



Revisión

ARTÍCULO EN INGLÉS

Maximum power, optimal load and optimal power spectrum for power training in upper-body (bench press): a review

F. Castillo^a, T. Valverde^b, A. Morales^c, A. Pérez-Guerra^a, F. de León^a and J.M. García-Manso^a

^aDepartamento de Educación Física. Universidad de Las Palmas de Gran Canaria. España.

^bFacultad de Ciencias de la Actividad Física. Universidad Católica de Valencia. España.

^cFacultad de Ciencias de la Actividad Física. Universidad de Granada. España.

History of the article:

Received June 9, 2011

Accepted November 17, 2011

Key words:

Bench press.

Maximal power.

Optimal load.

Optimal power spectrum.

Palabras clave:

Press de banca.

Potencia máxima.

Carga óptima.

Rango óptimo de potencia.

ABSTRACT

It is a fact that high performance sport has been characterized in recent years as a more specific training and in which coaches and athletes tend to use exercise and training loads which significantly resemble athletes' real actions during competition.

Principles of individuality and specificity are two aspects which best explain this trend. In that vein, this review analyzes and understands what specialized literature says to reach one of the most popular exercises used in upper-body power development: bench press in its different variants.

© 2012 Revista Andaluza de Medicina del Deporte.

RESUMEN

Potencia máxima, potencia óptima y espectro óptimo en el entrenamiento de la potencia del miembro superior (*bench press*): una revisión

Es un hecho que el deporte de alto rendimiento se ha caracterizado durante los últimos años por un entrenamiento cada vez más específico en el que técnicos y deportistas tienden a utilizar ejercicios y cargas de entrenamiento que se asemejan significativamente a las acciones que debe realizar el deportista durante la competición. Los principios de individualidad y especificidad son dos de los aspectos que mejor explican esta tendencia. En esa línea, esta revisión trata de analizar y entender lo que la bibliografía especializada señala con la realización de uno de los ejercicios más populares que se emplean en el desarrollo de la potencia del *upper-body*: *bench press* en sus diferentes variantes.

© 2012 Revista Andaluza de Medicina del Deporte.

Correspondence:

J.M. García Manso.

E-mail: jgarcia@def.ulpgc.es

Introduction

Resistance training plays a fundamental role in most of conditioning sports programs^{1,2} especially at high levels³. It is also known that most of sports actions, especially sport activities involving striking, throwing, jumping or rapid acceleration movements, sustain their performance in specific technical gestures implementation performed at maximum power⁴⁻⁷.

Power is the work amount produced per unit time or the product of force and velocity ($\text{Power} = \text{Force} \times \text{Displacement/Time} = \text{Force} \times \text{Velocity}$)⁷ and maximal power is the highest power level achieved in muscular contractions⁸. Maximal power output in a sport gesture varies with the load, contraction type and technique⁹.

Some papers suggest that, maximal power in single muscle fibres and single joint movements is reached approximately at 30% of maximum isometric strength and 30% of maximum isometric shortening velocity¹⁰⁻¹⁶. For multiple-joint muscle actions, optimal load varies with exercise. It is often said that, for lower body movements, optimal power appears at 0%¹⁷⁻²¹ and 55-59% 1RM²² in jump squat, 60-70% 1RM²³ and 40-65% 1RM²⁴ in half squat, and 56-78% 1RM²⁵ in leg press. Optimal load for weightlifting movements, such as clean and/or snatch has been identified at 70-80% 1RM²⁶. For upper-body movements, as for example bench press, countermovement bench press and bench press throw, optimal load is achieved between 30% and 70% of 1RM.

Aspects such as movement mechanics^{18,19,27-29} age⁹, gender²⁵, fibre type³⁰, muscle-tendon morphology³¹, muscular fatigue³², training level strength and training experience^{33,34} are some parameters that can affect the load percentage at which maximum power is reached in a technical gesture^{12,35}.

Consequently, the optimal load at which power output is reached is the load intensity in which the perfect combination between velocity and load displacement is produced¹⁶. This is known as *optimal load* (OL)^{7,18,26,35-37}. From a practice point of view, OL and similar power loads where there are no significant differences (optimal power spectrum) are considered as more appropriate loads to develop power at a specific technical gesture⁷.

Most of the studies related to OL determination have used three types of exercises (and its variants): total body (e.g. clean, snatch, hang power clean), lower body (e.g. squat, squat jump, leg press, leg extension) and upper body (e.g. bench press, bench pull and curl biceps). OL changes depending on the exercise and muscular group: Olympic lifters (80-100% 1RM) (e.g.^{26,38}); lower body (60%) (e.g.^{24,39}); upper body (40-70%) (e.g.^{13,40}).

In this review, we will focus on the upper-body OL analyzing published works in which bench press, in its different variants, is used. Articles were selected using US National Library of Medicine (PubMed), Scholar Google databases and indexed magazines in Spanish language (Redalyc, Dialnet LILAC y Latindex).

Bench press is one of the most common exercise used in training routines by most of athletes in every sport, being an optimal gesture to increase muscular force of the front of the chest (pectoralis major and pectoralis minor), arms (crural triceps: long, intern and extern portions) and shoulders (medial and anterior deltoids)⁴¹⁻⁴³.

Therefore, the aim of this paper is to review the optimal load for optimal power development spectrum in bench press (with and without countermovement) in different subjects, thinking of the influence on different kinetic and kinematic variables in maximal power output.

Methodological aspects which can affect muscular power assessment and optimal training load

In order to create force and muscular power assessment studies selection at which capabilities can be trained, we firstly created methodological criteria, which can allow right data interpretation of related studies. We should take into consideration that mostly all investigators make muscular power studies using kinematic systems, which enable muscular parameters assessment in terms of lifted load displacement during an exercise. From an external load on, and once known its displacement and the time to reach it, through specific designed softwares, optimal and mean power are estimated, as well as other kinematic parameters that can be useful to assess muscle or muscle groups mechanical characteristics during performance.

Frequently, there are significant changes in these studies because of the methods used, making difficult the results comparison. The main conflictive highlighted aspects are the few detailed information about morphofunctional sample characteristics, unclear description about exercise execution (e.g. slow or fast countermovement, stopping or not the movement at the end of the eccentric phase, etc.), few information about weight and size values corresponding to body segment displacement (arms, legs or whole body), different load increases used during performance, different exercises used to assess a same body segment (e.g. concentric bench press, bench press with countermovement, or bench press throwing), different measurement instruments and criteria used in the data interpretation (maximal power or mean power), etc. Before starting the bench press revision, we will explain some refinements about the previous mentioned points that, in case not being considered, will affect the revision interpretation.

Power peak and power mean

One of the main data interpretation and bibliography analysis difficulties were peak and mean power values not being clearly indicated. *Maximal power* is defined by Baker and Newton⁴⁴, as the maximal power production in the whole range or range of motion/concentric contraction. These authors refer to this value as *peak power* (PP), referring to a higher instantaneous production in a period of 1/ms without an apparent movement being observed. Some authors define this capability as the moment at which *threshold muscular performance is reached*, corresponding to a maximal mechanical performance that an athlete can produce in a concrete gest or movement⁴⁵. *Mean power* (MP) corresponds to mean values obtained from the sum of all the positive values developed during concentric phase performance divided by the number of data obtained in that gest or range of motion.

Free weights vs. resistance machine

In overweight training, common materials used are divided into two groups: machines and free-weights. The fact of using one or another will make a change in the final results. The term "machine" usually refers to resistance training devices with cables, pin loaded weight stacks or fixed lever arms. Free-weight encompass dumbbells and plates are typically loaded on to the end of a barbell. Free-weight exercises are performed usually at utility benches or squat racks.

Especially relevant is that barbell free force exercises highly controlled during the whole range motion optimize the gest and prevent from execution possible risks. A detailed analysis of 25 BP movements in

120 experienced weightlifters (candidate master: 30 people; Master of sports of Russia: 70 people; World-class athletes: 20 people) have allowed to establish Ivanovich et al⁴⁶ that the barbell displacement, even showing stable parameters in every weightlifter, can present significant individual differences, making changes during the seven phases in which lifting is divided and affecting any body plane (frontal, sagittal and transversal).

Using free-weights enhances stabilizing muscles group participation and level activation^{47,48}. Contrary, machine exercises cause opposite effect⁴⁸⁻⁵¹. Relative to bench press, it is especially relevant the free-weight effect on shoulders muscles (deltoids). This muscle has a stabilizing function, so that the anterior portion tends to resist a humerus lateral rotation at the same time that medial deltoid tends to resist abduction⁴⁸. This muscle relevancy has also been highlighted by Scheving and Pauly⁵² stating that its three portions (anterior, medial and posterior) are activated during this exercise to stabilize the humerus in glenoid cavity and as synergist movement structures.

Anthropometric characteristics

Athletes' anthropometric characteristics are determinant variables in all sports performance and, especially, in those where force is a discriminant capability. Relative to the studied exercise (BP), height and, fundamentally, upper-body (arm and forearm) length are two morphologic parameters which significantly influence in power levels reached during this movement. Generally, individuals with longer arms have greater advantage in BP power developing, rather than those with shorter arms or, even sometimes, higher force levels⁵³. Time and displacement to take the barbell to the chest depend on athletes' anthropometric particularities, width of the grip, height of the bridge, barrel displacement, lowering velocity and the barbell weight.

Incidence in the total mechanical system inertia

In order to assess kinematic parameters, the whole mechanical system inertia, must be carefully determined (i.e. mass of the lifted load plus the inertia of the levers and body segments) to be able to precisely calculate the load at which its power training is optimized⁵⁴. In case not making it real, results interpretation force to conclude erroneously, where there is a tendency to underestimate force levels⁵⁵ and power⁵⁴. Nelson and Duncan⁵ suggested that the gravity effect on the muscular performance should always be taken into consideration in force assessment. According to these authors, not considering these parameters takes, in isokinetic dynamometer (Cybex) assessments, into 4% errors for extensive knee muscles and to 15% for flexive muscles.

Sport gesture technical domain

A right domain technique execution is considered as the key for movement balance and stability, as well as to reach the right force application and power development. Load magnitude will represent the main factor that causes mechanical alterations in force exercises. Specifically, it is easy to prove how by making a PB at high loads ($>80\%$ 1RM) technical execution is seriously compromised, especially when training level and experience are low. Major changes are observed in load control and range motions. The movement magnitude in a BP decreases at higher loads because of a higher scapula protraction⁵⁶.

Individuals' training level

Power production depends on the subjects' maximal strength level, which frequently is determined by their training level. So, it doesn't seem strange to conclude that, the more trained the subjects are, the higher levels in force and PP will have.

Assessment methods and data collection

It is a fact that every instrument presents a different reliability degree, which affect directly to published results. This mechanism consisted of an optic encoder with a digital recorder connected (displacement error below 0.16%; 0.02% of time circuit error). Cronin et al⁵⁷ used a lineal position transducer (Unimeasure, Corvallis, OR), connected to a Smith Press machine, which would allow a velocity and movement variation assessment at a sampling frequency of 200 Hz (0.01 cm of precision). Siegel et al²⁹ used a chronoscopic light mechanism (model 63501 IR, Lafayette Instrument Company, Lafayette, IN) connected to a time mechanism (CLOCK Model 54050, Lafayette). On its part, Jandacka and Vaverka⁵⁸ used a rotational encoder at a 100 Hz sampling frequency (FitroDyne Premium - University of Komensky, Bratislava).

Optimal load, mean power and peak power in bench press

The following analysis is divided into PB_{cc} and PB_{ssc} movements from BT in its different variants (BT_{cc} , BT_{ssc} and BT). In each case, the following parameters are assessed: OL, PLS and power (PP and PM).

Concentric bench press and countermovement bench press

In order to make the OL analysis in PB_{cc} and PB_{ssc} , thirteen studies have been included. Nine of these examined B_{cc} , three studied B_{ssc} and the other two studied both movements ($B_{cc} + B_{ssc}$). Except in the study by Jandacka and Vaverka⁵⁸ and Naclerio and García⁵³ in which women were included in the sample ($n=52$), the rest of the evaluated subjects were men ($n=363$; men: 311; women: 52) of different level performing and force training experience. One group was formed by young elderly (≈ 40 years) and elderly (≈ 65 years)⁵⁹ and the rest were young adults ($\approx 20-25$ years), in which practitioners from different sports modalities were included (weightlifting, bodybuilding, basketball, handball, cyclists, volleyball, sprinters, middle distance runners and sailors)^{2,24,60-62}. The rest of the sample is formed by moderate active young health men volunteered^{29,53,57,58,63-65} (table 1).

As expected, most active subjects, especially the ones who practiced sports which required force conditions (powerlifting, weightlifting) or power (sprint), presented higher force levels, especially when results were expressed in relative values (1RM/BW) or were assessed according to their body weight and age (advanced or intermediate level). The strongest ones were powerlifting practitioners⁶⁰. In addition, sample subjects' high strength levels were highlighted in the study by Pearson et al (2009), which were elite-level sailors from the Emirates Team New Zealand America's Cup syndicate. Weakest subjects (novice or untrained) were the endurance modalities practitioners (cyclists and runners), subjects who didn't regularly practice sport and the oldest ones⁵⁹.

Table 1
Studies analyzing concentric bench press movement (BP_{cc}) and with countermovement (BP_{ss}). Sample characteristics are identified, age, body weight, maximal force (1-RM), relative force (RM/BW) and assessment instruments

Bench press with countermovement (BP _{cc}) and without it (BP _{ss})					
Article	Sample	Age (years)	BW (kg)	Performance 1-RM - RM/BW/L	MV
Mayhew et al ⁶³ BP _{SSC}	Male: 24 College Stud FA/TE (Smith Rack)	20.1±1.5	80.6±12.5	93.9±28.3/1.17 (I)	Optic Digital Encoder
Izquierdo et al ⁵⁹ BP _{CC}	G ₄₀ : 26 M-FA G ₆₅ : 21 M-FA	42.0±2.9 65.0±4.1	84.0±9.6 78.0±9.3	59.5±2.0/0.71 (N) 47.0±2.4/0.60 (I)	LPT
Cronin et al ⁵⁷ BP _{CC}	Male: 27 EE (U _{6M})	21.9±3.1	89.0±12.5	RM _L : 72.0±6.6/0.81 (N) RM _H : 100.9±7.2/1.13 (I)	LPT
Cronin et al ⁶⁴ BP _{CC} + BP _{SSC}	Male: 27 EE (U _{6M})	21.9±3.1	89.0±2.5	86.3±13.7/0.97 (N)	LPT
Izquierdo et al ²⁴ BP _{CC}	Male: 70 WL: 11 HP: 19 RR: 18 MD: 10 CG: 12	22.6±3.0 22.4±6.0 21.4±1.0 23.1±5.0 20.6±1.0	80.6±10.0 83.1±10.0 67.0±15.0 66.4±4.0 71.9±8.0	87.3±1.0/1.08 (I) 77.2±1.0/0.93 (I) 53.9±7.0/0.80 (N) 53.9±7.0/0.81 (N) 53.9±7.0/0.75 (UT)	LPT ER
Siegel et al ²⁹ BP _{SSC}	Male: 25 Student TE	23.0±4.0	89.0±30.0	79.3±14.1/0.89 (N)	CTL+TD
Naclerio and García ⁵³ BP _{CC}	UUTS: 37 Male: 33	22.0±2.1	U	76.0±10.2 (-)	LPT
Naclerio et al ⁶⁰ BP _{CC}	Male: 9 PL Female: 4	18.0-39.0	99.3±15.9	137.0±34.5/1.38 (A)	LPT
Asçi and Açıkkada ⁶² BP _{CC}	Male: 56 SP: 13 BP: 16 HP: 16 VP: 5 BB: 6	24.1±6.1 23.3±3.5 22.6±4.9 23.2±3.8 24.2±3.1	72.5±7.1 84.3±10.3 86.1±8.9 81.6±6.7 77.5±7.2	82.3±18.4/1.14 (I) 79.2±14.1/0.94 (N) 77.2±12.8/0.90 (N) 75.5±12.2/0.93 (N) 86.3±10.8/1.11 (I)	LPT
Marqués et al ⁶¹ BP _{CC}	Male: 14 HP	22.3±3.7	82.5±12.2	68.9±10.1/0.84	Rotary Encoder
Jandacka and Vaverka ⁵⁸ BP _{SSC}	Male: 55 Female: 48 PES	21.8±1.5 21.1±1.2	75.2±8.7 58.7±6.6	68.99±17.3/0.92 (N) 31.50±6.2/0.54 (N)	LPT FitroDyne
Pearson et al ² BP _{CC}	Male: 12 FA-Sailors	33.95±3.5	97.8±12.5	119.7±23.9/1.22 (I)	LPT
Sánchez et al ⁶⁵ BP _{CC}	Male: 100	25.1±5.0	79.4±8.3	98.7±12.5/1.24 (I)	LPT

A: advance level; BB: bodybuilders; BP: basketball players; BPCC: concentric bench press; BPSSC: bench press stretch-shorten cycle; BW: body weight; CG: control group; CTL+TD: chronoscopic timing lights connected to a timing device; EE: experienced subjects; F: female; FA: physically active; G: group; HP: handball players; I: Intermediate level; L: level; LPT: lineal position transducer; M: men; MD: middle distance runner; MV: valuation method; N: novice level; PES: physical education students; PL: powerlifters; RE: rotary encoder; RM: repetition maximum; RMH: strongest subjects; RML: weaker subjects; RR: road race; SP: sprinters; SP: sprinters; U: values not identified; U6M: untrained during the last 6 months; UT: untrained level; UUTS: not strength-trained university; VP: volleyball players.

It is well known that there is a strong relationship between maximal strength (1RM) and maximal power production^{20,39,66-75}. However, the strongest relationship between them occurs in heavier loading intensity⁷⁶. The reason for this is the fact that strongest subjects usually possess favorable neuromuscular characteristics^{15,77}. Also, it should be taken into consideration that strongest subjects usually present higher muscular development with a high cross-sectional area⁷⁸⁻⁸¹. In case of the most explosive athletes, hypertrophy mainly corresponds to fibers type II⁸¹⁻⁸³.

Optimal power training zones range between 30 and 70% of 1RM, with values always close to 50% for OL. However, OL does not seem to show a stable behavior in all analyzed studies, influenced by age, force level or training type of the subject. We can observe in the study by

Izquierdo et al²⁴ how strongest subjects present lower OL, while in the ones by Asçi and Açıkkada⁶² and Cronin et al⁵⁷, higher OL values correspond to bodybuilders (63%) or to subjects with a higher RM (60%) level. To our understanding, these differences have turned into a discussion topic between sports coaches, at the time of selecting the most efficient training loads to develop muscular power.

Some studies^{39,84,85} suggest that optimal load occur at higher loads in individuals with significantly greater maximal strength. We find the most highlighted case in the study by Propawski⁸², who proposes loads of approximately 70% 1RM for strongest subjects and 50% 1RM for weakest. However, Baker²² suggest that stronger athletes reach their maximal power output at lower loading rates in comparison to weaker

Table 2

Performance factors in concentric bench press movement (BPcc) and with countermovement (BPssc). Optimal load, optimal power spectrum, peak power and mean power values are shown.

Article	OL % - 1RM	OPS % - 1RM	PP - Watts (M-SD)	PM - Watts (M-SD)
Mayhew et al ⁶³ BP _{ssc}	40% Pre 40% Post	40-60% Pre 40-60% Pos	437.0±138.6 474.2±121.7	U
Izquierdo et al ²³ BP _{cc}	G ₄₀ : 45% G ₆₅ : 30%	30-45% (AG)	U	G ₄₀ : 293.0±11.0
Cronin et al ⁵⁷ BP _{cc} + BP _{ssc}	RM _i : 60% RM _h : 40% RM _i : 40% PP RM _i : 60% PM RM _h : 60% PP RM _h : 60% PM	40-60% 40-60% PP 40-60% PM	BP _{cc} RM _i : 501.8±55.3 RM _h : 572.7±79.8 BP _{ssc} RM _i : 444.9±66.5 RM _h : 556.0±80.9	BP _{cc} RM _i : 237.6±29.0 RM _h : 314.6±62.0 BP _{ssc} RM _i : 243.8±52.1 RM _h : 353.1±66.3
Cronin et al ⁶⁴ BP _{cc} + BP _{ssc}	50% PP: 50% MP: 70%	50-70% PP: 50-70% MP: 50-70%	BP _{cc} ≈555.0 ^{DG} BP _{ssc} ≈560.0 ^{DG}	BP _{cc} ≈275.0 ^{DG} BP _{ssc} ≈325.0 ^{DG}
Izquierdo et al ²⁴ BP _{cc}	WL: 30% HP: 30% RR: 45% MD: 45% CG: 45%	30-45% 30-45% 45-60% 45-60% 30-60%	U	486.0±10.0 468.0±76.0 272.0±52.0 269.0±45.0 266.0±30.0
Siegel et al ²⁹ BP _{ssc}	PP: 50%	PP: 40-60%	U	≈500.0
Naclerio and García ⁵³ BP _{cc}	MP: 53.3±10.7% PP: 47.1±10.7%	U	627.0±150.7	371.4±93.7
Naclerio et al ⁶⁰ BP _{cc}	45.5±10.6%	U	U	619.2±150.3
Asçi and Açıkada ⁶² BP _{cc}	SP: 52±12% BP: 50±14% HP: 58±16% VP: 54±12% BB: 63±16%	50-63%	U	SP: 227.0±115.0 BP: 232.0±201.0 HP: 190.0±98.0 VP: 300.0±307.0 BB: 221.0±96.0
Marques et al ⁶¹ BP _{cc}	PP: 38% MP: 52%	PP: 38-52% MP: 38-52%	≈820.0	≈450.0
Jandacka and Vaverka ⁵⁸ BP _{ssc}	M: 56±9% F: 63±8%	50-70% 50-70%	279.4±73.5 109.2±73.5	U
Pearson et al ² BP _{cc}	50%	U	≈600.0	306.0±75.0
Sánchez et al ⁶⁵ BP _{cc}	54.2%	40-65%	453.0±69.0	938.0±148.0

BB: bodybuilders; BP: basketball players; DG: graphic values; HP: handball players; OL: maximal power percentage; OPS: optimal power spectrum; PP: peak power; PM: mean power; RM: maximal repetition; RM_h: stronger subjects; RM_i: weaker subjects group; SP: sprinters; U: not shown; W: watts; M_{up}: male upward phase; M_{acc}: male acceleration phase.

athletes independently analyzed gesture. Other authors suggest that OL could always be the same independently of the subjects' force level²⁰. From the study by Izquierdo et al²³ we can also deduce that OL depends on the age, diminishing its value as it increases.

As shown in table 2, from the study by Izquierdo et al²³, higher [power values...] higher power values also correspond to strongest subjects, who present close values to 600 watts or more in PP and 300 watts in PM. Specifically, force velocity sports modalities athletes in which higher loads are moved (weightlifting), or those athletes to who add to the explosive gests a high height (basketball and volleyball players), are the ones who reach higher power levels.

Bodybuilders' low power level is highlighted in the study by Asçi and Açıkada⁶² (≈220W), which can be explained by the type of training, where work volume is significantly higher than the quality and execution high velocity adaptations in which hypertrophy has a general character and the same influence on slow and fast fibers.

Force importance and, more concrete, the way this is manifested, is especially relevant to power development. This capability is directly proportional to the peak acceleration and the mass of the object ($a=F/m$). Peak barbell acceleration is decreased as the intensity level is increased mainly being affected at the 2nd pull phase⁸⁶. We should take into consideration that, at constant resistance, non-ballistic movement involves two phases (acceleration and deceleration). The middle portion of the ascent is composed by the first deceleration phase and is defined as the effort portion where the applied force falls below the weight of load. The second acceleration phase, or the maximum strength region, is defined as the period where the applied force becomes greater than the load for the second period of time⁸⁷. Elliot et al⁸⁸, assessing the bench press, demonstrated that the deceleration phase corresponds to 23% of the last barbell range motion, when work at high loads is produced (1RM), increasing its value until 52% of the total displacement when loads were reduced to the 80%. However, we should take into

consideration that, when loads are especially high, (80-100% 1RM), instead of two phases, force presents four phases or regions (acceleration, sticking, maximum strength and deceleration)^{51,88,89}.

Sato et al⁸⁶, suggest that peak barbell acceleration does not change from 50 to 80% of 1RM in elite and experienced weightlifters, indicating that the force production becomes greater while the barbell mass is increased and the peak barbell acceleration remains relatively constant. Peak barbell acceleration is significantly decreased at increases ranging from 80 to 85% of 1RM. The results demonstrated that the force affecting barbell acceleration at the 2nd pull phase reaches near maximal level around 85% of 1RM.

Duration of the acceleration, during concentric phase, decrease with load. For example, acceleration phase change of 63.8% of the duration of the concentric phase, to 82.9 for 30 and 80% 1RM respectively⁵⁶. However, Sato et al⁸⁶ found that the peak barbell acceleration showed no changes from 50-80% 1RM among elite and experienced weightlifters, indicating that barbell acceleration remains relatively constant. Force affecting barbell acceleration at the 2nd pull phase reaches near maximal level around 85% 1RM. In other words, force production remains relatively the same while the peak acceleration decreases and the mass of the barbell increases. These showed that roughly 80% 1RM is the threshold for the elite level weightlifters to be able to maintain the peak barbell acceleration.

As previously mentioned, the way force is developed and applied to movement is the key of the barbell acceleration. In case peak force appears prematurely during concentric phase, a force decrease will occur during the last period of the range motion or, contrary, if velocity peak delays and acceleration phase is too long, end of a greater decline in force and a drastic deceleration at the concentric phase will occur⁸⁷. Barbell acceleration magnitude determines its velocity at the different displacement points. Its mean and maximum values will vary depending on the work load, decreasing velocity as the load is increased. González-Badillo and Sánchez-Medina⁹⁰ found a high relationship between mean velocity and relative load (1RM-%) that allows us to use one to estimate the OL with great precision. Also, these authors suggest that mean velocities attained with each 1RM (%) can differ very slightly due to differences in velocity at 1RM. During concentric phase an increase in mean velocity is associated with a decrease in concentric phase duration and the magnitude of the load lifted. A 100% increase in mean velocity necessitated a 37.5% load reduction, whereas a 50% reduction in load was required to attain an equivalent increase in the peak velocity⁵⁶.

In most frequently movements used for power training (e.g. bench press; squat, clean, etc.), we notice how barbell velocity decreases in the last part of the movement. This is probably due to a decrease in agonists muscle activation and possible increase in antagonist muscles activity, in order to stop the load at the end of the range of motion⁴. In ballistic actions (e.g. jump squat and bench press throw), a continued acceleration is observed throughout the range of motion, concentric velocity, force, power and muscle activation. These factors are higher during a ballistic movement in comparison to a similar traditional resistance training exercise^{4,16}.

A key in force development and muscular power is the kind of movement during the exercise execution. Countermovement actions take place to increase gest efficiency and enhance muscle mechanical answer. This hypothesis is true for PM in both studies^{57,64} including both movements (BP_{CC} and BP_{SSC}), as well as for PP^{58,64}, but not being the same in the study by Cronin et al⁵⁷, where peak power across the total

concentric phase was not affected by rebound action. These authors explained that rebound movement effect is produced to cause a shift phase in the power-time signal onto the left, peak power remaining unaffected in temporal terms. Consequently, these authors suggested that greater peak power would seem like a maximal strength function rather than individual's ability to utilize the SSC.

An eccentric muscle action stimulates the stretch reflex and builds up elastic energy allowing, mainly, force and power levels improvement during the subsequent concentric action^{71,91-93}. Mechanical source SSC bases were initially established by Cavagna et al⁹⁴ and have been analyzed by numerous subsequent studies aimed at analyzing its manifestation and effects magnitude. Potential SSC benefits are caused by energy stored in the elastic components (tendons and actin-myosin complex) utilization, reutilization and parallel (aponeurosis) of the musculotendinous system⁹⁴⁻¹⁰¹, spinal reflex⁹⁸⁻¹⁰⁰ and long latency responses¹⁰¹.

Elastic-reflex enhancement may be reflected in higher increases at 10-15% in power output^{54,57,97,102}. However, as seen in this review, SSC benefits vary considerably in each individual^{103,104}, especially when execution deficiencies differences are shown and force levels are low. Some authors^{64,99} suggested that a part of countermovement efficiency lost could be due to two phenomena: elastic energy loss caused by slow decreases and prolonged coupling phases, or by muscles inability to generate force at high muscle shortening velocities.

Peak velocity occurs later whereby the SSC effect has diminished^{13,64}. Some authors suggested that the elastic-reflex use only maximizes concentric movement initial part^{13,57,76,105-107}. Cronin et al⁵⁷ observed how enhancement is manifested in the first 200 ms of the concentric phase^{13,57}, linking the efficiency in the initial phase of the thrust stage with the athlete's maximal force. Bosco et al⁹⁹ suggested that during elongation phase above 500 ms, longer coupling is produced; causing an elastic energy decrease that could be stored in the muscle during the eccentric phase. However, Schmidtbleicher⁷⁶, doesn't fully accept this announcement, suggesting that if maximal strength is the main power performance factor, especially at the beginning of the push phase, everything will be conditioned by the external load used, diminishing their influence with decreasing load.

In countermovement muscle actions, peak acceleration and peak force have been shown to increase intensively. To this, it is necessary to have enough force to reduce the eccentric velocity of the load to zero prior to begin the concentric action. The change in momentum is directly proportional to the change in velocity and the mass of the load, increasing at fast and short eccentric phases and decreasing otherwise. This change in momentum is also proportional to the force which is causing such a change, and the duration over which the changes take place. The sum of external force in eccentric phase, supposedly, allows higher accelerations during the initial portion of the concentric phase. Perhaps, potential benefits depend on the ability to use the force increase quickly, via recruitment of a high number of motor units and a quickly elastic energy recovery.

Bench press throwing: concentric bench press and stretch-shorten cycle

In this section, six studies have been analyzed^{13,17,22,34,40,64}, of which three study only concentric bench press throw (BT_{CC}) and the other three assessed the same movement, adding the stretch-shorten cycle bench

press ($BT_{cc}+BT_{ssc}$) (table 3). In all of them, young adults (230 males), were used as sample, except in the study by Cronin et al⁶⁴, in which the sample presented a low performance (Novice level), the other five studies, assessed athletes with good force levels (Intermediate or Advanced level).

Exercises with barbell release, as the ones analyzed in this paragraph, are called *dynamic*³⁶, *explosive*¹⁰⁸ or *ballistic exercises*⁴. In any exercise, barbell throwing incorporation pretends to get closer to a competition behavior. This way, we can adapt the motor pattern used in competition, as well as.

The used motor pattern to real athlete's needs during performance. Also, it is pretended to eliminate, or minimize the characteristic deceleration phase at the end of the sports gest in which the bar (or implement) is not released during the last displacement part^{4,87}.

These reasons motivate athletes and coaches to train this kind of movement, especially when the training goal is power development. In the case of BP, its use makes significant changes in kinetic and kinematic traditional movements¹⁰⁹, with or without countermovement, where it is important that the barbell is controlled during the whole range of motion. We find the most important differences in concentric phase where a shorter time period is reached, as well as higher peaks and average velocities, which later will be traduced into average force enhancements, MP and PP in traditional movements⁴.

We can easily observe that the barbell velocity changes at any load intensity. Newton et al⁴ quantified these increases at 27.3% of mean velocity at 45% 1RM. This velocity will go on increasing or will keep maintained while the athlete keeps the barbell control. The point at which the load loses contact with the athlete, leads to any muscular force and therefore any change in velocity is not possible except for the gravity force causes, which is not included in any kinematic or kinetic calculation⁸⁷.

Commonly, most of every overweighed training exercise, forces the athlete to stop a substantial portion of the range and control the movement to guarantee the structural muscular integrity, as well as the

gestural involved joints (e.g. in BP we talk about elbow joint). This supposes that, during the barbell range, there is a decelerated gest phase prior to achieving zero velocity^{4,88,108}. This displacement part is accompanied by a muscular activity reduction, manifested in the agonist muscles electromyography activity. Deceleration results from shortening agonist activation and greater antagonist co-activation, especially at low loads performance⁴.

In these circumstances, load determinates the acceleration range and stops at concentric movement phase. Cronin et al⁶⁴ suggested that higher peak velocities will be reached later when PT is being performed, rather than when classic PB is performed. However, this phase will be determined by the load to move. Some studies demonstrated that during bench press with a light load (45% 1RM) deceleration phase was shorter ($\approx 40\%$ of concentric time)⁴ compared to heavy loads of 80% of 1RM (50% of concentric time)⁸⁸. Consequently, subjects with a higher force level will be in advantage during ballistic movements, when capability to develop force in a few period of time is high.

Results will turn into faster and more powerful gest. In assessed studies, we observe that except in the sample studied by Cronin et al⁶⁴, capability to apply force and power development is higher in mostly all the subjects (PP: 600-1,000 watts), than the one we see in traditional PB, where the barbell is not released (table 4). In the analyzed studies, maximal power was found in diverse power values. This way, while in the studies by Newton et al¹³ (OL: 15-30%) or Bevan et al¹⁷ (OL: 30%) optimal power was obtained at low work intensities, in studies by Baker^{22,34,40} and Cronin et al⁶⁴ OL was reached at loads between 50-60%. No significant differences were found between OL in PB_{cc} and PB_{ssc} . Optimal power spectrum was always detected at nearly or slightly higher OL intensities.

As noted above, in barrel released movements, force changes during the course, as it depends on the load (RM). Thus, at high loads, an initial peak force in the early stage of the movement is produced and then it decreases near the end of gesture. Different is the behavior at lower loads, where maximal force is produced at the beginning and then gradually decreased until the end of the movement.

Table 3

Performance factors in concentric bench press movement (BP_{cc}) and with countermovement (BP_{ssc}). Optimal load, optimal power espectrum, peak power and mean power values are shown

Concentric Bench Press Throw (BT_{cc}) SST Bench Press (BT_{ssc})					
Author	Sample	Age (years)	BW (Kg)	1-RM (Kg)	MV
Newton et al ¹³ BT_{cc} - BT_{ssc}	17 - Men TE (T_{6M})	20.6±1.9	83.7±8.2	104.0±16/1.24 (I)	ER
Cronin et al ⁶⁴ BT_{cc} - BT_{ssc}	27 - Men TE (U_{6M})	21.9±3.1	89.0±12.5	86.3±13.7/0.97 (N)	LPT (PPS)
Baker ³⁴ BT_{cc}	Male: 49 NRL: 22 SRL: 27	24.3±3.7 18.1±1.1	93.4±9.6 91.1±9.8	134.8±15.2/1.44 (A) 111.0±15.3/1.22 (I)	LPT (PPS)
Baker ⁴⁰ BT_{cc}	Male: 59 NL: 19 SL: 23 SRL: 17	NL: 25.1±3.4 SL: 19.7±2.0 SRL: 17.6±0.9	NL: 94.8±10.0 SL: 91.8±7.0 SRL: 91.8±10.0	NL: 140.1±14.0/1.48 (A) SL: 121.1±13.0/1.32 (I) SRL: 108.7±16.0/1.18 (I)	LPT (PPS)
Baker et al ²² BT_{cc}	Male: 31 Football	22.2±3.5	92.0±11.1	129.7±14.3/1.41 (I)	LPT (PPS)
Bevan et al ¹⁷ BT_{cc} - BT_{ssc}	Male: 47 Rugby	25.5±4.8	101.0±12.8	124.0±19.0 - 1.22 (I)	BMS LPT

A: advance level; BT_{cc} : concentric bench press throwing; BT_{ssc} : stretch-shorten cycle bench press throwing; BW: body weight; ER: encoder rotatory; I: Intermediate level; LPT: lineal position transducer; M: men; MV: valuation method; N: novice level; NRL: national rugby league; PES: physical education students; PPS: plyometric power system; RM: maximal repetition; RM_1 : strongest subjects; RM_2 : weaker subjects; SRL: college-aged rugby players; U_{6M} : untrained the last 6 months; U: values not identified.

Table 4

Performance factors influence in concentric Bench Press Throw movement (BP_{cc}) and Bench Press Throw with countermovement (BP_{ssc}). Optimal load, training optimal range and mean and peak power values are shown

Bench Press Throw Concentric (BT _{cc})				
Author	OL % - 1RM	OPS % - 1RM	PP Watts (M/SD)	MP Watts (M/SD)
Newton et al ¹³ BT _{cc} -BT _{ssc}	PP _{cc} : 15% MP _{cc} : 30% PP _{ssc} : 15% MP _{ssc} : 30%	PP _{cc} : 15-30% MP _{cc} : 30-45% PP _{ssc} : 15-30% MP _{ssc} : 30-45%	≈1.000.0 ^{DG} ≈1.050.0 ^{DG}	≈400 ^{DG} 563.0±104.0
Cronin et al ⁶⁴ BT _{cc} -BT _{ssc}	PP _{cc} : 50% MP _{cc} : 60% PP _{ssc} : 50% MP _{ssc} : 60%	PP _{cc} : 40-60% MP _{cc} : 50-70% PP _{ssc} : 40-60% MP _{ssc} : 50-70%	≈620 ^{DG} ≈625 ^{DG}	≈290 ^{DG} ≈340 ^{DG}
Baker ³⁴ BT _{cc}	NRL: 51.1±5.3% SRL: 54.9±5.6	44.4-59.2% 45.1-63.1%	U	610±79 515±78
Baker ⁴⁰ BT _{cc}	NL: 50% SL: 50% SRL: 55%	NL: ≈40-60% SL: ≈40-60% SRL: ≈40-60%	U	NL: 635±87 SL: 561±57 SRL: 499±81
Baker et al ²² BT _{cc}	54.9±5.3%	50-60%	U	588.0±95.0
Bevan et al ¹⁷ BT _{cc} -BT _{ssc}	30% 30%	30-50% 20-60%	U 873.0±24.2	873.0±23.7

DG: graphic values; MP: mean power; OL: optimal power load percentage; OPS: optimal power spectrum; PP: peak power; RM: maximal repetition; RMH: strongest subjects group; RML: less strong subjects; TE: training experience; U: not shown; W: watts.

This leads to changes in size, shape and timing in the acceleration phases. In PB, the acceleration phase is larger than in a single concentric movement. Along these lines, Newton et al⁴ found that for ballistic actions, working with a load of 45% 1RM (bench press), acceleration is generated during 96% of the course, compared to 60% of concentric actions. This represents an increase of ≈36 of peak velocity and significant changes in peak power⁴. As in traditional BP, when a countermovement is included in BT, MP is favored (with differences ranging between 15-30%), although it's not the same situation as in PP.

Conclusions and practical implications

The results in this review show how upper-body power training with bench press exercise passes through the optimal load and optimal power spectrum, allowing the maximal power output evaluation. Parameters such as age, training level and sport specialization marked differences in the optimal load and optimal power spectrum value. In addition, with the aim of optimizing bench press technical variants (BP_{cc}, BP_{ssc} or BT) it is necessary that, previously, have enough force levels and an appropriate execution technique.

References

- Harman E. Strength and power: a definition of terms. *Nat Strength Cond Assoc J.* 1993;15:18-20.
- Pearson SN, Cronin JB, Hume PA, Slyfield D. Kinematics and kinetics of the bench-press and bench-pull exercises in a strength-trained sporting population. *Sports Biomech.* 2009;8:245-54.
- Keogh JW, Wilson GJ, Weatherby RP. A cross-sectional comparison of different resistance training techniques in the bench press. *J Cond Strength Res.* 1999;13:247-58.
- Newton RU, Kraemer WJ, Häkkinen K, Humphries BJ, Murphy AJ. Kinematics, kinetics and muscle activation during explosive upper body movements. *J Appl Biom.* 1996;12:31-43.
- Sleivert G, Taingahue M. The relationship between maximal jump-squat power and sprint acceleration in athletes. *Eur J Appl Physiol.* 2004;91:46-52.
- Young WB, Newton RU, Doyle TL, Chapman D, Cormack S, Stewart G, et al. Physiological and anthropometric characteristics of starters and non-starters and playing positions in elite Australian Rules Football: a case study. *J Sci Med Sport.* 2005;8:333-45.
- Cronin J, Sleivert G. Challenges in understanding the influence of maximal power training on improving athletic performance. *Sports Med.* 2005;35:213-34.
- Gollnick PD, Bayley WM. Biochemical training adaptations and maximal power. In: Jones NL, McCartney N, McComas AJ, editors. *Human muscle power.* Champaign, IL: Human Kinetics; 1986. p. 255-67.
- Zatsiosky VM, Kraemer JW. *Science and practice of strength training.* Champaign, IL: Human Kinetics; 2006.
- Hill AV. The heat of shortening and the dynamic constants of muscle. *Proceedings of the Royal Society of London B.* 1938;126:136-95.
- Wilkie DR. The relation between force and velocity in human muscle. *J Physiol.* 1949;110:249-80.
- Kaneko M, Fuchimoto T, Toji H, Sueti K. Training effect of different loads on the force-velocity relationship and mechanical power output in human muscle. *Scand J Sports Sci.* 1983;5:50-5.
- Newton RU, Murphy AJ, Humphries BJ, Wilson GJ, Kraemer WJ, Häkkinen K. Influence of load and stretch shortening cycle on the kinematics, kinetics and muscle activation that occurs during explosive upper-body movements. *Eur J Appl Physiol Occup Physiol.* 1997;75:333-42.
- Bottinelli R, Pellegrino MA, Canepari M, Rossi R, Reggiani C. Specific contributions of various muscle fibre types to human muscle performance: an in vitro study. *J Electromyogr Kinesiol.* 1999;9:87-95.
- Toji H, Kaneko M. Effect of multiple-load training on the force-velocity relationship. *J Strength Cond Res.* 2004;18:792-5.
- Cormie P, McGuigan MR, Newton RU. Developing maximal neuromuscular power: Part 1-biological basis of maximal power production. *Sports Med.* 2011;41:17-38.
- Bevan HR, Bunce PJ, Owen NJ, Bennett MA, Cook CJ, Cunningham DJ, et al. Optimal loading for the development of peak power output in professional rugby players. *J Strength Cond Res.* 2010;24:43-7.
- Cormie P, McCaulley GO, Triplett NT, McBride JM. Optimal loading for maximal power output during lower-body resistance exercises. *Med Sci Sports Exerc.* 2007;39:340-9.
- Cormie P, McCaulley GO, McBride JM. Power versus strength-power jump squat training: influence on the load-power relationship. *Med Sci Sports Exerc.* 2007;39:996-1003.
- Cormie P, McGuigan MR, Newton RU. Influence of strength on magnitude and mechanisms of adaptation to power training. *Med Sci Sports Exerc.* 2010;42:1566-81.

21. Dayne AM, McBride JM, Nuzzo JL, Triplett NT, Skinner J, Burr A. Power output in the jump squat in adolescent male athletes. *J Strength Cond Res.* 2011;25:585-9.
22. Baker D, Nance S, Moore M. The load that maximizes the average mechanical power output during explosive bench press throws in highly trained athletes. *J Strength Cond Res.* 2001;15:20-4.
23. Izquierdo M, Häkkinen K, Antón A, Garrues M, Ibáñez J, Ruesta M, et al. Maximal strength and power, endurance performance, and serum hormones in middle-aged and elderly men. *Med Sci Sports Exerc.* 2001;33:1577-87.
24. Izquierdo M, Häkkinen K, González-Badillo JJ, Ibáñez J, Gorostiaga EM. Effects of long-term training specificity on maximal strength and power of the upper and lower extremities in athletes from different sports. *Eur J Appl Physiol.* 2002;87:264-71.
25. Thomas GA, Kraemer WJ, Spiering BA, Volek JS, Anderson JM, Maresh CM. Maximal power at different percentages of one repetition maximum: influence of resistance and gender. *J Strength Cond Res.* 2007;21:336-42.
26. Kawamori N, Crum AJ, Blumert PA, Kulik JR, Childers JT, Wood JA, et al. Influence of difference relative intensities on power output during the hang power clean: identification of the optimal load. *J Strength Cond Res.* 2005;19:698-706.
27. Burke ER. The control of muscle force: Motor Unit recruitment and firing patterns. In: Jones NL, McCartney N, McComas AJ, editors. *Human Muscle Power.* Champaign, IL: Human Kinetics; 1986. p. 98-109.
28. Komi PV. The stretch shortening cycle and human power output. In: Jones NL, McCartney N, McComas AJ, editors. *Human muscle power.* Champaign, IL: Human Kinetics; 1986. p. 20-34.
29. Siegel JA, Gilders RM, Staron RS, Hagerman FC. Human muscle power output during upper- and lower-body exercises. *J Strength Cond Res.* 2002;16:173-8.
30. Faulkner JA. Power output of the human diaphragm. *Am Rev Respir Dis.* 1986;134:1081-3.
31. Edgerton RV, Roy RR, Gregor JR, Rugg S. Morphological basis of skeletal muscle power output. In: Jones NL, McCartney N, McComas AJ, editors. *Human Muscle Power.* Champaign, IL: Human Kinetics; 1986. p. 43-64.
32. Green JH. Muscle power: fiber type recruitment, metabolism and fatigue. In: Jones NL, McCartney N, McComas AJ, editors. *Human Muscle Power.* Champaign, IL: Human Kinetics; 1986. p. 65-79.
33. Kraemer WJ, Newton RU. Training for muscular power. *Phys Med Rehabil Clin N Am.* 2000;11:341-68.
34. Baker D. Comparison of upper-body strength and power between professional and college-aged rugby league players. *J Strength Cond Res.* 2001;15:30-5.
35. McBride JM, Triplett-McBride T, Davie A, Newton RU. The effect of heavy- vs. light-load jump squats on the development of strength, power and speed. *J Strength Cond Res.* 2002;16:75-82.
36. Wilson GJ, Newton RU, Murphy AJ, Humphries BJ. The optimal training load for the development of dynamic athletic performance. *Med Sci Sports Exerc.* 1993;25:1279-86.
37. Kawamori N, Haff GG. The optimal training load for the development of muscular power. *J Strength Cond Res.* 2004;18:675-84.
38. Kilduff LP, Bevan H, Owen N, Kingsley MI, Bunce P, Bennett M, et al. Optimal loading for peak power output during the hang power clean in professional rugby players. *Int J Sports Physiol Perform.* 2007;2:260-9.
39. Stone MH, O'Bryant HS, McCoy L, Coglianese R, Lehmkuhl M, Schilling B. Power and maximal strength relationship during performance of dynamic and static weighted jumps. *J Strength Cond Res.* 2003;17:140-7.
40. Baker D. A series of studies on the training of high-intensity muscle power in rugby league football players. *J Strength Cond Res.* 2001;15:198-209.
41. Wilson GJ, Elliot BC, Kerr GK. Bar path and force profile characteristics for maximal loads in the bench press. *Int J Sport Biomech.* 1989;5:90-402.
42. Power G, Stratton S. Resistance training with machines. Champaign, IL: Stipes; 1989.
43. Barnett C, Kippers V, Turner P. Effects of variations of the bench press exercise on the EMG activity of five shoulder muscles. *J Strength Cond Res.* 1995;9:223-7.
44. Baker D, Newton RU. Acute effect on power output of alternating an agonist and antagonist muscle exercise during complex training. *J Strength Cond Res.* 2005;19:202-5.
45. González-Badillo JJ, Ribas J. Bases de la programación del entrenamiento de la fuerza. Barcelona: Editorial INDE; 2002.
46. Ivanovich SI, Georgievich LB, Stanislavovich FV. First results of the biomechanical analysis of the bench press technique. Available from: <http://erictalmant.com/pdf/Boris+Sheiko+Bench+Press+Technique+Article.pdf>
47. Garhammer J. Weight lifting and training. In: Vaughn CL, editors. *Biomechanics of Sport.* Boca Raton, FL: CRC Press; 1989. p. 169-211.
48. McCaw ST, Friday JJ. A comparison of muscle activity between a free weight and machine bench press. *J Strength Cond Res.* 1994;84:259-64.
49. Andrews JG, Hay JG, Vaughan CL. Knee shear forces during squat exercise using a barbell and a weight machine. In: Matsui H, Kabashi K, editors. *Biomechanics VIII-B.* Champaign, IL: Human Kinetics; 1983. p. 923-7.
50. Ariel GB. Resistive training. *Clin Sports Med.* 1983;21:55-69.
51. Lander JE, Bates BT, Sawhill JA, Hamill J. A comparison between free-weight and isokinetic bench pressing. *Med Sci Sports Exerc.* 1985;17:344-53.
52. Scheving LE, Pauly JE. An electromyographic study of some muscles acting on the upper extremity of man. *Anat Rec.* 1959;135:239-45.
53. Naclerio F, García S. Influencia de la longitud de los miembros superiores sobre la fuerza y la potencia producida en el Press de Banca. *PublICE Standard.* 13/11/2006. Pid: 740.
54. Rambaud O, Rahmani A, Moya B, Bourdin M. Importance of upper-limb inertia in calculating concentric bench press force. *J Strength Cond Res.* 2008;22:383-9.
55. Nelson SG, Duncan PW. Correction of isokinetic and isometric torque recordings for the effects of gravity. A clinical report. *Phys Ther.* 1983;63:674-6.
56. Cronin JB, McNair PJ, Marshall RN. Force-velocity analysis of strength-training techniques and load: implications for training strategy and research. *J Strength Cond Res.* 2003;17:148-55.
57. Cronin JB, McNair PJ, Marshall RN. The role of maximal strength and load on initial power production. *Med Sci Sports Exerc.* 2000;32:1763-9.
58. Jandačka D, Veverka F. A regression model to determine load for maximum power output. *Sports Biomech.* 2008;7:361-71.
59. Izquierdo M, Aguado X, González R, López JL, Häkkinen K. Maximal and explosive force production capacity and balance performance in men of different ages. *Eur J Appl Physiol Occup Physiol.* 1999;79:260-7.
60. Naclerio F, Jiménez A, Forte D. Relación del peso máximo con la fuerza aplicada y la potencia producida en un test creciente, en el ejercicio de press de banca plano con barra libre, en levantadores. *APUNTS Educación Física y Deportes.* 2006;4:42-52.
61. Marqués MC, Van den Tilaar R, Vescovi JD, González-Badillo JJ. Relationship between throwing velocity, muscle power, and bar velocity during bench press in elite handball players. *Int J Sports Physiol Perform.* 2007;2:414-22.
62. Aşçi A, Aşikada C. Power production among different sports with similar maximum strength. *J Strength Cond Res.* 2007;21:10-6.
63. Mayhew JL, Ware JS, Johns RA, Bemben MG. Changes in upper body power following heavy-resistance strength training in college men. *Int J Sports Med.* 1997;18:516-20.
64. Cronin J, McNair PJ, Marshall RN. Developing explosive power: a comparison of technique and training. *J Sci Med Sport.* 2001;4:59-70.
65. Sánchez-Medina L, Pérez CE, González-Badillo JJ. Importance of the propulsive phase in strength assessment. *Int J Sports Med.* 2010;31:123-9.
66. Moss BM, Refsnes PE, Abildgaard A, Nicolaysen K, Jensen J. Effects of maximal effort strength training with different loads on dynamic strength, cross-sectional area, load-power and load-velocity relationships. *Eur J Appl Physiol Occup Physiol.* 1997;75:193-9.
67. Blackburn JR, Morrissey MC. The relationship between open and closed kinetic chain strength of the lower limb and jumping performance. *J Orthop Sports Phys Ther.* 1998;27:430-5.
68. Carlock JM, Smith SL, Hartman MJ, Morris RT, Ciroslan DA, Pierce KC, et al. The relationship between vertical jump power estimates and weight-lifting ability: a field-test approach. *J Strength Cond Res.* 2004;18:534-9.
69. Wisløff U, Castagna C, Helgerud J, Jones R, Hoff J. Strong correlation of maximal squat strength with sprint performance and vertical jump height in elite soccer players. *Br J Sports Med.* 2004;38:285-8.
70. Ugrinowitsch C, Tricoli V, Rodacki AL, Batista M, Ricard MD. Influence of training background on jumping height. *J Strength Cond Res.* 2007;21:848-52.
71. Miyaguchi K, Demura S. Muscle power output properties using the stretch-shortening cycle of the upper limb and their relationships with one-repetition maximum bench press. *J Physiol Anthropol.* 2006;25:239-45.
72. Nuzzo JL, McBride JM, Cormie P, McCaulley GO. Relationship between countermovement jump performance and multijoint isometric and dynamic tests of strength. *J Strength Cond Res.* 2008;22:699-707.
73. Baker DG, Newton RU. Comparison of lower body strength, power, acceleration, speed, agility, and sprint momentum to describe and compare playing rank among professional rugby league players. *J Strength Cond Res.* 2008;22:153-8.
74. Baker DG, Newton RU. Effect of kinetically altering a repetition via the use of chain resistance on velocity during the bench press. *J Strength Cond Res.* 2009;23:1941-6.
75. Cormie P, McBride JM, McCaulley GO. Power-time, force-time, and velocity-time curve analysis of the countermovement jump: impact of training. *J Strength Cond Res.* 2009;23:177-86.
76. Schmidbleicher D. Training for power events. In: Komi PV, editors. *Strength and Power in Sport.* Boston: Blackwell Scientific Pub; 1992. p. 381-95.

77. Folland JP, Williams AG. The adaptations to strength training: morphological and neurological contributions to increased strength. *Sports Med*. 2007;37:145-68.
78. Costill DL, Coyle EF, Fink WF, Lesmes GR, Witzmann FA. Adaptations in skeletal muscle following strength training. *J Appl Physiol*. 1979;46:96-9.
79. Jones DA, Rutherford OM, Parker DF. Physiological changes in skeletal muscle as a result of strength training. *Q J Exp Physiol*. 1989;74:233-56.
80. Staron RS, Leonardi MJ, Karapondo DL, Malicky ES, Falkel JE, Hagerman FC, et al. Strength and skeletal muscle adaptations in heavy-resistance-trained women after detraining and retraining. *J Appl Physiol*. 1991;70:631-40.
81. Campos GE, Luecke TJ, Wendeln HK, Toma K, Hagerman FC, Murray TF, et al. Muscular adaptations in response to three different resistance-training regimens: specificity of repetition maximum training zones. *Eur J Appl Physiol*. 2002;88:50-60.
82. Thorstensson A, Hultén B, Von Döbeln W, Karlsson J. Effect of strength training on enzyme activities and fibre characteristics in human skeletal muscle. *Acta Physiol Scand*. 1976;96:392-8.
83. Dons B, Bollerup K, Bonde-Petersen F, Hancke S. The effect of weightlifting exercise related to muscle fiber composition and muscle cross-sectional area in humans. *Eur J Appl Physiol Occup Physiol*. 1979;40:95-106.
84. Poprawski B. Aspects of strength, power and speed in shot put training. *New Stud Athl*. 1988;3:89-93.
85. Driss T, Vandewalle H, Quièvre J, Miller C, Monod H. Effects of external loading on power output in a squat jump on a force platform: a comparison between strength and power athletes and sedentary individuals. *J Sports Sci*. 2001;19:99-105.
86. Sato K, Fleschler P, Sands W. Barbell acceleration analysis on various intensities of weightlifting. *ISBS-Conference Proceedings Archive, 27 International Conference on Biomechanics in Sports; 2009*.
87. Frost DM, Cronin JB, Newton RU. A comparison of the kinematics, kinetics and muscle activity between pneumatic and free weight resistance. *Eur J Appl Physiol*. 2008;104:937-56.
88. Elliot BC, Wilson GJ, Kerr GK. A biomechanical analysis of the sticking region in the bench press. *Med Sci Sports Exerc*. 1989;21:450-62.
89. Frost DM, Cronin J, Newton RU. A biomechanical evaluation of resistance: fundamental concepts for training and sports performance. *Sports Med*. 2010;40:303-26.
90. González-Badillo JJ, Sánchez-Medina L. Movement velocity as a measure of loading intensity in resistance training. *Int J Sports Med*. 2010;31:347-52.
91. Norman RW, Komi PV. Electrochemical delay in skeletal muscle under normal movement conditions. *Acta Physiol Scand*. 1979;106:241-8.
92. Komi PV. Physiological and biomechanical correlates of muscle functions: effects of muscle structure and stretch-shortening cycle on force and speed. *Exerc Sport Sci Rev*. 1984;12:81-121.
93. Walshe AD, Wilson GJ, Ettema GJ. Stretch-shorten cycle compared with isometric preload: contributions to enhanced muscular performance. *J Appl Physiol*. 1998;84:97-106.
94. Cavagna GA, Saibene FP, Margaria R. Effect of negative work on the amount of positive work performed by an isolated muscle. *J Appl Physiol*. 1965;20:157-8.
95. Asmussen E, Bonde-Petersen F. Storage of elastic energy in skeletal muscles in man. *Acta Physiol Scand*. 1974;91:385-92.
96. Cavagna GA, Citterio G. Effect of stretching on the elastic characteristics and the contractile component of frog striated muscle. *J Physiol*. 1974;239:1-14.
97. Komi PV. Stretch shortening cycle. In: Komi PV, editors. *The Encyclopaedia of Sports Medicine III. Strength and power in sport*. 1992. p. 169-79.
98. Dietz V, Noth J, Schmidtbleicher D. Interaction between pre-activity and stretch reflex in human triceps brachii during landing from forward falls. *J Physiol*. 1981;311:113-25.
99. Bosco C, Tihanyi J, Komi PV, Fekete G, Apor P. Store and recoil of elastic energy in slow and fast types of human skeletal muscles. *Acta Physiol Scand*. 1982;116:343-9.
100. Nicol C, Komi PV. Significance of passively induced stretch reflexes on achilles tendon force enhancement. *Muscle Nerve*. 1998;21:1546-8.
101. Melville-Jones G, Watt DG. Muscular control of landing from unexpected falls in man. *J Physiol*. 1971;219:729-37.
102. Enoka RM. *Neuromechanical basis of kinesiology*. 2nd ed. Champaign, IL: Human Kinetics; 1994. p. 1-466.
103. Wilson GJ, Murphy AJ, Pryor JF. Musculotendinous stiffness: its relationship to eccentric, isometric and concentric performance. *J Appl Physiol*. 1994;76:2714-9.
104. Walshe AD, Wilson GJ, Murphy AJ. The validity and reliability of a test of lower body musculotendinous stiffness. *Eur J Appl Physiol Occup Physiol*. 1996;73:332-9.
105. Thys H, Cavagna GA, Margaria R. The role played by elasticity in an exercise involving movements of small amplitude. *Pflugers Arch*. 1975;354:281-6.
106. Chapman AE, Caldwell GE, Selbie WS. Mechanical output following muscle stretch in forearm supination against inertial loads. *J Appl Physiol*. 1985;59:78-86.
107. Wilson GJ, Elliott BC, Wood GA. The effect on performance of imposing a delay during a stretch-shorten cycle movement. *Med Sci Sports Exerc*. 1991;23:364-70.
108. Häkkinen K, Komi PV. The effect of explosive type training on electromyographic and force production characteristics of the leg extensor muscles during concentric and various stretch-shortening cycle exercises. *Scand J Sports Scie*. 1985;7:65-76.
109. Claxton J. Kinematics of explosive upper body movements: A comparison of the traditional bench press, bench press throw and bungy resisted bench press. Auckland: Department of Sport and Health Science, Auckland Institute of Technology; 2001.