




Contents lists available at ScienceDirect

ISPRS Journal of Photogrammetry and Remote Sensing

journal homepage: www.elsevier.com/locate/isprsjprs

Review Article

Optical remotely sensed time series data for land cover classification: A review

Cristina Gómez^{a,b}, Joanne C. White^{c,*}, Michael A. Wulder^c^a INIA, Forest Research Centre, Dpt. of Silviculture and Forest Management, Crta. La Coruña km 7,5, 28040 Madrid, Spain^b Department of Geography and Environment, School of Geoscience, University of Aberdeen, Aberdeen AB24 3UE, Scotland, UK^c Canadian Forest Service (Pacific Forestry Centre), Natural Resources Canada, Victoria, British Columbia V8Z 1M5, Canada

ARTICLE INFO

Article history:

Received 19 January 2016

Received in revised form 7 March 2016

Accepted 8 March 2016

Available online 24 March 2016

Keywords:

Remote sensing

Landsat

Sentinel 2

Monitoring

Composite

Change detection

Mapping

Large area

ABSTRACT

Accurate land cover information is required for science, monitoring, and reporting. Land cover changes naturally over time, as well as a result of anthropogenic activities. Monitoring and mapping of land cover and land cover change in a consistent and robust manner over large areas is made possible with Earth Observation (EO) data. Land cover products satisfying a range of science and policy information needs are currently produced periodically at different spatial and temporal scales. The increased availability of EO data—particularly from the Landsat archive (and soon to be augmented with Sentinel-2 data)—coupled with improved computing and storage capacity with novel image compositing approaches, have resulted in the availability of annual, large-area, gap-free, surface reflectance data products. In turn, these data products support the development of annual land cover products that can be both informed and constrained by change detection outputs. The inclusion of time series change in the land cover mapping process provides information on class stability and informs on logical class transitions (both temporally and categorically). In this review, we present the issues and opportunities associated with generating and validating time-series informed annual, large-area, land cover products, and identify methods suited to incorporating time series information and other novel inputs for land cover characterization.

Crown Copyright © 2016 Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Land cover relates to the biophysical cover of the Earth's terrestrial surface, identifying vegetation, inland water, bare soil or human infrastructure. Distinct land cover types provide specific habitats and determine the exchange of energy and carbon between terrestrial and atmospheric regions (Houghton et al., 2012; Running et al., 1999). Biophysical processes occurring on the land surface have an impact on the climate system (Mahmood et al., 2014) and also on habitat diversity and the capacity of ecosystems to support human needs. Land cover, recognized as an essential climate variable (GCOS, 2003), is the most important element for description and study of the environment (Herold et al., 2006) and is a key input to models of ecosystem services (Andrew et al., 2014).

Characterizing and mapping land cover is essential for planning and managing natural resources (e.g. development, conservation), modeling environmental variables, and for understanding distribution of habitats. Remote sensing and digital image processing

enable observation, identification, mapping, assessment, and monitoring of land cover at a range of spatial, temporal, and thematic scales (Rogan and Chen, 2004). Identification of land cover types provides basic information for generation of other thematic maps, and establishes a baseline for monitoring activities.

Regional and global land cover products have been made possible by the synoptic and periodic observations of satellite Earth Observations (EO) missions, starting in 1972 with Landsat-1 (Belward and Skoien, 2015). Several national land cover mapping programs became operational in the 1990s and the early 2000s (e.g. Vogelmann et al., 2001; Wulder et al., 2008a). National and international agencies have produced regional land cover maps representing single-date conditions (e.g. EOSD 2000 in Canada; CORINE Land Cover 2000 in Europe), with Landsat observations acquired within an interval of the target year (e.g. ±1–3 years). Some programs repeat the land cover mapping periodically (e.g. NLCD 2001/2006/2011 in the US; Australian NCAS-LCCP) and even annually for certain land cover types (e.g. Brazilian PRODES) to enable change assessment.

Global maps have also been produced, typically based on cloud-free composites of coarse spatial resolution data from AVHRR,

* Corresponding author.

SPOT VEGETATION, MODIS or MERIS data (e.g. IGBP-DISCover, GLC2000, MODIS Collection 4 and 5 Land Cover product, GlobCover). Global land cover maps generated from coarse spatial resolution data have typically had low local accuracy (Frey and Smith, 2007; Fritz et al., 2010), particularly in regions with heterogeneous land cover (Herold et al., 2008). Also, due to disparities in classification schemes, spatial resolution, thematic detail, or estimated accuracy, comparison of these maps to inform on change over time is problematic (Bai et al., 2014; Pérez-Hoyos et al., 2012; Pflugmacher et al., 2011) and generally discouraged (Gebhardt et al., 2014; Homer et al., 2007), as inaccuracies in thematic classes compound when change is considered (Fuller et al., 2003). Global land cover maps have also been produced recently with multiple-year data from Landsat (Gong et al., 2013) and single-year Landsat-like imagery (Chen et al., 2014), with reported overall accuracies ranging from 65–80%.

Land cover maps can be updated by identification and mapping of changed areas, leaving unchanged areas in the original map intact. Some large-area land cover programs currently apply such a change-updating approach, for example European CORINE Land Cover (Büttner, 2014) or the US NLCD (Xian et al., 2009). Remote sensing is well suited to monitoring land cover change, and a myriad of approaches have been developed for the purpose (Coppin et al., 2004; Hussain et al., 2013; Lu et al., 2004; Singh, 1989; Tewkesbury et al., 2015). The change-updating approach for production of frequent land cover maps is conceptually appealing, operationally efficient, and economically convenient; however, a potential downside is the introduction of additional sources of error (i.e. via change detection), particularly in heterogeneous and dynamic regions, with frequent and diverse land cover changes. Alternative and robust methods to generate annual large-area land cover maps, such as from time series of EO imagery, require additional investigation and application development.

Until recently, land cover maps generated from coarse spatial resolution data (e.g. 1 km from AVHRR or MODIS) were deemed adequate for ingestion in global and large-area environmental models (Hansen and Loveland, 2012; Loveland et al., 2000). A confluence of data availability, technological capacity, and increasingly detailed information needs has motivated progression of research and operational efforts for generation of temporally, spatially, and thematically improved land cover databases.

Land cover changes can be related to natural processes (e.g. flooding, wildfire) and anthropogenic activities (e.g. urbanization, agriculture). The rate of change and the nature of land cover transitions can also differ in time and space. Some regions are relatively stable (e.g. permanent forest); whereas, other areas are subject to rapid and persistent transformation (e.g. urban expansion of previously vegetated areas). Increased human population and technological development has been found to accelerate land cover change (Goldewijk, 2001; Ramankutty and Foley, 1999). Assessing the location, extent, type, and frequency of land cover transitions, as well as identifying spatial and temporal patterns of change through interpretation and analysis of frequent land cover maps provide insights into underlying processes and drivers of change (Kennedy et al., 2014). Annual land cover information is valuable to aid in formulation of socio-economic policies (e.g. European Common Agriculture Policy) and data provision for environmental applications (e.g. vulnerability and risk assessment).

Maps that relate change between two dates typically lack information regarding underlying processes and do not enable insights on the nature of the transformations present, such as rate or persistence of change (Gillanders et al., 2008). In order to study changes in land cover, ideally both subtle modifications and rapid changes should be accounted for (Lambin et al., 2003). A temporal series of land cover maps can capture the complexities of Earth's changing surface (Liu and Cai, 2012; Sexton et al., 2013a) and can be used

to parameterize biogeochemical models (e.g. Feddema et al., 2005; Running et al., 1994). Furthermore, the level of detail, the definition of land cover categories, and the land cover types selected for generation of a map are often insufficient or inadequate for a broad range of uses, and specific semantics might be misunderstood and wrongly used (Comber et al., 2005). An alternative option is to generate information of essential land cover attributes (e.g. canopy cover) as a versatile approach for generation of widely usable land cover products. Elementary feature layers can support—via rule-based combinations of attributes—multiple classification schemes to suit a range of science communities and policy makers. Demonstrating progress in this direction, estimates of bare ground, woody and herbaceous vegetation proportions are provided globally using MODIS (Hansen et al., 2003), and in the US using Landsat Vegetation Continuous Field products (Hansen et al., 2011). Likewise, the European CORINE Land Cover project also aims to periodically generate five high resolution attribute layers: imperviousness, forest, grassland, wetland, and water bodies (Blanes Guàrdia et al., 2014).

Time series of medium spatial resolution optical data have demonstrated high capacity for characterization of environmental phenomena, describing trends as well as discrete change events. Time series of medium spatial resolution data have been used to map forest disturbance (Kennedy et al., 2010) and surface water bodies (Tulbure and Broich, 2013), to characterize land cover change (Zhu and Woodcock, 2014a) and to identify the nature of land cover changes (Olthof and Fraser, 2014). Time series enable modeling and estimation of ecosystem structural variables and have been used to estimate aboveground biomass (Gómez et al., 2014; Main-Korn et al., 2013), forest carbon sinks (Gómez et al., 2012), forest degradation (Shimabukuro et al., 2014), and forest disturbance (Schroeder et al., 2014). Strategies have been developed to deal with irregular and sparse time series of data (Gómez et al., 2011), but near-anniversary annual series are most appropriate for information extraction in vegetated ecosystems. Moreover, intra-annual time series have proven of great value to acquire phenological insights for land cover mapping (Melaas et al., 2013).

Herein, we investigate current opportunities and issues for generation and validation of temporally dense, large-area land cover products to meet current science, monitoring, and reporting information needs. Our focus is on data and land cover products with sufficient temporal frequency (i.e. annual) and spatial resolution to support monitoring efforts. Specifically, we focus on time series of medium spatial resolution remotely sensed data (e.g. 10–100 m), primarily Landsat (with full awareness of applicability to the forthcoming Sentinel 2 data stream), with time series being defined as a series of consecutive observations, acquired at regular time intervals. In addition, we focus on products that provide accurate estimates of land cover change (including anthropogenic impacts), enable land cover transitions to be characterized, and that are sufficiently flexible or adaptable to provide versatility for a range of applications. We place emphasis on the identification of data and approaches suited to incorporating time series information and novel inputs for land cover characterization.

2. Developments enabling progress: Data, technology, and analysis

2.1. Medium spatial resolution satellite observations

Medium spatial resolution (10–100 m) EO data are captured by an increasing number of satellite missions (Belward and Skøien, 2015). Synergies between operational programs like Landsat and Sentinel-2 (Wulder et al., 2015) will raise the frequency of geometrically and radiometrically compatible acquisitions. Combined, the

Landsat OLI and Sentinel-2 dual system are expected to provide global observations with 2–5 day frequency (Drusch et al., 2012; Irons et al., 2012; Wulder et al., 2015). Satellite sensors appropriate for land cover classification provide consistent and repeatable measurements at an appropriate spatial scale (Verbesselt et al., 2010). Landsat data (TM, ETM+, and OLI) are the standard for land cover classification (Cohen and Goward, 2004) and change detection (Wulder et al., 2008a), as a result of their spatial resolution (30 m), revisit period (16 days; 8 days with multiple Landsat sensors in orbit), and wide spatial coverage (185 × 185 km). Rigorous calibration and consistency in the radiometry of the Landsat sensors, in particular the Thematic Mapper (TM), Enhanced Thematic Mapper Plus (ETM+), and Operational Land Imager (OLI), makes the Landsat image archive a strong example for the benefits of calibration and data interoperability. Given operational imperatives, other data sources (e.g. SPOT, IRS, ASTER) can be availed upon to offer complementary image coverage (Wulder et al., 2008b).

2.2. Landsat pixel-based image composites

Despite long-term plans for periodic systematic acquisitions (Markham et al., 2004; Williams et al., 2006) and the improved accessibility to Landsat data through global archive consolidation efforts (Roy et al., 2014; Wulder et al., 2016) the amount of Landsat data available for many locations is less than desirable (Roy et al., 2010a). In persistently cloudy areas, the number of available observations may be inadequate for analysis (due to cloud, shadow and haze), both within and between years. To help address the limitations of scene-level data quality, pixel-based compositing with Landsat data increased in popularity soon after the opening of the USGS Landsat archive in 2008 (Woodcock et al., 2008). Image compositing is an established approach, common to global and regional studies using lower spatial resolution data (as reviewed in White et al., 2014; Hansen and Loveland, 2012). Prior to free and open access to the Landsat archive (Woodcock et al., 2008), compositing with higher resolution data was largely precluded due to data costs (Wulder et al., 2012) and a lack of image data sources with both large area and time series coverage.

Repeat observations over the same location increase the chance of cloud-free data, even in areas of persistent cloud coverage. Large-area and cloud-free surface reflectance composites can be generated from preferred Landsat observations (Griffiths et al., 2013a; White et al., 2014), selecting the best available observation subject to user defined criteria (e.g. sensor preference, acquisition day of year, free of cloud, etcetera). These best-available-pixel (BAP) composites offer opportunities for monitoring with frequent data in areas where restrictions exist due to cloud persistence. BAP image composites, produced with rule-based criteria and preferences, can be optimized for specific applications (White et al., 2014), including land cover. Furthermore, the spatiotemporal context of the time series provides a means for improvement of residual missing or noisy pixels (i.e. spectral spikes) in image composites (Hermosilla et al., 2015a; Weiss et al., 2014), with these gap-free, infilled composites referred to as proxy BAP composites.

BAP image compositing approaches can take advantage of the robust radiometric calibration associated with the Landsat program, and automated approaches for generating surface reflectance values (e.g. LEDAPS, Masek et al., 2006). Automated processes can also be implemented for cloud and shadow detection and subsequent masking (e.g. Fmask; Zhu and Woodcock, 2012; Zhu et al., 2015a). For efficient, consistent, and timely large-area land cover mapping, scene-based Landsat inputs are not practical (Hansen et al., 2014), and the feasibility of image composites as foundation for generation of large-area products is a topic of active

research (Franklin et al., 2015). To date, Landsat BAP image composites have been used for a range of applications (Table 1).

3. Towards annual land cover products derived from time-series data

The availability of analysis-ready data products, increased technological capacity, and robust time series analysis approaches are guiding the emergence of methods for generation of annual thematic products informing current and historical land cover dynamics. The greatest challenge for novel methods is the complete integration of the temporal dimension into modeling. Temporal series approaches have been demonstrated as superior to single-date methods for a range of applications (Broich et al., 2011; Gómez et al., 2014), including land cover (Franklin et al., 2015).

3.1. Operational strategies for generating annual land cover maps

Methods for generating large-area land cover products must be robust, consistent, and repeatable (Franklin and Wulder, 2002). Although automation can minimize the rate of random error and reduce economic costs, operational experiences have uncovered the difficulties for automated mapping of large areas with acceptable accuracy (Bontemps et al., 2012; Gong et al., 2013). Fundamentally, two approaches can be followed for generation of annual land cover products supported by time-series data. First, independent land cover maps can be generated for each time step (i.e. for each year), using pre- and post-change information derived from the time series (Fig. 1A). This approach assumes that the models used to generate the land cover information are portable, and that land cover signature extension is enabled through the use of surface reflectance values (Song et al., 2001; Gray and Song, 2013). MODIS Collection 5 Global Land Cover product (Friedl et al., 2010) is an example of an annual series of land cover maps produced independently using a time-series of intra-annual spectral data. A series of annual maps derived from inter-annual spectral time series, such as that proposed in Fig. 1A, was demonstrated by Franklin et al. (2015). In this study, the use of time-series metrics in the classification approach improved the overall accuracy by 6.38% relative to single-date results. A second approach for generating annual land cover products from time-series data would be to generate a base map representing reference date conditions for a single year, which could then be updated and backdated annually with change information obtained from the spectral time series (Fig. 1B). In this case annual spectral data provides information to identify the nature of change (e.g. subtle versus drastic), including the no-change option. As an example, Pouliot et al. (2014) mapped the forest area of Canada (2000–2011) annually with MODIS data and a change-updating approach. All efforts to generate time series informed land cover maps benefit from expert knowledge of ecologically feasible land cover transitions, as this contributes in building confidence, evaluating accuracy, and improving classification with an iterative strategy (Liu and Cai, 2012).

The complexity of the land cover characterization problem, together with increasingly detailed information needs for science, monitoring, and reporting, have advanced methods for mapping of single land cover classes (i.e. forest, water) (Frazier and Page, 2000; Schneider et al., 2010), in contrast to the more common multiple-class characterization required for simultaneous and spatially exhaustive mapping. In single-class mapping, binary classifiers segregate the target from all other cover types, thereby saving sampling resources (Foody et al., 2007). Building upon the single-class identification approach, cascade or hierarchical strategies progressively identify land cover types one by one (Chen et al.,

Table 1
Select applications using Landsat pixel-based image composites.

Study	Location	Type of composite	Application
Hansen et al. (2011)	Conterminous United States	Annual, seasonal (summer, autumn), 7 monthly WELD Period: climate year 2008	Land cover characterization
Potapov et al. (2011)	Boreal European Russia	Multi-year (target ± 2 years) Two epochs: 2000/2005	Quantification of gross forest cover loss
Broich et al. (2011)	Sumatra and Kalimantan (Indonesia)	Multi-year (target ± 1 year) Two epochs: 2000/2005	Quantification of gross forest cover loss
Griffiths et al. (2013b)	Carpathian ecoregion (Czech Republic, Austria, Poland, Hungary, Ukraine, Romania, Slovakia)	Annual Period: 2000–2005	Mapping forest disturbance and change in forest type
Sexton et al. (2013b)	Washington-Baltimore (US)	Multi-year (target ± 2 years) Six epochs: 1985/1990/1995/2000/2005/2010 Seasonal (winter and summer) Period: 1984–2010	
Peterson and Nelson (2014)	Alaska (US)	Annual WELD Period: 2003–2012	Estimation of forest height
Hansen et al. (2014)	Conterminous United States	Annual WELD Period: 2006–2010	Land cover monitoring (Forest cover loss and bare ground gain) Crop field extraction
Yan and Roy (2014)	Three tiles (150 \times 150 km) Texas, California, South Dakota (US)	Weekly WELD 5 \times 52 (2006–2010)	
Hermosilla et al. (2015a)	Saskatchewan (Canada)	Proxy annual (summer) Period: 1998–2012	Characterization of per-pixel change
Yan and Roy (2015)	Texas, Kansas, South Dakota (US)	Annual, seasonal (summer, autumn), 7 monthly, and 52 weekly WELD Period: climate year 2010	Land cover characterization
Thompson et al. (2015)	Saskatchewan (Canada)	Proxy annual BAP (2010)	Tree species mapping
Gómez et al. (2015)	Saskatchewan (Canada)	Proxy annual (summer) Period: 1998–2012	Description of landscape processes of change
Zald et al. (2016)	Saskatchewan (Canada)	Proxy annual (summer) Period: 1998–2012	Estimation of biomass




Fig. 1. Strategies for generation of annual land cover maps. (A) Generate an independent classification for each time step (i.e. each year)—assumes models are portable and signature extension is enabled through the use of surface reflectance values. (B) A single classification is generated on a reference year, and this classification is then “rolled” backwards or forwards through the time-series based on spectral/change information.

2014; Sulla-Menashe et al., 2011) minimizing effects of spectral confusion among land cover classes.

Intra-annual and inter-annual spectral series offer different opportunities for a wide range of classification approaches. Intra-annual series inform on the average phenology of individual land cover types, enabling identification of subtle differences and variations over multiple years. Inter-annual series provide medium- to long-term unique spectral profiles of land cover types and inform subtle modifications or drastic changes. Plus, inter-annual series provide evidence in support of possible changes to land cover. Several single date land cover maps have been generated with intra-annual time-series data over large areas (Table 3). Also, some series of annual land cover maps have been produced over large areas (Gebhardt et al., 2014; Friedl et al., 2010; Pouliot et al., 2014).

3.2. Variables derived from spectrotemporal feature space

Spectral signatures of distinct land cover types vary with soil moisture, sun elevation, topography, atmospheric conditions, and view angle (sensor dependent). Spectral variability and signature overlap increase for land cover over large areas and in complex environments (e.g. mixed forests versus coniferous forests). Much of the spectral variability of land cover types can be captured with repeat observations over time, and thus the value of time-series data for discriminating between land cover types. Intra-annual variability is particularly helpful for identification of land cover types dominated by vegetation of marked phenology (e.g. deciduous forest, annual pasture). Phenological differences within a single-year or over consecutive years has helped in discriminating

between cover types such as herbaceous crops and savannah (Senf et al., 2015) or crops and pasture (Müller et al., 2014). Inter-annual variations captured with anniversary measures inform processes of change or stability (Gómez et al., 2015; Lehmann et al., 2013); some processes are only feasible in certain cover types (e.g. stability of rock outcrop), while other processes indicate transitions from one land cover to another (Olthof and Fraser, 2014).

Multi-temporal spectral data encapsulates rich information (Kennedy et al., 2014) that requires specialized tools. Numerous approaches and perspectives are possible for disentangling this multidimensional feature space (Table 2). Multi-temporal spectral data captured at various dates of the same or consecutive years can be summarized by statistical metrics (e.g. average, variability), which in turn can function as descriptive or predictive variables. Spectrotemporal statistical metrics have shown to be a viable means for discrimination of land cover types (DeFries et al., 1995; Gebhardt et al., 2014; Petitjean et al., 2012) and for quantification of forest cover change (Hansen et al., 2014).

A multi-year series of radiometrically calibrated anniversary spectral data is known as a temporal trajectory. Temporal trajectories provide insights of the spectral behavior of corresponding land cover elements over time. A number of methods have been developed to unveil and interpret the rich information provided by temporal trajectories. One strategy partitions the temporal trajectory into linear segments and interprets individual components (LandTrendr; Kennedy et al., 2010). LandTrendr and similar algorithms that segment the temporal trajectory (e.g. C2C, Hermosilla et al., 2015a; DBEST, Jamali et al., 2015) derive descriptive measures of individual segments known as spectrotemporal change metrics

Table 2
Variables derived from the spectrotemporal feature space by different approaches.

Variable type	Characteristics
Statistical metric	<ul style="list-style-type: none"> • Statistical summary of spectral values over one or more periods • Typically data informing seasonality or phenology • <i>Example:</i> average, maximum, minimum
Change metric	<ul style="list-style-type: none"> • Descriptive attribute of a temporal segment • Typically from annual data • <i>Example:</i> magnitude, duration, slope
Shape non-stationary	<ul style="list-style-type: none"> • Pattern of anniversary data • Can/cannot be characterized by parameters • Typically annual data
Shape stationary	<ul style="list-style-type: none"> • Periodic pattern of multiple values per year • Can be described by parameters • <i>Example:</i> sine or cosine based curve
Trend	<ul style="list-style-type: none"> • Non stationary • Annual or longer interval • Admits irregular intervals

(e.g. magnitude, duration). Spectrotemporal change metrics have been used successfully for characterization of land cover (Franklin et al., 2015), as well as forest disturbance and recovery (Hermosilla et al., 2015a; Kennedy et al., 2012).

Temporal trajectories can also be treated with a continuous approach as stationary (e.g. periodic) or non-stationary shape variables (i.e. referring to the shape of the temporal trajectory). Deviations from modeled stationary patterns, derived from multiple observations per year, have been used for classification of land cover types (Zhu and Woodcock, 2014b). Idealized non-stationary shapes have been used for identification of land cover transitions like stand-replacing forest disturbance (Kennedy et al., 2007) and for characterizing the nature of land cover change (Olthof and Fraser, 2014). Temporal trajectories capturing processes of natural succession (e.g. growth) constitute a spectrotemporal signature of specific land cover types. The length or duration of the shape variables necessary to identify significant patterns is presumably variable, as in the estimation of biomass (Gómez et al., 2014; Pflugmacher et al., 2014), and dependent on land cover type. Different strategies for analysis of time series are the decomposition into trend, seasonal, and break elements (BFAST, Verbesselt et al., 2010) enabling analyses of components in relation to landscape processes of change and land cover (DeVries et al., 2015; Verbesselt et al., 2012), or the simple analysis of *trend* (Eastman et al., 2009; Parmentier and Eastman, 2014). The existing range and variety of approaches and more than 30 years of radiometrically and spatially consistent Landsat TM/ETM+ observations will enable retrospective land cover mapping based on the spectrotemporal feature space.

3.3. Classification methods incorporating time series data

The choice of classification algorithm requires considering multiple aspects of the problem: type of data, statistical distribution of classes, target accuracy, ease of use, speed, scalability, and interpretability of the classifier. Direct tradeoffs of some of these factors exist, and it is also important to establish a balance between acceptable accuracy and optimal use of resources. Algorithms that cluster elements by similarity of attributes without *a priori* human intervention (i.e. unsupervised classification) are typically used when scarce knowledge of the land cover types is available (Chen and Gong, 2013; Eva et al., 2004). Clustering algorithms (e.g. k-means, ISODATA) run iteratively until convergence of an optimal set of clusters. Because clusters produced automatically do not

necessarily correspond with the land cover types (Loveland et al., 2000), post-classification refinement techniques (e.g. merging and splitting clusters) are necessary before labelling (e.g. Wulder et al., 2008a). Also, to avoid classes with high internal variance (e.g. water, bare soil, snow) dominating the clustering (Loveland et al., 1991), prior stratification and masking are common practices (Furby et al., 2008; Wulder et al., 2008a). Despite the attractiveness of the automatic character of clustering algorithms, they become time consuming when the data dimension is high or the data volume large (Chen and Gong, 2013), and interpreting clusters properly is a challenging and intensive process. Alternatively, supervised classification approaches assimilate the data to a number of reference land cover samples labelled *a priori*. Selecting an adequate number of good quality training samples is crucial (Bruzzone and Demir, 2014; Foody and Mathur, 2006; Shao and Lunetta, 2012), a time consuming task typically done manually, although accumulated experience and improved databases enable semiautomatic selection in certain conditions (e.g. Radoux et al., 2014). Selecting and labeling samples is not error-free and potentially the cause of poor and biased classification performance (McIver and Friedl, 2002; Pal and Mather, 2006). Supervised methods require that the training data completely represent the classification problem, the classifier is incapable of identifying what is unknown to the training sample (Foody, 2000, 2004). The state of practice for large-area land cover mapping has evolved during the last decade, from a preponderance of unsupervised techniques (Franklin and Wulder, 2002) to an increased use of supervised techniques (Khatami et al., 2016), in part due to increasing ancillary data that facilitates sample collection for training data (e.g. Colditz et al., 2011; Homer et al., 2007; Radoux et al., 2014). Other approaches involve various classifiers used in parallel or in succession (e.g. Bauer et al., 1994; Bontemps et al., 2011; Chen et al., 2014; Tateishi et al., 2011) that can be both supervised and unsupervised. Table 3 illustrates how several of the common classification algorithms have been applied for large-area land cover mapping using time series of optical EO data. Table 4 synthesizes the strengths and weaknesses of algorithms included in Table 3, as identified in the literature.

Partitioning is the only clustering category (Han and Kamber, 2001) widely used for land cover mapping with remotely sensed data, although sporadic efforts of hierarchical clustering exist (Shenthilnath et al., 2011). For large datasets, k-means and ISO-DATA algorithms are preferred, as they are less time consuming than others. Parametric supervised classifiers (e.g. maximum likelihood, minimum distance, discriminant analysis) are difficult to use with multi-temporal data of many spectral features and multimodal distributions (Glanz et al., 2014). As a general rule, their reduced flexibility in decision boundaries makes parametric classifiers unsatisfactory for characterization of land cover in large areas and complex environments (Hubert-Moy et al., 2001). On the other hand, non-parametric classifiers (e.g. k-Nearest Neighbor, kNN; decision trees, DT; neural networks, NN; and Support Vector Machines, SVM) impose boundaries of arbitrary geometries, and provide higher flexibility at the expense of computationally intense iterative processes. In general, non-parametric classifiers that focus decision rules on class boundaries are proficient when the statistics and distribution of land cover types are unknown (Foody and Mathur, 2006; Hansen, 2012)—a common scenario for large areas—thereby making non-parametric classifiers more appropriate than parametric classifiers, which focus on central tendency statistics.

Significant effort has been dedicated to evaluating the performance of land cover classification algorithms, and identifying their relative strengths and shortcomings (Huang et al., 2002; Pal and Mather, 2003, 2005; Rodríguez Galiano et al., 2012). A few efforts have specifically compared algorithms using time-series data (Jia

Table 3

Examples illustrating the range of algorithms and type of variables used for classification of time-series optical data in large area land cover characterization and mapping.

Algorithm	Example	Data	Variables	OCA
Artificial Neural Network	Single date map (1987) Global (Gopal et al., 1999) <i>Fuzzy ARTMAP</i>	AVHRR (1°) Intra-annual series: monthly composites	Spectral parameters NDVI Latitude	0.85
	Single date map (2002) Regional: China (Bagan et al., 2005) <i>Self-Organizing Map</i>	MODIS (500 m) Intra-annual series: Growing season 16-day composites (17 dates)	Spectral parameters EVI	0.91
	Series of maps (1972–1988) Regional: Australia (Furby et al., 2008) <i>Bayesian Network</i>	Landsat (25 m)	Spectral parameters	Not reported
Clustering	Single date map (1992) Regional: Conterminous US NLCD 1992 (Vogelmann et al., 2001) <i>CLUSTER</i>	Landsat (30 m) (leaf-on and leaf-off mosaics)	Spectral parameters (reflectance, NDVI) Ancillary parameters (elevation, slope, aspect, pop. density)	0.6–0.8
	Single date map (2000) Regional: Africa GLC2000 (Mayaux et al., 2004)	SPOT 4VEGETATION (1000 m) Intra-annual series: 10-day and monthly composites	Statistical metrics (maximum monthly NDVI, third lowest albedo index)	Not reported
	Single date map (2000) Regional: Temperate East Asia (Boles et al., 2004) <i>ISODATA</i>	SPOT 4 VEGETATION (1000 m) Intra-annual series: 10-day composites during growing season (27 dates)	Spectral parameters (LSWI, EVI)	0.6
Decision Tree	Series of seven maps (1993–2008) Regional: Mexico (Gebhardt et al., 2014)	Landsat (30 m) All yearly images per scene (964–5706 images) 135 scenes	Statistical metrics NDVI, EVI, SR, ARVI, 6TCT (maximum, minimum, range, average, standard deviation)	0.76
	Single date map (2001) Regional: Conterminous US NLCD 2001 (Homer et al., 2004)	Landsat (30 m) 3 seasonal: spring, summer, fall Nominal year 2001	Spectral parameters (reflectance, TCT, thermal) Topographic parameters	0.85
	Single date map (1987) Global (Friedl and Broadley, 1997)	AVHRR (1°) Intra-annual series: monthly composites	Spectral parameters (maximum monthly NDVI) Latitude	0.88
Ensemble (same base classifier)	Single date map (2000) Global MODIS Collection 4 Land Cover product (Friedl et al., 2002) <i>Boosted decision tree</i>	MODIS (1000 m) Intra-annual series: 16-day composites	Spectral parameters (reflectance, EVI, LST, texture, water mask) Topographic parameters	~0.75
	Series of annual maps (2001–2012) Global MODIS Collection 5 Land Cover product (Friedl et al., 2010) <i>Boosted decision tree</i>	MODIS (500 m) Multiple intra-annual series: 32-day composites	Spectral parameters (reflectance, EVI, LST) Statistical metrics (minimum, maximum, mean annual)	0.75
	Series of annual maps (2001–2010) Regional: Latin America and Caribe (Clark et al., 2012) <i>Random Forest</i>	MODIS (250 m) Multiple intra-annual series: 16-day composites	Statistical metrics (maximum, minimum, range, standard deviation of 3-, 6-, and 12-month periods) (reflectance, NDVI, EVI)	0.85

(continued on next page)

Table 3 (continued)

Algorithm	Example	Data	Variables	OCA
Gaussian Maximum Likelihood	Area and product			
	Single date map Regional: Eurasia (Radoux et al., 2014)	MERIS (300 m) Multi-annual (2008–2012) series: 7-day composites	Spectral parameters (reflectance, summer and autumn composites)	0.82
	Single date map (2013) Regional: China (partial) (Jia et al., 2014a)	Fused Landsat – MODIS Intra-annual series: 16-day composites	Statistical metrics (phenology) NDVI (growing season: begin, end, length, amplitude, maximum)	0.95
	Single date map (1987) Global (Defries and Townshend, 1994)	AVHRR (1°) Intra-annual series: monthly composites	Spectral parameters NDVI Latitude	0.78
Support Vector Machine	Single date map (2000) Regional: Portugal (Gonçalves et al., 2005)	MODIS (500 m) Intra-annual series: 8-day composites	Shape stationary (parameters)	0.78
	Single date map (2000) Regional: Portugal (Carrão et al., 2008)	MODIS (500 m) Intra-annual series: 43 composites	Spectral parameters (Reflectance, NDVI, EVI)	0.88
	Single date map (2003) Regional: Wisconsin (Dash et al., 2007)	MERIS (300 m) Two images: August, September	Spectral parameters Vegetation indices: MGVI, MTCl	0.73

Note. EVI: Enhanced Vegetation Index. LST: Land Surface Temperature. LSWI: Land Surface Water Index. MGVI: MERIS Global Vegetation Index. MTCl: MERIS Terrestrial Chlorophyll Index. NDVI: Normalized Difference Vegetation Index. OCA: Overall Classification Accuracy.

et al., 2014b; Schneider, 2012). The strengths of an algorithm may be general (e.g. easiness of application and interpretation) or they may manifest only in certain scenarios (e.g. capacity to handle missing data) (Table 4). DT, which are based on recursive binary partitions complying with a set of optimized rules (Breiman et al., 1984), are an attractive option for large-area land cover classification for a number of reasons, primarily their ease of application and interpretation, and their capacity to handle data measured on different scales, non-linear relationships, and missing data (Friedl et al., 2002). DT can be trained quickly and perform rapidly (Pal and Mather, 2003); however, DT have been shown to perform less well than algorithms such as SVM and NN in feature spaces with high dimensionality (Hansen, 2012; Pal and Mather, 2003), and are also sensitive to noisy observations and over-fitting (Ghimire et al., 2012). Random Forest (RF) (Breiman, 1996) is an improved implementation of a DT, which votes for the best tree out of a forest of trees created by recursive modification of the sampling data (Belgiu and Drăguț, 2016). RF provides higher classification accuracy than other forms of DT and avoids over-fitting, at the cost of increased computational intensity and a black box nature that obfuscates decision rules (Rodríguez Galiano et al., 2012). SVM algorithms identify one or more hyperplanes separating target groups in a multi-dimensional space. SVM is less sensitive to the Hughes phenomenon (Hughes, 1968)—whereby classification accuracy decreases when the number of input features to the classifier is over a given threshold—than other algorithms, and performs well with a large number of variables relative to training size (Pal and Mather, 2005; Shao and Lunetta, 2012), a significant benefit when abundant remotely sensed data exists and little ground truth is available. Neural networks are accurate classifiers, but tend to overfit the data (Rodríguez Galiano et al., 2012) and remain a black box for interpretation. Both NN and SVM are computationally intense and require a number of parameters to be adjusted (Table 4).

Combining the strengths of various algorithms into an ensemble classifier tends to increase accuracy (Bauer and Kohavi, 1999) and provides information on classification uncertainty (Foody et al., 2007) or confidence (McIver and Friedl, 2001), but it also tends to reduce interpretability and increase computational complexity and overhead (Table 4). Ensemble learning algorithms use the same or a combination of base classifiers to produce multiple classifications of the same data (e.g. Random Forest, bagging, boosting) (Rodríguez Galiano et al., 2012). Ensemble approaches can be dependent, whereby the output of one classifier is used to inform the next classifier, or independent, whereby each classifier is run independently and the outputs are combined using some weighting or voting mechanism (Rokach, 2010). Bagging methods resample the original training set with replacement each time, while boosting methods resample or reweigh the training data by emphasizing more those instances that are misclassified by previous classifiers. Boosting has been found to be a convenient technique for large classification problems (Friedl et al., 2010).

The incorporation of complex temporal data into land cover classification has not been paired with novel classification algorithms and in most cases the same rules are applied as for classifying single-date data. A notable exception by Liu and Cai (2012) demonstrated the utility of considering spectrotemporal context in the classifier. It has been noted that the inclusion of time-series data can have a more positive impact on results (i.e. accuracy) than the choice of classification algorithm (Jia et al., 2014b). Novel algorithms are needed to leverage the predictive potential of time-series data; consider the case of hyperspectral and multispectral datasets, whereby techniques that function well with single-date data, may not offer the same performance when using time-series data (Bruzzone and Demir, 2014).

Table 4
Strengths and weaknesses of algorithms used for large-area land cover characterization with time-series optical data.

Algorithm	Strengths/characteristics	Weaknesses
Artificial Neural Networks Non-parametric	<ul style="list-style-type: none"> • Manage well large feature space • Indicate strength of class membership • Generally high classification accuracy • Resistant to training data deficiencies—requires less training data than DT 	<ul style="list-style-type: none"> • Needs parameters for network design • Tends to overfit data • Black box (rules are unknown) • Computationally intense • Slow training
Clustering (partitioning)	<ul style="list-style-type: none"> • Do not need previous knowledge • Do not need samples 	<ul style="list-style-type: none"> • Cluster-class correspondence not assured • Complex identification of classes • Computationally intense
Decision trees Non-parametric	<ul style="list-style-type: none"> • No need of any kind of parameter • Easy to apply and interpret • Handle missing data • Handle data of different types (e.g. continuous, categorical) and scales • Handle non-linear relationships • Insensitive to noise 	<ul style="list-style-type: none"> • Sensitive to noise • Tend to overfit • Not as good as others in large feature spaces • Large training sample needed
Gaussian Maximum likelihood Parametric	<ul style="list-style-type: none"> • Simple application • Easy to understand and interpret • Predicts class membership probability 	<ul style="list-style-type: none"> • Parametric • Assumes normal distribution of data • Large training sample needed
Support Vector Machines Non-parametric	<ul style="list-style-type: none"> • Manages well large feature space • Insensitive to Hughes effect • Works well with small training dataset • Does not overfit 	<ul style="list-style-type: none"> • Needs parameters: regularization and kernel • Poor performance with small feature space • Computationally intense • Designed as binary, although variations exist
Random Forests Non-parametric	<ul style="list-style-type: none"> • Capacity to determine variable importance • Robust to data reduction • Does not over-fit • Produces unbiased accuracy estimate • Higher accuracy than DT 	<ul style="list-style-type: none"> • Decision rules unknown (black box) • Computationally intense • Needs input parameters (#trees and #variables per node)
Bagging	<ul style="list-style-type: none"> • Provides measures of classification confidence • Does not overfit 	<ul style="list-style-type: none"> • Complex incomprehensible classifiers
Boosting	<ul style="list-style-type: none"> • Provides measures of classification confidence • Does not overfit • Robust to noise 	<ul style="list-style-type: none"> • Stops if a classifier achieves zero training set error • Complex incomprehensible classifiers • Ineffective if excessive error in training sample

4. Novel inputs for land cover classification

Progress in the development of robust methods for the generation of repeat land cover products over large areas is made possible with contributions from novel inputs that were not necessary or possible before. Knowledge of ecological succession can facilitate refinement of a time series of land cover maps, while information on land cover dynamics can be used to impose model restrictions on short- and medium-term land cover transitions, contributing to the temporal consistency of land cover products. Variables with an inherent temporal dimension and temporal trajectories of multiple vegetation indices provide complementary insights regarding the status and change of essential biophysical attributes associated with land cover types. Improved quality of training samples, and combinations of multi-scale and multi-sensor data for enhancement of temporal and spatial resolution has direct impacts on the accuracy of land cover characterization.

4.1. Land cover transitions and temporal stabilization of classification

The likely character of land cover transitions depends on the time interval considered and on the environmental context. Likely successional transitions (e.g. open to dense forest) may take just a few years to manifest in tropical areas but longer in boreal or Mediterranean regions. Possible land cover transitions by loss of vegetation (e.g. shrub to bare ground) could happen rapidly anywhere following a sudden disturbance. Furthermore, transitions related to subtle, long-term processes occur in different directions (e.g. canopy closure by natural growth, loss of density by disease-related mortality, change in dominant species). Understanding ecological processes is essential for production of reliable time

series land cover maps, regardless of the mapping strategy applied. Insights of possible and likely transitions at different time intervals can be used to impose logical restrictions for generation of consecutive maps (Liu and Cai, 2012; Pouliot and Latifovic, 2013) and also for refinement and verification. Land cover transitions can be identified by modifications in spectral seasonal curves over several years (Gutiérrez-Vélez and DeFries, 2013; Reed et al., 1994) and by specific patterns or trends in inter-annual spectral trajectories that can inform on change types (Olthof and Fraser, 2014).

Developing accurate and consistent annual land cover maps requires dedicated techniques for temporal stabilization of the classification results (Friedl et al., 2010). Furthermore, for some change assessment applications, temporal consistency is more important than the overall accuracy of individual maps (Radoux et al., 2014). Markov Random Field (MRF) models are particularly attractive techniques in temporal pattern recognition, as a result of their ability to identify relationships of temporally dependent data, and have been applied in land cover map series to identify and correct illogical class transitions (i.e. transitions that are ecologically impossible) (Wehmann and Liu, 2015), and to incorporate spatial context into classifications (Li et al., 2014). Liu and Cai (2012) constructed land cover change trajectories from time-series Landsat data with a transition probability model based on MRF. The temporal-contextual constraints imposed by the model improved temporal consistency of a series of seven Landsat-based land cover maps. Cai et al. (2014) applied the same model to a MODIS Collection 5 Global Land Cover product, detecting illogical transitions and improving the accuracy and temporal consistency of the product in five tiles with abundant illogical transitions. Similarly, Wang et al. (2015) applied the maximum a posteriori MRF in post-classification processing, as

	Snow/Ice	Rock/Rubble	Exposed land	Water	Shrub-Tall	Shrub-Low	Herb	Bryoids	Wetland-Treed	Wetland-Shrub	Wetland-Herb	Coniferous-Dense	Coniferous-Open	Coniferous-Sparse	Broadleaf-Dense	Broadleaf-Open	Broadleaf-Sparse	Mixed Wood-Dense	Mixed Wood-Open	Mixed Wood-Sparse	
	TO																				
FROM	31	32	33	20	51	52	100	40	81	82	83	211	212	213	221	222	223	231	232	233	
Snow/Ice	31	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Rock/Rubble	32	1	3	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
Exposed land	33	1	0	3	0	0	0	1	1	0	0	1	0	0	0	0	0	0	0	0	0
Water	20	1	0	0	3	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0
Shrub-Tall	51	1	0	1	0	3	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
Shrub-Low	52	1	0	1	0	1	3	1	0	0	0	0	0	0	0	0	0	0	0	0	0
Herb	100	1	0	1	0	0	1	3	0	0	0	1	0	0	0	0	0	0	0	0	0
Bryoids	40	1	1	1	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0
Wetland-Treed	81	1	0	0	1	0	0	0	0	3	1	1	0	0	0	0	0	0	0	0	0
Wetland-Shrub	82	1	0	0	1	0	0	0	0	1	3	1	0	0	0	0	0	0	0	0	0
Wetland-Herb	83	1	0	0	1	0	0	1	0	0	2	3	0	0	0	0	0	0	0	0	0
Coniferous-Dense	211	1	0	1	0	0	0	0	0	0	0	0	3	2	1	0	0	0	0	0	0
Coniferous-Open	212	1	0	1	0	0	0	0	0	0	0	0	2	3	2	0	0	0	0	0	1
Coniferous-Sparse	213	1	0	1	0	0	0	0	0	0	0	0	2	3	0	0	0	0	0	1	1
Broadleaf-Dense	221	1	0	1	0	0	0	0	0	0	0	0	0	0	0	3	2	1	0	0	0
Broadleaf-Open	222	1	0	1	0	0	0	0	0	0	0	0	0	0	0	2	3	1	0	0	0
Broadleaf-Sparse	223	1	0	1	0	0	0	0	0	0	0	0	0	0	0	2	3	0	0	0	1
Mixed Wood-Dense	231	1	0	1	0	0	0	0	0	0	0	0	1	1	0	1	1	3	1	1	1
Mixed Wood-Open	232	1	0	1	0	0	0	0	0	0	0	0	1	1	0	0	1	1	3	1	1
Mixed Wood-Sparse	233	1	0	1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	2	3	3

0 Not likely 1 Possible 2 Probable 3 Likely

Likely: stability
Probable: succession, can happen any time
Possible: not impossible but not normal
Not likely: illogical or rare transition

Fig. 2. Class transition matrix for the land cover types present over the forested area of Canada (after [Wulder et al., 2008a](#)). For a single year period four kinds of transition are considered according to their likelihood to occur: from “likely” (stability or no change) and “probable” (natural succession transition), through “possible” (transition that could happen) to “not likely” (ecologically illogical transition). “Not likely” transitions point to potential error in the prior or/and later land cover classification.

a spatial-temporal consistency model, improving the classification of single date maps produced with multi-year MODIS data. Other approaches incorporating transitions in the process of classification have also demonstrated an improvement to the temporal consistency of land cover maps series. For example, working with annual series of land cover class probabilities derived from MODIS via Random Forest classifiers, [Yin et al. \(2014\)](#) identified land cover transitions by the change in probability of occurrence for the five land cover types of their classification system. Post-classification approaches that apply temporal filters with transition rules ([Clark et al., 2010](#)) or change transition matrices ([Pouliot et al., 2014](#)) have also been shown to be useful for identification and amendment of illogical transitions. Utilizing the legend relating the land cover present over the forested area of Canada (after [Wulder et al., 2008a](#)) a class transition matrix is generated and presented ([Fig. 2](#)). The above logic is incorporated to create scores in the transition matrix from likely, through probable and possible, to not likely. Such a matrix can be used to reconcile the consistency across annual land cover maps and to incorporate and inform with knowledge of change (such as from spectral trends or change detection). Class stability in a non-changing pixel-series can be enforced through rules. More sophisticated approaches to use of the class transition matrix may incorporate disturbance and rules informed by knowledge of successional trends and spectral evidence.

4.2. Multi-scale and multi-sensor combination of spectral data

Bringing together medium spatial and high temporal data from complementary sensors like Landsat and MODIS or MERIS has been shown to be successful for land cover applications ([Amorós-López et al., 2013](#); [Gong et al., 2013](#)). Combining these data sources is increasingly frequent, either via concurrent use or by fusing data sets to generate synthetic images. Supported by the spatial detail of Landsat, the high temporal data from MODIS/MERIS informs phenological traits and helps identify difficult land cover types ([Jia et al., 2014a, 2014c](#)). In their Landsat-based global map, [Gong et al. \(2013\)](#) supported training data selection with MODIS Enhanced Vegetation Index (EVI) time series from 2010, aiming to improve spectral separation between cultivated bare land and natural barren lands, and to distinguish sparsely vegetated land from more persistent barren land.

Among data fusion algorithms, the Spatial and Temporal Adaptive Reflectance Fusion Model (STARFM) ([Gao et al., 2006](#)) with improvements ([Hilker et al., 2009](#); [Zhu et al., 2010](#)) stands out for the algorithm’s ability to generate Landsat-like images, although unmixing-based algorithms and the Spatial and Temporal Reflectance Unmixing Model (STRUM) are preferable when few Landsat data are available ([Gevaert and García-Haro, 2015](#)). In a land cover change updating approach context, [Chen et al. \(2015\)](#) employed downscaled intra-annual MODIS NDVI time series for

identification of real rather than spurious change, improving the overall land cover classification accuracy from 0.76 to 0.89. Jia et al. (2014a) fused Landsat and MODIS data via STARFM into 23 16-day composites and used statistical metrics (maximum, minimum, mean, and standard deviation) for supervised classification improving single date classification accuracies by ~4%. In this case no land cover change is assumed during the period of blended data.

Despite successful efforts blending MODIS and Landsat data, an important restriction for this approach is the limited availability of MODIS data, going back just to year 2000. Also, Senf et al. (2015) demonstrated this type of synthetic seasonal data improves single date land cover classification, but the authors show that the results from the synthetic data are inferior to those from the original Landsat data. Synthetic Landsat images free of clouds, shadows, snow and SLC-off gaps, and without phenological and sun orientation issues can also be generated from Landsat models in regions where abundant observations are already available (Zhu et al., 2015b). The capacity of the synthetic Landsat data developed by Zhu et al. (2015b) for land cover mapping has yet to be demonstrated. Another approach for improving land cover characterization is to combine data from different temporal scales acquired by a single sensor. For instance, both an annual composite and a time series of bimonthly MERIS composites were used to improve classification results of the GlobCover 2005 and 2009 products (Bontemps et al., 2012). The advantages of using data from a single sensor are in the consistency of the spatial registration and radiometric calibration.

4.3. Multiple temporal spectral variables

For accurate characterization of land cover types and optimal detection of land cover transitions multiple spectral variables are needed (Zhu and Woodcock, 2014a). The temporal trajectory of vegetation indices like the Normalized Burnt Ratio (NBR) is particularly useful for identification of land cover transitions associated with drastic loss of vegetation cover (e.g. forest to bare soil), because values are markedly changed after years of relative stability (Hermosilla et al., 2015b). Other spectral variables provide information of change causality (Hais et al., 2009; Schroeder et al., 2011) informing possible land cover transitions. Furthermore, subtle processes responsible of transitions between similar cover types (e.g. open forest to dense forest by recruitment or growth) can only be identified by temporal spectral trajectories robust to noise and sensitive to variations in vegetation density (e.g. Tasseled Cap Angle). Smoothing algorithms have also been applied to reduce noise inherent in the time series data (e.g. Shao et al., 2016). For generation of rich and detailed land cover dynamics databases over large areas, multiple spectral variables are required. Parmentier and Eastman (2014) combined time series of various vegetation indices and successfully identified land cover transitions, although the contribution of each variable remained unsolved.

Including multiple spectrottemporal variables in the classification problem increases the feature space, hampering some classifiers' performance. Dimensionality reduction by linear or non-linear techniques (Yan and Roy, 2015) facilitates data compression while retaining relevant information. Principal Component Analysis (PCA) is a well understood dimensionality reduction technique, but its components can be difficult to interpret (Homer et al., 2004). Vegetation indices are an efficient way to reduce the amount of data and some of them provide the additional advantage of being linked to vegetation's physical characteristics (e.g. Tasseled Cap Transformation).

4.4. Optimized training sample data

Reference data should represent all land cover types within the area of interest, and should be defined in the context of the classi-

fication approach to be applied. Classifiers based on statistical distributions (e.g. Maximum likelihood) perform better with pure samples (Lillesand and Kiefer, 2008), whereas classifiers with rules targeting boundary regions between classes (e.g. SVM, DT) benefit from critical border samples (Foody, 2004; Hansen, 2012). Non-parametric methods such as kNN require the reference data to represent the full range of variability in the population, as these methods cannot extrapolate beyond the range of the training data. The adequate size of training sample is also algorithm dependent (Foody and Mathur, 2006; Pal and Mather, 2003) and although no guidelines exist, a general assumption is that the size of the training sample should be proportional to the independent space size (i.e. the number of variables in the model). Classification algorithms frequently suffer from inability to handle high-dimensional data. Therefore, supervised classifications aim to target an adequate sample with a minimum number of elements that maximize accuracy. To improve the accuracy/sample size relation, dimensionality reduction via feature transformation (e.g. PCA; Tasseled Cap Transformation, TCT) and feature selection (Elghazel and Aussem, 2015) are useful tools.

Acquiring reference data is expensive and time consuming, but also critical for the accuracy of classification results (Shao and Lunetta, 2012). The use of time-series data for land cover classification presents additional challenges for reference data sampling. Signature extension by generalization of a single set of class spectral signatures across time and space might yield substantial inter-class mixing. Distributing samples in space and time with the objective of capturing spectral variability (Sexton et al., 2013a, 2013b) relies on good radiometric normalization between image dates. Alternatively, automatic and adaptive generalization of spectral signatures in temporally irregular (non-anniversary) time series (Gray and Song, 2013) is possible. Annual series of BAP image composites represent a seamless radiometrically corrected dataset that enables internally consistent and repeatable land cover characterization (Hansen et al., 2013; Pengra et al., 2015). Furthermore, datasets with a temporal dimension (e.g. inter-annual and intra-annual spectral series) provide opportunities to augment the sample size, assimilating patterns of identical spectrottemporal trajectories across space and time to a good reference (Xue et al., 2014) with reliable measures of time series similarity (e.g. Dynamic Time Warping or variations; Maus et al., in press). Active learning (semi-supervised) methods may also optimize the choice of training samples, proposing training samples to the operator in an iterative form until satisfactory classification is achieved (Tuia et al., 2011). Active learning can be beneficial at the initial phases of the mapping project, reducing the expensive phase of redundant sample labelling (Bruzzone and Demir, 2014), but can be difficult to implement over large areas.

5. Validation and accuracy assessment

Validation of land cover products is essential to demonstrate the quality of the products for operational applications (Cihlar, 2000; Foody, 2002). The accuracy of classified data has to be assessed and reported with informative metrics adequate for the user community. Although there are no standard methods to assess the accuracy of thematic maps, best practices have emerged in the literature, indicating the inclusion of an error or confusion matrix (Foody, 2002), which can aid in identifying confusion between classes, as well as potential sources of error. Furthermore, quantitative metrics that are derived from the confusion matrix, such as overall accuracy, as well as area-weighted metrics and confidence intervals are useful measures (Olofsson et al., 2014). Validation of large-area land cover products is challenging and requires appropriate sampling strategies for statistical assessment

of the product's accuracy. Moreover, the reference data must have sufficient accuracy to enable robust validation (Wulder et al., 2006). The reliability of large-area land cover maps is frequently tested against reference maps produced at a higher spatial resolution assuming good, although not perfect accuracy.

Validation of two or more consecutive land cover maps with reference datasets is a difficult task, because acquiring reference data for multiple years over large areas can be unfeasible, even if auxiliary maps exist at certain dates. Adequate reference data is particularly elusive if land cover products span a historical period. In such cases, manual interpretation of the Landsat time-series data is becoming an accepted approach for generating the required reference data (Cohen et al., 2010). In the change-based updating approach (Fig. 1B), if the area changed is proportionally small, the accuracy of land cover maps in a time series is often assumed to be close to that of the base map (Pouliot and Latifovic, 2013). In theory, validation of a time series of land cover and change detection results can be achieved with independent error matrices (Mertens and Lambin, 2000; Yuan et al., 2005). However, to ensure robustness, this approach requires no-change samples to be acquired as a component of the reference data (Olofsson et al., 2014). Post-classification comparison of a time series of land cover maps can aid in identifying illogical land cover transitions in space and time (Liu and Cai, 2012), which, in well-registered maps of the same spatial resolution, can be indicative of classification error (Townsend et al., 2009). Moreover, explicit information of change incorporated in land cover products provides a powerful tool for self-assessment and validation. The temporal consistency between consecutive maps can, to some extent, be evaluated against land cover changes. Acceptable land cover transitions in consecutive maps should conform to the time interval separating them, that is, short intervals impose tighter restrictions and provide more reliable judgement. For example, in an annual series of land cover maps, grassland can transit to pasture but not to open forest, but the same transitions would be more uncertain in a five year period. Permitted transitions between land cover classes provide a means for self-assessment, but only partially, because even ecologically logical class transitions could be incorrect.

Two main difficulties arise with validation via logical land cover transitions. First, transitions associated with subtle or progressive spectral variations are difficult to assess, because enough evidence or historical data are seldom available. However, a short delay in detection of such transitions should be considered a minor error compared with, for example, missing a forest to bare ground disturbance-related transition. Second, validation of transitions becomes even more challenging with a higher number or very similar classes (Olofsson et al., 2013). Likewise, confusion between discrete classes of a continuum (e.g. between herb and shrub) does not have the same implications as other confusion types (e.g. between herb and water). Adapted or new measures of error are also needed to report the accuracy of time series land cover maps (e.g. Tsutsumida and Comber, 2015).

6. Synthesis: Issues and opportunities

Despite progress in technology and data availability, generation of large-area land cover products is challenging. While some of these challenges relate to the migration of algorithms from a research to an operational phase (e.g. handling and processing massive amounts of data), we have identified some challenges specific to the use of time-series data, such as annual image BAP composites, for land cover mapping. We have also identified opportunities afforded by novel data analysis techniques developed by the remote sensing community and other scientific disci-

plines. These techniques merit further exploration for the purpose of land cover characterization with time-series data.

6.1. Time-series data and temporal variables

Generating Landsat annual BAP image composites (Roy et al., 2010b; White et al., 2014) and improved annual series of proxy image composites (Hermosilla et al., 2015a) over large areas has been demonstrated. Until now, single-date land cover maps have been produced with multiple intra-annual composites (Yan and Roy, 2014). When generating a series of annual land cover maps, consideration of phenological information from seasonal image composites (Roy et al., 2010b)—in areas where there is sufficient data to do so—would presumably facilitate better discrimination of land cover types. However, the irregular availability of medium spatial resolution data over large areas and across many years, as well as in high latitudes with persistent snow coverage, may jeopardize the robustness and spatial-temporal consistency of the land cover product. MODIS data can support Landsat data via combination or fusion, and inform on phenological traits for dates after 1999 only (Chen et al., 2015; Jia et al., 2014a).

Among the common approaches applied thus far for characterizing the spectrottemporal domain of Landsat time series (e.g. change metrics, trends), some are better suited for certain applications. Statistical metrics have demonstrated efficiencies for synthesizing temporal data to distinguish among various land cover types, but their specificity is not demonstrated: various land cover types could be represented by the same statistical metrics over a certain period if the metrics are not properly defined. Change metrics describe the evolution over time of biophysical parameters correlated with the spectral features observed. Change metrics are intuitive and easy to characterize and interpret. The choice of an optimal set of change metrics, including single or multiple spectral features, as well as the interval prior and after the target date, are crucial and application specific. Shape variables are visually appealing and easy to interpret, but characterizing and modeling the domain of *shape* variables is not an easy task. The relevance of slight pattern variations, which might be difficult to capture with parametric or functional approaches, is also application dependent; for example, a shortly delayed maximum in a density indicator does not modify biomass estimation (Gómez et al., 2014) but is a determinant of land cover change types (Olthoff and Fraser, 2014). Parameterized trends are not as intuitive as other variables, but have been effective in some land cover problems (Eastman et al., 2009; Parmentier and Eastman, 2014). The capacity of these approaches to provide temporal variables that are resilient to inter-annual reflectance variability and capable of robustly and consistently discriminating land cover types in annual series of large-area land cover maps needs to be demonstrated. The spatio-temporal context provided by an annual series of proxy images and insights of ecologically feasible land cover transitions facilitate development of meaningful rules with which to define spatially and temporally consistent maps. Hermosilla et al. (2015b) present an example for using Random Forests to label disturbance types combining change metrics, object-level spectral information, and the geometric characteristics of the objects. Disturbance year is assigned using a breakpoint analysis of the Landsat time series, with 90% accuracy found for the disturbance types labeled over the 30-year analysis period.

6.2. Emerging options for automatic classification of land cover

Despite the widespread incorporation of temporal data into approaches for characterizing the status and dynamics of ecosystems, and recognition of the superiority of models that integrate temporal data, development of novel land cover classification

methods that take full advantage of this information remains to be realized. Temporal data are operationally analyzed disregarding the full capacity of the time dimension and presumably missing the opportunity to unveil more detailed information. We mention here some techniques that are novel or are being used in an exploratory phase, and that could benefit the land cover mapping community, some of which incorporate the temporal dimension.

Unsupervised clustering is currently only marginally used in large-area land cover mapping (e.g. GlobCover 2009; Wulder et al., 2008a), despite the attractiveness of its automatic character and independence from field samples. In the context of time series informed land cover, clustering deserves renewed attention because the strengths of novel techniques could be decisive for the land cover problem. Traditional clustering focuses on discovering a single grouping of the data, which does not necessarily correspond with the land cover types, particularly when dealing with multidimensional feature domains. Some novel approaches define multiple sets of clusters that describe alternative aspects and characterize the data in different ways. Three novel clustering paradigms are attracting considerable research effort (Müller et al., 2015): alternative clustering, clustering ensemble, and subspace clustering. Alternative clustering and clustering ensemble search for an alternative to a given solution and a consensus of various clustering options respectively, and neither of them seem practical for the large-area land cover problem. In contrast, subspace clustering focuses on detecting multiple clusters in arbitrary, sometimes overlapping, subspace projections of the high dimensional data (Parsons et al., 2004). For land cover characterization with time-series features, subspace clustering is an attractive tool for exploration and identification of relevant subsets of variables that characterize essential land cover attributes individually. Subspace clustering also provides a means for interpretation of complex data spaces.

Clustering of time series differs from clustering of static data mainly in how the similarity between elements is computed (Liao, 2005). Dynamic Time Warping, Mahalanobis, or Triangle distances are measures generally preferred over Euclidean distance, but the adequacy of a time series similarity measure is application dependent (Gorecki, 2014; Lhermitte et al., 2008). Time-series spectral data can be directly clustered as raw data, or indirectly clustered via features (e.g. statistics, change metrics) or modeled series (e.g. shape, trends) (Liao, 2005). The second approach (i.e. features) facilitates reduction of noise and data size, as well as classification of sequences of unequal length (e.g. missing data). Techniques for clustering continuous series are used in medical (Aach and Church, 2001; Wismüller et al., 2002), pattern recognition (Liao, 2005; Maharaj, 1999), and finance applications, accommodating single and sometimes multiple variables (Ramoni et al., 2000), and with hidden Markov models frequently underlying time series models for clustering (Zhang et al., 2011).

In supervised classification, machine learning and ensemble algorithms are more accurate and efficient than conventional parametric classifiers when faced with large dimensional and complex data spaces, and are currently used for large-area land cover mapping. However, supervised classification methods can benefit from semi-automated techniques for speeding up training data collection (Schneider, 2012). Semi-supervised learning (SSL) (also known as *deep learning* and *active learning*) methods for classification exploit both training data and unlabeled samples in the learning phase of the classifier, in order to obtain a general decision function that considers the training set information and the structure of all data in the feature space (Bruzzone and Demir, 2014). SSL can therefore outperform other classifiers when few training samples are available. SSL deals with the intrinsic variability of spectral signatures in land cover classes and has already been used for land cover characterization with hyperspectral data (Camps-Valls et al., 2007; Jun and Ghosh, 2013).

Another relatively new classification technique with potential application in large-area land cover mapping with time-series data is the Relevance Vector Machine (RVM), a Bayesian extension of the SVM (Tipping, 2001). As a kernel method like SVM, RVM classifies large dimensional data with few training samples, achieving high accuracy. Unlike SVM, RVM provides an estimate of the posterior probability of class membership, which may be used to illustrate the uncertainty of the class allocations (Foody, 2008). RVM provides more flexibility for the choice of kernel type than SVM, and its implementation does not require the penalty and kernel parameters, saving considerable computational resources (Gómez-Chova et al., 2011). RVM is a technique frequently used in pattern recognition (Caesarendra et al., 2010; Wang et al., 2009), and to date, just a few studies can be found in the land cover literature, using optical (Foody, 2008) and hyperspectral (Camps-Valls and Bruzzone, 2005; Demir and Ertürk, 2007) data.

7. Summary

Characterization of large-area land cover dynamics is necessary to inform monitoring, reporting, science, and policy. The increased availability of EO data at a range of spatial and temporal resolutions, coupled with increasingly complex information needs for science and policy making, and technological improvements in image processing capacity and storage, guide the advance of methods to generate frequent and accurate land cover products. Time-series spectral data provides better characterization of land cover types, with intra-annual series informing on phenological differences and inter-annual series informing on land cover dynamics.

Techniques for data analysis and interpretation that fully incorporate the temporal dimension remain an area of intense research and represent an important challenge for operational mapping. For land cover characterization, intra-annual series of coarse spatial data have long informed on phenological traits and differences among land cover types. Medium spatial resolution intra-annual series with less frequency (i.e. Landsat) are feasible in certain areas, but not a global reality, due to variation in data availability. Inter-annual time series, on the other hand, are facilitated by BAP image composites, which compile the best possible observation at each pixel location, subject to user criteria (e.g. day of acquisition, sensor preference). Annual series of gap-free proxy image composites represent a valuable source of data for generation of time series land cover maps as well as other applications.

To date, information from Landsat time-series data has taken the form of statistical metrics, change metrics, or pattern/trend components. These metrics or components are subsequently used in applications, such as land cover mapping. Despite the temporal character of the data employed, classifiers used in operational mapping are those typically used for single date spectral data classification; however, the special traits of time-series data are quickly driving evolution of novel techniques, such as RVM. Improvement of existing approaches, as well as inclusion of novel techniques, imported and adapted from other disciplines, is important to make the most of data available and to produce annual land cover maps that meet a wide range of needs. Subspace clustering, time series clustering, and semi-supervised learning are techniques with potential to facilitate annual large-area land cover maps production.

Validation of annual series of land cover maps, and particularly historical maps, requires methods not fully dependent on field or ancillary reference data. Ecological insights of land cover transitions over different time intervals provide opportunities for assessment of land cover products. Incorporating spatiotemporal context into classification and validation improves the accuracy and temporal consistency of land cover time series. In this context, the

quality of annual series of proxy image composites, without data gaps or anomalous values, represents a real opportunity for generation of accurate and consistent annual land cover maps.

Through this review we have demonstrated the need and related utility of annual land cover products to support increasingly complex information needs for monitoring, reporting, science, and policy. Time-series data offers new opportunities and challenges for land cover mapping. There is currently a dearth of land cover classification methods that can take full advantage of the rich information available from time-series data. This review presented a range of options for source image data (via image archives, image compositing, synthetic data generation) as well as a number of classification routines that avail upon both the spatial and temporal information present. The production of land cover maps was among the first applications of satellite multispectral remotely sensed data. Over four decades later, new and improved classification algorithms continue to emerge. Furthermore, sole reliance on spectral information for classification purposes is not required with supplemental datasets available to inform on conditions that can further aid in differentiation between classes. Incorporation of the temporal element into classification approaches is already benefiting from sophisticated spatial and spectral algorithms, pointing to the potential for further improvements in land cover classification outcomes. It is unlikely that there will be a single, or optimal, method for land cover mapping incorporating time-series data. Rather, approaches that incorporate spatial, spectral, and temporal data, and the knowledge of the underlying processes (e.g. phenology, disturbance, and succession) of the classes of interest are poised to meet a wide range of information needs.

Acknowledgements

This research was undertaken as part of the “National Terrestrial Ecosystem Monitoring System (NTEMS): Timely and detailed national cross-sector monitoring for Canada” project jointly funded by the Canadian Space Agency (CSA) Government Related Initiatives Program (GRIP) and the Canadian Forest Service (CFS) of Natural Resources Canada. We appreciate the time, effort, and insight offered by the journal editors and reviewers.

References

- Aach, J., Church, G.M., 2001. Aligning gene expression time series with time warping algorithms. *Bioinformatics* 17 (6), 495–508.
- Amorós-López, J., Gómez-Chova, L., Alonso, L., Guanter, L., Zurita-Milla, R., Moreno, J., Camps-Valls, G., 2013. Multitemporal fusion of Landsat/TM and ENVISAT/MERIS for crop monitoring. *Int. J. Appl. Earth Obs. Geoinf.* 23, 132–143.
- Andrew, M.E., Wulder, M.A., Nelson, T.A., 2014. Potential contributions of remote sensing to ecosystem service assessments. *Prog. Phys. Geogr.* 38 (3), 328–353.
- Bagan, H., Wang, Q., Watanabe, M., Yang, Y., Ma, J., 2005. Land cover classification from MODIS EVI times-series data using SOM neural network. *Int. J. Remote Sens.* 26 (22), 4999–5012.
- Bai, Y., Feng, M., Jiang, H., Wang, J., Zhu, Y., Liu, Y., 2014. Assessing consistency of five global land cover data sets in China. *Remote Sens.* 6, 8739–8759.
- Bauer, M.E., Kohavi, R., 1999. An empirical comparison of voting classification algorithms: bagging, boosting, and variants. *Mach. Learn.* 36, 105–142.
- Bauer, M.E., Burk, T.E., Ek, A.R., Coppin, P.R., Lime, S.D., Walsh, T.A., Walters, D.K., Befort, W., Heinzen, D.F., 1994. Satellite inventory of Minnesota forest resources. *Photogramm. Eng. Remote Sens.* 60 (3), 287–298.
- Belgiu, M., Drăguț, L., 2016. Random forests in remote sensing: a review of applications and future directions. *ISPRS J. Photogramm. Remote Sens.* 114, 24–31.
- Belward, A.S., Sköien, J.O., 2015. Who launched what, when and why; trends in global land cover observation capacity from civilian earth observation satellites. *ISPRS J. Photogramm. Remote Sens.* 103, 115–128.
- Blanes Guàrdia, N., Green, T., Simón, A., 2014. The user's role in the European land monitoring context. In: Manakos, I., Braun, M. (Eds.), *Land Use and Land Cover Mapping in Europe*. Springer, Dordrech Heidelberg, New York, London, pp. 31–42.
- Boles, S.H., Xiao, X., Liu, J., Zhang, Q., Munkhutuya, S., Chen, S., Ojima, D., 2004. Land cover characterization of Temperate East Asia using multi-temporal VEGETATION sensor data. *Remote Sens. Environ.* 90, 477–489.
- Bontemps, S., Defourny, P., Bogaert, E.V., Arino, O., Kalogirou, V., Perez, J.R., 2011. Globcover 2009-Products Description and Validation Reports. Technical Report for ESA GlobCover project: UCLouvain & ESA Team pp. 53.
- Bontemps, S., Arino, O., Bicheron, P., Brockman, C.C., Leroy, M., Vancutsem, C., Defourny, P., 2012. Operational service demonstration for global land cover mapping: the GlobCover and GlobCorine experiences from 2005 and 2009. In: Giri, Ch P. (Ed.), *Remote Sensing of Land Use and Land Cover, Principles and Applications*. Taylor and Francis Group, CRC Press, Boca Raton/London/New York, pp. 243–264.
- Breiman, L., Friedman, J.H., Olshen, R.A., Stone, C.J., 1984. *Classification and Regression Trees*. Chapman and Hall/CRC, Boca Raton, FL, USA, p. 358.
- Breiman, L., 1996. Bagging predictors. *Mach. Learn.* 24, 123–140.
- Broich, M., Hansen, M.C., Potapov, P., Adusei, B., Lindquist, E., Stehman, S.V., 2011. Time-series analysis of multi-resolution optical imagery for quantifying forest cover loss in Sumatra and Kalimantan, Indonesia. *Int. J. Appl. Earth Obs. Geoinf.* 13, 277–291.
- Bruzzone, L., Demir, B., 2014. A review of modern approaches to classification of remote sensing data. In: Manakos, I., Braun, M. (Eds.), *Land use and land cover mapping in Europe*. Springer, Dordrech Heidelberg, New York, London, pp. 127–143.
- Büttner, G., 2014. CORINE Land cover and land cover change products. In: Manakos, I., Braun, M. (Eds.), *Land Use and Land Cover Mapping in Europe*. Springer, Dordrech Heidelberg, New York, London, pp. 31–42.
- Caesarendra, W., Widodo, A., Yang, B.S., 2010. Application of relevance vector machine and logistic regression for machine degradation assessment. *Mech. Syst. Signal Process.* 24 (4), 1161–1171.
- Cai, Sh., Liu, D., Sulla-Menashe, D., Friedl, M.A., 2014. Enhancing MODIS land cover product with a spatial-temporal modeling algorithm. *Remote Sens. Environ.* 147, 243–255.
- Camps-Valls, G., Bruzzone, L., 2005. Kernel-based methods for hyperspectral image classification. *IEEE Trans. Geosci. Remote Sens.* 43, 1351–1362.
- Camps-Valls, G., Bandos Marsheva, T.V., Zhou, D., 2007. Semi-supervised graph-based hyperspectral image classification. *IEEE Trans. Geosci. Remote Sens.* 45 (10), 3044–3054.
- Carrão, H., Gonçalves, P., Caetano, M., 2008. Contribution of multispectral and multitemporal information from MODIS images to land cover classification. *Remote Sens. Environ.* 112, 986–997.
- Chen, Y., Gong, P., 2013. Clustering based on eigenspace transformation – CBEST for efficient classification. *ISPRS J. Photogramm. Remote Sens.* 83, 64–80.
- Chen, J., Chen, J., Liao, A., Cao, X., Chen, L., Chen, X., He, Ch., Han, G., Peng, Sh., Lu, M., Zhang, W., Tong, X., Mills, J., 2014. Global land cover mapping at 30 m resolution: a POK-based operational approach. *ISPRS J. Photogramm. Remote Sens.*
- Chen, X., Yang, D., Chen, J., Cao, X., 2015. An improved automated land cover updating approach by integrating with downscaled NDVI time series. *Remote Sens. Lett.* 6 (1), 29–38.
- Cihlar, J., 2000. Land cover mapping of large areas from satellites: status and research priorities. *Int. J. Remote Sens.* 21, 1093–1114.
- Clark, M.L., Aide, T.M., Grau, H.R., Riner, G., 2010. A scalable approach to mapping annual land cover at 250 m using MODIS time-series data: a case study in the Dry Chaco ecoregion of South America. *Remote Sens. Environ.* 114, 2816–2832.
- Clark, M.L., Aide, T.M., Riner, G., 2012. Land change for all municipalities in Latin America and the Caribbean assessed from 250-m MODIS imagery (2001–2010). *Remote Sens. Environ.* 126, 84–103.
- Cohen, W.B., Goward, S., 2004. Landsat's role in ecological applications of Remote Sensing. *Bioscience* 54, 535–545.
- Cohen, W.B., Zhiqiang, Y., Kennedy, R., 2010. Detecting trends in forest disturbance and recovery using yearly Landsat time series: 2. TimeSync—Tools for calibration and validation. *Remote Sens. Environ.* 114 (12), 2911–2924.
- Colditz, R.R., Schmidt, M., Conrad, C., Hansen, M.C., Dech, S., 2011. Land cover classification with coarse spatial resolution data to derive continuous and discrete maps for complex regions. *Remote Sens. Environ.* 115, 3264–3275.
- Comber, A., Fisher, P., Wadsworth, R., 2005. What is land cover? *Environ. Plan. B: Plan. Des.* 32, 199–209.
- Coppin, P., Jonckherre, I., Nackaerts, K., Muys, B., 2004. Digital change detection methods in ecosystem monitoring: a review. *Int. J. Remote Sens.* 25, 1565–1596.
- Dash, J., Mathur, A., Foody, G.M., Curran, P.J., Chipman, J.W., Lillesand, T.M., 2007. Land cover classification using multi-temporal MERIS vegetation indices. *Int. J. Remote Sens.* 28 (6), 1137–1159.
- DeFries, S., Townshend, J.R.G., 1994. NDVI-derived land cover classifications at a global scale. *Int. J. Remote Sens.* 15 (17), 3567–3586.
- DeFries, S., Hansen, M., Townshend, J., 1995. Global discrimination of land cover types from metrics derived from AVHRR Pathfinder data. *Remote Sens. Environ.* 54, 209–222.
- Demir, B., Ertürk, S., 2007. Hyperspectral image classification using relevance vector machines. *IEEE Geosci. Remote Sens. Lett.* 4, 586–590.
- DeVries, B., Verbesselt, J., Kooistra, L., Herold, M., 2015. Robust monitoring of small-scale forest disturbances in a tropical montane forest using Landsat time series. *Remote Sens. Environ.* 161, 107–121.
- Drusch, M., Del Bello, U., Carlier, S., Colin, O., Fernandez, V., Gascon, F., Hoersch, B., Isola, C., Laberinti, P., Marimort, P., Meygret, A., Spoto, F., Sy, O., Marchese, F.,

- Bargellini, . Sentinel-2: ESA's optical high-resolution mission for GMES operational services. *Remote Sens. Environ.* 120, 25–36.
- Elghazel, H., Aussem, A., 2015. Unsupervised feature selection with ensemble learning. *Mach. Learn.* 98, 157–180.
- Eastman, J.R., Sangermano, F., Ghimire, B., Zhu, H., Chen, H., Neeti, N., Cai, Y., Machado, E.A., Crema, S.C., 2009. Seasonal trend analysis of image time series. *Int. J. Remote Sens.* 30 (10), 2721–2726.
- Eva, H., Belward, A.S., DeMiranda, E.E., Dibellaz, C.M., Gond, V., Hubber, O., Jonesk, S., Sgrenzaroli, M., Fritz, S., 2004. A land cover map of South America. *Glob. Change Biol.* 10, 731–744.
- Feddema, J.J., Oleson, K.W., Bonan, G.B., Mearns, L.O., Buja, L.E., Meehl, G.A., Washington, W.M., 2005. The importance of land-cover change in simulating future climates. *Science* 310, 1674–1678.
- Footy, G.M., 2000. Estimation of sub-pixel land cover composition in the presence of untrained classes. *Comput. Geosci.* 26, 469–478.
- Footy, G.M., 2002. Status of land cover accuracy assessment. *Remote Sens. Environ.* 80, 185–201.
- Footy, G.M., 2004. Supervised image classification by MLP and RBF neural networks with and without an exhaustively defined set of classes. *Int. J. Remote Sens.* 25 (15), 3091–3104.
- Footy, G.M., Mathur, A., 2006. The use of small training sets containing mixed pixels for accurate hard image classification: training on mixed spectral responses for classification by a SVM. *Remote Sens. Environ.* 103, 179–189.
- Footy, G.M., Boyd, D.S., Sánchez-Hernández, C., 2007. Mapping a specific class with an ensemble of classifiers. *Int. J. Remote Sens.* 28 (8), 1733–1746.
- Footy, G.M., 2008. RVM-based multi-class classification of remotely sensed data. *Int. J. Remote Sens.* 29 (6), 1817–1823.
- Franklin, S.E., Wulder, M.A., 2002. Remote sensing methods in medium spatial resolution satellite data land cover classification of large areas. *Prog. Phys. Geogr.* 26 (2), 173–205.
- Franklin, S.E., Ahmed, O.S., Wulder, M.A., White, J.C., Hermosilla, T., Coops, N.C., 2015. Large area mapping of annual land cover dynamics using multi-temporal change detection and classification of Landsat time-series data. *Can. J. Remote Sens.* 41, 293–314.
- Frazier, P., Page, K., 2000. Water body detection and delineation with Landsat TM data. *Photogramm. Eng. Remote Sens.* 66 (12), 1461–1467.
- Frey, K.E., Smith, L.C., 2007. How well do we know northern land cover? Comparison of four global vegetation and wetland products with a new ground-truth database for West Siberia. *Global Biogeochem. Cycles* 21, GB1016.
- Friedl, M.A., Brodley, C.E., 1997. Decision tree classification of land cover from remotely sensed data. *Remote Sens. Environ.* 61, 399–409.
- Friedl, M.A., McIver, D.K., Hodges, J.C.F., Zhang, X.Y., Muchoney, D., Strahler, A.H., Woodcock, C.E., Gopal, S., Schneider, A., Cooper, A., Baccini, A., Gao, F., Schaaf, C., 2002. Global land cover mapping from MODIS: algorithms and early results. *Remote Sens. Environ.* 83, 287–302.
- Friedl, M.A., Sulla-Menashe, D., Tan, B., Schneider, A.M., Ramakutty, N., Sibley, A., Huang, X., 2010. MODIS collection 5 global land cover: algorithm refinements and characterization of new datasets. *Remote Sens. Environ.* 114, 168–182.
- Fritz, S., See, L., Rembold, F., 2010. Comparison of global and regional land cover maps with statistical information for the agricultural domain in Africa. *Int. J. Remote Sens.* 31, 2237–2256.
- Fuller, R.M., Smith, G.M., Devereux, B.J., 2003. The characterisation and measurement of land cover change through *Remote Sensing*: problems in operational applications? *Int. J. Appl. Earth Obs. Geoinf.* 4, 243–253.
- Furby, S., Caccetta, P., Wu, X., Chia, J., 2008. Continental scale land cover change monitoring in Australia using Landsat imagery. In: *International Earth Conference: Studying, Modeling and Sense Making of Planet Earth*, Mytilene, Lesvos, Greece, June 2008, pp. 1–8.
- Gao, F., Masek, J., Schwaller, M., Hall, F., 2006. On the blending of the Landsat and MODIS surface reflectance: predicting daily Landsat surface reflectance. *IEEE Trans. Geosci. Remote Sens.* 44, 2207–2218.
- GCOS, 2003. The second report on the adequacy of the Global Observing System for Climate Support of the UNFCCC, GCOS-82. Secretariat of the World Meteorological Organization: Geneva, Switzerland, 74 pp.
- Gebhardt, S., Wehrmann, T., Muñoz Ruiz, M.A., Maeda, P., Bishop, J., Schramm, M., Kopeinig, R., Cartus, O., Kellndorfer, J., Ressler, R., Santos, L.A., Schmidt, M., 2014. MAD-MEX: automatic wall-to-wall land cover monitoring for the Mexican REDD-MRV program using all Landsat data. *Remote Sens.* 6, 3926–3943.
- Gevaert, C.M., García-Haro, F.J., 2015. A comparison of STARFM and an unmixing-based algorithm for Landsat and MODIS data fusion. *Remote Sens. Environ.* 156, 34–44.
- Ghimire, B., Rogan, J., Rodríguez Galiano, V., Panday, P., Neeti, N., 2012. An evaluation of bagging, boosting, and Random Forests for land cover classification in Cape Cod, Massachusetts, USA. *GIScience Remote Sens.* 49 (5), 623–643.
- Gillanders, S.N., Coops, N.C., Wulder, M.A., Gergel, S.E., Nelson, T., 2008. Multitemporal remote sensing of landscape dynamics and pattern change: describing natural and anthropogenic trends. *Prog. Phys. Geogr.* 35 (5), 502–528.
- Glanz, H., Carvalho, L., Sulla-Menashe, D., Friedl, M.A., 2014. A parametric model for classifying land cover and evaluating training data based on multi-temporal *Remote Sensing* data. *ISPRS J. Photogramm. Remote Sens.* 97, 219–228.
- Goldewijk, K.K., 2001. Estimating global land use change over the past 300 years: the HYDE database. *Global Biogeochem. Cycles* 15 (2), 417–433.
- Gómez-Chova, L., Muñoz Marí, J., Laparra, V., Malo-López, J., Camps-Valls, G., 2011. A review of kernel methods in *Remote Sensing* data analysis. In: Prasad, S., Bruce, L.M., Chaussoot, J. (Eds.), *Optical Remote Sensing. Advances in Signal Processing and Exploitation Techniques*. Springer-Verlag, Berlin, Heidelberg, pp. 171–206.
- Gómez, C., White, J.C., Wulder, M.A., 2011. Characterizing the state and processes of change in a dynamic forest environment using hierarchical spatio-temporal segmentation. *Remote Sens. Environ.* 115, 1665–1679.
- Gómez, C., Wulder, M.A., White, J.C., Montes, F., Delgado, J.A., 2012. Characterizing 25 years of change in the area, distribution, and carbon stock of Mediterranean pines in Central Spain. *Int. J. Remote Sens.* 33 (17), 5546–5573.
- Gómez, C., White, J.C., Wulder, M.A., Alejandro, P., 2014. Historical forest biomass dynamics modelled with Landsat spectral trajectories. *ISPRS J. Photogramm. Remote Sens.* 93, 14–28.
- Gómez, C., White, J.C., Wulder, M.A., Alejandro, P., 2015. Integrated object-based spatiotemporal characterization of forest change from an annual time series of Landsat image composites. *Can. J. Remote Sens.* 41 (4), 271–292.
- Gonçalves, P., Carrão, H., Pinheiro, A., Caetano, M., 2005. Land cover classification with support vector machine applied to MODIS. In: *Proceedings of the 25th EARSeL Symposium*, 2005.
- Gong, P., Wang, J., Yu, L., Zhao, Y., Zhao, Y., Liang, L., Niu, Z., Huang, X., Fu, H., Liu, S., Li, C., Li, X., Fu, W., Liu, C., Xu, Y., Wang, X., Cheng, Q., Hu, L., Yao, W., Zhang, H., Zhu, P., Zhao, Z., Zhang, H., Zheng, Y., Ji, L., Zhang, Y., Chen, H., Yan, A., Guo, J., Yu, L., Wang, L., Liu, X., Shi, T., Zhu, M., Chen, Y., Yang, G., Tang, P., Xu, B., Giri, C., Clinton, N., Zhu, Z., Chen, J., Chen, J., 2013. Finer resolution observation and monitoring of GLC: first mapping results with Landsat TM and ETM+ data. *Int. J. Remote Sens.* 34 (7), 2607–2654.
- Gopal, S., Woodcock, C.E., Strahler, A.H., 1999. Fuzzy Neural Network classification of global land cover from a 1° AVHRR Data Set. *Remote Sens. Environ.* 67, 230–243.
- Gorecki, T., 2014. Using derivatives in a longest common subsequence dissimilarity measure for time series classification. *Pattern Recogn. Lett.* 45, 99–105.
- Gray, J., Song, C., 2013. Consistent classification of image time series with automatic adaptive signature generalization. *Remote Sens. Environ.* 134, 333–341.
- Griffiths, P., Van der Linden, S., Kuemmerle, T., Hostert, P., 2013a. A pixel-based Landsat compositing algorithm for large area land cover mapping. *IEEE J. Sel. Top. Appl. Earth Observations Remote Sens.* 6 (5), 2088–2101.
- Griffiths, P., Kuemmerle, T., Baumann, M., Radeloff, V.C., Abrudan, I.V., Lieskovsky, J., Munteanu, C., Ostapowicz, K., Hostert, P., 2013b. Forest disturbances, forest recovery, and changes in forest types across the Carpathian ecoregion from 1985 to 2010 based on Landsat image composites. *Remote Sens. Environ.* 151, 72–88.
- Gutiérrez-Vélez, V.H., DeFries, R., 2013. Annual multi-resolution detection of land cover conversion to oil palm in the Peruvian Amazon. *Remote Sens. Environ.* 129, 154–167.
- Hais, M., Jonášová, M., Langhammer, J., Kučera, T., 2009. Comparison of two types of forest disturbance using multitemporal Landsat TM/ETM+ imagery and field vegetation data. *Remote Sens. Environ.* 113, 835–845.
- Han, J., Kamber, M., 2001. *Data Mining: Concepts and Techniques*. Morgan Kaufmann, San Francisco, pp. 384–466.
- Hansen, M.C., DeFries, R.S., Townshend, J., Carroll, M., Dimiceli, C., Sohlberg, R., 2003. Global percent tree cover at a spatial resolution of 500 m: first results of the MODIS Vegetation Continuous Fields algorithm. *Earth Interact.* 7, 1–15.
- Hansen, M.C., Egorov, A., Roy, D.P., Potapov, P., Ju, J., Turubanova, S., Kommareddy, I., Loveland, T.R., 2011. Continuous fields of land cover for the conterminous United States using Landsat data: first results from the Web-Enabled Landsat Data (WELD) project. *Remote Sens. Lett.* 2 (4), 279–288.
- Hansen, M.C., 2012. Classification trees and mixed pixel training data. In: Giri, Ch P. (Ed.), *Remote Sensing of Land Use and Land Cover, Principles and Applications*. Taylor and Francis Group, CRC Press, Boca Raton/London/New York, pp. 127–136.
- Hansen, M.C., Loveland, T.R., 2012. A review of large area monitoring of land cover change using Landsat data. *Remote Sens. Environ.* 122, 66–74.
- Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau, D., Stehman, S.V., Goetz, S.J., Loveland, T.R., Kommareddy, A., Egorov, A., Chini, L., Justice, C.O., Townshend, J.R.G., 2013. High-resolution global maps of 21st century forest cover change. *Science* 342 (6160), 850–853.
- Hansen, M.C., Egorov, A., Potapov, P.V., Stehman, S.V., Tyukavina, A., Turubanova, S. A., Roy, D.P., Goetz, S.J., Loveland, T.R., Ju, J., Kommareddy, A., Kovalsky, V., Forsyth, C., Bents, T., 2014. Monitoring conterminous United States (CONUS) land cover change with Web-Enabled Landsat Data (WELD). *Remote Sens. Environ.* 140, 466–484.
- Hermosilla, T., Wulder, M.A., White, J.C., Coops, N.C., Hobart, G.W., 2015a. An integrated Landsat time series protocol for change detection and generation of annual gap-free surface reflectance composites. *Remote Sens. Environ.* 158, 220–234.
- Hermosilla, T., Wulder, M.A., White, J.C., Coops, N.C., Hobart, G.W., 2015b. Regional detection, characterization, and attribution of annual forest change from 1984 to 2012 using Landsat-derived time-series metrics. *Remote Sens. Environ.* 170, 121–132.
- Herold, M., Latham, J.S., DiGregorio, A., Schmullius, C.C., 2006. Evolving standards in land cover characterization. *J. Land Use Sci.* 1 (2–4), 157–168.
- Herold, M., Mayaux, P., Woodcock, C.E., Baccini, A., Schmullius, C., 2008. Some challenges in global land cover mapping: an assessment of agreement and accuracy in existing 1 km datasets. *Remote Sens. Environ.* 112, 2538–2556.
- Hilker, T., Wulder, M.A., Coops, N.C., Linke, J., McDermid, G., Masek, J.G., Gao, F., White, J.C., 2009. A new data fusion model for high spatial- and temporal-resolution mapping of forest disturbance based on Landsat and MODIS. *Remote Sens. Environ.* 113 (8), 1613–1627.

- Homer, C., Huang, C., Yang, L., Wylie, B., Coan, M., 2004. Development of a 2001 national land cover database for the United States. *Photogramm. Eng. Remote Sens.* 70 (7), 829–840.
- Homer, C., Dewitz, J., Fry, J., Coan, M., Hossain, N., Larson, Ch., Herold, N., McKerrow, A., VanDriel, J.N., Wickham, J., 2007. Completion of the 2001 National Land Cover Database for the conterminous United States. *Photogramm. Eng. Remote Sens.* 73, 337–341.
- Houghton, R.A., House, J.I., Pongratz, J., van der Werf, G.R., DeFries, R.S., Hansen, M. C., LeQuere, C., Ramankutty, N., 2012. Carbon emissions from land use and land cover change. *Biogeosciences* 9, 5125–5142.
- Huang, C., Davis, L.S., Townshend, J.R.G., 2002. An assessment of support vector machines for land cover classification. *Int. J. Remote Sens.* 23, 725–749.
- Hubert-Moy, L., Cotonnec, A., Le Du, L., Chardin, A., Pérez, P., 2001. A comparison of parametric classification procedures of remotely sensed data applied on different landscape units. *Remote Sens. Environ.* 75, 174–187.
- Hughes, G.F., 1968. On the mean accuracy of statistical pattern recognizers. *IEEE Trans. Inform. Theory* 14 (1), 55–63.
- Hussain, M., Chen, D., Cheng, A., Wei, H., Stanley, D., 2013. Change detection from remotely sensed images: from pixel-based to object-based approaches. *ISPRS J. Photogramm. Remote Sens.* 80, 91–106.
- Irons, J.R., Dwyer, J.L., Barsi, J.A., 2012. The next Landsat satellite: the Landsat data continuity mission. *Remote Sens. Environ.* 122, 11–21.
- Jamali, S., Jönsson, P., Eklundh, L., Ardö, J., Seaquist, J., 2015. Detecting changes in vegetation trends using time series segmentation. *Remote Sens. Environ.* 156, 182–195.
- Jia, K., Liang, S., Zhang, N., Wei, X., Gu, X., Zhao, X., Yao, Y., Xie, X., 2014a. Land cover classification of finer remote sensing data integrating temporal features from time series coarser resolution data. *ISPRS J. Photogramm. Remote Sens.* 93, 49–55.
- Jia, K., Liang, S., Wei, X., Yao, Y., Su, Y., Jiang, B., Wang, X., 2014b. Land cover classification of Landsat data with phenological features extracted from time series MODIS NDVI data. *Remote Sens.* 6, 11518–11532.
- Jia, K., Liang, S., Wei, X.Q., Zhang, L., Yao, Y.J., Gao, S., 2014c. Automatic land cover update approach integrating iterative training sample selection and a Markov Random Field model. *Remote Sens. Lett.* 5, 148–156.
- Jun, G., Ghosh, J., 2013. Semisupervised learning of hyperspectral data with unknown land-cover classes. *IEEE Trans. Geosci. Remote Sens.* 51 (1), 273–282.
- Kennedy, R.E., Cohen, W.B., Schroeder, T.A., 2007. Trajectory-based change detection for automated characterization of forest disturbance dynamics. *Remote Sens. Environ.* 110, 370–386.
- Kennedy, R.E., Ziquiang, Y., Cohen, W., 2010. Detecting trends in forest disturbance and recovery using yearly Landsat time series: 1. LandTrendr – temporal segmentation algorithms. *Remote Sens. Environ.* 114, 2897–2910.
- Kennedy, R.E., Zhiqiang, Y., Cohen, W.B., Pfaff, E., Braaten, J., Nelson, P., 2012. Spatial and temporal patterns of forest disturbance and regrowth within the area of the Northwest Forest Plan. *Remote Sens. Environ.* 122, 117–133.
- Kennedy, R.E., Andréfouët, S., Gómez, C., Griffiths, P., Hais, M., Healey, S., Helmer, E. H., Hostert, P., Lyons, M., Meigs, G.W., Pflugmacher, D., Phinn, S., Powell, S., Scarth, P.F., Sen, S., Schroeder, T.A., Schneider, A.M., Sonnenschein, R., Vogelmann, J.E., Wulder, M.A., Zhu, Z., 2014. Bringing an ecological view of change to Landsat-based Remote Sensing. *Frontiers Ecol. Environ.* 12 (6), 339–346.
- Khatami, R., Mountrakis, G., Stehman, S.V., 2016. A meta-analysis of remote sensing research on supervised pixel-based land cover image classification processes: general guidelines for practitioners and future research. *Remote Sens. Environ.* 177, 89–100.
- Lambin, E.F., Geist, H.J., Lepers, E., 2003. Dynamics of land-use and land cover change in tropical regions. *Annu. Rev. Environ. Resour.* 28, 205–341.
- Lehmann, E.A., Wallace, J.F., Caccetta, P.A., Furby, S.L., Zdunic, K., 2013. Forest cover trends from time series Landsat data for the Australian continent. *Int. J. Appl. Earth Observation Geoinformation* 21, 453–462.
- Lhermitte, S., Verbesselt, J., Verstraeten, W.W., Coppin, P., 2008. A comparison of time series similarity measures for classification and change detection of ecosystem dynamics. *Remote Sens. Environ.* 115, 3129–3152.
- Li, M., Zang, Sh., Zhang, B., Li, Sh., Wu, Ch., 2014. A review of *Remote Sensing* image classification techniques: the role of spatio-contextual information. *Eur. J. Remote Sens.* 47, 389–411.
- Liao, T.W., 2005. Clustering of time-series data. *Pattern Recogn.* 38, 1857–1874.
- Lillesand, T.M., Kiefer, R.W., 2008. *Remote Sensing and Image Interpretation*. John Wiley and Sons, New York, p. 736.
- Liu, D., Cai, S., 2012. A spatial-temporal modeling approach to reconstructing land-cover change trajectories from multi-temporal satellite imagery. *Ann. Assoc. Am. Geogr.* 102, 1329–1347.
- Loveland, T.R., Merchant, J.W., Ohlen, D.O., Brown, J.F., 1991. Development of a land-cover characteristics database for the conterminous U.S. *Photogramm. Eng. Remote Sens.* 57 (11), 1453–1463.
- Loveland, T.R., Reed, B.C., Brown, J.F., Ohlen, D.O., Zhu, Z., Yang, L., Merchant, J.W., 2000. Development of a global land cover characteristics database and IGBP DISCover from 1 km AVHRR data. *Int. J. Remote Sens.* 21, 1303–1330.
- Lu, D., Mausel, P., Bronzdzio, E., Moran, E., 2004. Change detection techniques. *Int. J. Remote Sens.* 25, 2365–2407.
- Maharaj, E.A., 1999. Comparison and classification of stationary multivariate time series. *Pattern Recogn.* 32, 1129–1138.
- Mahmood, R., Pielke, R.A., Hubbard, K.G., Ngoyi, D., Dirmeyer, P.A., McAlpine, C., Carleton, A.M., Hale, R., Gameda, S., Beltran-Przekurat, A., Baker, B., McNider, R., Legates, D.R., Shepherd, M., Du, J., Blanken, P.D., Frauenfeld, O.W., Nair, U.S., Fall, S., . Land cover changes and their biogeophysical effects on climate. *Int. J. Climatol.* 34, 929–953.
- Main-Korn, M., Cohen, W.B., Kennedy, R.E., Grodzki, W., Pflugmacher, D., Griffiths, P., Hostert, P., 2013. Monitoring coniferous forest biomass change using a Landsat trajectory approach. *Remote Sens. Environ.* 139, 277–290.
- Markham, B.L., Storey, J.C., Williams, D.L., Irons, J.R., 2004. Landsat sensors performance: history and current status. *IEEE Trans. Geosci. Remote Sens.* 42 (12), 2691–2694.
- Masek, J.G., Vermote, E.F., Saleous, N.E., Wolfe, R., Hall, F.G., Huemmrich, K.F., Gao, F., Kutler, J., Lim, T.K., 2006. A Landsat surface-reflectance dataset for North America, 1990–2000. *IEEE Geosci. Remote Sens. Lett.* 3 (1), 68–72.
- Maus, V., Câmara, G., Cartaxo, R., Sanchez, A., Ramos, F.M., de Queiroz, G.R., in press. A time-weighted dynamic time warping method for land-use and land-cover mapping. *IEEE J. Sel. Topics Appl. Earth Observations Remote Sens.* <http://dx.doi.org/10.1109/JSTARS.2016.2517118>.
- Mayaux, P., Bartholomé, E., Fritz, S., Belward, A., 2004. A new land cover map of Africa for the year 2000. *J. Biogeogr.* 31, 861–877.
- McIver, D.K., Friedl, M.A., 2001. Estimating pixel-scale land cover classification confidence using nonparametric machine learning methods. *IEEE Trans. Geosci. Remote Sens.* 39, 1959–1968.
- McIver, D.K., Friedl, M.A., 2002. Using prior probabilities in decision-tree classification of remotely sensed data. *Remote Sens. Environ.* 81, 253–261.
- Melaas, E.K., Friedl, M.A., Zhu, Z., 2013. Detecting interannual variation in deciduous broadleaf forest phenology using Landsat TM/ETM+ data. *Remote Sens. Environ.* 132, 176–185.
- Mertens, B., Lambin, E., 2000. Land-cover-change trajectories in southern Cameroon. *Ann. Assoc. Am. Geogr.* 90 (3), 467–494.
- Müller, H., Rufin, Ph., Griffiths, P., Barros Siqueira, A.J., Hostert, P., 2014. Mining dense Landsat time series for separating cropland and pasture in a heterogeneous Brazilian savanna landscape. *Remote Sens. Environ.* 156, 490–499.
- Müller, E., Assent, I., Günemann, S., Seidl, T., Dy, J., 2015. MultiClust special issue on discovering, summarizing and using multiple clusterings. *Mach. Learn.* 98, 1–5.
- Olofsson, P., Foody, G.M., Herold, M., Stehman, S.V., Woodcock, C.E., Wulder, M.A., 2014. Good practices for estimating area and assessing accuracy of land change. *Remote Sens. Environ.* 148, 42–57.
- Olofsson, P., Foody, G.M., Stehman, S.V., Woodcock, C.E., 2013. Making better use of accuracy data in land change studies: estimating accuracy and area and quantifying uncertainty using stratified estimation. *Remote Sens. Environ.* 129, 122–131.
- Olthof, I., Fraser, R.H., 2014. Detecting landscape changes in high latitude environments using Landsat trend analysis: 2. Classification. *Remote Sens.* 6 (11), 11558–11578.
- Pal, M., Mather, P.M., 2003. An assessment of the effectiveness of decision tree methods for land cover classification. *Remote Sens. Environ.* 86, 554–565.
- Pal, M., Mather, P.M., 2005. Support vector machines for classification in Remote Sensing. *Int. J. Remote Sens.* 26 (5), 1007–1011.
- Pal, M., Mather, P.M., 2006. Some issues in the classification of DAIS hyperspectral data. *Int. J. Remote Sens.* 27 (14), 2895–2916.
- Parmentier, B., Eastman, J.R., 2014. Land transitions from multivariate time series: using seasonal trend analysis and segmentation to detect land cover changes. *Int. J. Remote Sens.* 35 (2), 671–692.
- Parsons, L., Haque, E., Liu, H., 2004. Subspace clustering for high dimensional data: a review. *SigKdd Explor.* 6 (1), 90–105.
- Pengra, B., Long, J., Dahal, D., Stehman, S.V., Loveland, T.R., 2015. A global reference database from very high resolution commercial satellite data and methodology for application to Landsat derived 30 m continuous field tree cover data. *Remote Sens. Environ.* 165, 234–248.
- Pérez-Hoyos, A., García-Haro, F.J., SanMiguel-Ayanz, J., 2012. Conventional and fuzzy comparisons of large scale land cover products: application to CORINE, GLC2000, MODIS and GlobCover in Europe. *ISPRS J. Photogramm. Remote Sens.* 74, 185–201.
- Peterson, B., Nelson, K.J., 2014. Mapping forest height in Alaska using GLAS, Landsat composites, and airborne LiDAR. *Remote Sens.* 6, 12409–12426.
- Petitjean, F., Inglada, J., Gancarski, P., 2012. Satellite image time series analysis under time warping. *IEEE Trans. Geosci. Remote Sens.* 50, 3081–3095.
- Pflugmacher, D., Krankina, O.N., Cohen, W.B., Friedl, M.A., Sulla-Menashe, D., Kennedy, R.E., Nelson, P., Loboda, T.V., Kuemmerle, T., Dyukarev, E., Elsakov, V., Kharuk, V.I., 2011. Comparison and assessment of coarse resolution land cover maps for Northern Eurasia. *Remote Sens. Environ.* 115, 3539–3553.
- Pflugmacher, D., Cohen, W.B., Kennedy, R.E., Yang, Z., 2014. Using Landsat-derived disturbance and recovery history and lidar to map forest biomass dynamics. *Remote Sens. Environ.* 151, 124–137.
- Potapov, P., Turubanova, S., Hansen, M.C., 2011. Regional-scale boreal forest cover and change mapping using Landsat data composites for European Russia. *Remote Sens. Environ.* 115, 548–561.
- Pouliot, D., Latifovic, R., 2013. Accuracy assessment of annual land cover time series derived from change-based updating. In: 7th International Workshop on the Analysis of Multitemporal Remote Sensing Image, Banff, Canada. 25–27 June 2013.
- Pouliot, D., Latifovic, R., Zabcic, N., Guindon, L., Olthof, I., 2014. Development and assessment of a 250 m spatial resolution MODIS annual land cover time series (2000–2011) for the forest region of Canada derived from change-based updating. *Remote Sens. Environ.* 140, 731–743.

- Radoux, J., Lamarche, C., Van Bogaert, E., Bontemps, S., Brockmann, C., Defourny, P., 2014. Automated training sample extraction for global land cover mapping. *Remote Sens.* 6, 3965–3987.
- Ramankutty, N., Foley, J.A., 1999. Estimating historical changes in global land cover: croplands from 1700 to 1992. *Global Biogeochem. Cycles* 13 (4), 997–1027.
- Ramoni, M., Sebastiani, P., Cohen, P.R., 2000. Multivariate clustering by dynamics. In: *Proceedings of the Seventeenth National Conference on Artificial Intelligence*, pp. 633–638.
- Reed, B.C., Brown, J.F., VanderZee, D., Loveland, T.R., Merchant, J.W., Ohlend, D.O., 1994. Measuring phenological variability from satellite imagery. *J. Veg. Sci.* 5, 703–714.
- Rodríguez Galiano, V.F., Chica Olmo, M., Abarca Hernández, F., Atkinson, P.M., Jeganathan, C., 2012. Random forest classification of mediterranean land cover using multi-seasonal imagery and multi-seasonal texture. *Remote Sens. Environ.* 121, 93–107.
- Rogan, J., Chen, D.M., 2004. Remote Sensing technology for mapping and monitoring land cover and land-use change. *Prog. Plan.* 61, 301–325.
- Rokach, L., 2010. Ensemble-based classifiers. *Artif. Intell. Rev.* 33, 1–39.
- Roy, D.P., Ju, J., Mbaw, Ch., Frost, Ph., Loveland, T., 2010a. Accessing free Landsat data via the internet: Africa's challenge. *Remote Sens. Lett.* 1 (2), 111–117.
- Roy, D.P., Ju, J., Kline, K., Scaramuzza, P.L., Kovalsky, V., Hansen, M., Loveland, T.R., Vermote, E., Zhang, Ch., 2010b. Web-enable Landsat data (WELD): Landsat ETM + composited mosaics of the conterminous United States. *Remote Sens. Environ.* 114, 35–49.
- Roy et al., 2014. Landsat-8: science and product vision for terrestrial global change research. *Remote Sens. Environ.* 145, 154–172.
- Running, S.W., Loveland, T.R., Pierce, L.L., 1994. A vegetation classification logic based on Remote Sensing for use in global scale biogeochemical models. *Ambio* 23, 77–81.
- Running, S.W., Baldocchi, D.D., Turner, D.P., Gower, S.T., Bakwin, P.S., Hibbard, K.A., 1999. A global terrestrial monitoring network integrating tower fluxes, flask sampling, ecosystem modeling and EOS satellite data. *Remote Sens. Environ.* 70, 108–127.
- Schneider, A., Friedl, M.A., Potere, D., 2010. Mapping global urban areas using MODIS 500-M data: new methods and datasets based on 'Urban Ecoregions'. *Remote Sens. Environ.* 114, 1733–1746.
- Schneider, A., 2012. Monitoring land cover change in urban and peri-urban areas using dense time stacks of Landsat satellite data and a data mining approach. *Remote Sens. Environ.* 124, 689–704.
- Schroeder, T.A., Healey, S.P., Moisen, G.G., Frescino, T.S., Cohen, W.B., Huang, Ch., Kennedy, R.E., Yang, Z., 2014. Improving estimates of forest disturbance by combining observations from Landsat time series with U.S. Forest Service Forest Inventory and Analysis data. *Remote Sens. Environ.* 154, 61–73.
- Schroeder, T.A., Wulder, M.A., Healey, S.P., Moisen, G.G., 2011. Mapping wildfire and clearcut harvest disturbances in boreal forests with Landsat time-series data. *Remote Sens. Environ.* 115, 1421–1433.
- Senf, C., Leitão, P.J., Pflugmacher, D., van der Linden, S., Hostert, P., 2015. Mapping land cover in complex Mediterranean landscapes using Landsat: improved classification accuracies from integrating multi-seasonal and synthetic imagery. *Remote Sens. Environ.* 156, 527–536.
- Sexton, J.O., Urban, D.L., Donohue, M.J., Song, C., 2013a. Long-term dynamics by multi-temporal classification across the Landsat-5 record. *Remote Sens. Environ.* 128, 246–258.
- Sexton, J.O., Song, X.P., Huang, Ch., Channan, S., Baker, M.E., Townshend, 2013b. Urban growth of the Washington, D.C.–Baltimore, MD metropolitan region from 1984 to 2010 by annual, Landsat-based estimates of impervious cover. *Remote Sens. Environ.* 129, 42–53.
- Shao, Y., Lunetta, R.S., 2012. Comparison of support vector machine, neural network, and CART algorithms for the land-cover classification using limited training data points. *ISPRS J. Photogramm. Remote Sens.* 70, 78–87.
- Shao, Y., Lunetta, R.S., Wheeler, B., James, J.S., Campbell, J.B., 2016. An evaluation of time-series smoothing algorithms for land-cover classifications using MODIS-NDVI multi-temporal data. *Remote Sens. Environ.* 174, 258–265.
- Shenthilnath, J., Omkar, S.N., Mani, V., Tejovanth, N., Diwakar, P.G., Shenoy, A., 2011. Hierarchical clustering algorithm for land cover mapping using satellite images. *IEEE J. Sel. Topics Appl. Earth Observations Remote Sens.* 5 (3), 762–768.
- Shimabukuro, Y.E., Beuchle, R., Grecchi, R.C., Achard, F., 2014. Assessment of forest degradation in Brazilian Amazon due to selective logging and fires using time series of fraction images derived from Landsat ETM+ images. *Remote Sens. Lett.* 5 (9), 773–782.
- Singh, 1989. Digital change detection techniques using remotely-sensed data. *Int. J. Remote Sens.* 10 (6), 989–1003.
- Song, C., Woodcock, C.E., Seto, K.C., Lenney, M.P., Macomber, S.A., 2001. Classification and change detection using Landsat TM data: when and how to correct atmospheric effects? *Remote Sens. Environ.* 75, 230–244.
- Sulla-Menashe, D., Friedl, M.A., Krankina, O.N., Baccini, A., Woodcock, C.E., Sibley, A., Sun, G., Kharuk, V., Elsakov, V., 2011. Hierarchical mapping of northern Eurasian land cover using MODIS data. *Remote Sens. Environ.* 115, 392–403.
- Tateishi, R., Uriyangqai, B., Al-Bilbisi, H., Ghar, M.A., Tsend-Ayush, J., Kobayashi, T., Kasimu, A., Thanh Hoan, N., Shalaby, A., Alsaaidh, B., Enkhzaya, T., Geggantana, Sato, H.P., 2011. Production of global land cover, GLCNMO. *Int. J. Digital Earth* 4, 22–49.
- Tewkesbury, A.P., Comber, A.J., Tate, N.J., Lamb, A., Fisher, P.F., 2015. A critical synthesis of remotely sensed optical image change detection techniques. *Remote Sens. Environ.* 160, 1–14.
- Thompson, S.D., Nelson, T.A., White, J.C., Wulder, M.A., 2015. Large area mapping of tree species using composited Landsat imagery. *Can. J. Remote Sens.* 41 (3), 203–218.
- Tiping, M.E., 2001. Sparse Bayesian learning and the relevance vector machine. *J. Mach. Learn. Res.* 1, 211–244.
- Townsend, P.A., Helmers, D.P., Kingdon, C.C., McNeil, B.F., deBeurs, K.M., Eshleman, K.N., 2009. Changes in the extent of surface mining and reclamation in the Central Appalachians detected using a 1976–2006 Landsat time series. *Remote Sens. Environ.* 113, 62–72.
- Tsutsumida, N., Comber, A., 2015. Measures of spatio-temporal accuracy for time series land cover data. *Int. J. Appl. Earth Obs. Geoinf.* 41, 46–55.
- Tuia, D., Pasolli, E., Emery, W.J., 2011. Using active learning to adapt *Remote Sensing* image classifiers. *Remote Sens. Environ.* 115, 2232–2242.
- Tulbure, M.G., Broich, M., 2013. Spatiotemporal dynamic of surface water bodies using Landsat time-series data from 1999 to 2011. *ISPRS J. Photogramm. Remote Sens.* 79, 44–52.
- Verbesselt, J., Hyndman, R., Zeileis, A., Culvenor, D., 2010. Phenological change detection while accounting for abrupt and gradual trends in satellite image time series. *Remote Sens. Environ.* 114, 2970–2980.
- Verbesselt, J., Zeileis, A., Herold, M., 2012. Near real-time disturbance detection using satellite image time series. *Remote Sens. Environ.* 123, 98–108.
- Vogelmann, J.E., Howard, S.M., Yang, L., Larson, C.R., Wylie, B.K., Van Driel, J.N., 2001. Completion of the 1990's national land cover data set for the conterminous United States. *Photogramm. Eng. Remote Sens.* 67, 650–661.
- Wang, X., Ye, M., Duanmu, C.J., 2009. Classification of data from electronic nose using relevance vector machines. *Sens. Actuators B: Chem.* 140, 143–148.
- Wang, J., Zhao, Y., Li, C., Yu, C., Liu, D., Gong, P., 2015. Mapping global land cover in 2001 and 2010 with spatial-temporal consistency at 250 m resolution. *ISPRS J. Photogramm. Remote Sens.* 103, 38–47.
- Wehmann, A., Liu, D., 2015. A spatial-temporal contextual Markovian kernel method for multi-temporal land cover mapping. *ISPRS J. Photogramm. Remote Sens.* 107, 77–89.
- Weiss, D.J., Atkinson, P.M., Bhatt, S., Mappin, B., Hay, S.I., Gething, P.W., 2014. An effective approach for gap-filling continental scale remotely sensed time-series. *ISPRS J. Photogramm. Remote Sens.* 98, 106–118.
- White, J.C., Wulder, M.A., Hobart, G.W., Luther, J.E., Hermosilla, T., Griffiths, P., Coops, N.C., Hall, R.J., Hostert, P., Dyk, A., Guindon, L., 2014. Pixel-based image compositing for large-area dense time series applications and science. *Can. J. Remote Sens.* 40, 192–212.
- Williams, D.L., Goward, S., Arvidson, T., 2006. Landsat: yesterday, today, and tomorrow. *Photogramm. Eng. Remote Sens.* 72 (10), 1171–1178.
- Wismüller, A., Lange, O., Dersch, D.R., Leisinger, G.L., Hahn, K., Pütz, B., Auer, D., 2012. Cluster analysis of biomedical image time-series. *Int. J. Comput. Vision* 46 (2), 103–128.
- Woodcock, C.E., Allen, R., Anderson, M., Belward, A., Bindschadler, R., Cohen, W.B., Gao, F., Goward, S.N., Helder, D., Helmer, E., Nemani, R., Oreopoulos, L., Schott, J., Thenkabail, P.S., Vermote, E.F., Vogelmann, J., Wulder, M.A., Wynne, R., 2008. Free access to Landsat imagery. *Science* 320, 1011.
- Wulder, M.A., Franklin, S.E., White, J.C., Linke, J., Magnussen, S., 2006. An accuracy assessment framework for large-area land cover classification products derived from medium-resolution satellite data. *Int. J. Remote Sens.* 27 (4), 663–683.
- Wulder, M.A., White, J.C., Cranny, M., Hall, R.J., Luther, J.E., Beaudoin, A., Goodenough, D.G., Dechka, J.A., 2008a. Monitoring Canada's forests. Part 1: completion of the EOSD land cover project. *Can. J. Remote Sens.* 34 (6), 549–562.
- Wulder, M.A., White, J.C., Goward, S.M., Masek, J.G., Irons, J.R., Herold, M., Cohen, W. B., Loveland, T.R., Woodcock, C.E., 2008b. Landsat continuity: issues and opportunities for land cover monitoring. *Remote Sens. Environ.* 112, 955–969.
- Wulder, M.A., Masek, J.G., Cohen, W.B., Loveland, T.R., Woodcock, C.E., 2012. Opening the archive: how free data has enabled the science and monitoring promise of Landsat. *Remote Sens. Environ.* 122, 2–10.
- Wulder, M.A., Hilker, T., White, J.C., Coops, N.C., Masek, J.G., Pflugmacher, D., Crevier, Y., 2015. Virtual constellations for global terrestrial monitoring. *Remote Sens. Environ.* 170, 62–76.
- Wulder, M.A., White, J.C., Loveland, T.R., Woodcock, C.E., Belward, A.S., Cohen, W.B., Fosnight, G., Shaw, J., Masek, J.G., Roy, D.P., 2016. The global Landsat archive: status, consolidation, and direction. *Remote Sens. Environ.* <http://dx.doi.org/10.1016/j.rse.2015.11.032>.
- Xian, G., Homer, C., Fry, J., 2009. Updating the 2001 national land cover database land cover classification to 2006 using Landsat change detection methods. *Remote Sens. Environ.* 113, 1133–1147.
- Xue, Z., Du, P., Feng, L., 2014. Phenology driven land cover classification and trend analysis based on long-term *Remote Sensing* image series. *IEEE J. Sel. Topics Appl. Earth Observations Remote Sens.* 7 (4), 1142–1156.
- Yan, L., Roy, D.P., 2014. Automated crop field extraction from multi-temporal Web Enabled Landsat Data. *Remote Sens. Environ.* 144, 42–64.
- Yan, L., Roy, D.P., 2015. Improved time series land cover classification by missing-observation-adaptive nonlinear dimensionality reduction. *Remote Sens. Environ.* 158, 478–491.
- Yin, H., Pflugmacher, D., Kennedy, R.E., Sulla-Menashe, D., Hostert, P., 2014. Mapping annual land use and land cover changes using MODIS time series. *IEEE J. Sel. Topics Appl. Earth Observations Remote Sens.* 7 (8), 3421–3427.
- Yuan, F., Sawaya, K.E., Loeffelholz, B.C., Bauer, M.E., 2005. Land cover classification and change analysis of the Twin Cities (Minnesota) Metropolitan area by multitemporal Landsat remote sensing. *Remote Sens. Environ.* 98, 317–328.
- Zald, H.S.J., Wulder, M.A., White, J.C., Hilker, T., Hermosilla, T., Hobart, G.W., Coops, N.C., 2016. Integrating Landsat pixel composites and change metrics with lidar

- plots to predictively map forest structure and aboveground biomass in Saskatchewan, Canada. *Remote Sens. Environ.* 176, 188–201.
- Zhang, X., Liu, J., Du, Y., Lv, T., 2011. A novel clustering method on time-series data. *Expert Syst. Appl.* 38, 11891–11900.
- Zhu, X., Chen, J., Gao, F., Chen, X., Masek, J.G., 2010. An enhanced spatial and temporal adaptive reflectance fusion model for complex heterogeneous regions. *Remote Sens. Environ.* 114 (11), 2610–2623.
- Zhu, Z., Woodcock, C.E., 2012. Object-based cloud and cloud shadow detection in Landsat imagery. *Remote Sens. Environ.* 118, 83–94.
- Zhu, Z., Woodcock, C.E., 2014a. Automate cloud, cloud shadow, and snow detection in multitemporal Landsat data: an algorithm designed specifically for monitoring land cover change. *Remote Sens. Environ.* 152, 217–234.
- Zhu, Z., Woodcock, C.E., 2014b. Continuous change detection and classification of land cover using all available Landsat data. *Remote Sens. Environ.* 144, 152–171.
- Zhu, Z., Wang, Sh., Woodcock, C.E., 2015a. Improvement and expansion of the Fmask algorithm: cloud, cloud shadow, and snow detection for Landsats 4–7, 8, and Sentinel 2 images. *Remote Sens. Environ.* 159, 269–277.
- Zhu, Z., Woodcock, C.E., Holden, C., Yang, Z., 2015b. Generating synthetic Landsat images based on all available Landsat data: predicting Landsat surface reflectance at any given time. *Remote Sens. Environ.* 162, 67–83.