



NORTH-HOLLAND

## The Maximum Determinant of $\pm 1$ Matrices

M. G. Neubauer\* and A. J. Radcliffe†

*Department of Mathematics and Statistics  
University of Nebraska  
Lincoln, Nebraska 68588-0323*

Submitted by Richard A. Brualdi

---

### ABSTRACT

We give a new proof for the bound on the value of the determinant of a  $\pm 1$  matrix of dimension  $n \equiv 1 \pmod{4}$  first given by Barba. Adapting a construction of A. E. Brouwer, we give examples to show that the bound is sharp for infinitely many values of  $n$ . This in turn gives an infinite family of examples which attain the bound given by H. Ehlich and by M. Wojtas for the determinant of a  $\pm 1$  matrix of dimension  $n \equiv 2 \pmod{4}$ . For  $n \equiv 3 \pmod{4}$  we construct an infinite family of examples which attain slightly more than  $\frac{1}{3}$  of the bound given by Ehlich. © Elsevier Science Inc., 1997

---

### 1. INTRODUCTION AND MAIN RESULT

Let  $\mathcal{N}_n$  be the set of  $\pm 1$  matrices of dimension  $n$ . The question of the maximum value of the determinant of an element  $N \in \mathcal{N}_n$  is an old one which goes back to the beginnings of matrix theory. It is a simple conse-

---

\*Current address: Department of Mathematics, California State University Northridge, Northridge, CA 91330.

†Research partially supported by NSF 9401351.

*LINEAR ALGEBRA AND ITS APPLICATIONS* 257:289–306 (1997)

© Elsevier Science Inc., 1997  
655 Avenue of the Americas, New York, NY 10010

0024-3795/97/\$17.00  
PII S0024-3795(96)00147-4

quence of Hadamard’s inequality [16] that for all  $M \in \mathcal{H}_n$

$$\det M \leq \left( \prod_{i=1}^n \sum_{j=1}^n m_{ij}^2 \right)^{1/2} \leq n^{n/2}. \tag{1}$$

There is a large body of work addressing the question of when (1) is sharp.

**DEFINITION 1.1.** Matrices in  $\mathcal{H}_n$  for which equality holds in (1) are called *Hadamard matrices*.

For an  $n \times n$  Hadamard matrix to exist it is necessary that  $n$  be either 1, 2, or a multiple of 4, and it is conjectured that this condition is also sufficient. According to [2] the smallest value for which the existence of a Hadamard matrix is in question is  $n = 4 \times 107 = 428$ .

Hadamard matrices do exist for many infinite families of values of  $n$ ; e.g., Sylvester [26] proved that Sylvester’s matrix,

$$S = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix},$$

tensored with itself  $t$  times, gives a Hadamard matrix of dimension  $2^t$ . The Paley construction (see e.g. [2, pp. 271–275] or [20]) for an odd prime power  $p$  gives a Hadamard matrix of size  $p + 1$  if  $p \equiv 3 \pmod{4}$  and size  $2p + 2$  if  $p \equiv 1 \pmod{4}$ , thus providing another infinite family of Hadamard matrices.

For further discussion of Hadamard matrices the reader might consult [4] or one of the surveys [1] or [25].

*1.1. (0, 1) Matrices*

There is a strong connection between  $\mathcal{H}_n$  and  $\mathcal{Z}_{n-1}$ , the set of (0, 1) matrices of dimension  $n - 1$ . Specifically, there exists an injection  $\phi$  from  $\mathcal{Z}_{n-1}$  into  $\mathcal{H}_n$  which preserves the relative sizes of determinants. The function  $\phi$  can be described as follows:

Given  $M$  in  $\mathcal{Z}_{n-1}$ , let  $M'$  be the  $n \times n$  block matrix

$$M' = \begin{pmatrix} 1 & 1 \\ 0 & -2M \end{pmatrix}.$$

$\phi(M)$  is obtained by adding the first row of  $M'$  to each other row. The resulting matrix is clearly a  $\pm 1$  matrix, and the passage from  $M'$  to  $\phi(M)$

doesn't change the value of the determinant. Thus it is clear that

$$\det \phi(M) = (-2)^{n-1} \det M.$$

Let  $J_n$  denote the  $n \times n$  matrix all of whose entries are 1. The map  $\phi$  can also be described as follows:

$$\phi(M) = J_n - \begin{pmatrix} 0 & 0 \\ 0 & 2M \end{pmatrix}.$$

The image of  $\phi$  is the set  $\mathcal{X}'_n$  of matrices  $N = (n_{ij})$  in  $\mathcal{X}_n$  such that  $n_{1j} = 1$  for all  $1 \leq j \leq n$  and  $n_{i1} = 1$  for all  $1 \leq i \leq n$ . The set  $\mathcal{X}'_n$  is a sufficiently large subset of  $\mathcal{X}_n$  for our purposes, since for every matrix  $M \in \mathcal{X}_n$  we can find suitable diagonal matrices  $P = (p_{ij})$  and  $Q = (q_{ij})$  with  $p_{ii}, q_{ii} \in \{\pm 1\}$  such that  $PMQ \in \mathcal{X}'_n$ . Since  $\det P = \pm 1 = \det Q$ , we have  $|\det M| = |\det PNQ|$ .

If we write  $m(\mathcal{E})$  for  $\max\{\det M \mid M \in \mathcal{E}\}$ , then the above remarks show that

$$m(\mathcal{X}_n) = m(\mathcal{X}'_n) = 2^{n-1}m(\mathcal{Z}_{n-1}).$$

In particular,  $m(\mathcal{Z}_{n-1}) \leq n^{n/2}/2^{n-1}$ , and the  $(0, 1)$  matrices which attain this bound are the preimages under  $\phi$  of Hadamard matrices. From another point of view they can be regarded as the incidence matrices of Hadamard designs; a class of symmetric designs with parameters  $(4m - 1, 2m - 1, m - 1)$ . (See [2], [4], or [1] for more details.)

1.2. *What Is Known When  $n \not\equiv 0 \pmod{4}$ ?*

The first reference to the case  $n \not\equiv 0 \pmod{4}$  seems to be Colucci [10] in 1926. The following lemma, concerning the cases  $n \equiv 1 \pmod{2}$ , was discussed by Barba [3] and proved, independently of Barba, by Ehlich [13, Satz 4.1].

LEMMA A (Barba, Ehlich). *Suppose that  $n \equiv 1 \pmod{2}$ . For all  $N \in \mathcal{X}_n$ ,*

$$\det N \leq (2n - 1)^{1/2}(n - 1)^{(n-1)/2}. \tag{2}$$

*That is,  $m(\mathcal{X}_n) \leq (2n - 1)^{1/2}(n - 1)^{(n-1)/2}$ . Equivalently, for all  $M \in \mathcal{Z}_{n-1}$ ,*

$$2^{n-1} \det M \leq (2n - 1)^{1/2}(n - 1)^{(n-1)/2}. \tag{3}$$

In 1937 T. Popoviciu [22], apparently unaware of [3], gave a weaker bound. Curiously enough, J. Brenner [5] claims Popoviciu's bound to be sharper than (2). Neither [3] nor [22] addresses the case  $n \equiv 2 \pmod{4}$  separately; rather than use Hadamard's bound for  $n \equiv 0 \pmod{4}$  in this case as well, though it was known that Hadamard's bound can only be attained for  $n = 1, 2$  or  $n \equiv 0 \pmod{4}$ .

It seems that H. Ehlich and, independently, M. Wojtas were the first to address the case  $n \equiv 2 \pmod{4}$  [13, 28]. They also seem to have been the first to address the question of the structure of matrices of maximal determinants.

**PROPOSITION A** (Ehlich, Wojtas). *Suppose that  $n \equiv 1 \pmod{4}$ . For all  $N \in \mathcal{H}_n$  the inequality (2) holds, and in order for equality to hold in (2) it is necessary that  $2n - 1$  be a square and that there exists an  $N \in \mathcal{H}_n$  with  $NN^T = (n - 1)I_n + J_n$ .*

**PROPOSITION B** (Ehlich, Wojtas). *Assume  $n \equiv 2 \pmod{4}$ . For all  $N \in \mathcal{H}_n$*

$$\det N \leq (2n - 2)(n - 2)^{\frac{1}{2}n - 1}. \tag{4}$$

*Moreover, equality in (4) holds if and only if there exists  $N \in \mathcal{H}_n$  such that*

$$NN^T = N^T N = \begin{pmatrix} L & 0 \\ 0 & L \end{pmatrix},$$

*where  $L = (n - 2)I + 2J$  is an  $(n/2) \times (n/2)$  matrix. A further necessary condition for equality to hold is that  $2n - 2$  is the sum of two squares.*

H. S. E. Cohn gave an independent proof of Lemma B above and provided further information on the structure of maximal examples [12, Theorem 2].

In a sequel to [13] Ehlich investigated the case  $n \equiv 3 \pmod{4}$  [14].

**PROPOSITION C** [Ehlich]. *Assume  $n \equiv 3 \pmod{4}$  and  $n \geq 63$ . For all  $N \in \mathcal{H}_n$ ,*

$$\det N \leq \sqrt{\frac{4 \times 11^6}{7^7} (n - 3)^{n-7} n^7}. \tag{5}$$

Moreover, for equality to hold it is necessary that  $n = 7m$  and that there exists  $N \in \mathcal{X}_n$  with

$$NN^T = I_7 \otimes [(n - 3)I_m + 4J_m] - J_n.$$

The corresponding bounds for all values  $n \equiv 3 \pmod{4}$ ,  $n < 63$ , are also given in [14], as are structures of  $NN^T$  for normalized maximal examples  $N$ . The theme for values  $n < 63$  is the same as for the above example: A  $\pm 1$  matrix  $N$  has maximal determinant if  $NN^T$  has block structure with the blocks along the diagonal of the form  $(n - 3)I + 3J$  and the off-diagonal blocks equal to  $-J$ .

Though it is well known that the Hadamard bound in the case  $n \equiv 0 \pmod{4}$  is attained infinitely often and has to be considered sharp in this sense, it was not known if the bounds given in Proposition A, Proposition B, and Proposition C are sharp in this sense. The object of this paper is to show that this is indeed the case for  $n \equiv 1 \pmod{4}$  and  $n \equiv 2 \pmod{4}$ . For  $n \equiv 3 \pmod{4}$  we provide an infinitely family of examples which asymptotically have determinants slightly larger than  $\frac{1}{3}$  of the bound of Proposition C. The case  $n \equiv 1 \pmod{4}$  is crucial to everything, and so we provide our own version and proof of Proposition A to keep the exposition self-contained. We then extend the known results to show that the bound of Proposition A is sharp, i.e., it is attained infinitely often.

**THEOREM A.** *Suppose that  $n \equiv 1 \pmod{4}$ . For all  $N \in \mathcal{X}_n$  the inequality (2) holds. In order for equality to hold in (2) it is necessary that  $2n - 1$  be a square and that  $NN^T = (n - 1)I_n + R$  where  $\text{rank } R = 1$  and  $|r_{ij}| = 1$  for all  $i, j$ .*

*Furthermore, if  $n = 2(q^2 + q) + 1$  for some odd prime power  $q$ , then there exist matrices  $N \in \mathcal{X}_n$  for which equality holds in (2), i.e.,*

$$m(\mathcal{X}_n) = (2n - 1)^{1/2}(n - 1)^{(n-1)/2} \tag{6}$$

*for all  $n = 2(q^2 + q) + 1$ ,  $q$  an odd prime power. In particular the bound in (2) is sharp for infinitely many values of  $n$ .*

The proof of Theorem A naturally divides into two parts. In Section 2 we prove the inequality (2). In Section 3 we build an infinite family of examples to prove that

$$\mathcal{B}_n = \{N \in \mathcal{X}_n \mid n \equiv 1 \pmod{4} \text{ and } N \text{ attains equality in (2)}\}$$

is nonempty whenever  $n = 2q^2 + 2q + 1$  and  $q$  is an odd prime power.

I. Kaplansky [18] claims that the bound in (2) can be attained only if there exists a design with parameters  $(2q^2 + 2q + 1, q^2, (q^2 - q)/2)$ .

The particular values  $n = 17$  and  $n = 21$  for which the bound cannot be attained are discussed in [21] and [8] respectively.

The examples that we construct can be used to give an infinite family of examples which attain the bound of Proposition B.

**THEOREM B.** *If  $n = 4(q^2 + q) + 2$  for some odd prime power  $q$ , then there exists a matrix  $N \in \mathcal{X}_n$  for which equality holds in the inequality (4) of Theorem A. In particular, equality holds in (4) for infinitely many values of  $n$ .*

This was already observed by Whiteman [27], who gave the same proof as given below. He presented his results in the context of  $D$ -optimal designs. For a definition of  $D$ -optimal designs and other optimality criteria see [24]. In [19] another infinite family of matrices is given whose determinants attain the bound in (4).

*Proof of Theorem B.* Suppose  $n \equiv 1 \pmod{4}$  and that  $N \in \mathcal{X}_n$ . After a suitable change of basis we may assume, by Theorem A, that  $NN^T = (n - 1)I_n + J_n$ . Now define

$$\tilde{N} = S \otimes N = \begin{pmatrix} N & N \\ N & -N \end{pmatrix}, \tag{7}$$

where  $S$  is Sylvester's matrix defined above. It is easy to show that

$$\tilde{N}\tilde{N}^T = \begin{pmatrix} (2n - 2)I_n + 2J_n & 0 \\ 0 & (2n - 2)I_n + 2J_n \end{pmatrix}, \tag{8}$$

which implies that  $\det \tilde{N} = (2n - 2)(n - 2)^{\frac{1}{2}n - 1}$  ([13], [28], or [12]). ■

Unfortunately, we were unable to construct an infinite family of examples which attain the bound of Proposition C. In fact, we do not even know of a single example for which the bound in Proposition C is attained. Nevertheless, our efforts yielded an infinite family of examples whose determinants asymptotically take on values slightly larger than  $\frac{1}{3}$  of the bound of Proposition C. This shows that the bound of Proposition C is of the correct order. This is a significant improvement over the lower bound given in [9].

THEOREM C. *There exists an infinite family of examples  $N_n \in \mathcal{X}_n$  such that*

$$\lim_{n \rightarrow \infty} \frac{\det N_n}{h_n} = \frac{1}{2} \left(\frac{7}{11}\right)^3 \sqrt{7} \approx 0.34,$$

where  $h_n^2 = (4 \times 11^6 / 7^7)(n - 3)^{n-7} n^7$ .

A proof of Theorem C is given in Section 4.

At this point it remains unresolved if the bound in (5) can ever be attained. We have no doubt that for a particular value of  $n \equiv 3 \pmod{4}$  an efficient computer search is likely to produce examples whose determinants have larger values than the ones given by us in Section 4. The real challenge, however, is to find an infinite family of examples whose determinants take on the bound in (5) or to show that the bound in (5) can be improved. It is conceivable that it is not sharp. The methods used by Ehlich in [14] study the maximum of the values of the determinant on a subset of positive definite matrices. This subset is defined by certain necessary condition its elements have to satisfy if they are to be of the form  $NN^T$  for some  $N \in \mathcal{X}_n$ . It is certainly possible that this subset contains many positive definite matrices which are not of the form  $NN^T$  for some  $N \in \mathcal{X}_n$ . Hence the bound derived from this subset might be larger than the actual bound.

Having said that, we mention that B. Solomon [23] has provided us with examples for  $n = 11$ ,  $n = 15$ , and  $n = 19$ , believed on the basis of computer experiments to be maximal, which seem to show that maximal examples tend to exhibit the behavior forecast in [14].

Some interesting comments on determinants of  $\pm 1$  matrices in the context of  $D$ -optimal designs can be found in [15].

## 2. THE PROOF OF THE INEQUALITY OF THEOREM A

The next lemma is proved in [10]. For completeness we include a (slightly shorter) proof.

LEMMA 2.1. *Suppose that  $T \in M_n(\mathbf{R})$  is a symmetric, diagonalizable matrix with the following properties: it has 0 on the diagonal,  $I_n + T$  is positive definite, and the root mean square of its off-diagonal elements is  $c$ . Then*

$$\det(I_n + T) \leq [1 + c(n - 1)](1 - c)^{n-1}. \tag{9}$$

*Proof.* Let  $z_1, z_2, \dots, z_n$  be the eigenvalues of  $T$ . We know that  $\sum_{i=1}^n z_i = 0$  and

$$\sum_{i=1}^n z_i^2 = \sum_{i,j} t_{ij}^2 = n(n-1)c^2. \tag{10}$$

Split the eigenvalues into two classes according to their sign; let  $x_1, x_2, \dots, x_k$  be the nonnegative ones,  $y_1, y_2, \dots, y_{n-k}$  the negative ones. Set  $\bar{x} = (\sum_1^k x_i)/k$ , and  $\bar{y} = (\sum_1^{n-k} y_i)/(n-k)$ . Now

$$\begin{aligned} \det(I_n + T) &= \prod_{i=1}^k (1 + x_i) \cdot \prod_{i=1}^{n-k} (1 + y_i) \\ &\leq (1 + \bar{x})^k (1 + \bar{y})^{n-k}, \end{aligned} \tag{11}$$

where the inequality is just the arithmetic-mean-geometric-mean inequality. Thus if we define  $g : (k, x, y) \mapsto k \ln(1 + x) + (n - k) \ln(1 + y)$ , we have that  $\ln \det(I_n + T) \leq g(k, \bar{x}, \bar{y})$ . We want a bound on  $g(k, \bar{x}, \bar{y})$  given that  $k\bar{x} + (n - k)\bar{y} = 0$  and  $k\bar{x}^2 + (n - k)\bar{y}^2 = n(n - 1)c^2$ . For convenience we denote this last quantity by  $S$ . Solving for  $\bar{x}$  and  $\bar{y}$  in terms of  $k$ , we get

$$\bar{x} = \sqrt{S} \frac{\sqrt{n - k}}{\sqrt{n} \sqrt{k}}, \quad \bar{y} = -\sqrt{S} \frac{\sqrt{k}}{\sqrt{n} \sqrt{n - k}}.$$

Therefore the quantity we wish to estimate is

$$g(k) = k \ln \left( 1 + \sqrt{S} \frac{\sqrt{n - k}}{\sqrt{n} \sqrt{k}} \right) + (n - k) \ln \left( 1 - \sqrt{S} \frac{\sqrt{k}}{\sqrt{n} \sqrt{n - k}} \right).$$

The function  $g(k)$  is of course defined for all real  $k \in (0, n)$ , but condition (10) ensures that  $k \in [1, n - 1]$ . We will show that  $g(k)$  is decreasing on  $(0, n)$ . Indeed, differentiating we get (writing  $\bar{x}$  and  $\bar{y}$  for the above functions of  $k$ ),

$$g'(k) = \ln(1 + \bar{x}) - \ln(1 + \bar{y}) - \frac{1}{2}(\bar{x} - \bar{y}) \left( \frac{1}{1 + \bar{x}} + \frac{1}{1 + \bar{y}} \right). \tag{12}$$

To show that  $g'(k) < 0$  we need only prove that

$$\frac{1}{\bar{x} - \bar{y}} \int_{\bar{y}}^{\bar{x}} \frac{dt}{1 + t} < \frac{1}{2} \left( \frac{1}{1 + \bar{x}} + \frac{1}{1 + \bar{y}} \right).$$



This is an immediate consequence of the strict convexity of the function  $t \mapsto 1/(1+t)$ . Therefore, for all  $k \in [1, n-1]$  we have  $g(k) \leq g(1)$ , where

$$\begin{aligned}
 g(1) &= \ln\left(1 + \sqrt{S} \sqrt{1 + \frac{1}{n}}\right) + (n-1) \ln\left(1 + \sqrt{S} \sqrt{\frac{1}{n(n-1)}}\right) \\
 &= \ln[1 + c(n-1)] + (n-1) \ln(1 - c).
 \end{aligned}
 \tag{13}$$

The result follows immediately. ■

Next we prove a result which is analogous to Lemma 1 of [12]. It is slightly more general than is needed here but the proof makes the result natural. Define the function

$$f : (n, h) \mapsto (n + h - 1)(h - 1)^{n-1}.
 \tag{14}$$

LEMMA 2.2. *Suppose that  $M$  is a positive definite symmetric matrix of the form  $(h - 1)I_n + R \in M_n(\mathbf{R})$ , where  $h > 1$  and  $R = (r_{ij})$  has  $|r_{ij}| \geq 1$  and  $r_{ii} = 1$ . Then:*

1.  $|\det M| \leq f(n, h)$ .
2. *Equality holds if and only if  $|r_{ij}| = 1 \forall 1 \leq i, j \leq n$  and  $\text{rank } R = 1$ .*

*Proof.* First notice that if  $M = (h - 1)I_n + R$  with  $R$  as in part 2, then

$$\begin{aligned}
 \det M &= (h - 1)^n \left(1 + \frac{1}{h - 1} \text{trace } R\right) \\
 &= (h - 1)^n \left(1 + \frac{n}{h - 1}\right) \\
 &= (h - 1)^{n-1} (h - 1 + n) = f(n, h).
 \end{aligned}$$

We prove the theorem by induction on the dimension  $n$ . The case  $n = 2$  is a straightforward computation. Assume then that  $n > 2$  and that the result holds for smaller values of  $n$ .

Now  $\det M = \det[(h - 1)I_n + R] = \det[hI_n + (R - I_n)] = h^n \det[I_n + h^{-1}(R - I_n)]$ , and the matrix  $T = h^{-1}(R - I_n)$  satisfies the conditions of

Lemma 2.1 with  $c^2 = [1/n(n - 1)]\sum_{i \neq j} r_{ij}^2/h^2 \geq 1/h^2$ . Thus

$$\begin{aligned} \det M &= h^n \det(I_n + T) \\ &\leq h^n \left(1 + \frac{n - 1}{h}\right) \left(1 - \frac{1}{h}\right)^{n-1} \\ &= (n + h - 1)(h - 1)^{n-1} \\ &= f(n, h). \end{aligned} \tag{15}$$

It is clear that if  $|m_{ij}| > 1$  for some pair  $(i, j)$ ,  $i \neq j$ , then  $c > 1/h$  and the inequality becomes strict. Thus for equality to occur it is necessary that  $|m_{ij}| = 1$  for all  $1 \leq i, j \leq n$ .

After a suitable base change we may assume that the entries of the last row and the last column of  $M$  are all equal to 1, i.e.,  $m_{in} = m_{nj} = 1$  for all  $1 \leq i, j \leq n$ .

Now subtract  $1/h$  times the last row from the first  $n - 1$  rows and then subtract  $1/h$  times the last column from the first  $n - 1$  columns. The resulting matrix is

$$M' = \begin{pmatrix} M_1 & 0 \\ 0 & h \end{pmatrix},$$

where  $M_1 = (m'_{ij})$ ,  $1 \leq i, j \leq n - 1$ , is a symmetric, positive definite matrix [of size  $(n - 1) \times (n - 1)$ ] such that

$$\left(1 - \frac{1}{h}\right)^{-1} m'_{ij} = \begin{cases} h + 1 & \text{if } i = j, \\ 1 & \text{if } m_{ij} = 1, \\ -\frac{h + 1}{h - 1} & \text{if } m_{ij} = -1. \end{cases} \tag{16}$$

Note that  $(h + 1)/(h - 1) > 1$ . Since  $\det M = f(n, h)$  and  $\det M = h \det M_1$ , we see that

$$\det M_1 = \frac{f(n, h)}{h} = \frac{(h + n - 1)(h - 1)^{n-1}}{h}. \tag{17}$$

On the other hand, applying the induction hypothesis to the matrix  $\tilde{M}_1 = (1 - 1/h)^{-1}M_1$ , we get

$$\det M_1 \leq \left(1 - \frac{1}{h}\right)^{n-1} (h + n - 1)h^{n-2} \tag{18}$$

$$= \frac{(h + n - 1)(h - 1)^{n-1}}{h}. \tag{19}$$

Thus  $\det \tilde{M}_1 = f(n - 1, h + 1)$ . By induction this implies that  $\tilde{M}_1 = hI_{n-1} + \tilde{R}$  where  $\text{rank } \tilde{R} = 1$  and  $|\tilde{r}_{ij}| = 1$ . Hence, by (16), we conclude that  $m_{ij} = 1$  for all  $1 \leq i, j \leq n$ , i.e.,  $P = (h - 1)I_n + R$  with  $\text{rank } R = 1$  and  $|r_{ij}| = 1$ . ■

The statements of Theorem A pertaining to the inequality follow from Lemma 2.2. We summarize this in the next result.

**COROLLARY 2.3.** *Let  $N \in \mathcal{X}_n$ . Then  $|\det N| \leq \sqrt{f(n, n)}$  with equality if and only if  $NN^T = (n - 1)I_n + R$  with  $|r_{ij}| = 1$  and  $\text{rank } R = 1$ .*

*Proof.* Instead of working with the  $\pm 1$  matrix  $N$ , we consider the matrix  $M = NN^T$ . If  $N$  is singular, there is nothing to prove. If  $N$  is invertible, then, because  $n$  is odd, it follows that

$$M = (n - 1)I_n + R$$

is symmetric, positive definite, with  $|r_{ij}| \geq 1$ . Since  $\det M = (\det N)^2$ , the corollary now follows from Lemma 2.2. ■

### 3. AN INFINITE FAMILY OF MAXIMAL EXAMPLES

In this section we construct an infinite family of  $\pm 1$  matrices which yield equality in (2). Using the function  $\phi$ , we can construct from these an infinite family of  $(0, 1)$  matrices which yield equality in (3). The construction is very close to one by A. E. Brouwer [6]; we include the details for completeness.

Recall that, by Proposition A, we are trying to construct, for some  $n \equiv 1 \pmod{4}$ , a matrix  $N \in \mathcal{X}_n$  such that  $N^TN = (n - 1)I_n + J_n$ . We will show that this is possible whenever  $n = 2q^2 + 2q + 1$  and  $q$  is an odd prime power.

Down to work. Let  $q$  be an odd prime power. By the Paley construction [20] there exists a Hadamard matrix  $H_{2q+2}$  of size  $2q + 2$ , which we may assume is normalized; i.e., its first row and column are all 1's. We use  $H_{2q+2}$  to define matrices  $E$  and  $A$ , of dimension  $q \times 2q$  and  $(q + 1) \times 2q$  respectively: after a suitable rearrangement of rows we may assume that

$$H_{2q+2} = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ 1 & 1 & & & \\ \vdots & \vdots & & E & \\ 1 & 1 & & & \\ 1 & -1 & & & \\ \vdots & \vdots & & A & \\ 1 & -1 & & & \end{pmatrix}.$$

The matrices  $E$  and  $A$  are  $\pm 1$  matrices satisfying the following (where  $J_{m,l}$  is an  $m \times l$  matrix of 1's):

$$J_{m,q}E = -J_{m,2q}, \tag{20}$$

$$EJ_{2q,m} = -2J_{q,m}, \tag{21}$$

$$J_{m,q+1}A = 0, \tag{22}$$

$$AJ_{2q,m} = 0, \tag{23}$$

$$E^TE + A^TA = (2q + 2)I_{2q} - J_{2q}. \tag{24}$$

Let  $C$  be a  $(q + 1) \times (q + 1)$  conference matrix. That is,  $C$  is a  $(0, \pm 1)$  matrix with  $C_{ij} = 0 \iff i = j$ , and

$$C^TC = CC^T = qI. \tag{25}$$

Such a matrix can be constructed, for instance, by enumerating the elements  $\mathbf{F}_q = \{x_1, x_2, \dots, x_q\}$ , the field with  $q$  elements, then adding a row and column of 1's to the matrix  $(\chi(x_i - x_j))_{i,j=1}^q$ , and finally changing the first entry on the diagonal back to 0. Here  $\chi : \mathbf{F}_q \rightarrow \{0, \pm 1\}$  is the quadratic character.

Let  $L$  be the  $q^2 \times q(q + 1)$  incidence matrix of points and lines in affine  $\mathbf{F}_q^2$ . Then lines in  $\mathbf{F}_q^2$  can be split into  $q + 1$  parallel classes according to their slope, and two lines are disjoint if they are parallel, and meet in one point if they are not. Thus if we group the columns of  $L$  by their slope we find that  $L$



properties of  $E$ ,  $A$ ,  $C$ , and  $L$  established above in (20) through (30):

$$\begin{aligned}
 M_q^T M_q &= (I_{q+1} \otimes E^T) L^T L (I_{q+1} \otimes E) \\
 &\quad + (C^T \otimes A^T - I_{q+1} \otimes J_{2q, q+1}) (C \otimes A - I_{q+1} \otimes J_{q+1, 2q}) \\
 &= (I_{q+1} \otimes E^T) [(J_{q+1} - I_{q+1}) \otimes J_q + I_{q+1} \otimes qI_q] (I_{q+1} \otimes E) \\
 &\quad + C^T C \otimes A^T A - C \otimes (J_{2q, q+1} A) - C^T \otimes (A^T J_{q+1, 2q}) \\
 &\quad + I_{q+1} \otimes (q+1) J_{2q} \\
 &= (J_{q+1} - I_{q+1}) \otimes (E^T J_q E) + I_{q+1} \otimes qE^T E + qI_{q+1} \otimes (A^T A) \\
 &\quad + I_{q+1} \otimes (q+1) J_{2q} \\
 &= (J_{q+1} - I_{q+1}) \otimes (-E^T J_{q, 2q}) + qI_{q+1} \otimes (E^T E + A^T A) \\
 &\quad + (q+1) I_q \otimes J_{2q} \\
 &= (J_{q+1} - I_{q+1}) \otimes J_{2q} + qI_{q+1} \otimes [(2q+2)I_{2q} - J_{2q}] \\
 &\quad + (q+1) I_q \otimes J_{2q} \\
 &= qI_{q+1} \otimes [(2q+2)I_{2q} - J_{2q}] + (qI_{q+1} + J_{q+1}) \otimes J_{2q} \\
 &= (2q^2 + 2q) I_{2q^2+2q} + J_{2q^2+2q}.
 \end{aligned}$$

From this it follows immediately that  $N_q^T N_q = (2q^2 + 2q) I_{2q^2+2q+1} + J_{2q^2+2q+1}$  and thence that  $N_q \in \mathcal{B}_{2q^2+2q+1}$ .

#### 4. THE PROOF OF THEOREM C

Let  $N \in \mathcal{B}_k$ ,  $k = 2q^2 + 2q + 1$ , be one of the examples constructed Section 3, i.e., we may assume  $NN^T = (k-1)I_k + J_k$ . Recall that each row and each column of such an  $N$  has  $q^2$  entries 1 and  $(q+1)^2$  entries  $-1$ .

Let

$$B = \begin{pmatrix} 1 & 1 & \cdots & 1 & 1 & \cdots & 1 \\ 1 & & & & & & \\ \vdots & & N & & & & N \\ 1 & & & & & & \\ -1 & & & & & & \\ \vdots & & N & & & & -N \\ -1 & & & & & & \end{pmatrix}. \tag{31}$$

Then  $B \in \mathcal{K}_n$ , where  $n = 2k + 1 = 4q^2 + 4q + 3$ . We note that  $n \equiv 3 \pmod{4}$ . It is easy to see that

$$BB^T = \begin{pmatrix} n & -4q - 1 & \cdots & -4q - 1 & -1 & \cdots & -1 \\ -4q - 1 & & & & & & \\ \vdots & & (n - 3)I_k + 3J_k & & & & -J_k \\ -4q - 1 & & & & & & \\ -1 & & & & & & \\ \vdots & & -J_k & & & & (n - 3)I_k + 3J_k \\ -1 & & & & & & \end{pmatrix}. \tag{32}$$

Let

$$h_n = \sqrt{\frac{4 \times 11^6}{7^7} (n - 3)^{n-7} n^7} \tag{33}$$

be the bound in (5) of Proposition C.

LEMMA 4.1. *Assume that  $B$  is as in (31). Then:*

1.

$$\begin{aligned} d_n &= \det BB^T \\ &= [n - y - kt(4q + y + 1)][n + k(3 - y) - 2] \\ &\quad \times (n + 3k - 2)(n - 3)^{n-3}, \end{aligned}$$

where

$$y = \frac{k}{5(k-1)}$$

and

$$t = \frac{4q + y + 1}{n + k(3 - y) - 3}.$$

2.

$$\lim_{n \rightarrow \infty} \frac{\det B}{\sqrt{n^n}} = \sqrt{e^{-3}}.$$

3.

$$\lim_{n \rightarrow \infty} \sqrt{\frac{d_n}{h_n}} = \frac{1}{2} \left( \frac{7}{11} \right)^3 \sqrt{7} \approx 0.34.$$

*Proof.* 1: The formula for  $d_n$  is obtained by first zeroing out the  $-J$  matrix in the upper right-hand corner of  $BB^T$  and secondly zeroing out the first row of the matrix that remains in the upper left-hand corner.

2: The critical term in the formula for  $d_n$  occurs in the first factor. We see that

$$\lim_{q \rightarrow \infty} y = \frac{1}{5}$$

and

$$\lim_{q \rightarrow \infty} qt = \frac{5}{12}.$$

The analysis of the remaining terms is straightforward.

3: This follows directly from part 2 of this lemma and (1.2) of [14]. ■

Theorem C is now a consequence of Lemma 4.1.

*We would like to thank Bruce Solomon for providing us with examples of large-determinant  $(0, 1)$  matrices for many values of  $n \leq 20$ .*



After having read a preprint of this paper, I. Kaplansky pointed out more recent work on the problem, and he provided us with the references [7], [8], [15], [17], [19], [21], and [27]. This improved the paper considerably.

## REFERENCES

- 1 S. S. Aghaian, *Hadamard Matrices and Their Application*, Lecture Notes in Math. 1168, Springer-Verlag, Berlin, 1985.
- 2 E. F. Assmus, Jr., and J. D. Key, *Designs and Their Codes*, Cambridge Tracts in Math. 103, Cambridge U.P., Cambridge, 1992.
- 3 G. Barba, Intorno al teorema di Hadamard sui determinanti a valore massimo, *Giorn. Mat. Battaglia* **71**:70–86 (1933).
- 4 B. Bollobás, *Combinatorics, Set Systems, Hypergraphs, Families of Vectors, and Combinatorial Probability*, Cambridge U.P., Cambridge, 1986.
- 5 J. Brenner, The Hadamard maximum determinant problem, *Amer. Math. Monthly* **79**:626–629 (1972).
- 6 A. E. Brouwer, An Infinite Series of Symmetric Designs, Report ZW202/83, Math. Centrum Amsterdam, 1983.
- 7 F. Bussemaker, I. Kaplansky, B. McKay, and J. Seidel, Determinants of matrices of conference type, preprint.
- 8 T. Chadjipantelis, C. Moysiadis, and S. Kounias, The maximum determinant of  $21 \times 21$   $(\pm 1, -1)$ -matrices and D-optimal designs, *J. Statist. Plann. Inference* **16**:167–178 (1987).
- 9 G. F. Clements and B. Lindstrom, A sequence of  $(\pm 1)$ -determinants with large values, *Proc. Amer. Math. Soc.* **16**:548–550 (1965).
- 10 A. Colucci, Sui valore massimi dei determinanti ad elementi  $+1$  e  $-1$ , *Giorn. Mat. Battaglini* **64**:217–221 (1926).
- 11 J. H. E. Cohn, Determinants with elements  $\pm 1$ , *J. London Math. Soc.* **42**:436–442 (1967).
- 12 J. H. E. Cohn, On determinants with elements  $\pm 1$ , II, *Bull. London Math. Soc.* **21**:36–42 (1989).
- 13 H. Ehlich, Determinantenabschätzungen für binäre Matrizen, *Math. Z.* **83**:123–132 (1964).
- 14 H. Ehlich, Determinantenabschätzungen für binäre Matrizen mit  $n \equiv 3 \pmod{4}$ , *Math. Z.* **84**:438–477 (1964).
- 15 Z. Galil and J. Kiefer, D-optimum weighing designs, *Ann. Statist.* **8**(6):1293–1306 (1980).
- 16 J. Hadamard, Résolution d'une question relative aux déterminants, *Bull. Sci. Math.* **2**:240–246 (1893).
- 17 I. Kaplansky, A theorem on doubling quadratic forms and a theorem of Ehlich, preprint.
- 18 I. Kaplansky, private communication.
- 19 C. Koukouvinos, S. Kounias, and J. Seberry, Supplementary difference sets and optimal designs, *Discrete Math.* **88**:49–58 (1991).

- 20 J. H. van Lint and R. M. Wilson, *A Course in Combinatorics*, Cambridge U.P., 1992.
- 21 C. Moyssiadis and S. Kounias, The exact  $D$ -optimal first order saturated design with 17 observations, *J. Statist. Plann. Inference* **7**:13–27 (1982).
- 22 T. Popoviciu, Remarques sur le maximum d'un déterminant dont torus les éléments sont non-négatifs, *Soc. Sci. Cluj* **8**:572–582 (1937).
- 23 B. Solomon, private communication.
- 24 K. Shah and B. Sinha, *Theory of Optimal Designs*, Lecture Notes in Statist. 54, Springer-Verlag, 1989.
- 25 J. R. Seberry and M. Yamada, Hadamard matrices, sequences, and block designs, in *Contemporary Design Theory—A Collection of Surveys* (J. H. Dinitz and D. R. Stinson, Eds.), Wiley.
- 26 J. J. Sylvester, Thoughts on inverse orthogonal matrices, simultaneous sign-succession, and tessellated pavements in two or more colours, with applications to Newton's rule, ornamental tile-work, and the theory of numbers, *London Edinburgh and Dublin Philos. Mag. and J. Sci.* **34**:461–475 (1867).
- 27 A. Whiteman, A family of  $D$ -optimal designs, *Ars Combin.* **30**:23–26 (1990).
- 28 M. Mojtas, On Hadamard's inequality for the determinants of order non-divisible by 4, *Colloq. Math.* **12**:73–83 (1964).

*Received 21 July 1995; final manuscript accepted 18 January 1996*