

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Biochimica et Biophysica Acta 1662 (2004) 102–112



Review

# Gap junctions and connexin-mediated communication in the immune system

Ernesto Oviedo-Orta<sup>a,1</sup>, W. Howard Evans<sup>b,\*</sup><sup>a</sup>*Bristol Heart Institute, Bristol Royal Infirmary, Level 7, Upper Maudlin Street, Bristol BS2 8HW, UK*<sup>b</sup>*Department of Medical Biochemistry and Immunology, University of Wales, College of Medicine, Heath Park, Cardiff CF14 4XN, UK*

Received 4 July 2003; received in revised form 20 October 2003; accepted 20 October 2003

## Abstract

Gap junctions and connexins are present in the immune system. In haematopoiesis, connexin 43, the most widely distributed gap junction protein, appears to be a key player in the development of progenitor cells and their communication with stromal cells. Connexin 43 is expressed by macrophages, neutrophils and mast cells. Lymphocytes also express connexin 43, and inhibition of gap junction channels in these cells by using highly specific connexin mimetic reagents has profound effects on immunoglobulin secretion and synthesis of cytokines. Lymphocytes and leukocytes also communicate directly *in vitro* with endothelial cells via gap junctions. Connexins are implicated in inflammatory reactions in a range of tissues. Their involvement in atherosclerotic plaque formation in the vascular system is also a current growth point in research, and could lead to the development of therapeutic interventions.

© 2004 Elsevier B.V. All rights reserved.

**Keywords:** Connexin; Inflammation; Lymphocyte; Endothelium; Leukocyte; Monocyte; Macrophage; Mast cell; Intercellular communication

## 1. Introduction

Multicellular organisms are characterised by the operation of extensive direct intercellular communication networks that link and balance the metabolic and mechanical activities of the individual functional units. Such networks connecting and co-ordinating attached cells are found not only in compact tissues and organs but also in the various cell types of the immune system comprising the lymphoid organs as well as peripheral lymphocytes migrating through the blood and lymphatic networks (Fig. 1). Direct intercellular communication and strong adhesion are effected by gap junctions (see Evans and Martin for a comprehensive review) [1]. The present article collates information on the distribution and possible roles of gap junctions and the channel building blocks, the connexins, in the various branches of the immune system. Connexins, the proteins assembled into gap junction channels, are tetra-pass membrane proteins that oligomerise into hex-

americ hemichannels called connexons [2]. These proteins range in size from about 25 to 60 kDa with the main amino acid sequence differences found in the variable-length cytoplasmically oriented carboxyl tail. Over 20 different connexins have been identified in murine and human tissues [1]. After the recruitment of hexameric connexons to the plasma membrane, they carry out two functional options.

In the first, they may operate as unapposed hemichannels. In the second, they dock with partner connexons in neighbouring cells, to form gap junctions that allow direct transfer of molecules and ions up to approximately 1.2 kDa between the cytoplasm of the co-operating cells. Connexins have been detected by a variety of techniques in all nucleated cells, excepting adult striated muscle and sperm. In the present article, we review data demonstrating the widespread distribution of connexins in the component parts of immune system and outline and speculate on some functional implications. We also discuss the evidence pointing to the probable roles of connexin-mediated intercellular signalling by cells of the immune system in inflammation. The distribution and likely roles of gap junctions in the immune system have been addressed in previous reviews [3–5].

\* Corresponding author. Tel./fax: +44-2920-743133.

E-mail addresses: [e.oviedo-orta@bristol.ac.uk](mailto:e.oviedo-orta@bristol.ac.uk) (E. Oviedo-Orta), [wmbwhe@cardiff.ac.uk](mailto:wmbwhe@cardiff.ac.uk) (W. Howard Evans).

<sup>1</sup> Tel.: +44-117-928-4139; fax: +44-117-928-3581.




Fig. 1. Diagrammatic overview of the distribution of connexins in the immune system. The diagram shows the distribution of connexins in primary, secondary lymphoid organs and the various permutations of connexin-mediated interactions that may operate between monocytes (M) and lymphocytes (L) and endothelium cells of the vascular wall. Monocytes and lymphocytes cross the endothelial wall and may communicate with it and each other by gap junctions (double barrel) and/or by release of ATP or NAD<sup>+</sup> through connexin hemichannels (single barrel). In peripheral tissues, communication via gap junctions or hemichannels by lymphocytes (L) or macrophages (M $\phi$ ) with, for example, stromal cells or fibroblasts may also occur.

In view of the fundamental requirement for the cooperative behaviour of cells comprising the various branches of the immune system (Fig. 1), it is perhaps not surprising that gap junctions and their connexin protein building blocks have been identified in lymphoid tissues and cells (Table 1). Complex homocellular and heterocellular type cross-talk occurring between lymphocytes involves multiple receptor–ligand complexes and ion channels and these cellular interactions, for example, have recently been conceptualised as comprising an immunological synapse [6–9]. Although there is presently no direct evidence showing that connexins reside at such immunological synapses, there is indirect evidence showing that blockage of connexon channel interactions using highly specific peptide reagents disrupts immunoglobulin release and cytokine production by lymphocytes [10]. Furthermore, since cells may express multiple connexins that are likely to be assembled into a permutation of homo/heteromeric connexons, thus providing the basis for homo/heterotypic communication [1,11], these properties can equip cells to fulfil the basic requirements for the extensive and complex cross-talk that characterises the operation of cells in the immune system. Thus, in the various branches of the immune system, connexins and their assembled

channels should be considered as components that may complement the complex cascade of molecular interactions that underpin, for example, the rolling of leukocytes on the endothelial wall (Fig. 1), a process involving receptor–ligand interactions that implicate selectins, integrins, sialomucins, etc. [12].

Undoubtedly delaying the growing appreciation of functional roles for connexins in the immune system has been the paucity of morphological evidence for gap junctions in lymphoid organs. This is despite functional evidence for direct communication in cells populating the immune system suggested by the low electrical resistance detected between contacting cells, and especially the direct intercellular transfer between lymphocytes and leukocytes of small highly fluorescent probes.

Nevertheless, the low incidence of morphologically recognised gap junctions and the failure by immunologists to recognise and appreciate the consequences of specific properties highly indicative of direct and rapid intercellular communication in cells of the immune system should not exclude completely the operation of a system that is widespread in almost all tissues and organ systems. It is also worth pointing out at this stage that a connexin-dependent mechanism of intercellular communication that

Table 1  
Gap junction intercellular communication and connexin expression in cells of the immune system

Cell type	Morphology	Dye transfer	Connexin (Cx) isoform	References
<i>Homotypic interaction</i>				
T lymphocytes <sup>a</sup>	+	+	Cx43, Cx40	[59]
B lymphocytes <sup>a</sup>	+	+	Cx43, Cx40	[59]
Bone marrow stromal cells (BMSC)	+	+	Cx43	[37, 105–109]
Follicular dendritic cells (FDC)	+	+	Cx43	[42,43, 110]
Thymic epithelial cells (TEC)	+	+	Cx43	[5,38]
Monocytes	+	–	Cx43	[78]
Macrophages	+	–	Cx43, Cx37	[48,82, 111]
Polymorphonuclear neutrophils (PMN)	+	+	Cx43	[54]
Mast cells	+	–	Cx43	[18]
<i>Heterotypic interaction</i>				
T–B cells <sup>a</sup>	+	+	Cx43, Cx40	[59]
T cells–endothelium	+	+	Cx43	[75,110]
T–TEC <sup>b</sup>	+	+	Cx43	[34]
B–FDC	+	+	Cx43	[42,110]
Haematopoietic stem cells–BMSC	+	+	Cx43	[36,37, 112]
FDC–endothelium	+	–	Cx43	[110]
PMN–endothelium	+	+	Cx43	[54,100]
Monocytes–endothelium	+	–	Cx43	[70,78]
Macrophages–endothelium	+	+	Cx43	[70,113]
Macrophages–smooth muscle cells	+	–	Cx43	[70,78, 79,82]
Mast cells–fibroblast	+	–	Cx43, Cx32	[18]

<sup>a</sup> Derived from peripheral blood and/or secondary lymphoid organs.

<sup>b</sup> Thymocytes.

features the propagation of calcium waves between cells featuring unopposed connexin hemichannels in the plasma membrane is now being studied in an increasing number of mammalian cells (Fig. 1). This connexin-dependent mechanism of intercellular communication does not appear to utilise morphologically identifiable gap junctions, and features the release by cells of adenine nucleotides, especially ATP, that induce calcium signalling in close neighbours and implicating plasma membrane purinergic receptors [13–16]. Also this mechanism appears to be sensitive to cellular injury, e.g. in hypoxic conditions. Although the precise mechanism of ATP liberation by cells and the contribution of connexin hemichannels are unclear, the probable involvement of connexin hemichannels in this release process has been highlighted recently by studies showing that short connexin mimetic peptides, which correspond to amino acid sequences in the two extracellular loops of connexin 43, inhibit the release of ATP by cells induced either by photoliberation of intracellular IP<sub>3</sub> or by reducing extracellular calcium levels [17–19]. The connexin mimetic peptides are well-studied

inhibitors that prevent Ca<sup>2+</sup> wave propagation via hemichannels as well as gap junctional communication in tracheal epithelial cells [20,21]. This variation suggests that the inhibitory action of connexin mimetics on connexin hemichannels and gap junctions may vary between different tissues. Connexin hemichannels were predicted to exist on the plasma membrane from gap junction biosynthetic considerations [1,22,23], the penetration into cells of small dyes and the blockage of this dye uptake by cells by connexin antibodies [24] and from electrophysiological approaches [25–28] using *Xenopus* oocytes [29,30] and cells of the retina [31]. Their gating and regulation are being studied [32]. The potential functional importance of these unapposed connexin hemichannels is increasingly being defined in several cell types and these results clearly broaden the scope of intercellular communicative action by plasma membrane channels constructed of connexin subunits.

### 1.1. Gap junctions and connexins during haematopoiesis

Detailed characterisation of connexin expression has been carried out in the haematopoietic system where stem cells have the ability to renew themselves and give rise to the nine or more types of circulating blood cells. Earlier studies showed that connexins 30.3, 31, 31.1, 43, and 45 mRNA are expressed [33] but analysis of freshly isolated bone marrow cells as well as studies in derived cell lines concluded that connexin 43 was by far the major gap junction protein expressed by the specialised microenvironmental stromal cells as well as stem cells (Fig. 1) [33–36]. The regulatory mechanisms operational during haematopoiesis are complex [35]. Nevertheless, present evidence strongly points to important roles for connexins and gap junctions in effecting communication between haematopoietic stem cells and stromal cells [35,37]. The generation of Cx43 knockout mice has allowed more definitive studies that have further emphasised the critical role of this connexin in the intercellular networks that operate during haematopoiesis [33,35].

### 1.2. Gap junctions and connexins in lymphoid organs

T-cell lymphoid progenitors differentiate in the bone marrow and then migrate into the thymus where tight intercellular contact and adhesion with stromal cells contribute to their maturation. The thymus, with its compact meshwork of cells ensuring direct exchange of intercellular information, provides the organ platform where T cell clones differentiate to provide ultimately the peripheral repertoire of circulating cells. Thymic epithelial cells and thymocytes have been shown to form functional gap junction channels in both homotypic and heterotypic modes [38,39]. Again, these gap junctions are constructed exclusively of connexin 43, and many studies have shown that their function is differentially modulated by a variety of soluble factors

including growth hormones [5]. Effector and memory cell clones migrate from primary (thymic) to secondary (lymph node) lymphoid organs where they mature further and differentiate upon antigen stimulation, a process that is acutely dependent on direct cell-to-cell contact and co-operation. In this crucial differentiation step, other antigen-presenting cells and epithelial supporting cells play important roles. [40,41]. Connexin 43 appears to be the sole connexin expressed in lymphoid tissue (Fig. 1), and supporting its implication in antigen processing and presentation are studies showing that connexin 43 mRNA and protein expression are directly up-regulated by antigenic challenge [42]. Such studies suggest that connexin 43 is highly likely to participate in antigen processing and presentation.

Follicular dendritic cells are also key players in antigen presentation in secondary lymphoid organs with evidence from dye transfer studies for direct communication between each other and with B-lymphocytes implicating connexins [42]. It has been proposed that gap junctions contribute to the development and metabolism of germinal centres by controlling the growth follicular dendritic cells [43].

### 1.3. Gap junctions in the innate immune system

The innate immune system provides the first line of defence against pathogen infection, and consists mainly of phagocytic cells (neutrophils and macrophages) as well as dendritic and natural killer cells that act in an integrated fashion to eliminate infectious agents [44,45].

A number of studies have pointed to the presence of gap junctions in contacting macrophages [46–48] and between macrophages and intestinal epithelial cells [49], although some of these studies are controversial [48]. Specifically, electron microscopic analysis of bone marrow has shown gap junction-like structures in contacting macrophages and between macrophages and reticular cells, neutrophils, eosinophils, monocytes and erythroblasts [50]. These studies suggest a role for macrophages in haematopoiesis. The identification of connexin 43 as the constituent protein of macrophage gap junctions has been confirmed [51–54]. Continuing studies have focussed on the regulation of these gap junctions by a range of pro-inflammatory factors such as cytokines as well as growth factors [51–53]. Connexins have emerged as key proteins in polymorphonuclear neutrophil functions after these cells are activated by lipopolysaccharide *in vivo* and *in vitro* [54]. Corroborative studies have identified gap junctions at leukocyte–leukocyte and leukocyte–endothelial cell contacting areas, and gap junctions are implicated in their trans-endothelial migration and its modification under inflammatory conditions [42,51,54,55]. Both connexins 43 and 40 have been detected in these cells [56]. Coupling/attachment of neutrophils to endothelium by gap junctions is a time-dependent process that is modulated by the cytokine tumour necrosis factor and is facilitated by heterotypic cell adhesion involving a range of well-studied adhesion molecules [56].

Mast cells, located in mucosa and connective tissue, account for less than 0.2% of leukocytes, and their proliferation is dependent on direct interactions with T lymphocytes although mastocytes in connective tissue interact extensively with fibroblasts [18]. These cells are major participants in allergic interactions and in providing immunity to parasitic infections. Gap junctions connecting mast cells and fibroblasts were shown by electron microscopy in a chick embryo model [19], and they appear to be constructed of connexins 32 and 43. Such studies strengthen the universal roles played by gap junctions in the physiology of tissue microenvironments, but the effects of reagents that inhibit and disrupt gap junctions in these systems are awaited.

### 1.4. Connexins and communication in lymphocytes and between lymphocytes and endothelial cells

The pioneering electrophysiological studies of Hulser and Peters [57,58], demonstrating low intercellular resistance properties indicative of the presence of gap junctions linking attached lymphocytes, drew little attention for over 20 years. No doubt, this lack of recognition of a role for connexins in lymphocyte signalling persisted because of the perceived limited requirement for such a mechanism in cells that spend much of their time migrating through the lymphatic and blood vessels. However, several lines of evidence, accrued in many laboratories, have independently corroborated these important early studies.

Peripheral T, B and NK lymphocytes purified from healthy humans express connexin 43. Lymphocytes derived from secondary organs express, in addition, relatively low levels of Cx40 [59]. Flow cytometry approaches using anti-peptide connexin antibodies that bind to the extracellular loops confirmed that the connexins are located on the cell surface (Fig. 2) [59]. These observations are also highly suggestive of the presence of unapposed connexon hemichannels on lymphocytes with independent alternative communication functions such as the ATP-mediated propagation of calcium waves [60–62]. Calcein, a gap junction permeant fluorescent dye, was observed to transfer between contacting lymphocytes; importantly, this direct transfer was inhibited by two categories of gap junction inhibitors [59] and thus implying direct communication across these junctions. 18- $\alpha$ -Glycyrrhetic acid, a widely used inhibitor of gap junctional communication, is likely to enter cells and modify protein kinases that feature in the regulation of connexin 43 [63,64] that is extensively phosphorylated at multiple serine residues and a few threonine residues on the cytoplasmic tail [65]. In contrast, the connexin mimetic peptides that bind to two specific amino acid regions on the extracellular loops of connexin 43 are unlikely to penetrate into cells, undoubtedly a property accounting for their high specificity and the absence of inhibition of cellular metabolism [66]. Connexin mimetic peptides are thus emerging as useful tools




Fig. 2. Connexins 43 and 40 are expressed by cell extracts (A) and at the plasma membrane (B) of T, B and NK human lymphocytes. In A, lymphocytes were treated with phytohaemagglutinin (PHA-L) (lanes 1, 3 and 5) and lipopolysaccharide (LPS) (lanes 2, 4 and 6) for 0, 6 and 12 h, respectively; cellular proteins were separated by SDS-polyacrylamide gel electrophoresis; and connexin expression analysed by Western blotting using antibodies to connexins 43 or 40. Note the increase in connexin expression following addition of the two polyclonal stimulators. In B, cell surface expression of connexins was studied by FACS analysis by double labelling with T cell (anti-CD3), B cell (anti-CD19) and NK cell (anti-CD56) antibodies and a connexin antibody generated to the extracellular loops of connexin 43 [59].

in the functional dissection of gap junctions in the immune system and in unravelling roles for connexon hemichannels in cell–cell signalling [17,61,62]. The adoption of two means of connexin-mediated  $\text{Ca}^{2+}$  signalling between cells involving both hemichannels and gap junctions and the variable contribution of each route to overall communication is also likely to be influenced by the partner cells and the extracellular matrix as exemplified by the roles of integrins in modulating gap junctional communication in alveolar cells [21,67,68].

Connexin mimetic peptides have emerged as important reagents in revealing roles for connexins as functional players in lymphocytes. Exposure of lymphocytes for 10–50 h to GAP26 and GAP27, short peptides that correspond to specific amino acid sequences in the extracellular loops of connexin 43, markedly reduced the production of immunoglobulins (Ig) G, M and A in mixed cultures of human T and B lymphocytes [10] (Fig. 3). A small degree of intercellular channel rectification was observed in heterotypic and homotypic T and B cell cultures suggesting a role for lymphocyte gap junctions in the polarisation of the immune response [10]. Addition to cell media of connexin mimetics also demonstrated complex temporal inhibitory effects on the synthesis of cytokines, but less so on IL-2 and especially IL-10 (Fig. 3). FACS analysis using extracellular loop antibodies to connexin 43 of lymphocyte populations revealed differ-

ences indicating that CD4+, CD8+ and CD19+ cells displayed a characteristic surface patterns under mitogenic stimulation [59].

Interaction between lymphocytes and endothelial cells is a central feature of vascular physiology. Blood-borne lymphocytes leave the circulation via specialised postcapillary vessels, the high endothelial venules (HEV) [12]. The active processes of adhesion and transmigration across the endothelial cell layer are complex and involve a series of time-dependent molecular mechanisms which guarantee not only adhesion but also direct intercellular communication. Gap junctions are present in varying degrees of complexity along the vascular tree [69], and connexin43 [55,70], connexin40 [71,72] and connexin37 [73,74] have been found to be consistently expressed in the vascular wall as determined by mRNA and protein analysis. Lymphocytes establish functional gap junction channels with endothelial cells *in vitro* during active transendothelial migration in a time-dependent manner; these studies were reinforced by others showing that the process was inhibited by connexin mimetic peptides (Fig. 4) [75].

### 1.5. Connexins as possible inflammatory response markers

Inflammation is a local or systemic response of a tissue to infection and usually leads to structural or functional damage. Although the overall inflammatory response is




Fig. 3. Gap junction inhibitors decrease production by mixtures of T and B lymphocytes of (A) various immunoglobulins and (B) cytokines. Panel A shows the inhibition of IgG, IgM and IgA by connexin mimetic peptide GAP27 or 18- $\alpha$ -glycyrrhetic acid (alpha-GA) over 60 h. Panel B shows the inhibition of IL-2 and IL-10 by GAP27 and 18- $\alpha$ -glycyrrhetic acid over 72 h.

causally diverse, leukocyte migration, activation and maturation are key components. Leukocytes gain access to inflammation sites by interacting with parenchymal cells and the extracellular matrix, and an important aspect of their involvement in inflammation involves transmigration across endothelium.

There is increasing evidence for the implication of connexins, especially connexin 43 during inflammation. Endothelial and smooth muscle cells express connexin 43 with endothelial cells also expressing connexin 37 [1]. Normally, mononuclear cells are rarely found in subendothelial spaces but during tissue or local vascular inflammation, such as in atherosclerosis, there is a large increase in monocyte and lymphocyte numbers that associate with the vascular wall and permeate extensively into the subendothelial spaces [76,77]. In vitro studies have shown that coculturing endothelial and smooth muscle cells resulted in increased production of IL-1 and IL-6 transforming growth factor, and, importantly in the present context, significantly higher expression of connexin 43 [78]. Connexin 43 expression by leukocytes is also susceptible to LPS stimulation and increased during ischaemia reperfusion injury [54]. These observations combine to suggest a connexin involvement in the overall inflammatory response in the arterial wall. A range of supporting studies, carried out mainly in vitro, have demonstrated that gap junction-mediated communication operates as assessed by direct dye transfer between peripherally derived monocytes and endothelial and smooth muscle cells or between macrophage

foam cells [55,79]. Furthermore, lipid loading of monocytes/macrophages markedly reduced gap junctional communication [80]. Expression of connexin 43 is increased in areas of vessel intimal thickening at atheromatous lesions where foam cells aggregate [55,79]. However, a direct functional role specifically for connexin 43 is still controversial and difficult to establish unequivocally since other connexins including connexins 30, 37, and 45 are identified in these lesions [81]. Further supporting the role for connexin 43 in the inflammatory response are recent studies by Kwak et al. [82] who studied expression patterns of connexins 37, 40 and 43 in low-density-lipoprotein-deficient mice and in human atherosclerotic plaques where a redistribution of these connexins in the arterial wall occurred during the development of atheroma [82]. Analysis of the progression of atheroma indicated that it was much reduced in the aorta of low-density-lipoprotein-deficient mice as compared to controls, and these mice exhibited fewer fibrous caps. Furthermore, and of direct clinical relevance, are results showing that statins, drugs that reduce cholesterol synthesis by inhibition of HMG CoA reductase, reduced connexin 43 expression by cultured endothelial cells [83]. These diverse observations strongly indicate that connexin 43 and therefore channels constructed of this protein are implicated in inflammatory events leading to atheroma.

Besides involvement in inflammatory events in the cardiovascular system, connexin expression is modified in inflammatory events in other organ systems. When inflam-




Fig. 4. Direct intercellular communication via gap junctions between lymphocytes and endothelial cells. (A) During transendothelial migration in an in vitro assay, the transfer of calcein (a low molecular weight green fluorescent dye) from endothelial cells (E) to lymphocytes (L) was analysed over time by two photon laser scanning microscopy. In B, the specificity of transfer by a connexin-mediated mechanism was determined by FACS analysis of the dye transfer occurring in the trans-well migration system. Connexin mimetic peptide GAP27 inhibited calcein transfer between endothelial cells and lymphocytes; as a control, GAP18 (a peptide incorporating a short amino acid sequence in the intracellular loop of connexin 43) was used [75].

matory cells invade and damage tubular and interstitial cells in kidneys of patients with glomerulitis, connexin expression increases in parallel with ICAM-1 and VCAM-1 [84,85]. Connexins and gap junctions were also modified when damage occurred in the central and peripheral nervous systems. In the central nervous system, connexins are widely distributed with connexin expression varying during development and in injury processes. In summary, neurons express mainly connexins 32, 36 and 43, oligodendrocytes express connexins 29, 32 and 47, and astrocytes express connexins 26, 30 and 43 [86–90]. Leukocytes have been implicated in cerebral ischemia [91], where a role for connexins and especially connexin hemichannels in astrocytes has been investigated in stroke models [92]. In peripheral nerves, connexin 43 is mainly associated with perineural fibroblasts and Schwann cell cytoplasm [93–95], and there is high expression of connexin 32 at reflexive gap junctions in Schwann cells [96].

Connexin expression is modified in the inflammatory response to peripheral nerve damage and this has been investigated in mouse and rat models of sciatic nerve damage [97–99]. After injury there are changes in connexin 32 and 46 expression. Interestingly, there appears to be a linkage between the down regulation of connexin 32 and up-regulation of connexin 43 in endoneurial fibroblasts

and infiltrating macrophages. It has been proposed that many of these changes in various cell types are induced by the release of inflammatory cytokines by endothelial cells and transmigrating leukocytes [78,100]. The changes in expression of connexins 32 and 43 appear to vary according to the tissues examined. In liver, for example after LPS induced inflammation, connexin 32 is rapidly degraded [101–103], an event that coincides with a decrease in cell communication ascertained by lowered dye coupling [104].

Connexin 26 is one of the smallest connexins with a short 16-amino-acid cytoplasmic tail (connexin 43 has a tail of 156 amino acids). Connexin 26 is associated with inflammation in the cochlea where cells of the stria vascularis are connected by gap junctions. It has been suggested that local release of proinflammatory cytokines such as tumour necrosis factor alpha or interleukin 1 beta by immune system cells may contribute to the lowering of connexin 26 expression. This decrease in connexin 26 levels in the ear contrasts with the lack of change in connexin 26 expression in liver following LPS treatment [104].

The above observations combine to indicate that acute inflammation involving cells of the immune system can trigger changes in the expression and probably the functional involvement of gap junctions communication chan-

nels in a number of tissue/organ systems infiltrated by leukocytes.

However, it is apparent that these changes monitored at the protein or RNA levels are complex and vary between various tissue/organ systems. However, one may generalise and comment that changes in connexin 43 are mainly associated with early inflammatory events involving macrophage infiltration and endothelial cell activation. In contrast, down-regulation of connexin 32 seems to point to tissue damage and the consequential functional disruption. Various systems are thus likely to be affected in different ways probably because of the presence of different connexin expression profiles allowing channels linking similar and dissimilar cells with varying selectivity to be established.

## References

- [1] W.H. Evans, P.E. Martin, Gap junctions: structure and function, *Mol. Membr. Biol.* 19 (2002) 121–136.
- [2] S. Ahmad, P.E. Martin, W.H. Evans, Assembly of gap junction channels: mechanism, effects of calmodulin antagonists and identification of connexin oligomerization determinants, *Eur. J. Biochem.* 268 (2001) 4544–4552.
- [3] L.A. Alves, A.C. de Carvalho, W. Savino, Gap junctions: a novel route for direct cell–cell communication in the immune system? *Immunol. Today* 19 (1998) 269–275.
- [4] E. Oviedo-Orta, W.H. Evans, Gap junctions and connexins: potential contributors to the immunological synapse, *J. Leukoc. Biol.* 72 (2002) 636–642.
- [5] L.A. Alves, O.K. Nihei, P.C. Fonseca, A.C. Carvalho, W. Savino, Gap junction modulation by extracellular signaling molecules: the thymus model, *Braz. J. Med. Biol. Res.* 33 (2000) 457–465.
- [6] M.F. Krummel, M.M. Davis, Dynamics of the immunological synapse: finding, establishing and solidifying a connection, *Curr. Opin. Immunol.* 14 (2002) 66–74.
- [7] M.I. Dustin, P.M. Allen, A.S. Shaw, Environmental control of immunological synapse formation and duration, *Trends Immunol.* 22 (2001) 192–194.
- [8] S.K. Bromley, W.R. Burack, K.G. Johnson, K. Somersalo, T.N. Sims, C. Sumen, M.M. Davis, A.S. Shaw, P.M. Allen, M.L. Dustin, The immunological synapse, *Annu. Rev. Immunol.* 19 (2001) 375–396.
- [9] M.I. Dustin, J.A. Cooper, The immunological synapse and the actin cytoskeleton: molecular hardware for T cell signaling, *Nat. Immunol.* 1 (2000) 23–29.
- [10] E. Oviedo-Orta, P. Gasque, W.H. Evans, Immunoglobulin and cytokine expression in mixed lymphocyte cultures is reduced by disruption of gap junction intercellular communication, *FASEB J.* 15 (2001) 768–774.
- [11] W.H. Evans, S. Ahmad, J. Diez, C.H. George, J.M. Kendall, P.E. Martin, Trafficking pathways leading to the formation of gap junctions, *Novartis Found. Symp.* 219 (44–54) (1999) 54–59 (discussion).
- [12] J.P. Girard, T.A. Springer, High endothelial venules (HeVs)—specialized endothelium for lymphocyte migration, *Immunol. Today* 16 (1995) 449–457.
- [13] K. Paemeleire, P.E. Martin, S.L. Coleman, K.E. Fogarty, W.A. Carrington, L. Leybaert, R.A. Tuft, W.H. Evans, M.J. Sanderson, Intercellular calcium waves in HeLa cells expressing GFP-labeled connexin 43, 32, or 26, *Mol. Biol. Cell* 11 (2000) 1815–1827.
- [14] M.L. Cotrina, J.H. Lin, A. Alves-Rodrigues, S. Liu, J. Li, H. Azmi-Ghadimi, J. Kang, C.C. Naus, M. Nedergaard, Connexins regulate calcium signaling by controlling ATP release, *Proc. Natl. Acad. Sci. U. S. A.* 95 (1998) 15735–15740.
- [15] P.B. Guthrie, J. Knappenberger, M. Segal, M.V. Bennett, A.C. Charles, S.B. Kater, ATP released from astrocytes mediates glial calcium waves, *J. Neurosci.* 19 (1999) 520–528.
- [16] C.E. Stout, J.L. Costantin, C.G. Naus, A.C. Charles, Intercellular calcium signaling in astrocytes via ATP release through connexin hemichannels, *J. Biol. Chem.* 277 (2002) 10482–10488.
- [17] K. Braet, W. Vandamme, P.E. Martin, W.H. Evans, L. Leybaert, Photoliberating inositol-1,4,5-trisphosphate triggers ATP release that is blocked by the connexin mimetic peptide gap 26, *Cell Calcium* 33 (2003) 37–48.
- [18] H. Vliagoftis, A.M. Hutson, S. Mahmudi-Azer, H. Kim, V. Rumsaeng, C.K. Oh, R. Moqbel, D.D. Metcalfe, Mast cells express connexins on their cytoplasmic membrane, *J. Allergy Clin. Immunol.* 103 (1999) 656–662.
- [19] S.M. Oliani, A.P. Girol, R.L. Smith, Gap junctions between mast cells and fibroblasts in the developing avian eye, *Acta Anat. (Basel)* 154 (1995) 267–271.
- [20] S. Boitano, W.H. Evans, Connexin mimetic peptides reversibly inhibit Ca(2+) signaling through gap junctions in airway cells, *Am. J. Physiol., Lung Cell. Mol. Physiol.* 279 (2000) L623–L630.
- [21] B.E. Isakson, W.H. Evans, S. Boitano, Intercellular Ca<sup>2+</sup> signaling in alveolar epithelial cells through gap junctions and by extracellular ATP 280, *Am. J. Physiol., Lung Cell. Mol. Physiol.* 280 (2001) L221–L228.
- [22] G. Gaietta, T.J. Deerinck, S.R. Adams, J. Bouwer, O. Tour, D.W. Laird, G.E. Sosinsky, R.Y. Tsien, M.H. Ellisman, Multicolor and electron microscopic imaging of connexin trafficking, *Science* 296 (2002) 503–507.
- [23] W.H. Evans, P.E. Martin, Lighting up gap junction channels in a flash, *BioEssays* 24 (2002) 876–880.
- [24] R.A. Meyer, D.W. Laird, J.P. Revel, R.G. Johnson, Inhibition of gap junction and adherens junction assembly by connexin and A-CAM antibodies, *J. Cell Biol.* 119 (1992) 179–189.
- [25] S. Haubrich, H.J. Schwarz, F. Bukauskas, H. Lichtenberg-Frate, O. Traub, R. Weingart, K. Willecke, Incompatibility of connexin 40 and 43 Hemichannels in gap junctions between mammalian cells is determined by intracellular domains, *Mol. Biol. Cell* 7 (1996) 1995–2006.
- [26] G. Pfahnl A Dahl, G. Dahl, Localization of a voltage gate in connexin46 gap junction hemichannels, *Biophys. J.* 75 (1998) 2323–2331.
- [27] E.B. Trexler, F.F. Bukauskas, M.V. Bennett, T.A. Bargiello, V.K. Verselis, Rapid and direct effects of pH on connexins revealed by the connexin46 hemichannel preparation, *J. Gen. Physiol.* 113 (1999) 721–742.
- [28] R.P. Kondo, S.Y. Wang, S.A. John, J.N. Weiss, J.I. Goldhaber, Metabolic inhibition activates a non-selective current through connexin hemichannels in isolated ventricular myocytes, *J. Mol. Cell. Cardiol.* 32 (2000) 1859–1872.
- [29] D.L. Paul, L. Ebihara, L.J. Takemoto, K.I. Swenson, D.A. Goodenough, Connexin46, a novel lens gap junction protein, induces voltage-gated currents in nonjunctional plasma membrane of *Xenopus* oocytes, *J. Cell Biol.* 115 (1991) 1077–1089.
- [30] G.A. Zampighi, D.D. Loo, M. Kreman, S. Eskandari, E.M. Wright, Functional and morphological correlates of connexin50 expressed in *Xenopus laevis* oocytes, *J. Gen. Physiol.* 113 (1999) 507–524.
- [31] M. Kamermans, I. Fahrenfort, K. Schultz, U. Janssen-Bienhold, T. Sjoerdsma, R. Weiler, Hemichannel-mediated inhibition in the outer retina, *Science* 292 (2001) 1178–1180.
- [32] J.E. Contreras, J.C. Saez, F.F. Bukauskas, M.V. Bennett, Gating and regulation of connexin 43 (Cx43) hemichannels, *Proc. Natl. Acad. Sci. U. S. A.* 16 (2003) 12960393.
- [33] J.A. Cancelas, W.L. Koevoet, A.E. de Koning, A.E. Mayen, E.J. Rombouts, R.E. Ploemacher, Connexin-43 gap junctions are in-



- involved in multiconnexin-expressing stromal support of hemopoietic progenitors and stem cells, *Blood* 96 (2000) 498–505.
- [34] E. Montecino-Rodriguez, H. Leathers, K. Dorshkind, Expression of connexin 43 (Cx43) is critical for normal hematopoiesis, *Blood* 96 (2000) 917–924.
- [35] E. Montecino-Rodriguez, K. Dorshkind, Regulation of hematopoiesis by gap junction-mediated intercellular communication, *J. Leukoc. Biol.* 70 (2001) 341–347.
- [36] T. Krenacs, M. Rosendaal, Connexin43 gap junctions in normal, regenerating, and cultured mouse bone marrow and in human leukemias: their possible involvement in blood formation, *Am. J. Pathol.* 152 (1998) 993–1004.
- [37] M.M. Rosendaal, T.T. Krenacs, Regulatory pathways in blood-forming tissue with particular reference to gap junctional communication, *Pathol. Oncol. Res.* 6 (2000) 243–249.
- [38] L.A. Alves, A.C. Campos de Carvalho, E.O. Cirne Lima, C.M. Rocha e Souza, M. Dardenne, D.C. Spray, W. Savino, Functional gap junctions in thymic epithelial cells are formed by connexin 43, *Eur. J. Immunol.* 25 (1995) 431–437.
- [39] L.A. Alves, A.C. de Carvalho, L. Parreira-Martins, M. Dardenne, W. Savino, Intrathymic gap junction-mediated communication, *Adv. Exp. Med. Biol.* 355 (1994) 155–158.
- [40] F.W. Lusinskas, S. Ma, A. Nusrat, C.A. Parkos, S.K. Shaw, The role of endothelial cell lateral junctions during leukocyte trafficking, *Immunol. Rev.* 186 (2002) 57–67.
- [41] R.A. Worthylyake, K. Burridge, Leukocyte transendothelial migration: orchestrating the underlying molecular machinery, *Curr. Opin. Cell Biol.* 13 (2001) 569–577.
- [42] T. Krenacs, M. van Dartel, E. Lindhout, M. Rosendaal, Direct cell/cell communication in the lymphoid germinal center: connexin43 gap junctions functionally couple follicular dendritic cells to each other and to B lymphocytes, *Eur. J. Immunol.* 27 (1997) 1489–1497.
- [43] T. Krenacs, M. Rosendaal, Gap-junction communication pathways in germinal center reactions, *Dev. Immunol.* 6 (1998) 111–118.
- [44] A. Mantovani, M. Muzio, C. Garlanda, S. Sozzani, P. Allavena, Macrophage control of inflammation: negative pathways of regulation of inflammatory cytokines, *Novartis Found. Symp.* 234 (2001) 120–131.
- [45] A. Matsukawa, M. Yoshinaga, Neutrophils as a source of cytokines in inflammation, *Histol. Histopathol.* 14 (1999) 511–516.
- [46] J.A. Levy, R.M. Weiss, E.R. Dirksen, M.R. Rosen, Possible communication between murine macrophages oriented in linear chains in tissue culture, *Exp. Cell Res.* 103 (1976) 375–385.
- [47] M. Porvaznik, T.J. MacVittie, Detection of gap junctions between the progeny of a canine macrophage colony-forming cell in vitro, *J. Cell Biol.* 82 (1979) 555–564.
- [48] L.A. Alves, R. Coutinho-Silva, P.M. Persechini, D.C. Spray, W. Savino, A.C. Campos de Carvalho, Are there functional gap junctions or junctional hemichannels in macrophages? *Blood* 88 (1996) 328–334.
- [49] C.A. Martin, F.R. Homaidan, T. Palaia, R. Burakoff, M.E. el-Sabban, Gap junctional communication between murine macrophages and intestinal epithelial cell lines, *Cell Adhes. Commun.* 5 (1998) 437–449.
- [50] F.R. Campbell, Gap junctions between cells of bone marrow: an ultrastructural study using tannic acid, *Anat. Rec.* 196 (1980) 101–107.
- [51] E.A. Eugenin, M.C. Branes, J.W. Berman, J.C. Saez, TNF-alpha plus IFN-gamma induce connexin43 expression and formation of gap junctions between human monocytes/macrophages that enhance physiological responses, *J. Immunol.* 170 (2003) 1320–1328.
- [52] N. Rouach, C.F. Calvo, J. Glowinski, C. Giaume, Brain macrophages inhibit gap junctional communication and downregulate connexin 43 expression in cultured astrocytes, *Eur. J. Neurosci.* 15 (2002) 403–407.
- [53] R. Huang, Y. Lin, C.C. Wang, J. Gano, B. Lin, Q. Shi, A. Boynton, J. Burke, R.P. Huang, Connexin 43 suppresses human glioblastoma cell growth by down-regulation of monocyte chemotactic protein 1, as discovered using protein array technology, *Cancer* (2002) 2806–2812.
- [54] P.I. Jara, M.P. Boric, J.C. Saez, Leukocytes express connexin 43 after activation with lipopolysaccharide and appear to form gap junctions with endothelial cells after ischemia-reperfusion, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 7011–7015.
- [55] D. Polacek, F. Bech, J.F. McKinsey, P.F. Davies, Connexin43 gene expression in the rabbit arterial wall: effects of hypercholesterolemia, balloon injury and their combination, *J. Vasc. Res.* 34 (1997) 19–30.
- [56] M.C. Branes, J.E. Contreras, J.C. Saez, Activation of human polymorphonuclear cells induces formation of functional gap junctions and expression of connexins, *Med. Sci. Monit.* 8 (2002) BR313–BR323.
- [57] D.F. Hulser, J.H. Peters, Contact cooperation in stimulated lymphocytes: II. Electrophysiological investigations on intercellular communication, *Exp. Cell Res.* 74 (1972) 319–326.
- [58] D.F. Hulser, J.H. Peters, Intercellular communication in phytohemagglutinin-induced lymphocyte agglutinates, *Eur. J. Immunol.* 1 (1971) 494–495.
- [59] E. Oviedo-Orta, T. Hoy, W.H. Evans, Intercellular communication in the immune system: differential expression of connexin40 and 43, and perturbation of gap junction channel functions in peripheral blood and tonsil human lymphocyte subpopulations, *Immunology* 99 (2000) 578–590.
- [60] D.A. Goodenough, D.L. Paul, Beyond the gap: functions of unpaired connexon channels, *Nat. Rev., Mol. Cell Biol.* 4 (2003) 285–294.
- [61] K. Braet, S. Aspeslagh, W. Vandamme, K. Willecke, P.E.M. Martin, W.H. Evans, L. Laybaert, Pharmacological sensitivity of ATP release triggered by photoliberation of inositol-1,4,5-trisphosphate and zero extracellular calcium in brain endothelial cells, *J. Cell. Physiol.* 197 (2003) 205–213.
- [62] K. Braet, K. Paemeleire, K. D’Herde, M.J. Sanderson, L. Leybaert, Astrocyte–endothelial cell calcium signals conveyed by two signalling pathways, *Eur. J. Neurosci.* 13 (2001) 79–91.
- [63] R.B. Stagg, W.H. Fletcher, The hormone-induced regulation of contact-dependent cell–cell communication by phosphorylation, *Endocr. Rev.* 11 (1990) 302–325.
- [64] V. Cruciani, S.O. Mikalsen, Connexins, gap junctional intercellular communication and kinases, *Biol. Cell* 94 (2002) 433–443.
- [65] P.D. Lampe, A.F. Lau, Regulation of gap junctions by phosphorylation of connexins, *Arch. Biochem. Biophys.* 384 (2000) 205–215.
- [66] W.H. Evans, S. Boitano, Connexin mimetic peptides: specific inhibitors of gap-junctional intercellular communication, *Biochem. Soc. Trans.* 29 (2001) 606–612.
- [67] M. Koval, Sharing signals: connecting lung epithelial cells with gap junction channels, *Am. J. Physiol., Lung Cell. Mol. Physiol.* 283 (2002) L875–L893.
- [68] B.E. Isakson, G.J. Seedorf, R.L. Lubman, W.H. Evans, S. Boitano, Cell–cell communication in heterocellular cultures of alveolar epithelial cells, *Am. J. Respir. Cell Mol. Biol.* 14 (2003) 14.
- [69] E. Dejana, M. Corada, M.G. Lampugnani, Endothelial cell-to-cell junctions, *FASEB J.* 9 (1995) 910–918.
- [70] D. Polacek, R. Lal, M.V. Volin, P.F. Davies, Gap junctional communication between vascular cells. Induction of connexin43 messenger RNA in macrophage foam cells of atherosclerotic lesions, *Am. J. Pathol.* 142 (1993) 593–606.
- [71] R. Bruzzone, J.A. Haefliger, R.L. Gimlich, D.L. Paul, Connexin40, a component of gap junctions in vascular endothelium, is restricted in its ability to interact with other connexins, *Mol. Biol. Cell* 4 (1993) 7–20.
- [72] D. Gros, T. Jarry-Guichard, I. Ten Velde, A. de Maziere, M.J. van Kempen, J. Davoust, J.P. Briand, A.F. Moorman, H.J. Jongsma,

- Restricted distribution of connexin40, a gap junctional protein, in mammalian heart, *Circ. Res.* 74 (1994) 839–851.
- [73] D.M. Larson, M.J. Wroblewski, G.D. Sagar, E.M. Westphale, E.C. Beyer, Differential regulation of connexin43 and connexin37 in endothelial cells by cell density, growth, and TGF-beta1, *Am. J. Physiol.* 272 (1997) C405–C415.
- [74] H.V. van Rijen, M.J. van Kempen, S. Postma, H.J. Jongsma, Tumour necrosis factor alpha alters the expression of connexin43, connexin40, and connexin37 in human umbilical vein endothelial cells, *Cytokine* 10 (1998) 258–264.
- [75] E. Oviedo-Orta, R.J. Errington, W.H. Evans, Gap junction intercellular communication during lymphocyte transendothelial migration, *Cell Biol. Int.* 26 (2002) 253–263.
- [76] L.A. Salamonsen, J. Zhang, M. Brasted, Leukocyte networks and human endometrial remodelling, *J. Reprod. Immunol.* 57 (2002) 95–108.
- [77] W.A. Muller, Leukocyte–endothelial cell interactions in the inflammatory response, *Lab. Invest.* 82 (2002) 521–533.
- [78] M. Navab, F. Liao, G.P. Hough, L.A. Ross, B.J. Van Lenten, T.B. Rajavashisth, A.J. Lusis, H. Laks, D.C. Drinkwater, A.M. Fogelman, Interaction of monocytes with cocultures of human aortic wall cells involves interleukins 1 and 6 with marked increases in connexin43 message, *J. Clin. Invest.* 87 (1991) 1763–1772.
- [79] J.P. Blackburn, N.S. Peters, H.I. Yeh, S. Rothery, C.R. Green, N.J. Severs, Upregulation of connexin43 gap junctions during early stages of human coronary atherosclerosis, *Arterioscler. Thromb. Vasc. Biol.* 15 (1995) 1219–1228.
- [80] E.R. Andreeva, V.N. Serebryakov, A.N. Orekhov, Gap junctional communication in primary culture of cells derived from human aortic intima, *Tissue Cell* 27 (1995) 591–597.
- [81] N.J. Severs, S. Rothery, E. Dupont, S.R. Coppen, H.I. Yeh, Y.S. Ko, T. Matsushita, R. Kaba, D. Halliday, Immunocytochemical analysis of connexin expression in the healthy and diseased cardiovascular system, *Microsc. Res. Tech.* 52 (2001) 301–322.
- [82] B.R. Kwak, F. Mulhaupt, N. Veillard, D.B. Gros, F. Mach, Altered pattern of vascular connexin expression in atherosclerotic plaques, *Arterioscler. Thromb. Vasc. Biol.* 22 (2002) 225–230.
- [83] B.R. Kwak, N. Veillard, G. Pelli, F. Mulhaupt, R.W. James, M. Chanson, F. Mach, Reduced connexin43 expression inhibits atherosclerotic lesion formation in low-density lipoprotein receptor-deficient mice, *Circulation* 107 (2003) 1033–1039.
- [84] G.S. Hillis, L.A. Duthie, R. Mlynski, N.G. McKay, S. Mistry, A.M. MacLeod, J.G. Simpson, N.E. Haites, The expression of connexin 43 in human kidney and cultured renal cells, *Nephron* 75 (1997) 458–463.
- [85] G.S. Hillis, L.A. Duthie, P.A. Brown, J.G. Simpson, A.M. MacLeod, N.E. Haites, Upregulation and co-localization of connexin43 and cellular adhesion molecules in inflammatory renal disease, *J. Pathol.* 182 (1997) 373–379.
- [86] J. Li, E.L. Hertzberg, J.I. Nagy, Connexin32 in oligodendrocytes and association with myelinated fibers in mouse and rat brain, *J. Comp. Neurol.* 379 (1997) 571–591.
- [87] B. Nadarajah, A.M. Jones, W.H. Evans, J.G. Parnavelas, Differential expression of connexins during neocortical development and neuronal circuit formation, *J. Neurosci.* 17 (1997) 3096–3111.
- [88] B. Nadarajah, J.G. Parnavelas, Gap junction-mediated communication in the developing and adult cerebral cortex, *Novartis Found. Symp.* 219 (1999) 157–170.
- [89] R. Dermietzel, Diversification of gap junction proteins (connexins) in the central nervous system and the concept of functional compartments, *Cell Biol. Int.* 22 (1998) 719–730.
- [90] R. Dermietzel, Gap junction wiring: a ‘new’ principle in cell-to-cell communication in the nervous system? *Brain Res. Brain Res. Rev.* 26 (1998) 176–183.
- [91] G. Stoll, S. Jander, M. Schroeter, Inflammation and glial responses in ischemic brain lesions, *Prog. Neurobiol.* 56 (1998) 149–171.
- [92] J.H. Lin, H. Weigel, M.L. Cotrina, S. Liu, E. Bueno, A.J. Hansen, T.W. Hansen, S. Goldman, M. Nedergaard, Gap-junction-mediated propagation and amplification of cell injury, *Nat. Neurosci.* 1 (1998) 494–500.
- [93] S. Zhao, A. Fort, D.C. Spray, Characteristics of gap junction channels in Schwann cells from wild-type and connexin-null mice, *Ann. N. Y. Acad. Sci.* 883 (1999) 533–537.
- [94] T. Yoshimura, M. Satake, T. Kobayashi, Connexin43 is another gap junction protein in the peripheral nervous system, *J. Neurochem.* 67 (1996) 1252–1258.
- [95] E.T. Mambetisaeva, V. Gire, W.H. Evans, Multiple connexin expression in peripheral nerve, Schwann cells, and Schwannoma cells, *J. Neurosci. Res.* 57 (1999) 166–175.
- [96] J. Bergoffen, S.S. Scherer, S. Wang, M.O. Scott, L.J. Bone, D.L. Paul, K. Chen, M.W. Lensch, P.F. Chance, K.H. Fischbeck, Connexin mutations in X-linked Charcot–Marie–Tooth disease, *Science* 262 (1993) 2039–2042.
- [97] C.Y. Lin Sh Lu, R. Lu Cy Muhammad, W.Y. Muhammad R Chou, F.C. Chou Wy Lin, P.C. Lin Fe Wu, C.R. Wu Pc Lin, L.C. Lin Cr Yang, L.C. Yang, Induction of connexin 37 expression in a rat model of neuropathic pain, *Brain Res. Mol. Brain Res.* 99 (2002) 134–140.
- [98] Z. Sahenk, Abnormal Schwann cell–axon interactions in CMT neuropathies. The effects of mutant Schwann cells on the axonal cytoskeleton and regeneration-associated myelination, *Ann. N. Y. Acad. Sci.* 883 (1999) 415–426.
- [99] P. Anzini, D.H. Neuberger, M. Schachner, E. Nelles, K. Willecke, J. Zielasek, K.V. Toyka, U. Suter, R. Martini, Structural abnormalities and deficient maintenance of peripheral nerve myelin in mice lacking the gap junction protein connexin 32, *J. Neurosci.* 17 (1997) 4545–4551.
- [100] S. Zahler, A. Hoffmann, T. Gloe, U. Pohl, Gap-junctional coupling between neutrophils and endothelial cells: a novel modulator of transendothelial migration, *J. Leukoc. Biol.* 73 (2003) 118–126.
- [101] N.G. Theodorakis, A. De Maio, Cx32 mRNA in rat liver: effects of inflammation on poly(A) tail distribution and mRNA degradation, *Am. J. Physiol.* 276 (1999) R1249–R1257.
- [102] A. Temme, T. Ott, F. Dombrowski, K. Willecke, The extent of synchronous initiation and termination of DNA synthesis in regenerating mouse liver is dependent on connexin32 expressing gap junctions, *J. Hepatol.* 32 (2000) 627–635.
- [103] A. Temme, T. Ott, T. Haberberger, O. Traub, K. Willecke, Acute-phase response and circadian expression of connexin26 are not altered in connexin32-deficient mouse liver, *Cell Tissue Res.* 300 (2000) 111–117.
- [104] A. De Maio, C. Gingalewski, N.G. Theodorakis, M.G. Clemens, Interruption of hepatic gap junctional communication in the rat during inflammation induced by bacterial lipopolysaccharide, *Shock* 14 (2000) 53–59.
- [105] K. Yamazaki, *Sl/Slid* mice have an increased number of gap junctions in their bone marrow stromal cells, *Blood Cells* 13 (1988) 421–435.
- [106] A. Umezawa, K. Harigaya, H. Abe, Y. Watanabe, Gap-junctional communication of bone marrow stromal cells is resistant to irradiation in vitro, *Exp. Hematol.* 18 (1990) 1002–1007.
- [107] A. Umezawa, J. Hata, Expression of gap-junctional protein (connexin 43 or alpha 1 gap junction) is down-regulated at the transcriptional level during adipocyte differentiation of H-1/A marrow stromal cells, *Cell Struct. Funct.* 17 (1992) 177–184.
- [108] K. Dorshkind, L. Green, A. Godwin, W.H. Fletcher, Connexin-43-type gap junctions mediate communication between bone marrow stromal cells, *Blood* 82 (1993) 38–45.
- [109] J. Durig, C. Rosenthal, K. Halfmeyer, M. Wiemann, J. Novotny, D. Bingmann, U. Dührsen, K. Schirmacher, Intercellular communication between bone marrow stromal cells and CD34+ haematopoietic progenitor cells is mediated by connexin 43-type gap junctions, *Br. J. Haematol.* 111 (2000) 416–425.
- [110] T. Krenacs, M. Rosendaal, Immunohistological detection of gap junctions in human lymphoid tissue: connexin43 in follicular den-

- dritic and lymphoendothelial cells, *J. Histochem. Cytochem.* 43 (1995) 1125–1137.
- [111] E.C. Beyer, T.H. Steinberg, Evidence that the gap junction protein connexin-43 is the ATP-induced pore of mouse macrophages, *J. Biol. Chem.* 266 (1991) 7971–7974.
- [112] R.E. Ploemacher, A.E. Mayen, A.E. De Koning, T. Krenacs, M. Rosendaal, Hematopoiesis: gap junction intercellular communication is likely to be involved in regulation of stroma-dependent proliferation of hemopoietic stem cells, *Hematology* 5 (2000) 133–147.
- [113] N. Takasu, H. Kurihara, K. Uchida, Early changes in fine structures of the aortic arch in hamsters fed a high cholesterol diet, *Arch. Histol. Cytol.* 53 (1990) 211–218.