

# The importance of the negative charge of $\beta$ -lactam compounds for the inactivation of the active-site serine DD-peptidase of *Streptomyces* R61

Louis Varetto, Jean-Marie Frère and Jean-Marie Ghuysen

Service de Microbiologie, Université de Liège, Institut de Chimie, B6, Sart Tilman, B-4000 Liège, Belgium

Received 9 October 1987



The interaction between the *Streptomyces* R61 penicillin-sensitive DD-peptidase and deacetyl-cephalosporin C or its lactone derivative has been studied at different pH values. The results show the importance of an enzyme group of  $pK \cong 9$  which might form an ion pair with the free carboxylate of the former compound. This electrostatic interaction is shown to contribute to the formation of the first, non-covalent enzyme-inactivator complex by a factor of at least 50.

$\beta$ -Lactam; pH dependence; Enzyme inactivation; Deacetyl-cephalosporin C; DD-peptidase

## 1. INTRODUCTION

The study of the pH dependence of the active-site serine DD-peptidase of *Streptomyces* R61 [1] has revealed that the deprotonation of an enzyme group of  $pK \cong 9.3-9.6$  induces a sharp decrease in the rates of enzyme-catalysed hydrolysis of the peptide substrate and inactivation by various  $\beta$ -lactams. The group concerned has been hypothesized to be the side chain of a lysine residue. Indeed, the active-site serine residue of penicillin-binding proteins and  $\beta$ -lactamases is invariably followed by an -X-X-Lys sequence [2-5], suggesting that charge pairing between an enzyme positive group and the C-terminal carboxylate of the substrate or the carboxylate on C<sub>3</sub> of penicillins (or C<sub>4</sub> of cephalosporins) might be important for recognition of substrate and inactivator by the enzyme. Accordingly, esterification of the carboxylate on C<sub>3</sub> of penicillins strongly decreases the

antibacterial activity [6]. However, with cephalosporins, lactonization of the C<sub>4</sub> carboxylate with an alcohol function borne by the C<sub>3</sub> substituent yields active compounds. Moreover, the lactone of deacetyl-cephalosporin C (I) is a reasonably good substrate of  $\beta$ -lactamase I from *Bacillus cereus* [7]. In preliminary experiments, inactivation of *Streptomyces* R61 DD-peptidase by I was found to proceed at a non-negligible rate.



The present work compares deacetyl-cephalosporin C (II) to the corresponding lactone (I) as inactivators of *Streptomyces* R61 DD-peptidase.

Correspondence address: L. Varetto, Service de Microbiologie, Université de Liège, Institut de Chimie, B6, Sart Tilman, B-4000 Liège, Belgium

## 2. MATERIALS AND METHODS

All experiments were performed at 25°C.

### 2.1. Enzymes

*Streptomyces* R61 DD-peptidase and *B. licheniformis*  $\beta$ -lactamase were purified as described [8,9].

### 2.2. $\beta$ -Lactam compounds

Compounds I and II were kind gifts from Dr J.B. Ward (Glaxo Research Group, Greenford, Middlesex, England).

### 2.3. Buffers


The following buffers were used: pH 7.0, 10 mM sodium phosphate; pH 8.0, 7 mM sodium phosphate; pH 9, 12 mM glycine + NaOH; pH 9.5, 15 mM glycine + NaOH; pH 10 and 10.5, 25 mM glycine + NaOH. The ionic strength of all buffers was adjusted to a conductance of 0.1 mS using a Methrom E527 conductometer ( $c = 11.6 \text{ cm}^{-1}$ ) either by modifying the buffer concentration or by addition of NaCl in the case of the glycine-NaOH buffer, pH 9.

### 2.4. DD-peptidase activity

Hydrolysis of the synthetic substrate  $N^\alpha, N^\epsilon$ -diacetyl-L-lysyl-D-alanyl-D-alanine was stopped by heating the sample at 100°C for 1 min and the amount of released D-alanine was measured by the D-amino acid oxidase procedure [10].

### 2.5. Model and equations for the interaction with $\beta$ -lactam compounds [11]

The results were analysed on the basis of the three-step model



where E denotes the enzyme, I the  $\beta$ -lactam, EI\* the acyl enzyme and K the dissociation constant of EI. If  $[I] \gg [E]$ , the rate of accumulation of EI\*,  $[EI^*]_t - [EI^*]_{ss}$  (where  $[EI^*]_t$  and  $[EI^*]_{ss}$  are the concentrations of EI\* at time  $t$  and at the steady state, respectively) is proportional to  $e^{-k_a t}$  where  $k_a = (k_{+3} + k_f)$  and  $k_f = (k_{+2}[I]) / (K + [I])$ .

### 2.6. Determination of the kinetic parameters

#### 2.6.1. Value of $k_{+3}$

After complete inactivation of the enzyme, excess  $\beta$ -lactam was eliminated by addition of  $\beta$ -lactamase. The solutions were further incubated and, after increasing time periods, samples were taken to which was added substrate for activity estimation.

#### 2.6.2. Value of $k_f$

With deacetyl-cephalosporin C,  $k_{+3}$  was small (see below) and, in all experiments,  $[EI^*]_{ss}$  remained negligible. The value of  $k_f$  at low  $[I]$  was computed by measuring the enzyme inactivation as a function of time. Faster reactions were monitored by measuring the quenching of the enzyme fluorescence at 320 nm upon formation of the acyl enzyme [11]. These experiments were performed using a Kontron SFM23 spectrofluorometer. Excitation was at 280 nm. A 20  $\mu$ l aliquot of  $\beta$ -lactam solution was rapidly mixed with 980  $\mu$ l enzyme solution (5.3  $\mu$ M) and the exponential decrease in fluorescence was analysed as in [1]. At pH 10.5, the decrease in fluorescence was too small to allow utilization of this method, and the enzyme was too unstable for monitoring the decrease in activity. It was only possible to obtain a rough approximation of  $k_f$  by following the decrease in absorbance at 260 nm after adding concentrated enzyme to a 10  $\mu$ M solution of deacetyl-cephalosporin C. The final enzyme concentration was also 10  $\mu$ M so that  $k_f$  could be computed by using the second-order rate law.

With the lactone, the fluorescence method was used, but  $k_{+3}$  was not always negligible when compared to  $k_a$  and  $k_f$  was obtained from  $k_a - k_{+3}$ . The instability of the lactone precluded any measurement above pH 10.

The susceptibility of the compounds to nucleophilic attack by  $\text{OH}^-$  was determined by following the decrease in absorbance of a 150  $\mu$ M solution in 1 M  $\text{K}_2\text{HPO}_4$  adjusted to pH 11.25 (for the lactone) or 12.25 (for deacetyl-cephalosporin C) with 10 M KOH [12].

## 3. RESULTS

### 3.1. Values of $k_{+3}$ (table 1)

Reactivation was markedly more rapid with the

Table 1  
pH dependence of  $k_{+3}$

pH	$10^5 \times k_{+3} \text{ (s}^{-1}\text{)}$	
	Deacetyl- cephalosporin C	Lactone
7.0	2 ± 0.4	31 ± 4
8.0	2.4 ± 0.5	53 ± 9
9.0	6.1 ± 0.9	310 ± 50
10.0	12 ± 2	550 ± 80

Deacetyl-cephalosporin C: enzyme (9.4  $\mu\text{M}$ ) and  $\beta$ -lactam (625  $\mu\text{M}$ ) were incubated at pH 7.0 in a total volume of 170  $\mu\text{l}$  of 1 mM sodium phosphate, pH 7.0. After 15 min, inactivation was complete and 10- $\mu\text{l}$  aliquots were 20-fold diluted in a suitable buffer containing 23  $\mu\text{g}$   $\beta$ -lactamase. The recovery of activity was measured on 10- $\mu\text{l}$  aliquots over a period of 60 min. Lactone: enzyme (0.44  $\mu\text{M}$ ) and the lactone (616  $\mu\text{M}$ ) were incubated for 10 min in 0.1 mM sodium phosphate pH 7.0, which resulted in complete inactivation. Samples (50  $\mu\text{l}$ ) were then supplemented with 150  $\mu\text{l}$  of suitable buffer containing 5  $\mu\text{g}$   $\beta$ -lactamase. The reactivation was measured on 10- $\mu\text{l}$  aliquots over a period of 10 (pH 9 and 10) or 30 min (pH 7 and 8)

lactone and, as observed before, the value of  $k_{+3}$  sharply increased at high pH [1]. This phenomenon has been shown to be due to direct attack of the acyl enzyme by  $\text{OH}^-$  [1].

### 3.2. Values of $k_f$

In a first series of experiments, only one  $\beta$ -

lactam concentration was used at each pH (table 2). The decrease in  $k_f/[I]$  observed at pH 10 appeared to be significantly larger with deacetyl-cephalosporin C than with the lactone ( $3.75 \pm 0.6$  vs  $1.7 \pm 0.25$ -fold decrease between pH 8.0 and 10.0). The value of  $k_f$  was then measured at various concentrations of  $\beta$ -lactams and plots of  $1/k_f$  vs  $1/[I]$  allowed the computation of the individual values of  $k_{+2}$  and  $K$  (fig.1), in the pH range 7–10 for deacetyl-cephalosporin C and at pH 7.0 for the lactone. With this latter compound, experiments were performed at concentrations ranging between 140 and 412  $\mu\text{M}$ . Plots of  $1/k_f$  vs  $1/[I]$  yielded the following results:  $k_{+2}/K = 194 \pm 17 \text{ M}^{-1} \cdot \text{s}^{-1}$ ,  $k_{+2} = 0.30 \pm 0.18 \text{ s}^{-1}$  and  $K = 1.5 \pm 0.9 \text{ mM}$ . The large errors in the individual values of  $k_{+2}$  and  $K$  were due to the fact that the lactone concentrations were too low when compared to  $K$ . However, it was not possible to use greater concentrations of the lactone, because the reaction became too rapid and reabsorption of the emitted radiation (in the fluorescence experiments) became too significant. Consequently, it seemed that, for the lactone, the ratio  $k_f/[I]$  provided a good approximation of  $k_{+2}/K$  at all pH values, since it was very unlikely that  $K$  would decrease at high pH values. At pH 10.5, a  $k_f$  value of  $1.2 \pm 0.3 \times 10^{-3} \text{ s}^{-1}$  was measured with deacetyl-cephalosporin C, indicating a  $k_2/K$  value of  $120 \pm 30 \text{ M}^{-1} \cdot \text{s}^{-1}$ . The individual values of  $k_2$  and  $K$  could not be obtained.

### 3.3. Hydrolysis by $\text{OH}^-$

At pH 12.25 and 30°C, the pseudo-first-order rate constant for the degradation of deacetyl-

Table 2  
pH dependence of  $k_f/[I]$

pH	Deacetyl-cephalosporin C			Lactone		
	[I] ( $\mu\text{M}$ )	$k_f$ ( $\text{s}^{-1}$ )	$k_f/[I]^a$ ( $\text{M}^{-1} \cdot \text{s}^{-1}$ )	[I] ( $\mu\text{M}$ )	$k_f$ ( $\text{s}^{-1}$ )	$k_f/[I]$ ( $\text{M}^{-1} \cdot \text{s}^{-1}$ )
7.0	30	0.035 ± 0.0012	1170 ± 40	412	0.074 ± 0.004	180 ± 10
8.0	30	0.0405 ± 0.0012	1350 ± 40	412	0.082 ± 0.004	200 ± 10
9.0	30	0.0237 ± 0.0012	790 ± 40	412	0.075 ± 0.004	183 ± 10
9.5	50	0.0250 ± 0.002	500 ± 40		N.D.	
10	50	0.018 ± 0.002	360 ± 40	412	0.048 ± 0.004	117 ± 10

<sup>a</sup> The  $\beta$ -lactam concentrations were not much smaller than the  $K$  values, so  $k_f/[I]$  was not equivalent to  $k_{+2}/K$





Fig.1. pH dependence of  $K$  (A),  $k_{+2}/K$  (B) and  $k_{+2}$  (C) for deacetyl-cephalosporin C.  $\beta$ -Lactam concentrations varied from 20 to 100  $\mu\text{M}$  (pH 7.0), 30 to 150  $\mu\text{M}$  (pH 8.0), 50 to 150  $\mu\text{M}$  (pH 9.0) and 50 to 500  $\mu\text{M}$  (pH 10). The enzyme concentration was 5.3  $\mu\text{M}$ . At pH 10.5, both enzyme and  $\beta$ -lactam were 10  $\mu\text{M}$  and only the value of  $K$  at that pH was computed assuming that  $k_{+2}$  remained constant ( $0.06 \text{ s}^{-1}$ ). The following simple model was used:



The apparent dissociation constant of the non-covalent complex is


$$K_{\text{app}} = (K_{\text{EH}^+}) \frac{[1 + \frac{K}{[\text{H}^+]}]}{[1 + \frac{KK_{\text{EH}^+}}{K_{\text{E}}[\text{H}^+]}]}$$

The solid lines were drawn on the basis of this equation with the following values for the constants:  $\text{p}K = 9.0$ ;  $K_{\text{EH}^+} = 25 \mu\text{M}$ ;  $K_{\text{E}} = 1250 \mu\text{M}$ . If one assumes that  $K_{\text{E}} = \infty$ , the dashed curve (A) is obtained. The corresponding curve is not shown in (B), since it is nearly indistinguishable from that displayed.


cephalosporin C was  $1.6 \pm 0.2 \times 10^{-3} \text{ s}^{-1}$ , which yielded a value of  $0.09 \pm 0.01 \pm 0.2 \text{ M}^{-1} \cdot \text{s}^{-1}$  for the second-order rate constant, in good agreement with the values obtained before [12] for similar compounds. The lactone was about 200-fold more unstable: at pH 11.25, the pseudo-first-order rate constant was  $34 \pm 3 \times 10^{-3} \text{ s}^{-1}$ , yielding a second-order rate constant of  $19 \pm 2 \text{ M}^{-1} \cdot \text{s}^{-1}$ .

4. DISCUSSION


The relative instability of the acyl enzyme formed upon inactivation of *Streptomyces* R61 DD-peptidase by the lactone seemed to indicate that the lactone c ring remained intact (structure III)



Indeed, hydrolysis of the lactone would have yielded the same adduct as that obtained with deacetyl-cephalosporin C (IV), since expulsion of the OH group of cephalosporins containing a  $\text{CH}_2\text{OH}$  substituent on  $\text{C}_3$  does not seem to occur upon formation of the acyl enzyme [13].



Alternatively, rearrangement with expulsion of the leaving group attached to the methylene group on C<sub>3</sub> would have yielded V.



In the first case, the value of  $k_{+3}$  for the lactone would have been the same as that observed with deacetyl-cephalosporin C and, in the second, it should have been even lower [13].

As demonstrated in fig.1, the decrease in  $k_f/[I]$  as the pH increased was due to an increased value of  $K$ . It could be hypothesized that charge pairing between an enzyme positive group (of  $pK \cong 9$ ) and the cephalosporin carboxylate enhanced the strength of the non-covalent interaction by a large factor, which could be estimated to be close to or greater than 50. Unfortunately, the instability of the enzyme precluded the obtaining of accurate data at pH 10.5 or above. With the lactone and at pH 7.0, the  $K$  value was  $60 \pm 36$ -fold larger than that observed with deacetyl-cephalosporin C under similar conditions and an increase in pH to 10 had a much less dramatic effect on the  $k_{+2}/K$  value. Indeed, the value of the second-order rate constant was  $60 \pm 10\%$  of that observed at pH 7.0. This slight decrease was probably due to a corresponding increase in  $K$  (a larger increase in  $K$  would imply a corresponding increase in  $k_{+2}$ , a very unlikely phenomenon). The pH dependence of  $K$  for the lactone might be due to the presence of a partial negative charge on the carbonyl group of the latter compound.

It is also interesting to note that the first-order rate constants for acylation ( $k_{+2}$ ) and deacylation ( $k_{+3}$ ) were greater with the lactone. These observations might be correlated with the fact that the lactone is also a better substrate of  $\beta$ -lactamase I from *B. cereus* [7]. The value of  $k_{+2}$  for the lactone was  $5 \pm 3$ -fold greater than that for deacetyl-cephalosporin C, while the susceptibility to nucleophilic attack of the former compound was about 200-fold larger than that of the latter. Since both compounds have the same substituent on C<sub>7</sub>, the large discrepancy must be attributed to a less productive location of the  $\beta$ -lactam carbonyl near the enzyme active serine, probably due to a poor electrostatic interaction or to steric factors such as a lack of free rotation around the C<sub>3</sub>-CH<sub>2</sub>O bond.

In conclusion, the present data indicate that charge pairing between the  $\beta$ -lactam carboxylate and an enzyme positive group of  $pK \cong 9$  contribute by a factor of at least 50 in the initial, non-covalent interaction between the two reagents.

#### ACKNOWLEDGEMENTS

This work has been supported in part by the Fonds de la Recherche Scientifique Médicale (FRSM, Brussels, contract no. 3.4507.83), an Action concertée with the Belgian Government (convention no.86/91-90), a convention with the Région wallonne (C2/C16/CONV.246/20428), the Fonds de Recherche de la Faculté de Médecine de l'Université de Liège and a contract with the EEC (BAP-0197-B). We wish to thank Dr S.G. Waley (University of Oxford) for his interest in this work.

#### REFERENCES

- [1] Varetto, L., Frère, J.M., Nguyen-Distèche, M., Ghuysen, J.M. and Houssier, C. (1987) *Eur. J. Biochem.* 162, 525-531.
- [2] Ambler, R.P. (1980) *Phil. Trans. R. Soc. Lond. B* 289, 321-331.
- [3] Dale, J.W., Godwin, D., Mossakowska, D., Stephenson, P. and Wall, S. (1985) *FEBS Lett.* 191, 39-44.
- [4] De Meester, F., Joris, B., Lenzini, M.V., Dehottay, P., Ercicum, T., Dusart, J., Klein, D., Ghuysen, J.M., Frère, J.M. and Van Beeumen, J. (1987) *Biochem. J.* 244, 427-432.
- [5] Duez, C., Piron-Fraipont, C., Joris, B., Dusart, J., Urdea, M.S., Martial, J.A., Frère, J.M. and Ghuysen, J.M. (1987) *Eur. J. Biochem.* 162, 509-518.
- [6] Jaszberenyi, J.C. and Gunda, T.E. (1975) *Prog. Med. Chem.* 12, 395-477.
- [7] Kubarawa, S. and Abraham, E.P. (1967) *Biochem. J.* 103, 27c-29c.
- [8] Frère, J.M., Ghuysen, J.M., Perkins, H.R. and Nieto, M. (1973) *Biochem. J.* 135, 463-468.
- [9] Thatcher, D.R. (1975) *Methods Enzymol.* 43, 653-664.
- [10] Frère, J.M., Leyh-Bouille, M., Ghuysen, J.M., Nieto, M. and Perkins, H.R. (1976) *Methods Enzymol.* 45B, 610-636.
- [11] Frère, J.M., Ghuysen, J.M. and Iwatsubo, M. (1975) *Eur. J. Biochem.* 57, 343-351.
- [12] Frère, J.M., Kelly, J.A., Klein, D., Ghuysen, J.M., Claes, P. and Vanderhaeghe, H. (1982) *Biochem. J.* 203, 223-234.
- [13] Faraci, W.S. and Pratt, R.F. (1986) *Biochem. J.* 238, 309-312.