

Myosin light chain 2 modulates calcium-sensitive cross-bridge transitions in vertebrate skeletal muscle

Joseph M. Metzger and Richard L. Moss

Department of Physiology, School of Medicine, University of Wisconsin, Madison, Wisconsin 53706 USA

ABSTRACT We investigated the mechanism of the Ca^{2+} sensitivity of cross-bridge transitions that limit the rate of force development in vertebrate skeletal muscle. The rate of force development increases with Ca^{2+} concentration in the physiological range. We show here that at low concentrations of Ca^{2+} the rate of force development increases after partial extraction of the 20-kD light chain 2 subunit of myosin, whereas reconstitution with light chain 2 fully restores native sensitivity to Ca^{2+} in skinned single skeletal fibers. Furthermore, elevated free Mg^{2+} concentration reduces Ca^{2+} sensitivity, an effect that is reversed by extraction of the light chain but not by disruption of thin-filament activation by partial removal of troponin C, the Ca^{2+} binding protein of the thin filament. Our findings indicate that the Ca^{2+} sensitivity of the rate of force development in vertebrate skeletal muscle is mediated in part by the light chain 2 subunit of the myosin cross-bridge.

INTRODUCTION

Force development and shortening in muscle result from interaction of the contractile proteins myosin and actin. In all muscle cells, the extent of interaction between actin and myosin is regulated by the concentration of cytosolic Ca^{2+} , although the specific proteins that confer Ca^{2+} sensitivity differ among muscle types. For example, Ca^{2+} regulation of contraction in vertebrate striated muscle is mediated by troponin and tropomyosin associated with the thin filament (1), whereas striated muscles of various invertebrate species are regulated by Ca^{2+} binding directly to myosin (2–5).

The molecular basis of Ca^{2+} activation of contraction in myosin-regulated muscle is not well understood. The Ca^{2+} specific binding site responsible for activation of myosin-regulated muscle is located in the globular head region of myosin. This binding site is lost on removal of the light chain 2 (LC_2) protein subunit of myosin; however, LC_2 itself does not contain the Ca^{2+} specific binding site (4, 5). Recent evidence suggests that the Ca^{2+} specific binding site of myosin involves a three-peptide domain that consists of the regulatory LC (i.e., LC_2), essential LC, and a 10-kD myosin heavy chain fragment (6). It is possible that the Ca^{2+} specific binding site may be defined by a pocket formed by these peptides. In addition, it is known that myosin LC_2 exhibits significant primary sequence homology with Ca^{2+} binding proteins troponin C, parvalbumin, and calmodulin (7–9). Indeed, a high-affinity $\text{Ca}^{2+}/\text{Mg}^{2+}$ binding site has been localized to the N-terminal region of LC_2 (for review see reference 5). The role of this divalent cation binding site in Ca^{2+} activated contraction of myosin-regulated muscle is unclear. Interestingly, LC_2 is present in the myosin molecule of thin filament-regulated muscle. Myosin LC_2 of vertebrate skeletal muscle also contains the high-affin-

ity $\text{Ca}^{2+}/\text{Mg}^{2+}$ binding site, although its possible function in the Ca^{2+} sensitivity of contraction has not been established (4). However, in myosin-regulated muscle the inhibitory effect of LC_2 was lost in LC_2 extracted fibers that were reconstituted with a mutant form of vertebrate LC_2 that lacked a functional $\text{Ca}^{2+}/\text{Mg}^{2+}$ binding site (10). This suggests that the high-affinity binding site of LC_2 is somehow important in regulating the interaction of myosin with actin in these fibers, but whether a similar role exists in vertebrate muscle has not been established.

The idea that vertebrate skeletal muscle contains some form of myosin-based regulation of contraction has been inferred from various studies (11, 12). The applicability of these findings to intact muscle is uncertain because the studies typically used isolated proteins, which eliminated some factors that influence contraction *in vivo*, such as structural constraints on cross-bridge interaction imposed by the intact filament lattice. Thus, although the results of the earlier studies are convincing for the conditions used, there is as yet no physiological evidence for a myosin component of Ca^{2+} regulation of contraction in vertebrate striated muscle.

Here, we investigated the basis of the Ca^{2+} sensitivity of weak to strong cross-bridge transitions in mammalian skeletal muscle (13–15). Experiments were designed to determine whether myosin LC_2 is involved in conferring Ca^{2+} sensitivity to cross-bridge transitions. Single fibers from rabbit and rat fast-twitch skeletal muscles were treated chemically to permeabilize surface membranes, thereby allowing direct control of intracellular solutions and permitting specific extraction and readdition of contractile and regulatory protein subunits. Results showed that during submaximal Ca^{2+} activation LC_2 is repressive to cross-bridge transitions and that repression is relieved by increasing the concentration of Ca^{2+} or by removal of endogenous LC_2 . Furthermore, increased Mg^{2+} concentration decreased Ca^{2+} sensitivity, an effect

Address correspondence to Dr. Joseph M. Metzger, Department of Physiology, School of Medicine, University of Michigan, 7730 Medical Science II, Ann Arbor, Michigan 48109-0622.




FIGURE 1 (A) Records of sarcomere length (*upper traces*) and tension (*lower traces*) obtained in the determination of k_{tr} during maximal activation (pCa 4.5) of a psoas fiber before and after extraction of 25% LC₂. k_{tr} was 20 s⁻¹ both before and after partial extraction of LC₂. Sarcomere length was clamped to 2.44 μm during tension redevelopment. P_0 was 128 kN/m². To extract LC₂, the fiber was placed in a solution containing, in mM, 20 KCl, 20 EDTA, 5 imidazole, pH 7.00, for 3 h at 30°C. hs, half sarcomere. All tension traces are normalized to the maximum tension generated in each fiber under the particular experimental conditions (i.e., control, extracted; high Ca²⁺, low Ca²⁺, etc.). (B) Records of tension obtained during the protocol to determine k_{tr} in a psoas fiber during submaximal Ca²⁺ activation both before and after partial extraction of LC₂. k_{tr} was 3 s⁻¹ before extraction of LC₂ and 10 s⁻¹ after extraction. Sarcomere length was clamped as in A (records not shown). Results were obtained from the same fiber as in A.

that was reversed by extraction of LC₂. Our findings suggest a role for LC₂ in modulating Ca²⁺ sensitivity of contraction in mammalian skeletal muscle.

MATERIALS AND METHODS

Skinned fiber preparations and experimental apparatus

Fast-twitch skeletal muscle fibers were obtained from the superficial portion of the vastus lateralis (svl) muscles of adult female Sprague-

Dawley rats and from psoas muscles of adult male New Zealand rabbits. Results obtained using psoas and svl fibers were qualitatively similar. Bundles of ~50 fibers were dissected from each muscle while in relaxing solution (below) and were then tied with surgical silk to glass capillary tubes. Bundles were stored for up to 3 wk at -23°C in relaxing solution containing 50% (vol/vol) glycerol. Before each experiment, bundles were placed in relaxing solution containing 0.5% (wt/vol) Brij-58 for 30 min to disrupt the sarcoplasmic reticulum (16). Individual fibers were carefully pulled free from one end of the fiber bundle and mounted between a force transducer (model 407; Cambridge Technology, Inc., Cambridge, Massachusetts; sensitivity, 0.2 mV/μN; 1-99% response time, 100 μs; resonant frequency, ~5 kHz; noise level at the output equivalent to 1 mg peak-to-peak) and DC torque motor (model 300s; Cambridge Technology, Inc.). The fiber was viewed through an inverted microscope (model IM; Carl Zeiss, Inc., Thornwood, New York), and its overall length was adjusted with a mechanical translator to set resting sarcomere length. Complete details of the mounting procedure and experimental set-up have been reported elsewhere (16, 17).

Solutions

Relaxing and activating solutions contained (mM) 7 ethyleneglycol-bis(β-aminoethyl ether)-N,N'-tetraacetic acid, 1 or 10 free Mg²⁺, 4.4 total ATP, 14.5 creatine phosphate, 20 imidazole, and sufficient KCl to yield a total ionic strength of 180 mM. Solution pH was adjusted to 7 with KOH. Relaxing solution had a pCa (i.e., -log [Ca²⁺]) of 9.0, whereas the pCa of the solution for maximal activation was 4.5. The computer program of Fabiato and Fabiato (18) was used to calculate the final concentrations of each metal, ligand, and metal-ligand complex, using the stability constants listed by Godt and Lindley (19). The apparent stability constant for Ca²⁺-EGTA was corrected for ionic strength, pH, and an experimental temperature of 15°C (18).

Rate constant of tension redevelopment and sarcomere length control system

The experimental protocol for measuring the rate constant of tension redevelopment (k_{tr}) was a modification of the multistep protocol developed by Brenner and Eisenberg (20). Measurement of k_{tr} involved a mechanical maneuver to dissociate myosin cross-bridges from actin in a steadily activated fiber so that the subsequent rate of tension redevelopment reflects the forward and reverse rate constants for the rate limiting transition(s) in the cross-bridge cycle leading to formation of strongly bound force-generating states. The fiber was first transferred from relaxing solution to an activating solution, and steady isometric tension was developed. The fiber was then rapidly (within 0.5 ms) shortened by ~200-300 nm per half-sarcomere, resulting in an abrupt reduction of force to zero, and the fiber shortened for 5-40 ms under unloaded conditions (i.e., at maximum velocity). Before the imposed slack was taken up, the fiber was rapidly reextended to its initial length. Coincident with the restretch, force transiently increased and then rapidly declined to zero or very nearly zero. The redevelopment of force after this maneuver reflects the rate of reattachment of cross-bridges and the transition to strongly bound force-producing states. During redevelopment of force, sarcomere length was held constant, because in the absence of sarcomere length control k_{tr} would be underestimated due to end compliance (20). Sarcomere length was clamped to within 0.5 nm per half-sarcomere by servocontrol of the position of the first order line of the laser diffraction pattern (Fig. 1). Records of tension redevelopment were best fit by a single exponential equation. Complete details of the experimental protocol, curve fitting procedure, mechanical set-up, and sarcomere length control system are described elsewhere (17).

Tension-pCa relationship

At each pCa, steady isometric tension was allowed to develop, after which the fiber was slackened to obtain the tension baseline. The fiber




FIGURE 2 SDS-polyacrylamide gels and densitometric scans of gels obtained from segments of a psoas fiber before and after the procedure to extract LC₂. To quantitate the amount of LC₂ extracted, the ratio LC₂/(LC₁ + LC₃) was determined for both control and extracted fibers by measuring the areas under the peaks corresponding to these proteins. The extracted ratio was then divided by the control ratio to determine the reduction in LC₂ content. Calculated in this way, 61% of LC₂ was extracted from this fiber, which was the largest amount removed in this study. Importantly, exogenous troponin C was first added back to the fiber before mechanical measurements were made because the LC₂ extraction procedure also removes troponin C. Comparison of troponin C/(LC₁ + LC₃) ratios from control and extracted segments showed that readdition of troponin C to extracted fiber was stoichiometric. TnC, troponin C; MHC, myosin heavy chain.

was then relaxed. The difference between steady tension and the tension baseline after the slack step was measured as total tension. To obtain active tension, resting tension measured at pCa 9.0 (~1% of total tension) was subtracted from total tension. The fiber was transferred to relaxing solution after each activation at a given pCa. Tension-pCa relations were determined in each fiber by expressing tensions (P) at various submaximal Ca²⁺ concentrations as fractions of the maximum value, P_0 (i.e., isometric tension at pCa 4.5 and 1 mM free Mg²⁺) obtained in each fiber. Every fourth contraction was at pCa 4.5 to monitor any decline in fiber performance (16).

Protocols for extraction and readdition of LC₂

Methods to extract LC₂ from skinned skeletal muscle fibers have been published previously and briefly described here (21). To extract LC₂, the fiber was transferred from relaxing solution to a solution containing (mM): 20 KCl, 20 ethylenediaminetetraacetate (EDTA), 5 imidazole,

pH 7.00, for 1–4 h at 20–30°C. After extraction, the fiber was transferred to relaxing solution and temperature was returned to 15°C. Because there was a variable amount of extraction of troponin C during this procedure, exogenous troponin C was added back to the fibers before data collection. In this study, maximum isometric tension averaged $0.98 \pm 0.02 P_0$ after extraction of LC₂ and add back of troponin C.

Readdition of exogenous LC₂ involved transferring the fiber to a relaxing solution containing 0.5 mg/ml purified LC₂ for 2–3 h at 15°C.

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE)

Before attachment of the fiber to the experimental apparatus, a 0.5–1-mm segment of the fiber was removed for analysis of the control protein composition by SDS-PAGE. The remaining portion of the fiber, ~2–3 mm in length, was mounted into connectors (16). Contractile properties were determined in the control fiber and after extraction of




FIGURE 3 Summaries of the effects of partial extraction of LC₂ on the k_{tr} -pCa (upper), isometric tension-pCa (middle), and k_{tr} -isometric tension (lower) relationships in rabbit psoas fibers. Maximum isometric tension averaged $0.98 \pm 0.02 P_0$ after extraction of LC₂ and add back of troponin C (see Fig. 2). Values are mean \pm SE with 6–10 observations per point. In some instances, error bars were smaller than the symbol size. Asterisks indicate experimental values that were significantly greater than control ($P < 0.05$). Qualitatively similar results were obtained from svl fibers.

LC₂. After extraction, a 0.5–1-mm segment of the fiber was removed for protein analysis. Fibers were subsequently reconstituted by bathing in relaxing solution containing purified LC₂. Mechanical measurements were obtained after readdition of LC₂. Each fiber segment was placed in a 0.5-ml microfuge tube containing SDS sample buffer (10 μ l/mm length of fiber segment) and stored at -80°C for subsequent analysis of contractile and regulatory protein content by SDS-PAGE

and scanning densitometry, as described previously (21, 22). Thus, gels obtained from segments of the same fiber were analyzed to quantitate protein composition at the different stages of an experiment (Fig. 2). Gels of LC₂ reconstituted fibers have been published (22).

RESULTS AND DISCUSSION

To probe cross-bridge transitions that limit the rate of formation of the strongly bound force-bearing state, we used a mechanical maneuver to determine the rate constant of tension redevelopment (k_{tr}) after rapid release and reextension of fiber length during steady Ca²⁺ activation (13–15, 20) (Fig. 1). k_{tr} is sensitive to Ca²⁺ in the physiological range (i.e., 10^{-6} – 10^{-4} M) and is thought to be determined by the step or steps in the actin-myosin ATP hydrolysis reaction that limit the rate of formation of the strongly bound force-generating cross-bridge state (13–15). In earlier studies, the Ca²⁺ sensitivity of k_{tr} was unchanged when thin filament activation was disrupted by partial extraction of troponin C, whereas the Ca²⁺ sensitivity of steady-state isometric tension was markedly reduced (23). Partial extraction of troponin C results in a rightward shift (i.e., to higher [Ca²⁺]) and reduction in the steepness of the tension-pCa relationship due to a disruption of near-neighbor cooperativity between adjacent functional groups (defined structurally as 1 tropomyosin, 1 troponin complex, and 7 actin monomers) on the thin filament (24, 25). This dissociation of Ca²⁺ sensitivities of k_{tr} and isometric tension provides evidence that k_{tr} is significantly affected by an apparent rate constant that characterizes the kinetics of transitions




FIGURE 4 Records of tension obtained during determination of k_{tr} at a submaximal concentration of Ca²⁺ in a psoas fiber before and after extraction of LC₂ and after the readdition of exogenous LC₂. k_{tr} was 4.8 s⁻¹ in control, 6.5 s⁻¹ after partial extraction of LC₂, and 4.4 s⁻¹ after the readdition of purified LC₂ into the fiber. In each trial, sarcomere length was clamped to 2.58 μm during tension redevelopment (records not shown). P_0 was 131 kN/m². After extraction of LC₂ and readdition of troponin C, isometric tension at pCa 4.5 was 1.01 P_0 . The fiber was placed in LC₂ extracting solution (Fig. 1) for 2.5 h at 20°C. To reconstitute the extracted fiber with LC₂, the fiber was bathed for 3.5 h at 15°C in relaxing solution containing 0.58 mg/ml purified LC₂. SDS-PAGE gels of segments obtained from a single fiber before and after reconstitution with LC₂ have been published (22).




FIGURE 5 Effects on the Ca^{2+} sensitivity of k_{tr} due to elevated $[\text{Mg}^{2+}]$ in control and partially LC_2 extracted fibers. Results in *B–D* were obtained from the same fiber. *A* and *B* are controls, *C* and *D* are after partial extraction of LC_2 . (*A*) Records of tension obtained during determination of k_{tr} at maximal concentration of Ca^{2+} (pCa 4.5) at pMg 3 and 2 in a svl fiber. k_{tr} was 20 s^{-1} at pMg 3 and 19.5 s^{-1} at pMg 2. Qualitatively similar results were obtained from psoas fibers. $P_0 = 115 \text{ kN/m}^2$. (*B*) Records of tension obtained during determination of k_{tr} at a submaximal concentration of Ca^{2+} (pCa 5.6) at pMg 3 and 2 in a svl fiber. k_{tr} was 17 s^{-1} at pMg 3 and 2.8 s^{-1} at pMg 2. Relative tensions were $0.84 P_0$ at pMg 3 and $0.33 P_0$ at pMg 2. k_{tr} at pCa 4.5 was 23 s^{-1} . Average pCa₅₀'s from k_{tr} -pCa and tension-pCa data were 5.80 and 6.03 at pMg 3 and 5.46 and 5.55 at pMg 2, respectively. Qualitatively similar results were obtained from psoas fibers. (*C*) Records of tension obtained during determination of k_{tr} at pCa 5.6 and pMg 3 and 2 after the partial extraction of LC_2 from the same fiber as in *B*. k_{tr} was 17 s^{-1} at pMg 2 and pMg 3. (*D*) Records of tension obtained during determination of k_{tr} at pCa 6.0 and pMg 3 and 2 after partial extraction of LC_2 from the same fiber as in *B* and *C*. k_{tr} was 5 s^{-1} at pMg 3 and 4.5 s^{-1} at pMg 2 (noisy trace). For all records, sarcomere length (records not shown) was clamped during tension redevelopment. LC_2 extraction solution and protocol were as described in Fig. 1.

among cross-bridge states. Furthermore, k_{tr} increases in the presence of added phosphate, suggesting that the specific cross-bridge transition(s) that underlies k_{tr} , at least in part, includes the phosphate release step of the actomyosin ATP hydrolysis reaction and/or a step in rapid equilibrium with phosphate release (26, 27). Based on analysis of tension transients resulting from photogeneration of phosphate from caged phosphate, it has been proposed that phosphate release from actomyosin is a two-step process involving an isomerization step followed by the phosphate release step (28, 29).

Because LC_2 is essential in conferring Ca^{2+} sensitivity to thick filament regulated muscles (2–5), the Ca^{2+} sensitivity of k_{tr} was examined after partial extraction of myosin LC_2 from skeletal muscle fibers (Fig. 1). Partial extraction of LC_2 had no effect on k_{tr} at maximal $[\text{Ca}^{2+}]$, but k_{tr} increased markedly at submaximal $[\text{Ca}^{2+}]$ (Fig. 1).

In the example shown, there was an approximate three-fold increase in k_{tr} at pCa 6.0 (where pCa is $-\log[\text{Ca}^{2+}]$) after extraction of 25% of LC_2 , determined from SDS-polyacrylamide gels of segments of the same fiber obtained before and after extraction. Densitometric scans of SDS gels of these fiber segments showed that extraction also removed troponin C, but other contractile and regulatory proteins were unchanged by the protocol. It is important to note that before experimental measurements fibers were reconstituted with troponin C (see Fig. 2).

Fig. 3 summarizes the effects of partial extraction of LC_2 on the relationships between k_{tr} and pCa, steady-state tension and pCa, and k_{tr} and steady-state tension. At pCa 4.5, k_{tr} averaged $16.7 \pm 0.7 \text{ s}^{-1}$ ($n = 10$) in control fibers and $17.1 \pm 0.8 \text{ s}^{-1}$ ($n = 10$) after extraction of an average $28 \pm 7\%$ of endogenous LC_2 . We have inter-




FIGURE 6 Summary of effects on the k_{tr} -tension relationship (data obtained from Fig. 5) due to altered $[Mg^{2+}]$ before (*Control*) and after partial extraction of LC₂ (*Extracted*). ●, pMg 3; ○, pMg 2. Tension and k_{tr} values are scaled to values obtained at pCa 4.5. For the control and extracted data, the pMg 2 values (○ at point 1.0, 1.0 in both plots) were partially covered by the pMg 3 data point (●). For the extracted data, the line marked *a* connects data points obtained at pCa 5.6; at *b* the line connects data points obtained at pCa 5.7; at *c* the line connects data points obtained at pCa 6.0. Qualitatively similar findings obtained in both psoas ($n = 2$) and svl fibers ($n = 5$).

interpreted our results (below) based on the assumption that the extraction of LC₂ is random along the thick filament. This assumption is based on the observation that the extraction of troponin C is random along the length of the thin filament (30). Isometric tension and k_{tr} values at pCa 4.5 were not significantly different than control values (Fig. 3), and this is taken as evidence that the extraction procedure did not adversely affect contractile function. At pCa above 5.0, k_{tr} increased significantly after extraction of LC₂. The pCa required for half maximal k_{tr} , that is pCa₅₀, increased from an average of 5.71 in control fibers to 6.07 after partial extraction of LC₂, indicating that the sensitivity of k_{tr} to Ca²⁺ increased due to extraction of the light chain. For example, at pCa 6.0, k_{tr} was $3.8 \pm 0.7 \text{ s}^{-1}$ ($n = 9$) in controls and $9.3 \pm 1.0 \text{ s}^{-1}$ ($n = 9$) after partial extraction of LC₂.

An additional effect of LC₂ extraction was an increase in the minimum value of k_{tr} obtained at low $[Ca^{2+}]$. This suggests that partial removal of LC₂ altered the fundamental (i.e., apparent Ca²⁺ independent) value of an apparent rate constant, as well as affecting the sensitivity of the rate constant to Ca²⁺. In agreement with previous work (21), partial extraction of LC₂ increased the Ca²⁺ sensitivity of steady-state tension, with pCa₅₀ increasing from 6.10 in control fibers to 6.25 after extraction.

Taking these results together, partial extraction of LC₂ resulted in a left-shift of the relationship between k_{tr} and steady-state tension (Fig. 3). Thus, the effects on k_{tr} subsequent to extraction of LC₂ are not simply due to enhanced activation of the thin filament as a result of increased cross-bridge binding (21) but rather indicate that LC₂ has a direct effect to modulate k_{tr} . For example, our earlier study showed that tension and instantaneous stiffness were increased at submaximal concentrations of Ca²⁺ after partial extraction of LC₂ (21). This suggested that one effect of removal of LC₂ was to increase the activation of the thin filament at submaximal concentra-

tions of Ca²⁺. If the effects of LC₂ extraction on k_{tr} were solely related to the increase in thin filament activation, then the k_{tr} -relative tension relationship should be unaffected by extraction. However, we found the k_{tr} -tension relationship to be altered by extraction, which we take to indicate that removal of LC₂ additionally affects cross-bridge kinetics.

That the effects on k_{tr} are specific to the extraction of LC₂ is clearly demonstrated in experiments in which LC₂ readdition to extracted fibers resulted in complete restoration of native Ca²⁺ sensitivity of k_{tr} (Fig. 4). Similar findings were obtained in five separate experiments and provide strong evidence that the observed effects on k_{tr} can be directly attributed to LC₂ rather than a nonspecific effect of the extraction procedure. It should be noted here that these effects of LC₂ are not dependent on phosphorylation of the light chain, because as we showed earlier the phosphate content in these fibers is low (<0.1 mol phosphate/mol LC₂). Effects on k_{tr} and tension due to alterations in phosphorylation of LC₂ have been reported previously (17).

If altered Ca²⁺ sensitivity of k_{tr} is directly related to the extraction of LC₂, then there should be a graded effect of extraction on k_{tr} such that at low amounts of extraction the effect should be small, whereas with greater amounts of extraction the alteration in the Ca²⁺ sensitivity of k_{tr} should correspondingly be increased. To address this point, we examined the relationship between the Ca²⁺ sensitivity of k_{tr} , determined as the ratio of k_{tr} values (extracted/control; $k_{tr,ext/con}$) at a particular pCa, and the amount of LC₂ extracted determined from the same fiber by SDS-PAGE. The data were fit by a straight line using least-squares fit linear regression to obtain the slope of the relationship between $k_{tr,ext/con}$ and percent of LC₂ extracted. In this analysis, the slope was 0.07 and the correlation coefficient was 0.83. Thus, for each 1% of LC₂ extracted, the $k_{tr,ext/con}$ value increased by 7%. This indicates that the Ca²⁺ sensitivity of k_{tr} relates to LC₂ extraction in a graded manner; however, caution must be taken in interpreting the magnitude of this effect because the value of $k_{tr,ext/con}$ is highly dependent on the pCa value used in the comparison. For example, at low pCa values that yield near maximal values of k_{tr} , there would be relatively little or no effect of LC₂ extraction on

TABLE 1 Effects on k_{tr} due to partial extraction of troponin C

	pMg 3		pMg 2	
	Control	Extracted	Control	Extracted
k_{tr} (s ⁻¹)	11	11	3.5	3.5
P/P_0	0.89	0.30	0.46	0.11

All measurements were made at pCa 5.6. After partial extraction of troponin C, isometric tension at pCa 4.5 was $0.54 P_0$. The solution and procedure for extracting troponin C have been published (23, 36). Results were obtained from an svl fiber, with sarcomere length servocontrolled to $2.63 \mu\text{m}$ during measurement of k_{tr} .

the $k_{tr,ext/con}$ value (see Fig. 3). At higher pCa values, the $k_{tr,ext/con}$ value increases so that comparison among fibers with differing extents of LC₂ extraction is complex and involves two factors: the extent of LC₂ extracted and the pCa value used in determining the $k_{tr,ext/con}$ value. This complicating feature was not present in our earlier work that related extent of troponin C extraction to altered contractile function because we were able to use tension at maximal Ca²⁺ concentration as a general marker for extent of protein extraction (25).

LC₂ of vertebrate striated muscle belongs to a large family of Ca²⁺ binding proteins and has considerable sequence homology with the regulatory proteins calmodulin, troponin C, and scallop LC₂ (7–9). Vertebrate skeletal LC₂ has a high affinity divalent cation binding site that preferentially binds Ca²⁺ over Mg²⁺ (K_{Ca} 3×10^7 M⁻¹, K_{Mg} $2.5\text{--}3 \times 10^5$ M⁻¹) (31). However, under resting conditions this site would be occupied predominantly by Mg²⁺ because the concentration of Mg²⁺ is about four orders of magnitude greater than Ca²⁺ (31, 32). Also, due to the slow dissociation rate of Mg²⁺, this site is not usually thought to be involved in regulation of contraction, because Ca²⁺ would not significantly displace Mg²⁺ during the time course of a twitch (32). However, under conditions that raise the mean Ca²⁺ concentration for prolonged times, such as during successive twitches or tetany, Ca²⁺ binding to LC₂ may be increased to a value of ~ 0.70 mol Ca²⁺/mol myosin in the presence of physiological levels of Mg²⁺ and pH 7.00 (31). Presumably, during the protocol to determine k_{tr} in skinned fibers, Ca²⁺ is in equilibrium with the cation binding site of LC₂ because several seconds are required to achieve steady activation of the fibers. If the Ca²⁺ sensitivity of k_{tr} involves this site, increased [Mg²⁺] should reduce bound Ca²⁺ and decrease k_{tr} . Fig. 5 shows results of the effects on k_{tr} due to increasing free Mg²⁺ from 1 mM, which is near physiological, to 10 mM. Increases in free [Mg²⁺] had no effect on k_{tr} at pCa 4.5; however, the Ca²⁺ sensitivity of k_{tr} was markedly reduced in that pCa₅₀ decreased by 0.34 pCa units at pMg 2 compared with pMg 3. We next examined the possible role of LC₂ in mediating the effect of Mg²⁺ on k_{tr} . In fibers treated to partially extract LC₂, effects on k_{tr} due to increased [Mg²⁺] were reversed (Figs. 5 and 6). This suggests that LC₂ is critical in conferring Mg²⁺ sensitivity to the modulation of k_{tr} by Ca²⁺. This effect could involve the Ca²⁺-Mg²⁺ site of LC₂, although other mechanisms involving LC₂ (see below) are not excluded.

The possibility that the effects on Ca²⁺ sensitivity of k_{tr} due to increased Mg²⁺ are mediated by thin filament regulatory proteins also has been considered. It is known that Mg²⁺ depresses the Ca²⁺ sensitivity of steady-state tension (33), thought to be due to an effect of Mg²⁺ to reduce Ca²⁺ binding to the low-affinity sites of troponin C (34). Results in Fig. 6 show that in control fibers the k_{tr} -relative tension relationships are similar at pMg 3 and 2. This might be expected if the Ca²⁺ sensitivity of k_{tr}

is due to variations in thin filament activation when the concentrations of Ca²⁺ and Mg²⁺ are altered. However, after partial extraction of LC₂, high Mg²⁺ depressed tension to a disproportionately greater extent than k_{tr} (Fig. 6). Similar results were observed in a total of seven experiments. These findings argue against the idea that thin filament activation is the sole determinant of k_{tr} , but instead suggest that the effect of increased Mg²⁺ to reduce the Ca²⁺ sensitivity of k_{tr} is mediated by LC₂. It is possible, however, that effects of Ca²⁺ and Mg²⁺ on k_{tr} also involve the thin filament regulatory system, but in this case the expression of the thin filament component appears to depend on the presence of LC₂ on the myosin molecule.

The marked reduction in the effect of high Mg²⁺ concentration to alter the Ca²⁺ sensitivity of k_{tr} in LC₂-extracted fibers is at first glance surprising, because in our experiments only $\sim 30\%$ of total LC₂ was extracted. However, previous work suggests that for any given myosin molecule one LC₂ can be removed relatively easily with EDTA, but removal of the second LC₂ is more difficult and requires harsher conditions (35). Thus, in our experiments, an average 30% reduction in LC₂ content could indicate that as much as 60% of myosin molecules have lost an LC₂ subunit.

As a control, we investigated the possibility that troponin C mediates the effects of Mg²⁺ on k_{tr} . Partial removal of troponin C resulted in reduced Ca²⁺ sensitivity of tension due to the disruption of near-neighbor molecular cooperativity within the thin filament. In partially troponin C-extracted fibers, the effect of Mg²⁺ to depress the Ca²⁺ sensitivity of steady-state isometric tension remained, i.e., effects on tension due to troponin C extraction and to high [Mg²⁺] were additive (Table 1). In contrast, at both 1 and 10 mM Mg²⁺, the Ca²⁺ sensitivity of k_{tr} was unchanged due to the partial extraction of troponin C. These findings are consistent with the idea that LC₂ mediates the effect of altered [Mg²⁺] on the k_{tr} -pCa relationship.

The detailed mechanism of LC₂ modulation of the Ca²⁺ sensitivity of k_{tr} is not clear. There is evidence to indicate that binding of divalent metals to the high-affinity Ca²⁺/Mg²⁺ site of LC₂ induces a conformational change in myosin. In the presence of Ca²⁺, there is a decrease in the Stokes radius of LC₂ in solution (37), providing evidence of a marked alteration in the tertiary structure of this protein due to Ca²⁺ binding. Furthermore, a conformational change in myosin due to Ca²⁺ binding is suggested in experiments where Ca²⁺, but not Mg²⁺, altered the sedimentation coefficient and viscosity of isolated native and synthetic myosin filaments in solution (11). It is possible that a conformational change on Ca²⁺ binding to LC₂ underlies modulation of a step or steps that limit the rate of formation of the strongly bound force-bearing cross-bridge state. Myosin LC₂ is noncovalently bound to the globular head of myosin near the head-tail junction or "hinge" region (38). A po-

tential, although speculative, consequence of Ca^{2+} binding to LC_2 could be altered flexibility of the hinge region, an effect that may underlie modified cross-bridge kinetics. Another possibility is that the thin filament regulatory system mediates the effect of Ca^{2+} on k_{tr} , but the inhibition of cross-bridge kinetics by the thin filament requires that LC_2 be present on myosin so that the cross-bridge somehow senses or modifies the state of thin filament activation. Thus, after removal of LC_2 , the inhibition of cross-bridge kinetics due to the thin filament is removed and the rate of the force generating transition is disinhibited. Consistent with this idea, Wagner (39) showed in solution biochemical studies that LC_2 was required for Ca^{2+} sensitive binding of myosin to regulated thin filaments. However, such a mechanism does not easily explain our finding that in troponin C-extracted, but LC_2 -replete fibers, k_{tr} was unchanged even though thin filament activation, which was evident as a marked reduction in the Ca^{2+} sensitivity of tension, was substantially decreased.

In conclusion, based on our findings we propose that LC_2 represses Ca^{2+} -sensitive weak to strong transitions of cross-bridges and that this repression can be relieved by added Ca^{2+} , reduced Mg^{2+} , or removal of LC_2 . In this respect, vertebrate striated muscle may be somewhat similar to thick-filament regulated systems where removal of LC_2 or addition of Ca^{2+} activates myosin-actin interactions (2–5). Our working hypothesis is that in vertebrate skeletal muscle there are at least two effects of Ca^{2+} : regulation of cross-bridge formation mediated by Ca^{2+} binding to thin filament regulatory proteins and modulation of the rate of formation of force-bearing cross-bridges somehow involving LC_2 . Ca^{2+} modulation of the kinetics of force development that are mediated by LC_2 could provide a basis for frequency-dependent potentiation of twitch tension in living muscle.

We thank Dr. J. Graham for SDS-PAGE analysis of fiber segments.

This study was supported by a grant from the National Institutes of Health.

Received for publication 21 June 1991 and in final form 19 March 1992.

REFERENCES

1. Ebashi, E., and M. Endo. 1968. Calcium ion and muscle contraction. *Prog. Biophys. Mol. Biol.* 18:123–183.
2. Szent-Györgyi, A. G., E. M. Szentkiralyi, and J. Kendrick-Jones. 1973. The light chains of scallop myosin as regulatory subunits. *J. Mol. Biol.* 74:179–203.
3. Kendrick-Jones, J., and J. M. Scholey. 1981. Myosin-linked regulatory systems. *J. Muscle Res. Cell Motil.* 2:347–372.
4. Bagshaw, C. R. 1980. Divalent metal ion binding and subunit interactions in myosin: a critical review. *J. Muscle Res. Cell Motil.* 1:255–277.
5. Bagshaw, C. R., and J. Kendrick-Jones. 1979. Characterization of homologous divalent metal ion binding sites of vertebrate and molluscan myosins using electronic paramagnetic resonance spectroscopy. *J. Mol. Biol.* 130:317–336.
6. Kwon, H., E. B. Goodwin, L. Nyitray, E. Berliner, E. O'Neill-Hennessey, F. D. Melandri, and A. G. Szent-Györgyi. 1990. Isolation of the regulatory domain of scallop myosin: role of the essential light chain in calcium binding. *Proc. Natl. Acad. Sci. USA* 87:4771–4775.
7. Kretsinger, R. H. 1980. Structure and evolution of calcium-modulated proteins. *CRC Crit. Rev. Biochem.* 8:119–174.
8. Collins, J. H. 1976. Homology of myosin DTNB light chain with alkali light chains, troponin C and parvalbumin. *Nature (Lond.)* 259:699–700.
9. Cheung, W. Y. 1980. Calmodulin plays a pivotal role in cellular regulation. *Science (Wash. DC)* 207:19–27.
10. Reinach, F. C., K. Nagai, and J. Kendrick-Jones. 1986. Site-directed mutagenesis of the regulatory light-chain $\text{Ca}^{2+}/\text{Mg}^{2+}$ binding site and its role in hybrid myosins. *Nature (Lond.)* 322:80–83.
11. Morimoto, K., and W. F. Harrington. 1974. Evidence for structural changes in vertebrate thick filaments induced by calcium. *J. Mol. Biol.* 88:693–709.
12. Lehman, W. 1978. Thick-filament-linked calcium regulation in vertebrate striated muscle. *Nature (Lond.)* 274:80–81.
13. Brenner, B. 1988. Effect of Ca^{2+} on cross-bridge turnover kinetics in skinned single psoas fibers: implications for regulation of muscle contraction. *Proc. Natl. Acad. Sci. USA* 85:3265–3269.
14. Metzger, J. M., and R. L. Moss. 1990. pH modulation of the kinetics of a Ca^{2+} -sensitive cross-bridge state transition in mammalian single skeletal muscle fibres. *J. Physiol. (Lond.)* 428:751–764.
15. Metzger, J. M., and R. L. Moss. 1990. Calcium-sensitive cross-bridge transitions in mammalian fast and slow skeletal muscle fibers. *Science (Wash. DC)* 247:1088–1090.
16. Moss, R. L. 1979. Sarcomere length-tension relations of frog skinned muscle fibres during calcium activation at short lengths. *J. Physiol. (Lond.)* 292:177–192.
17. Metzger, J. M., M. L. Greaser, and R. L. Moss. 1989. Variations in cross-bridge attachment rate with phosphorylation of myosin in mammalian skinned skeletal muscle fibers. *J. Gen. Physiol.* 93:855–883.
18. Fabiato, A., and F. Fabiato. 1979. Calculator programs for computing the composition of the multiple metals and ligands used for experiments in skinned muscle cells. *J. Physiol. (Paris)* 75:463–505.
19. Godt, R. E., and B. D. Lindley. 1982. Influence of temperature upon contractile activation and isometric force production in mechanically skinned muscle fibers of the frog. *J. Gen. Physiol.* 80:279–297.
20. Brenner, B., and E. Eisenberg. 1986. Rate of force generation in muscle: correlation with actomyosin ATPase activity in solution. *Proc. Natl. Acad. Sci. USA* 83:3542–3546.
21. Hofmann, P. A., J. M. Metzger, M. L. Greaser, and R. L. Moss. 1990. Effects of partial extraction of light chain 2 on the Ca^{2+} sensitivities of isometric tension, stiffness, and velocity of shortening in skinned skeletal muscle fibers. *J. Gen. Physiol.* 95:477–498.
22. Moss, R. L., G. G. Giulian, and M. L. Greaser. 1983. Effects of EDTA treatment upon the protein subunit composition and mechanical properties of mammalian skinned skeletal muscle fibers. *J. Cell Biol.* 96:970–978.
23. Metzger, J. M., and R. L. Moss. 1991. Kinetics of a Ca^{2+} -sensitive cross-bridge state transition in skeletal muscle fibers. *J. Gen. Physiol.* 98:233–248.
24. Brandt, P. W., M. S. Diamond, and F. H. Schachat. 1984. The thin

- filament of vertebrate skeletal muscle co-operatively activates as a unit. *J. Mol. Biol.* 180:379–384.
25. Moss, R. L., G. G. Giulian, and M. L. Greaser. 1985. The effects of partial extraction of TnC upon the tension-pCa relationship in rabbit skinned skeletal muscle fibers. *J. Gen. Physiol.* 86:585–600.
26. Burton, K., and J. Sleep. 1988. The effect of phosphate on the rate of force recovery in rabbit psoas muscle fibers. *Biophys. J.* 53:564a. (Abstr.)
27. Metzger, J. M., and R. L. Moss. 1991. Phosphate and the kinetics of force generation in skinned skeletal muscle fibers. *Biophys. J.* 59:418a. (Abstr.)
28. Walker, J. W., Z. Lu, D. R. Swartz, and R. L. Moss. 1991. Thin filament modulation of cross-bridge transitions measured by photogeneration of P_i in skeletal muscle fibers. *Biophys. J.* 59:418a. (Abstr.)
29. Homsher, E., and N. C. Millar. 1989. Tension transients induced by photolysis of caged phosphate in glycerinated rabbit soleus fibres. *J. Physiol. (Lond.)* 418:P62.
30. Brandt, P. W., M. S. Diamond, J. S. Rutchik, and F. H. Schachat. 1987. Co-operative interactions between troponin-tropomyosin units extend the length of the thin filament in skeletal muscle. *J. Mol. Biol.* 195:885–896.
31. Holroyde, M. J., J. D. Potter, and R. J. Solaro. 1979. The calcium binding properties of phosphorylated and unphosphorylated cardiac and skeletal myosins. *J. Biol. Chem.* 254:6478–6482.
32. Bagshaw, C. R., and G. H. Reed. 1977. The significance of the slow dissociation of divalent metal ions from myosin 'regulatory' light chains. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 81:386–390.
33. Donaldson, S. K. B., and W. G. L. Kerrick. 1975. Characterization of the effects of Mg^{2+} on Ca^{2+} - and Sr^{2+} -activated tension generation of skinned skeletal muscle fibers. *J. Gen. Physiol.* 66:427–444.
34. Zot, A. S., and J. D. Potter. 1987. The effect of $[Mg^{2+}]$ on the Ca^{2+} dependence of ATPase and tension development of fast skeletal muscle. *J. Biol. Chem.* 262:1966–1969.
35. Kendrick-Jones, J., E. M. Szentkiralyi, and A. G. Szent-Györgyi. 1976. Regulatory light chains in myosins. *J. Mol. Biol.* 104:747–775.
36. Cox, J. A., M. Compte, and E. A. Stein. 1981. Calmodulin-free skeletal-muscle troponin C prepared in the absence of urea. *Biochem. J.* 195:205–211.
37. Alexis, M. N., and W. B. Gratzer. 1978. Interaction of skeletal muscle light chains with calcium ions. *Biochemistry.* 17:2319–2325.
38. Winkelman, D. A., and S. Lowey. 1986. Probing myosin head structure with monoclonal antibodies. *J. Mol. Biol.* 188:595–612.
39. Wagner, P. D. 1984. Effect of skeletal muscle myosin light chain 2 on the Ca-sensitive interaction of myosin and heavy meromyosin with regulated actin. *Biochemistry.* 23:5950–5956.