



Alexandria University Faculty of Medicine
Alexandria Journal of Medicine

<http://www.elsevier.com/locate/ajme>



Prevalence of intestinal parasites in newly diagnosed HIV/AIDS patients in Ilorin, Nigeria

O.A. Obateru^a, B.J. Bojuwoye^b, A.B. Olokoba^{b,*}, A. Fadeyi^c, A. Fowotade^d, L.B. Olokoba^e

^a Department of Medicine, Federal Medical Centre, Lokoja, Nigeria

^b Department of Medicine, University of Ilorin Teaching Hospital, Ilorin, Nigeria

^c Department of Microbiology, University of Ilorin Teaching Hospital, Ilorin, Nigeria

^d Department of Microbiology, University College Hospital, Ibadan, Nigeria

^e Department of Ophthalmology, General Hospital, Ilorin, Nigeria

Received 3 January 2016; revised 11 April 2016; accepted 12 April 2016

KEYWORDS

Intestinal parasites;
 HIV/AIDS;
 Newly diagnosed;
 Treatment naive;
 Adults

Abstract *Background:* Human immune-deficiency virus/acquired immune-deficiency syndrome predisposes to opportunistic parasitic infestations of the gastrointestinal tract. This study aimed to determine the prevalence of intestinal parasites in newly diagnosed treatment naïve HIV/AIDS patients.

Methods: This hospital-based cross-sectional study was carried out from December 2010 to June 2011. Questionnaires were administered to 238 HIV/AIDS subjects, and 238 age and sex-matched controls. CD4⁺ T cell count was carried out on HIV-positive subjects. Stool samples were examined using direct microscopic and modified Ziehl-Neelsen methods. Positivity of intestinal parasites was taken as the presence of worms, oocyst, cyst, ova or larvae in the stool samples.

Results: Ninety males and 148 females were studied for the HIV-positive and HIV-negative controls respectively. Intestinal parasitic infestation in HIV-positive subjects was 68.5%, and was significantly higher than in the HIV-negative controls 49.2% ($P < 0.05$). In HIV-positive subjects, *Cryptosporidium* spp. was the commonest (55.0%) parasite isolated. Others were *Cyclospora cayotensis* (41.2%), *Isospora belli* (3.0%), *Entamoeba histolytica* (8.4%), *Giardia lamblia* (3.7%), *Ascaris lumbricoides* (2.5%), *Strongyloides stercoralis* (1.7%), *Trichuris trichiura* (0.8%) and *Schistosoma mansoni* (0.4%). HIV-positive patients with CD4⁺ T cell count of less than 200 cells/ul were more at risk of opportunistic parasites compared to the HIV-negative controls.

Conclusion: The prevalence of intestinal parasites in newly diagnosed HIV/AIDS individuals was high, and its association with CD4⁺ T cell count was demonstrated. Routine screening for parasitic infestations at diagnosis is indicated to reduce the burden of the disease.

© 2016 Alexandria University Faculty of Medicine. Production and hosting by Elsevier B.V This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author. Tel.: +234 8038050480.

E-mail address: drabolokoba@yahoo.com (A.B. Olokoba).

Peer review under responsibility of Alexandria University Faculty of Medicine.

<http://dx.doi.org/10.1016/j.ajme.2016.04.001>

2090-5068 © 2016 Alexandria University Faculty of Medicine. Production and hosting by Elsevier B.V

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Human immune-deficiency virus/acquired immune-deficiency syndrome (HIV/AIDS) is a global public health problem.¹ The disease predisposes individuals to various opportunistic infections of which parasitic infestations of the gastrointestinal (GI) tract are prominent.² Several studies have shown that some intestinal parasites such as *Cryptosporidium* spp., *Cyclospora cayetanensis*, *Isospora belli*, *Entamoeba histolytica/dispar*, *Strongyloides stercoralis*, and *Giardia lamblia* are responsible for 60–80% of infestations.³ Thus infestation of the GI tract by these organisms plays a crucial role in HIV/AIDS pathogenesis, and diarrhoea diseases assume a prominent role reaching up to 50% in developing countries.² The decrease in immunity by attack on the immune system especially the cluster of differentiation (CD4⁺ T cells) component, macrophages, and defect in the production of immunoglobulin A (IgA) increases susceptibility to these parasites.⁴ Gut-associated lymphatic tissue (GALT) in the GI tract is an important site for early HIV replication and CD4⁺ T cell replication.⁵

Helper T cells which are very important in adaptive immunity, activate B cells to secrete antibodies, activate macrophages to destroy ingested microbes, and also activate cytotoxic T cells to kill infected target cells. These functions are impaired in patients with HIV/AIDS.⁶ Furthermore, HIV/AIDS decrease the circulating pool of effector, and memory CD8⁺ T cells which combat viral infections. The end result is immunosuppression.⁷ There have been reports of the prevalence of intestinal parasites in HIV/AIDS patients in other parts of Nigeria and the world.^{8–14} Ibrahim et al.⁸, and Babatunde et al.⁹, in North central, Nigeria, found an increase in the prevalence of intestinal parasites in HIV-positive patients compared to HIV-negative controls. Similarly, Akinbo et al.¹⁰, Hailemariam et al.¹¹, Alemu et al.¹², Jegede et al.¹³, and Oyediji et al.¹⁴ all found an increase in the prevalence and multiple intestinal parasitosis in HIV-positive Nigerian and Ethiopian patients compared with HIV-negative controls. These studies were mostly on patients already on highly active anti-retroviral therapy (HAART). There is a paucity of data on similar studies on treatment naïve HIV/AIDS patients. It is thus important to find out the common intestinal parasites in newly diagnosed treatment naïve adult HIV/AIDS patients in Ilorin, Nigeria, since these parasites play a prominent role in diarrhoeal diseases, and HIV/AIDS pathogenesis.

This study therefore aimed to determine the prevalence of intestinal parasites in newly-diagnosed treatment naïve HIV/AIDS patients.

2. Subjects, materials and methods

The study was a hospital-based cross-sectional study carried out over a six-month period (December 2010–June 2011), at the Infectious disease clinic of University of Ilorin Teaching Hospital (UIH), Ilorin, Nigeria. A standard structured questionnaire was administered to consecutive 238 newly diagnosed treatment naïve adult HIV/AIDS patients, and 238 age and sex-matched HIV-negative controls whose blood and stool samples were tested at first enrolment. Individuals who had taken antibiotics, anthelmintic drugs and steroids prior to enrolment were excluded from the study.

2.1. Specimen collection

Blood sample: Ten millilitres of venous blood samples were collected aseptically into labelled EDTA specimen bottles. Rapid Enzyme-linked immunosorbent assay (ELISA) was carried out to detect antibodies to HIV 1 and 2 using rapid ELISA test kit (AcuuBio Tech Co., Ltd., China). CD4⁺ T cell count was carried out with Partec 4000 cyflow counter Germany (2006) at the HIV laboratory of UIH.¹⁵ The range of CD4⁺ T cell count was classified using the 1993 Center for Disease Control and Prevention USA, immunological criteria.¹⁶

Stool sample: Stool samples were collected into sterile containers, labelled, and analysed within 24 h of collection. Stool samples were examined using direct microscopy for wet preparation, and iodine preparation.^{17,18} Modified Ziehl-Neelsen methods, and Faecal concentration technique (using formal ether concentration method) were also used to increase the yield of intestinal parasites. Positivity of intestinal parasites was taken as the presence of worms, ova, trophozoites, oocysts and cysts in the stool samples.^{17–19}

Data analysis: Epi-info™ version 3.5.1(CDC, USA 2008) software package was used for analysis. Statistical significance was taken as *p*-value less than 0.05.

Ethical approval was obtained from the Ethics and Research committee of UIH.

3. Results

There were 90 males (37.8%) and 148 females (62.2%) in both groups, with a male to female ratio of 0.6:1. The age range for both groups (HIV-positive subjects and HIV-negative controls) was from 16 to 65 years. The mean age of respondents was 36.1 ± 9.6 years and 35.9 ± 9.4 years for the HIV-positive subjects and HIV-negative controls respectively. The commonest age group was 26–35 years for both HIV-positive subjects and HIV-negative controls. The mean age of the respondents was similar (*P* > 0.05). Majority of the respondents were married, and secondary school level of education was the highest level of education attained by them. Most of the respondents were of Yoruba ethnic origin, and trading was the major occupation of the HIV-positive subjects, while most of the HIV-negative controls were self-employed (Table 1).

The prevalence of intestinal parasites in HIV-positive subjects and HIV-negative controls was 68.5% and 49.2% respectively. HIV-positive subjects had increased risk of intestinal parasites compared to HIV-negative controls (*P* < 0.05); OR = 2.25 CI (1.52–3.32).

The most frequently detected parasites in the HIV-positive subjects and HIV-negative controls were *Cryptosporidium* spp. and *C. cayetanensis*. These were significantly higher in the HIV-positive subjects than in HIV-negative controls (55% vs 16.8%) ($X^2 = 73.97$; *P* < 0.05) and (41.2% vs 8.4%) ($X^2 = 66.81$; *P* < 0.05) respectively. There were no statistically significant differences in the prevalence of *I. belli* (3.0% vs 0.8%; $X^2 = 1.81$; *P* > 0.05), *E. histolytica* (8.4% vs 8.8%) ($X^2 = 1.0$; *P* > 0.05), *G. lamblia* (3.7% vs 0%), *Hookworm* spp. (0% vs 0.8%), *S. stercoralis* (1.7% vs 0.8%; *P* > 0.05), *Ascaris lumbricoides* (2.5% vs 0%), *Trichuris trichiura* (0.8% vs 0%), *Schistosoma mansoni* (0.4% vs 0%) in both groups (Table 2).

Table 1 Socio-demographic characteristics of respondents.

	HIV-positive subjects frequency <i>n</i> (%)	HIV-negative controls frequency <i>n</i> (%)
<i>Age (years)</i>		
16–25	42(17.6)	42(17.6)
26–35	89(37.4)	89(37.4)
36–45	68(28.6)	68(28.6)
46–55	33(13.9)	33(13.9)
56–65	6(2.5)	6(2.5)
<i>Gender</i>		
Female	148(62.2)	148(62.2)
Male	90(37.8)	90(37.8)
<i>Marital status</i>		
Single	59(24.8)	89(37.4)
Married	169(71.0)	148(62.2)
Divorced	7(2.9)	1(0.4)
Widowed	3(1.3)	0(0)
<i>Ethnicity</i>		
Hausa	10(4.2)	12(5.0)
Yoruba	171(71.9)	191(80.3)
Ibo	12(5.0)	9(3.8)
Others	45(18.9)	26(10.9)
<i>Level of education</i>		
No formal education	65(27.3)	27(11.3)
Primary	70(29.4)	52(21.8)
Secondary	75(31.5)	83(34.9)
Tertiary	19(8.0)	68(28.6)
Quranic	9(3.8)	8(3.4)
<i>Occupation</i>		
Unemployed	37(15.6)	30(12.6)
Self employed	67(28.1)	61(25.6)
Students	7(2.5)	45(18.9)
Traders	77(32.4)	37(15.6)
Civil servants	42(17.6)	44(18.5)
Professionals	9(3.8)	21(8.8)

Some respondents were infected with up to four different parasites. HIV-positive subjects had more multiple parasitic infestation rates (more than one) than the HIV-negative controls 33.2% (79/238) and 25.6% (61/238) respectively ($\chi^2 = 34.13$; $P < 0.05$) (Fig. 1).

The range of CD4⁺ T cell count in HIV-positive patients was 13–883 cells/ul. The mean CD4⁺ T cell count was 257 ± 201.7. HIV-positive patients with CD4⁺ T cell count < 200 cells/ul had increased risk of intestinal parasitic infestation ($p = 0.017$; OR 1.95; CI = 1.09–3.55) compared to those with CD4⁺ T cell count of 200–499 ($p = 0.760$; OR = 1.09, CI = 0.59–2.04) and those with CD4⁺ T cell count > 500 ($p = 1.00$; OR = 0.24, CI = 0.10–0.54) (Table 3).

4. Discussion

Intestinal parasites, especially the opportunistic ones are a common cause of morbidity and mortality in HIV/AIDS patients.⁸ The prevalence of intestinal parasites in HIV-positive subjects in this study was found to be high (68.5%). Studies carried out in other parts of Nigeria showed diversity in findings. The figure of 68.5% is lower than the 87.8% reported in an earlier study carried out in our centre by Babatunde et al.⁹ Ibrahim et al. in North-central Nigeria, documented a figure of (50.0%).⁸ Akinbo et al. in South-south Nigeria reported 15.3%.¹⁰ This variation in prevalence is not different from that documented in Africa and other parts of the world. Hailemariam et al.¹¹ and Alemu et al.¹² in Ethiopia reported a prevalence of 52.6% and 80.3% respectively. Kipyegen et al. in Kenya documented a parasitic prevalence in HIV-positive patients which is to be 50.0%.²⁰

Da silva et al. in Brazil found a prevalence rate of 27.0%.²¹ Mohandas et al. in India found out that 30.0% of HIV-positive patients harbour intestinal parasites.²²

HIV-positive subjects in this study were all newly diagnosed and treatment naïve in contrast to the other studies mentioned above. Furthermore, the study subjects in aforementioned studies, were a combination of those on treatment and treatment naïve. They may have used anthelmintic drugs such as Albendazole, Piperazine, Levamisole, and HAART. This may have boosted their immunity and subsequently altered the intestinal flora in the gut.¹⁴ Sampling techniques and observer differences in the examination of stool samples may be another factor responsible for these differences in prevalence even though most of the studies involved the use of direct wet mount, formalin-ether sedimentation concentration, and modified Ziehl Neelsen methods. However, Hailemariam et al.¹¹ did not use Ziehl Neelsen method.

Table 2 Distribution of intestinal parasitic species in respondents.

Parasites	HIV-positive	HIV-negative	Chi-squared	<i>p</i> -value	Comment
(1) <i>Cryptosporidium</i> spp.	131(55.0)	40(16.8)	73.97	< 0.05	S
(2) <i>Cyclospora</i> spp.	98(41.2)	20(8.4)	66.81	< 0.05	S
(3) <i>Isospora belli</i>	7(3.0)	2(0.8)	1.81	> 0.05*	NS
(4) <i>E. histolytica</i>	20(8.4)	21(8.8)	1.0	> 0.05	NS
(5) <i>Giardia lamblia</i>	9 (3.7)	0(0)	–	–	–
(6) Hookworm spp.	0(0)	2(0.8)	–	–	–
(7) <i>S. stercoralis</i>	4(1.7)	2(0.8)	–	> 0.05*	–
(8) <i>A. lumbricoides</i>	6(2.5)	0(0)	–	–	–
(9) <i>Trichuris trichiura</i>	2(0.8)	0(0)	–	–	–
(10) <i>S. mansoni</i>	1(0.4)	0(0)	–	–	–

S-significant.

NS-not significant.

* *p* value from Fishers exact test.




Figure 1 Multiple parasitic infestation rate in respondents.

Table 3 Distribution of intestinal parasites in HIV-positive subjects at different values of CD4⁺ T cell count (cells/ul).

CD4 ⁺ T Cell count	n (%) n = 238	Intestinal parasites n (%) n = 163	Odds ratio (95% CI)	p-value
< 200	125(52.5)	94(57.7)	1.95(1.09–3.55)	0.017
200–499	79(33.2)	55(33.7)	1.09(0.59–2.04)	0.760
> 500	34(14.3)	14(8.6)	0.24(0.10–0.54)	1.000

Cryptosporidium spp. was found to be the most predominant intestinal parasite in HIV-positive subjects (55.0%). Studies carried out in Ile-Ife, South-west Nigeria, by Oyedeji et al.¹⁴ and Adesiji et al.²³ found the most predominant intestinal parasite in HIV-positive subjects to be *Cryptosporidium* spp. (19.2% and 52.7%), and Thomas et al. in Lagos, South-west Nigeria, documented similar findings.²⁴ However, this is in contrast to the findings by Nwokediuko et al. in Enugu, South-east Nigeria²⁵ and Oyerinde et al. in Lagos²⁶, who reported a rarity of *Cryptosporidium* spp. This is unusual, as *Cryptosporidium* spp. infection has been documented even in apparently normal individuals by Ikeh et al.²⁷, Nwabuisi²⁸, and Banwat et al.²⁹ all in North-central Nigeria. The prevailing environmental conditions may not be favourable for the transmission of these parasites in that part of the country, which may have accounted for the reported low incidence of *Cryptosporidium* spp. in those regions. These studies were noted to have been carried out on patients with diarrhoea; hence, they may have taken anthelmintic drugs such as Albendazole, Piperazine, and Levamisole prior to enrolment into the studies. The findings of Oyedeji et al.¹⁴ were influenced by the prior use of ant-helminthics and Co-trimoxazole. Jegede et al.¹³ in North central, Nigeria, did not find any statistical difference in the spectrum of intestinal parasites between HIV-positive and HIV-negative controls.

However, studies in Nigeria by Wokem et al.³⁰ and Ethiopia by Hailemariam et al.¹¹ found *A. lumbricoides* to be the most predominant parasites in HIV-positive subjects. *S. stercoralis* was found to be most predominant in HIV-positive subjects in Thailand.³¹ Further studies need to be carried out in those geographical areas to validate these findings as opportunistic parasites are expected to thrive more in HIV-positive patients.

From this study, *C. cayetanensis* was detected in a significant number of HIV-positive subjects compared to

HIV-negative controls (41.2% vs 8.4%). This differs from reports in some parts of Nigeria and the world at large, in which the prevalence was found to be low or undetected.^{23,24,32,33} The geographical variation and the remarkable similarity in morphologic features with *Cryptosporidium* spp. could account for these differences. The presence of *Cryptosporidium* spp. and *C. cayetanensis* in the HIV-negative controls may be due to the fact that asymptomatic carrier state can occur in immune-competent individuals and they only become symptomatic with diarrhoeal disease, when they become immune-deficient.^{28,30,34,35} Reports by Nwabuisi²⁸ in Ilorin, and Banwat et al.²⁹ in Jos, North-central Nigeria, have shown that infection with *Cryptosporidium* spp. and *C. cayetanensis* can occur in immuno-competent individuals. Poor personal, and environmental hygiene which could facilitate transmission of the cyst and ova of parasites may be responsible in this part of the world.

The rate of multiple parasitic infestations was 33.2% and 25.6% in HIV-positive and HIV-negative controls respectively. A report from Jos by Ibrahim et al.⁸, found multiple parasitic infection rates in HIV-positive subjects to be lower (13%), while Hailemariam et al.¹¹, and Alemu et al. documented a high multiple parasitic infestation rate of 91.3%.¹² This may be attributed to depressed immunity in the HIV-positive subjects, which predisposed them to mixed infections including the opportunistic and non-opportunistic parasites.

HIV-positive patients with CD4⁺ T cell count < 200 cells/ul had increased risk of intestinal parasitic infestation ($P < 0.05$) compared to those with CD4⁺ T cell count of > 200. Majority of HIV-positive subjects with CD4⁺ T cell count of < 200 cells/ul had more than double the risk of developing diarrhoea ($P < 0.05$) compared to those with CD4⁺ T cell count of > 200. This is in keeping of reports from earlier studies carried out by Babatunde et al.⁹, Wiwanitkit³¹ and Assefa et al.³² All these findings have shown that immunity plays a major role in

the survival and proliferation of intestinal parasites in HIV-positive patients.

A major limitation in this study, is the fact that other staining techniques such as modified trichome staining technique were not used which could have detected Microsporidia spores.

In conclusion, the prevalence of intestinal parasites in treatment naive adult HIV/AIDS individuals was found to be high. HIV-positive individuals with CD4⁺ T cell count less than 200 cells/ul were more at risk of infestations with intestinal parasites, and multiple intestinal parasites than those with more than 200 cell/ul. Apparently normal individuals also harbour opportunistic parasites.

Conflict of interest

None.

Source of funding

None

References

- Kallings LO. The first postmodern pandemic: 25 years of HIV/AIDS. *J Intern Med* 2008;**263**(3):218–43.
- Cimerman S, Cimerman B, Levi DS. Enteric parasites and AIDS. *Sao Paulo Med J* 1999;**117**(6):70–7.
- Bolinger RC, Quinn TC. Tropical diseases in the HIV-infected traveller. In: Broder S, Merigan Jr, Bolognesi D, editors. Textbook of aids medicine. New York: Williams & Williams; 1994. p. 311–22, 2nd ed..
- Osmanov S, Pathouy C, Walker N, Schward-lander B, Esparz J. WHO-UNAIDS network for HIV isolation and characterization. Estimated global distribution and regional spread of HIV-1 genetic subtypes in the year 2000. *J Acq Immune Def Syndr* 2002;**29**:184–90.
- Dandekar S. Pathogenesis of HIV in the gastrointestinal tract. *Curr HIV/AIDS Rep* 2007;**4**(1):10–5.
- Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Walker P. Helper T cells and lymphocyte activation. *Molecular biology of the cell*. 4th ed. New York: Garland Science; 2002. p. 561–2.
- Gulzar N, Copeland KF. CD8 + T-cells: function and response to HIV infection. *Curr HIV Res* 2004;**2**(1):23–37.
- Ibrahim A, Ikeh E, Malu A, Okeke E, Damen J. Intestinal parasitosis in Human Immunodeficiency Virus (HIV) infected adults with chronic diarrhoea at Jos University Teaching Hospital, Nigeria. *The Int J Par Dis* 2007;**2**(1).
- Babatunde SK, Salami AK, Fabiyi JP, Agbede OO, Desalu OO. Prevalence of intestinal parasitic infestations in HIV- seropositive and HIV- seronegative patients in Ilorin, Nigeria. *Ann Afr Med* 2010;**9**(3):123–8.
- Akinbo FO, Okaka CE, Omeregic R. Prevalence of intestinal parasitic infections among HIV patients in Benin city, Nigeria. *Libyan J Med* 2010;**5**:10.
- Hailemariam G, Kassu A, Abebe G, Abate E, Damte D, Mekonnen E, et al. Intestinal parasitic infections in HIV/AIDS and HIV- seronegative individuals in a Teaching Hospital in Ethiopia. *Jpn J Infect Dis* 2004;**57**(2):41–3.
- Alemu A, Siferaw Y, Getnet G, Yalew A, Addis Z. Opportunistic and other intestinal parasites among HIV/AIDS patients attending Gambi higher clinic in Bahir Dar city, North West Ethiopia. *Asian Pac J Med* 2011;**4**(8):661–5.
- Jegede EF, Oyeyi ETI, Bichi AH, Mbah HA, Torpey K. Prevalence of intestinal parasites among HIV/AIDS patients attending Infectious Disease Hospital, Kano, Nigeria. *Pan Afr Med J* 2014;**17**:295.
- Oyediji OA, Adejuyigbe E, Oninla SO, Akindele AA, Adedokun E, Agelebe E. Intestinal parasitosis in HIV infected children in a Nigerian Tertiary Hospital. *J Clin Diag Res* 2015;**9**(11):SC01–5.
- Partec Cyflow. Count of CD+4 cell with cyflow counter and partec CD+4 count essay count kit. Application note on procedures partec code number 05-8401 partec GmbH Germany; 2007.
- Center for Disease Control and Prevention. Revised classification system for HIV infection and expanded surveillance classification system case definition for AIDS among adolescence and adults. *JAMA* 1993;**269**(1993):729–30.
- Cheesbrough M. Parasitological tests. *District laboratory practice in tropical countries*. 2nd ed. Cambridge: Cambridge University Press; 2005. p. 191–200.
- World Health Organisation. Guidelines on standard operating procedures for laboratory diagnosis of HIV-opportunistic infections, Geneva; 2001.
- Said DE. Detection of parasites in commonly consumed raw vegetables. *Alexandria J Med* 2012;**48**:345–52.
- Kipyegen CK, Shivairo RS, Odhiambo RO. Prevalence of intestinal parasites among HIV patients in Baringo, Kenya. *Pan Afr Med J* 2012;**12**:37.
- Da Silva CV, Ferreira MS, Borges AS, Costa-Cruz JM. Intestinal parasitic infections in HIV/AIDS patients: experience at a teaching hospital in central Brazil. *Scan J Inf Dis* 2005;**37**(3):211–5.
- Mohandas K, Sehgal R, Sud A, Malla N. Prevalence of intestinal parasitic pathogens in HIV-positive individuals in Northern India. *Jpn J Infect Dis* 2002;**55**(3):83–4.
- Adesiji YO, Lawal RO, Taiwo SS, Fayemiwo SA, Adeyeba OA. Cryptosporidiosis in HIV infected patients with diarrhoea in Osun State south-western Nigeria. *Eur J Gen Med* 2007;**4**(3):119–22.
- Thomas B, Fagbenro-Beyioku AF, Ukegbuy CB. Prevalence of enteric opportunistic pathogens among HIV/AIDS patients in Lagos, Nigeria and comparative assessment of three staining techniques. Int Conf AIDS 2000 July 9–14; 13: abstract no. MoPeB2266.
- Nwokediuko SC, Bojuwoye BJ, Onyenekwe B. Apparent rarity of cryptosporidium in HIV related diarrhoea in Enugu, Nigeria. *Nig Postgrad Med J* 2002;**9**(2):70–3.
- Oyerinde JP, Odugbemi T, Benson RI, Alonge AA. Investigation of cryptosporidium in relation to other intestinal parasites at the Lagos university teaching hospital, Lagos. *West Afr J Med* 1989;**8**(4):264–9.
- Ikeh EI, Obadofin MO, Brindeiro B, Baugherb C, Frost F, Vanderjargt D. Intestinal parasitism in Magama Gumau rural village and Jos township in North Central Nigeria. *Nig Postgrad Med J* 2007;**14**(4):290–5.
- Nwabuisi C. Childhood cryptosporidiosis and intestinal parasitosis in association with diarrhoea in Kwara State, Nigeria. *West Afr J Med* 2001;**20**:165–84.
- Banwat EB, Egah DZ, Onile BA, Angyo IA, Audu ES. Prevalence of cryptosporidium infection among under-nourished children in Jos, Central Nigeria. *Niger Postgrad Med J* 2003;**10**:84–7.
- Wokem GN, Chukwu C, Nwachukwu BC. Prevalence of intestinal parasites seen in HIV sero-positive subjects in Port-Harcourt, Nigeria. *Niger. J Parasitol* 2008;**29**(2):115–20.
- Wiwanitkit V. Intestinal parasitic infections in Thai HIV-infected patients with different immunity status. *BMC Gastroenterol* 2001;**1**, 3e-pub.
- Assefa S, Erko B, Medhen G, Assefa Z, Shimeli T. Intestinal parasitic infection in relation to HIV, AIDS status, diarrhoea and CD4 cell-count. *BMC Inf Dis* 2009;**9**:155.

33. Brandonosio O, Maggi MA, Panaso S, Lisi A. Intestinal protozoa in HIV infected patients in Apulia, south Italy. *Epidemiol Infect* 1999;**123**:457–62.
34. Alakpa G, Fagbenro-Beyioku A, Clarke S. Cyclospora cayetansensis in stool submitted to hospitals in Lagos. *Int J Infect Dis* 2001;**6**(4):314–7.
35. Helmy MM, Rasheed LA, Abdel-Fattah HC. Co-infection with cryptosporidium and cyclospora in immunocompetent and immunocompromised patients. *J Egypt Soc Parasitol* 2003;**36**(2):613–27.