




ELSEVIER

Contents lists available at [ScienceDirect](http://ScienceDirect)

# Journal of Clinical Tuberculosis and Other Mycobacterial Diseases

journal homepage: [www.elsevier.com/locate/jctube](http://www.elsevier.com/locate/jctube)

## Corrigendum

### Corrigendum to “The nematode *Caenorhabditis elegans* displays a chemotaxis behavior to tuberculosis-specific odorants” [J. Clin. Tuberc. Mycobact. Dis. 4 (2016) 44–49]

Mário F. Neto<sup>a,1</sup>, Quan H. Nguyen<sup>b,1</sup>, Joseph Marsili<sup>b</sup>, Sally M. McFall<sup>a</sup>, Cindy Voisine<sup>b,\*</sup><sup>a</sup> Center for Innovation in Global Health Technologies (CIGHT), Department of Biomedical Engineering, Northwestern University, Evanston, IL 60208, USA<sup>b</sup> Department of Biology, Northeastern Illinois University, Chicago, IL 60625, USA

The authors regret that in the initially published version of the article, an error in calculating the highest concentration (reported as 100 ng/ul) used for each of the 4 TB specific VOCs was made. To generate the initial concentrations used in Fig. 2 A and B and Fig. 3 B, 1 ml of *o*-phenylanisole or methyl phenylacetate was added to 9 ml of ethanol. A ten fold serial dilution of this starting concentration was performed to generate the lower concentrations tested. For methyl *p*-anisate and methyl nicotinate, the highest concentration used was 100 ug/ul in ethanol. A ten fold serial dilution of this starting concentration was performed to generate the lower concentrations tested. The conclusions of the paper do not change but the initial concentration values are reported incorrectly. The authors would like to apologize for any inconvenience caused.

DOI of original article: [10.1016/j.jctube.2016.06.001](https://doi.org/10.1016/j.jctube.2016.06.001)

\* Corresponding author.

E-mail address: [c-voisine@neu.edu](mailto:c-voisine@neu.edu) (C. Voisine).<sup>1</sup> These authors contributed equally.

<http://dx.doi.org/10.1016/j.jctube.2016.11.002>  
2405-5794