


Contents lists available at [ScienceDirect](http://ScienceDirect.com)

## Physics Letters B

[www.elsevier.com/locate/physletb](http://www.elsevier.com/locate/physletb)The  $\pi\rho$  cloud contribution to the  $\omega$  width in nuclear matterD. Cabrera<sup>a,b</sup>, R. Rapp<sup>c</sup><sup>a</sup> Institute for Theoretical Physics, Frankfurt University, 60438 Frankfurt am Main, Germany<sup>b</sup> Frankfurt Institute for Advanced Studies, Frankfurt University, 60438 Frankfurt am Main, Germany<sup>c</sup> Cyclotron Institute and Department of Physics & Astronomy, Texas A&M University, College Station, TX 77843-3366, United States

## ARTICLE INFO

## Article history:

Received 17 July 2013

Received in revised form 9 December 2013

Accepted 26 December 2013

Available online 31 December 2013

Editor: J.-P. Blaizot

## ABSTRACT

The width of the  $\omega$  meson in cold nuclear matter is computed in a hadronic many-body approach, focusing on a detailed treatment of the medium modifications of intermediate  $\pi\rho$  states. The  $\pi$  and  $\rho$  propagators are dressed by their self-energies in nuclear matter taken from previously constrained many-body calculations. The pion self-energy includes  $Nh$  and  $\Delta h$  excitations with short-range correlations, while the  $\rho$  self-energy incorporates the same dressing of its  $2\pi$  cloud with a full 3-momentum dependence and vertex corrections, as well as direct resonance-hole excitations; both contributions were quantitatively fit to total photo-absorption spectra and  $\pi N \rightarrow \rho N$  scattering. Our calculations account for in-medium decays of type  $\omega N \rightarrow \pi N^{(*)}$ ,  $\pi\pi N(\Delta)$ , and 2-body absorptions  $\omega NN \rightarrow NN^{(*)}$ ,  $\pi NN$ . This causes deviations of the in-medium  $\omega$  width from a linear behavior in density, with important contributions from spacelike  $\rho$  propagators. The  $\omega$  width from the  $\rho\pi$  cloud may reach up to 200 MeV at normal nuclear matter density, with a moderate 3-momentum dependence. This largely resolves the discrepancy of linear  $T-Q$  approximations with the values deduced from nuclear photoproduction measurements.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>). Funded by SCOAP<sup>3</sup>.

## 1. Introduction

The low-mass vector mesons  $\rho$ ,  $\omega$  and  $\phi$  play a special role in the study of hot and dense nuclear matter, as their dilepton decay channel ( $I^+I^-$ ) provides a pristine window on their in-medium properties. This feature has been extensively and successfully exploited in the measurement of dilepton spectra in heavy-ion collisions [1–3]. In these reactions, the thermal emission of low-mass dileptons is dominated by the  $\rho$  meson, due to its much larger dilepton width compared to the  $\omega$ ,  $\Gamma_{\rho \rightarrow \ell\ell} \simeq 10 \Gamma_{\omega \rightarrow \ell\ell}$ . Dilepton data from the SPS and RHIC can now be consistently understood by a strong broadening (“melting”) of the  $\rho$  meson, as computed from hadronic many-body theory in the hot and dense system [4,5]. This approach also yields a good description [6,7] of the  $\rho$  broadening observed in nuclear photoproduction, if the data are corrected with absolute background determination [8,9]. As a further test of the validity and generality of the hadronic in-medium approach, the  $\omega$  meson, as the isospin zero pendant of the  $\rho$ , is a natural candidate.

The small dilepton decay width of the  $\omega$  led the CB-TAPS Collaboration to pursue the  $\pi^0\gamma$  decay channel in photon-induced production off nuclei. Early results for invariant-mass spectra reported significant downward mass shifts [10], seemingly in line with proton-induced dilepton production off nuclei [11]. However, with improved background determination these results were not

confirmed [12,13], leaving no evidence for a mass drop. As an alternative method, absorption measurements have been performed for  $\phi$  and  $\omega$  mesons in  $e^+e^-$  [14,15] and  $\pi^0\gamma$  [16] channels. These data are not directly sensitive to possible mass shifts, but they can be used to assess the in-medium (absorptive) widths. For both  $\phi$  and  $\omega$ , large in-medium widths have been deduced, e.g.,  $\Gamma_{\omega}^{\text{med}} \simeq 130\text{--}150$  MeV [16], or even above 200 MeV [15], for the  $\omega$  at normal nuclear matter density. These values exceed the free  $\omega$  width by a factor of  $\sim 20$ , posing a challenge for theoretical models [17–25].

Most of the calculations thus far are based on the so-called  $T-Q$  approximation, where the in-medium  $\omega$  self-energy is computed from the vacuum scattering amplitude and therefore depends linearly on nuclear density,  $\rho_N$  (see, however, Refs. [26,27]). In the present work we go beyond this approximation by simultaneously dressing the  $\pi$  and  $\rho$  propagators in the  $\pi\rho$  loop of the  $\omega$  self-energy. In the vacuum, the  $\omega$  decay into  $\pi\rho$  has a nominal threshold of  $m_\pi + m_\rho \simeq 910$  MeV and only proceeds through the low-mass tail of the  $\rho$  resonance, which is suppressed and possibly responsible for the small width of  $\Gamma_{\omega \rightarrow 3\pi} \simeq 7.5$  MeV. A broadening of the  $\rho$  in the medium enhances this decay channel, further augmented if the pion is dressed as well. This is a key point we aim to convey and elaborate quantitatively in this Letter by utilizing realistic in-medium  $\pi$  and  $\rho$  propagators.

<http://dx.doi.org/10.1016/j.physletb.2013.12.056>

0370-2693/© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>). Funded by SCOAP<sup>3</sup>.

Our Letter is organized as follows. In Section 2 we set up the  $\omega \rightarrow \pi\rho$  self-energy in vacuum (Section 2.1) and discuss the implementation of the  $\pi$  and  $\rho$  propagators in nuclear matter (Section 2.2). In Section 3 we quantitatively evaluate the consequences of the in-medium propagators on the density and 3-momentum dependence of the  $\omega$  width. We summarize and give an outlook in Section 4.

## 2. $\omega$ self-energy

### 2.1. $\omega$ width in vacuum

In vacuum we describe the coupling of the  $\omega$  to a pion and a  $\rho$  meson with the chiral anomalous interaction Lagrangian introduced, e.g., in the work by Jain et al. [28],

$$\mathcal{L}_{\omega\rho\pi}^{\text{int}} = g_{\omega\rho\pi} \epsilon_{\mu\nu\sigma\tau} \partial^\mu \omega^\nu \partial^\sigma \vec{\rho}^\tau \cdot \vec{\pi}. \quad (1)$$

The value of the coupling constant,  $g_{\omega\rho\pi}$ , determines the partial decay width  $\Gamma_{\omega \rightarrow \rho\pi}$  and will be discussed below. A straightforward application of Feynman rules for the  $\pi\rho$  loop yields the polarization-averaged self-energy of an  $\omega$  of 4-momentum  $P = (P^0, \vec{P})$  as

$$\begin{aligned} -i\Pi_\omega(P) = IF \frac{1}{3} \sum_{\lambda,\delta} \epsilon_\lambda^\nu(P) \epsilon_\delta^{\nu'}(P) i g_{\omega\rho\pi} i g_{\omega\rho\pi} \epsilon_{\mu\nu\alpha\beta} \epsilon_{\mu'\nu'\alpha'\beta'} \\ \times \int \frac{d^4q}{(2\pi)^4} P^\mu q^\alpha P^{\mu'} q^{\alpha'} i D_\rho^{\beta\beta'}(q) i D_\pi(P-q), \end{aligned} \quad (2)$$

where the isospin factor  $IF = 3$  accounts for the different  $\pi\rho$  charge states. Using standard representations of the polarization sum and of the spin-1  $\rho$  propagator,  $D_\rho^{\beta\beta'}$ , which we decompose in transverse ( $T$ ) and longitudinal ( $L$ ) modes [29], one finds

$$\begin{aligned} -i\Pi_\omega(P) = -\frac{4}{3} IF g_{\omega\rho\pi}^2 \int \frac{d^4q}{(2\pi)^4} D_\pi(P-q) \{v_1(q, P) D_\rho^T(q) \\ + v_2(q, P) [D_\rho^T(q) - D_\rho^L(q)]\} \end{aligned} \quad (3)$$

where  $D_\pi(P-q) = 1/[i(P-q)^2 - m_\pi^2 - \Pi_\pi]$  and  $D_\rho^{T,L}(q) = 1/[q^2 - M_\rho^2 - \Pi_\rho^{T,L}]$  are the scalar parts of the meson propagators with complex self-energies. The two vertex functions arise from the Lorentz contractions with the  $T$  and  $L$  projectors of the  $\rho$  propagator,  $v_1(q, P) = P^2 q^2 - (Pq)^2$  and  $v_2(q, P) = q^2(\vec{P}^2 - \vec{P} \cdot \vec{q}/\vec{q}^2)/2$ . The above expression is valid both in vacuum and in medium and incorporates the  $\omega$  3-momentum dependence. Using the Lehmann representation for the propagators one finds

$$\begin{aligned} \Pi_\omega(P) = -2\frac{4}{3} IF g_{\omega\rho\pi}^2 \int_0^\infty d\omega \int_0^\infty d\omega' \frac{\omega + \omega'}{(P^0)^2 - (\omega + \omega')^2 + i\eta} \\ \times \int \frac{d^3q}{(2\pi)^3} S_\pi(\omega', \vec{P} - \vec{q}) \{v_1(q, P) S_\rho^T(q) \\ + v_2(q, P) [S_\rho^T(q) - S_\rho^L(q)]\}_{q^0=\omega} \end{aligned} \quad (4)$$

with  $S_\rho^{T,L} = -\frac{1}{\pi} \text{Im} D_\rho^{T,L}$ ,  $S_\pi = -\frac{1}{\pi} \text{Im} D_\pi$  denoting the  $\rho$  and  $\pi$  spectral functions, respectively. The  $\omega$  width follows from the imaginary part of the self-energy as  $\Gamma_{\omega \rightarrow \rho\pi}(P) = -\text{Im} \Pi_\omega(P)/P^0$ . In vacuum, free spectral functions for the pion and the  $\rho$  meson are utilized,

$$\begin{aligned} S_\pi^{\text{vac}}(\omega', \vec{q}) = \delta(\omega'^2 - \vec{q}^2 - m_\pi^2), \\ S_\rho^{\text{vac}}(\omega, \vec{q}) = -\frac{1}{\pi} \frac{\text{Im} \Pi_{\rho\pi\pi}^{\text{vac}}(q^2)}{|\omega^2 - \vec{q}^2 - M_\rho^2 - \Pi_{\rho\pi\pi}^{\text{vac}}(q^2)|^2}. \end{aligned} \quad (5)$$

The  $\rho \rightarrow \pi\pi$  self-energy is often approximated by reabsorbing the real part into the physical  $\rho$  mass,  $m_\rho^2 \equiv M_\rho^2 - \text{Re} \Pi_{\rho\pi\pi}^{\text{vac}}$ , and an imaginary part

$$\text{Im} \Pi_{\rho\pi\pi}^{\text{vac}}(q^2) = -\frac{g_{\rho\pi\pi}^2}{48\pi\sqrt{q^2}} (q^2 - 4m_\pi^2)^{\frac{3}{2}} \Theta(q^2 - 4m_\pi^2) \quad (6)$$

with  $g_{\rho\pi\pi} \simeq 6$  to obtain  $\Gamma_{\rho \rightarrow \pi\pi} = -\text{Im} \Pi_{\rho\pi\pi}^{\text{vac}}(q^2 = m_\rho^2)/m_\rho \simeq 150$  MeV. Here, we use the microscopic vacuum spectral function underlying our in-medium model [29], which describes the low-mass tail of the  $\rho$  resonance more accurately, incorporating an energy dependence of  $\text{Re} \Pi_{\rho\pi\pi}^{\text{vac}}$ . With  $g_{\omega\rho\pi} = 1.9/f_\pi$  ( $f_\pi = 92$  MeV) [28,30], one obtains  $\Gamma_{\omega \rightarrow \rho\pi} = 3.6$  MeV, i.e., about 1/2 of the total  $3\pi$  width (2/3 when including interference effects [31]). Using a schematic Breit–Wigner  $\rho$  spectral function,  $\Gamma_{\omega \rightarrow \rho\pi}(m_\omega)$  is reduced by approximately 30%. In Ref. [31] the partial  $\pi\rho$  width was found to be 2.8 MeV. Rescaling our  $g_{\omega\rho\pi}$  to obtain that value would entail an according 22% reduction of our in-medium widths reported below. Some of this would be recovered by medium effects of the accompanying increase in the direct  $3\pi$  channel.

### 2.2. $\rho$ and $\pi$ propagators in nuclear matter

Before proceeding to calculate the  $\omega$  meson width in nuclear matter caused by the dressing of the propagators in the  $\pi\rho$  loop,  $\Gamma_{\omega \rightarrow \pi\rho}^{\text{med}}$ , two comments are in order.

We first note that the unnatural-parity coupling in the  $\omega\rho\pi$  Lagrangian (1) implies transversality of any contribution to the  $\omega$  self-energy with at least one  $\omega\rho\pi$  vertex with an external  $\omega$  [26]. Thus, in-medium vertex corrections, as required to ensure transversality for the pion cloud of the  $\rho$  meson [29,32,33] (or chiral symmetry in the  $\sigma$  channel [34]), are not dictated here, but correspond to contributions to  $\omega N \rightarrow \pi N, \pi\pi N$  scattering unrelated to the anomalous decay process. We will not include these in the present work.

Second, at finite 3-momentum relative to the nuclear medium, the  $\rho$  propagator splits into transverse and longitudinal modes. At  $\vec{P} = \vec{0}$ , the  $\omega$  self-energy only depends on the transverse modes of the  $\rho$ , since the vertex function  $v_2$  in Eq. (3) vanishes. However, for  $\vec{P} \neq \vec{0}$ ,  $v_2$  becomes finite and proportional to  $S_\rho^T - S_\rho^L$ . This contribution turns out to be appreciable due to the splitting of the in-medium  $T$  and  $L$  modes of the  $\rho$  [29] within the kinematics of the  $\omega \rightarrow \rho\pi$  decay.


Let us turn to briefly reviewing the main ingredients to the evaluation of  $\Gamma_{\omega \rightarrow \pi\rho}^{\text{med}}$  from Eq. (4), which are the microscopic calculations of the in-medium pion and  $\rho$  propagators.

The pion spectral function is evaluated with standard  $P$ -wave nucleon–hole ( $NN^{-1}$ ) and Delta–hole ( $\Delta N^{-1}$ ) excitations [35,36]. The corresponding irreducible  $P$ -wave pion self-energy,

$$\begin{aligned} \Pi_\pi(q^0, \vec{q}; \varrho) \\ = \frac{(f_N/m_\pi)^2 F_\pi(\vec{q}^2) \vec{q}^2 [U_{NN} + U_{\Delta N} - (g'_{11} - 2g'_{12} + g'_{22}) U_{NN} U_{\Delta N}]}{1 - (f_N/m_\pi)^2 [g'_{11} U_{NN} + g'_{22} U_{\Delta N} - (g'_{11} g'_{22} - g'_{12}^2) U_{NN} U_{\Delta N}]}, \end{aligned} \quad (7)$$

is given by the Lindhard functions  $U_\alpha$  for the loop diagrams [37]; they include transitions between the two channels through short-range correlations represented by Migdal parameters  $g'$ . The  $\pi NN$  and  $\pi N\Delta$  coupling constants,  $f_N \simeq 1$  and  $f_\Delta/f_N \simeq 2.13$  (absorbed in the definition of  $U_{\Delta N}$ ), are determined from pion–nucleon and pion–nucleus reactions. Finite-size effects on the  $\pi NN$  and  $\pi N\Delta$  vertices are simulated via hadronic monopole form factors,

$$F_\pi(\vec{q}^2) = \Lambda_\pi^2 / (\Lambda_\pi^2 + \vec{q}^2). \quad (8)$$



**Fig. 1.** Left: differential decay momentum distribution of the  $\omega \rightarrow \rho\pi$  width (for  $m_\omega = 782$  MeV) in vacuum (dotted line) and at saturation density when dressing either the pion (short-dashed line) or the  $\rho$  (long-dashed line), or both (dash-dotted line), without spacelike  $\rho$  modes. The solid line includes spacelike  $\rho$ 's, where the two maxima beyond  $q \simeq 0.4$  GeV correspond to  $\Delta N^{-1}$  and  $NN^{-1}$  excitations ( $\Lambda_{\rho NN} = 0.3$  GeV). Right: Energy dependence of  $\Gamma_{\omega \rightarrow \rho\pi}$  at saturation density for different contributions as in the left panel.

Consistency with our model for the in-medium  $\rho$  discussed below dictates a soft cutoff,  $\Lambda_\pi = 0.3$  GeV, following from constraints of  $\pi N \rightarrow \rho N$  scattering data and the non-resonant continuum in nuclear photo-absorption [38] (e.g., with  $\Lambda_\pi = 0.5$  GeV one overestimates the measured  $\pi N \rightarrow \rho N$  cross section by a factor of  $\sim 2$ ). Especially the former probe similar kinematics of the virtual  $\pi NN$  vertex as figuring into  $\omega N \rightarrow \rho N$  processes. The Migdal parameters are  $g'_{11} = 0.6$  and  $g'_{12} = g'_{22} = 0.2$ .

The in-medium  $\rho$  spectral function is taken from Refs. [29, 39], which start from a realistic description of the  $\rho$  in free space (reproducing  $P$ -wave  $\pi\pi$  scattering and the pion electromagnetic form factor). The self-energy in nuclear matter contains two components: pions ( $NN^{-1}$ ,  $\Delta N^{-1}$ ) in the two-pion cloud,  $\Pi_{\rho\pi\pi}$ , and direct baryon resonance excitations in  $\rho N$  scattering,  $\Pi_{\rho BN^{-1}}$  (“ $\rho$ -sobars”). The latter have been evaluated using effective Lagrangians in hadronic many-body theory (in analogy to the pion) [29,40,41], including ca. 10 baryonic resonances. In  $\Pi_{\rho\pi\pi}$ , the in-medium pion propagator described above is supplemented with vertex corrections to preserve the Ward-Takahashi identities of the  $\rho$  propagator; it extends to finite 3-momentum of the  $\rho$  which is essential for the  $\pi\rho$  loop in  $\Pi_\omega$ . The total  $\rho$  self-energy is quantitatively constrained by nuclear photo-absorption and  $\pi N \rightarrow \rho N$  scattering, dictating the soft  $\pi NN(\Delta)$  form factor quoted above [38]. The resulting  $\rho$  spectral function in nuclear matter is substantially broadened, with a (non-Breit-Wigner) shoulder around  $M \simeq 0.5$  GeV; this is precisely the region where most of the free  $\omega \rightarrow \rho\pi$  decays occur. Note that spacelike parts of the  $\pi$  and  $\rho$  spectral functions (i.e., with negative 4-momenta squared,  $q^2 < 0$ ) contribute to  $\Gamma_{\omega \rightarrow \pi\rho}^{\text{med}}$ ; they correspond to  $t$ -channel exchanges in  $\omega N$  scattering (e.g.,  $\rho$  exchange in  $\omega N \rightarrow \pi N^*$ ). For the pion these are encoded in the Lindhard functions in the self-energy, Eq. (7). For the  $\rho$  they also turn out to be dominated by the low-lying  $P$ -wave  $\rho$ -sobars,  $\rho NN^{-1}$  and  $\rho\Delta N^{-1}$ . The latter is well constrained by nuclear photo-absorption ( $f_{\rho\Delta N}^2/4\pi = 16.2$ ,  $\Lambda_{\rho\Delta N} = 0.7$  GeV), but the purely spacelike  $NN^{-1}$  mode (generating Landau damping of the exchanged  $\rho$ ) is not. An analysis of  $\rho$  photo-production cross sections,  $\gamma p \rightarrow \rho p$  [42], gave indications for a rather soft form factor,  $\Lambda_{\rho NN} \simeq 0.6$  GeV ( $f_{\rho NN}^2/4\pi = 6.0$ ), but it might be as soft as the  $\pi NN$  form factor in the pion cloud of the  $\rho$ . This needs to be investigated in future analysis of  $\omega N$  scattering data. Here, we bracket the uncertainty by varying  $\Lambda_{\rho NN} = 0.3\text{--}0.6$  GeV and  $g'_{NN} = 0\text{--}0.6$ . We find that the  $\omega$  coupling to spacelike  $S$ -wave rhosobars (e.g.,  $N^*(1520)N^{-1}$ , corresponding to  $\omega N \rightarrow \pi N^*(1520)$ ) is already much less important.

In addition to modifications of the  $\pi\rho$  cloud, pion dressing in the direct  $\omega \rightarrow \pi\pi\pi$  channel and  $\omega N^*N^{-1}$  excitations occur.

The direct  $3\pi$  decay has considerable phase space in vacuum, and thus we expect its in-medium modification to be smaller than for the  $\pi\rho$  channel, especially if the latter dominates in vacuum and with our soft form factors for the pion dressing; for  $\Lambda_{\pi NN(\Delta)} = 0.3$  GeV we estimate  $\Gamma_{\omega \rightarrow 3\pi}^{\text{med}}(Q_0) < 20$  MeV based on recent work in Ref. [43]. For the  $\omega$ -sobars, e.g.,  $N^*(1535)$ ,  $N^*(1520)$  or  $N^*(1650)$  [19,21], we cannot simply adopt the couplings from the literature, since they were adjusted to fit  $\omega N$  scattering data without the inclusion of  $\pi\rho$  cloud effects. If the latter are present, the direct-resonance contributions need to be suppressed to still describe  $\omega N$  scattering, and thus their contribution to the in-medium width will be (much) smaller than in Refs. [19,21].

### 3. $\omega$ width in nuclear matter

Let us first examine the differential distribution of the  $\omega$  width,  $d\Gamma_\omega/dq$ , over the center-of-mass decay momentum,  $|\vec{q}|$ , of the  $\pi$  and  $\rho$  spectral functions, recall Eq. (4). In vacuum, the fixed pion mass uniquely determines the (off-shell)  $\rho$  mass ( $M$ ) at given  $q$ . The maximum of the distribution occurs at  $q_{\text{max}} \simeq 0.2$  GeV, corresponding to  $M \simeq 0.5$  GeV (see Fig. 1 left). Consequently, the enhancement of the in-medium  $\rho$  spectral function around this mass strongly increases the phase space and thus  $\Gamma_{\omega \rightarrow \pi\rho}^{\text{med}}$ . A similar, albeit less pronounced effect is caused by the in-medium pion. A further remarkable increase in decay width is generated by spacelike  $\rho$ -sobars above  $q \simeq 0.4$  GeV, which, for a free pion ( $m = m_\pi$ ), marks the  $M = 0$  boundary. The low-lying collective excitations are sensitive to the  $\rho NN$  form factor. For a conservative choice of  $\Lambda_{\rho NN} = 0.3$  GeV, about 40% of the in-medium  $\omega$  width is generated by the spacelike  $\rho$  modes.

The energy dependence of  $\Gamma_{\omega \rightarrow \pi\rho}^{\text{med}}$  is rather pronounced (Fig. 1 right), a remnant of the (nominal) vacuum  $\pi\rho$  threshold together with the  $\vec{q}^2$  dependence of the  $\omega\pi\rho$  vertex. The density dependence of  $\Gamma_{\omega \rightarrow \pi\rho}^{\text{med}}$  (Fig. 2 left) exhibits significant nonlinearities. At normal nuclear matter density, the dominant uncertainty is due to the  $\rho NN$  form factor, quantified as  $\Gamma_{\omega \rightarrow \pi\rho}^{\text{med}} = 130\text{--}200$  MeV.

The 3-momentum dependence of the on-shell  $\omega$  width (i.e., for  $p^2 = (P^0)^2 - \vec{P}^2 = m_\omega^2$ ), relative to the nuclear rest frame, turns out to be moderate (Fig. 2 right), as generally expected from cloud effects with soft form factors counter-acting the momentum dependence of the vertices. A fair agreement with CBELSA/TAPS data [16] is found, apparently preferring the lower values of  $\Lambda_{\rho NN}$ , leaving room for (smaller) contributions from direct  $3\pi$  and interference terms, as well as from  $\omega$ -sobars which are expected to come in at higher 3-momenta [21]. However, we recall the somewhat larger in-medium width of  $\sim 200$  MeV found by CLAS [15].




Fig. 2. Left: Density dependence of the  $\omega \rightarrow \rho\pi$  width at  $P^0 = m_{\omega}$ ,  $\vec{P} = \vec{0}$ , and for different  $\rho NN$  form factors and short-range correlations. Right: Three-momentum dependence of  $\Gamma_{\omega \rightarrow \rho\pi}$  at saturation density for on-shell  $\omega$  mesons ( $P^2 = m_{\omega}^2$ , i.e.,  $E_{\omega}^2 = m_{\omega}^2 + P_{\omega}^2$ ), compared to CBELSA/TAPS data [16].

In the very recent work of Ref. [43], the total  $\omega$  width in nuclear matter is computed with similar methods. At  $\rho_N = \rho_0$  and  $\vec{P} = \vec{0}$ ,  $\Gamma_{\omega}^{\text{med}} = 129 \pm 10$  MeV is reported, predominantly due to the  $\rho\pi$  cloud modification and with a more pronounced momentum dependence. The  $\rho$  spectral function employed in there exhibits a factor of  $\sim 2$  less broadening than in our input, while the pion modifications are stronger due to a harder  $\pi NN$  form factor. We recall that the latter is fixed in our approach as part of the quantitatively constrained  $\rho$  spectral function. It was also argued in Ref. [43] that medium effects in interference terms of  $3\pi$  final states from direct  $3\pi$  and  $\rho\pi$  decays, which we neglected here, are small. Thus both our work and Ref. [43] identify the  $\pi\rho$  cloud as the main agent for the  $\omega$ 's in-medium broadening, albeit with some differences in the partitioning into  $\pi$  and  $\rho$  modifications, and in the 3-momentum dependence.

#### 4. Summary

We have studied the width of the  $\omega$  meson in cold nuclear matter focusing on the role of its  $\pi\rho$  cloud. We have employed hadronic many-body theory utilizing pion and  $\rho$  propagators evaluated with the same techniques, constrained and applied previously in both elementary and heavy-ion reactions. The low-mass shoulder in the in-medium  $\rho$  spectral function, together with spacelike contributions in the  $\pi\rho$  intermediate states, induce large effects, along with non-linear density dependencies, not captured in previous calculations based on  $T$ - $Q$  approximations. For an  $\omega$  at rest at saturation density, we find  $\Gamma_{\omega}^{\text{med}} = 130$ – $200$  MeV, where the uncertainty is largely due to the  $\rho NN$  vertex form factor which could not be accurately constrained before from  $\rho$  properties alone. Together with a rather weak 3-momentum dependence of the on-shell  $\omega$  width, our calculations compare favorably with data from recent absorption experiments. The present uncertainties can be reduced by systematic analyses of vacuum  $\omega$  scattering data (similar to the  $\pi NN$  form factor in the  $\rho$  cloud), where also contributions from direct  $3\pi$  couplings and  $\omega N$  resonances ( $\omega$ -sobars) need to be included. Work in this direction is in progress. The emergence of a large  $\omega$  width from  $\rho$  and pion propagators in nuclear matter is encouraging, and corroborates the quantum many-body approach as a suitable tool to assess the properties of hadrons in medium.

#### Acknowledgements

This work has been supported by the US National Science Foundation under grant No. PHY-1306359, the Humboldt Foundation, the BMBF (Germany) under project No. 05P12RFFCQ, the

Ministerio de Economía y Competitividad (Spain) under grant FPA2011-27853-C02, the Centro Nacional de Física de Partículas, Astropartículas y Nuclear (Consolider-Ingenio 2010) and the EU Integrated Infrastructure Initiative Hadron Physics Project under Grant Agreement No. 227431.

#### References

- [1] I. Tserruya, in: R. Stock (Ed.), *Relativistic Heavy-Ion Physics*, in: Landolt Börnstein (Springer), New Series, vol. 1/23A, 2010, 4-2, arXiv:0903.0415 [nucl-ex].
- [2] H.J. Specht, for the NA60 Collaboration, *AIP Conf. Proc.* 1322 (2010) 1.
- [3] F. Geurts, et al., STAR Collaboration, *Nucl. Phys. A* 904–905 (2010) 217c.
- [4] R. Rapp, J. Wambach, H. van Hees, in: R. Stock (Ed.), *Relativistic Heavy-Ion Physics*, in: Landolt Börnstein (Springer), New Series, vol. 1/23A, 2010, 4-1, arXiv:0901.3289 [hep-ph].
- [5] R. Rapp, *PoS CPOD 2013* (2013) 008.
- [6] S. Leupold, V. Metag, U. Mosel, *Int. J. Mod. Phys. E* 19 (2010) 147.
- [7] F. Riek, R. Rapp, Y. Oh, T.-S.H. Lee, *Phys. Rev. C* 82 (2010) 015202.
- [8] G.M. Huber, et al., TAGX Collaboration, *Phys. Rev. C* 68 (2003) 065202.
- [9] M.H. Wood, et al., CLAS Collaboration, *Phys. Rev. C* 78 (2008) 015201.
- [10] D. Trnka, et al., CBELSA/TAPS Collaboration, *Phys. Rev. Lett.* 94 (2005) 192303.
- [11] M. Naruki, et al., E325 Collaboration, *Phys. Rev. Lett.* 96 (2006) 092301.
- [12] M. Nanova, et al., CBELSA/TAPS Collaboration, *Phys. Rev. C* 82 (2010) 035209.
- [13] M. Kaskulov, E. Hernandez, E. Oset, *Eur. Phys. J. A* 31 (2007) 245.
- [14] T. Ishikawa, et al., LEPS Collaboration, *Phys. Lett. B* 608 (2005) 215.
- [15] M.H. Wood, et al., CLAS Collaboration, *Phys. Rev. Lett.* 105 (2010) 112301.
- [16] M. Kotulla, et al., CBELSA/TAPS Collaboration, *Phys. Rev. Lett.* 100 (2008) 192302.
- [17] F. Klingl, N. Kaiser, W. Weise, *Nucl. Phys. A* 624 (1997) 527.
- [18] M. Post, U. Mosel, *Nucl. Phys. A* 688 (2001) 808.
- [19] M.F.M. Lutz, G. Wolf, B. Friman, *Nucl. Phys. A* 706 (2002) 431; M.F.M. Lutz, G. Wolf, B. Friman, *Nucl. Phys. A* 765 (2006) 431 (Erratum).
- [20] S. Zschocke, O.P. Pavlenko, B. Kämpfer, *Phys. Lett. B* 562 (2003) 57.
- [21] P. Muehlich, V. Shklyar, S. Leupold, U. Mosel, M. Post, *Nucl. Phys. A* 780 (2006) 187.
- [22] A.T. Martell, P.J. Ellis, *Phys. Rev. C* 69 (2004) 065206.
- [23] F. Eichstaedt, S. Leupold, U. Mosel, P. Muehlich, *Prog. Theor. Phys. Suppl.* 168 (2007) 495.
- [24] T.E. Rodrigues, J.D.T. Arruda-Neto, *Phys. Rev. C* 84 (2011) 021601.
- [25] S. Ghosh, S. Sarkar, *Eur. Phys. J. A* 49 (2013) 97.
- [26] M. Wachs, PhD thesis, TU Darmstadt, 2000, <http://tuprints.ulb.tu-darmstadt.de/epda/000050/>.
- [27] F. Riek, J. Knoll, *Nucl. Phys. A* 740 (2004) 287.
- [28] P. Jain, R. Johnson, U.G. Meissner, N.W. Park, J. Schechter, *Phys. Rev. D* 37 (1988) 3252.
- [29] M. Urban, M. Buballa, R. Rapp, J. Wambach, *Nucl. Phys. A* 641 (1998) 433.
- [30] F. Klingl, N. Kaiser, W. Weise, *Z. Phys. A* 356 (1996) 193.
- [31] D.G. Gudino, G.T. Sanchez, *Int. J. Mod. Phys. A* 27 (2012) 1250101.
- [32] G. Chanfray, P. Schuck, *Nucl. Phys. A* 555 (1993) 329.
- [33] M. Herrmann, B.L. Friman, W. Nörenberg, *Nucl. Phys. A* 560 (1993) 411.
- [34] H.C. Chiang, E. Oset, M.J. Vicente-Vacas, *Nucl. Phys. A* 644 (1998) 77.
- [35] E. Oset, H. Toki, W. Weise, *Phys. Rep.* 83 (1982) 281.
- [36] A.B. Migdal, E.E. Saperstein, M.A. Troitsky, D.N. Voskresensky, *Phys. Rep.* 192 (1990) 179.

- [37] E. Oset, P. Fernandez de Cordoba, L.L. Salcedo, R. Brockmann, *Phys. Rep.* 188 (1990) 79.
- [38] R. Rapp, J. Wambach, *Adv. Nucl. Phys.* 25 (2000) 1.
- [39] R. Rapp, M. Urban, M. Buballa, J. Wambach, *Phys. Lett. B* 417 (1998) 1.
- [40] M. Urban, M. Buballa, R. Rapp, J. Wambach, *Nucl. Phys. A* 673 (2000) 357.
- [41] R. Rapp, J. Wambach, *Eur. Phys. J. A* 6 (1999) 415.
- [42] F. Riek, R. Rapp, T.-S.H. Lee, Y. Oh, *Phys. Lett. B* 677 (2009) 116.
- [43] A. Ramos, L. Tolos, R. Molina, E. Oset, arXiv:1306.5921v3 [nucl-th].