

A metacaspase of *Trypanosoma brucei* causes loss of respiratory competence and clonal death in the yeast *Saccharomyces cerevisiae*

Alexander Szallies^a, Bruno K. Kubata^c, Michael Duszenko^{a,b,*}

^aVerfügungsgebäude der Universität Tübingen, Auf der Morgenstelle 15, 72076 Tübingen, Germany

^bPhysiologisch-chemisches Institut, Universität Tübingen, Hoppe-Seyler-Strasse 4, 72076 Tübingen, Germany

^cDepartment of Molecular Behavioral Biology, Osaka Bioscience Institute, Osaka 565-0874, Japan

Received 14 February 2002; revised 11 March 2002; accepted 15 March 2002

First published online 27 March 2002

Edited by Vladimir Skulachev

Abstract Metacaspases constitute a new group of cysteine proteases homologous to caspases. Heterologous expression of *Trypanosoma brucei* metacaspase *TbMCA4* in the budding yeast *Saccharomyces cerevisiae* resulted in growth inhibition, mitochondrial dysfunction and clonal death. The metacaspase orthologue of yeast, *ScMCA1* (*YOR197w*), exhibited genetic interaction with *WWMI* (*YFL010c*), which encodes a small WW domain protein. *WWMI* overexpression resulted in growth arrest and clonal death, which was suppressed by concomitant overexpression of *ScMCA1*. GFP-fusion reporters of *WWMI*, *ScMCA1* and *TbMCA4* localized to the nucleus. Taken together, we suggest that metacaspases may play a role in nuclear function controlling cellular proliferation coupled to mitochondrial biogenesis. © 2002 Published by Elsevier Science B.V. on behalf of the Federation of European Biochemical Societies.

Key words: Metacaspase; Cysteine protease; WW domain; Mitochondrion; *Saccharomyces cerevisiae*; *Trypanosoma brucei*

1. Introduction

Cysteiny proteases of Clan CD [1] serve diverse functions in various organisms. Legumains are involved in lysosomal degradation, caspases control apoptosis, a form of programmed cell death (PCD), and modulate signal transduction, Gpi8 proteins are a core component of the endoplasmic reticulum glycosylphosphatidylinositol (GPI) transamidase complex, gingipains are virulence factors of pathogenic bacteria [1] and separins trigger anaphase progression [2]. All these proteases share a typical secondary and, where known, tertiary core structure composed of several consecutive β -strands and α -helices around a catalytic dyad of histidine and cysteine constituting the active center [1,3,4]. Metacaspases have been denoted as putative Clan CD peptidases by iterative sequence analyses and seem to be homologous to caspases and paracaspases [5]. They occur in virtually all non-metazoan eukary-

otic lineages and might be functional homologues of metazoan caspases, which are primarily involved in PCD [5].

PCD is thought to be a product of the coevolution of the eukaryotic cell and its mitochondrial endosymbiont [6–10]. It is anticipated to occur not only in metazoans, but also in plants and unicellular eukaryotes [10–12]. Conclusive genetic evidence for the existence of PCD in plants and in *Dictyostelium* was presented [13,14]. Interestingly, plant PCD could be inhibited by expression of various metazoan antiapoptotic genes such as p35 of baculovirus [15], which acts as a substrate upon diverse caspases [16], and Bcl-xL, a protector of mitochondrial function and integrity [13].

On unicellular eukaryotes, such as the yeast *Saccharomyces cerevisiae*, cellular death can be inflicted by expression of diverse PCD mediating metazoan gene products. Bax for instance caused clonal death of yeast by perturbation of their mitochondria [17,18].

We performed studies on the metacaspases of *S. cerevisiae* and *Trypanosoma brucei* in budding yeast to take advantage of the versatility with which gene function can be assessed in this model organism. Recent comprehensive two-hybrid screens of yeast protein interactions [19,20] revealed interactions of the yeast metacaspase (Mca1p, YOR197p) with several proteins, amongst others Wwm1p (YFL010p) [19].

WWMI (*YFL010c*) encodes a protein of 211 amino acids and is characterized by a high glycine content (14%) and a high hydrophilicity. It was categorized as a potential hydrophilin, which are often involved in osmotic stress response [21]. Wwm1p contains a 40 amino acid N-terminal domain with two signature tryptophan (W) residues (WW domain), a widespread module mediating protein–protein interactions in a variety of cellular processes [22].

2. Materials and methods

2.1. Cloning of *TbMCA1–5*

Genes of *T. brucei* metacaspases *TbMCA1–5* were cloned by reverse transcription based polymerase chain reaction (RT-PCR) as described previously [23] using *Pwo* polymerase (Roche, Mannheim, Germany) into pBluescript KS⁺ (Stratagene, La Jolla, CA, USA). Primer sequences were TGT GTG GAC TGA TCA CTT CCG ACT GCG C (*TbMCA1* sense), GTG TCA ACA CCG ACT TAT GCA CAT (*TbMCA1* antisense), TAA TTT TCC ACG CAC ATC CGC AGA CAG TAG (*TbMCA2* sense), GTG GTA GAA GCT CTT GCT ACA ACA (*TbMCA3* sense), AAC CCT TCT GCA GCT CCC AGG CAC (*TbMCA2/3* antisense), GCG AAT TCC TGA AAA CAT GGG AGG C (*TbMCA4* sense), CAC ATC TTC ATT CCA GGC AAA GAG (*TbMCA4* antisense), AGC ATT TAA GAT AAA ATA CAG AAA GC (*TbMCA5* sense), GAG GCA AAT CCT GCA

*Corresponding author. Fax: (49)-7071-295565.

E-mail address: michael.duszenko@uni-tuebingen.de (M. Duszenko).

Abbreviations: GFP, green fluorescent protein; GPI, glycosylphosphatidylinositol; PCD, programmed cell death; RT-PCR, reverse transcription based polymerase chain reaction; WW domain, domain with two conserved tryptophan (W) residues

CCA GGC TTG (*TbMCA5* antisense). Sequencing was carried out by GATC GmbH (Konstanz, Germany) according to the Sanger-dideoxynucleotide method. Sequences have been submitted to EMBL (accession numbers AJ437301, AJ437302, AJ437303, AJ437304, AJ437305).

2.2. Metacaspase sequence retrieval and analysis

The sequence of *Rhodobacter sphaeroides* was obtained from the *Rhodobacter* genome project by the University of Texas (<http://mmg.uth.tmc.edu/sphaeroides>), designated putative ORF 'OR0035'. The sequence of *Geobacter sulfurreducens* was obtained from the TIGR database (<http://www.tigr.org>), designated contig 2947. Sequences of fission and budding yeast as well as of *Arabidopsis* were obtained from the NCBI database (<http://www.ncbi.nlm.nih.gov>; accession numbers AAG38593, Z75105, AAC24380, AAD11574 and CAB79420). Metacaspase sequences were aligned with DNAMAN software (version 4.15, Lynnon BioSoft). Each one of the primary sequences was calculated for putative secondary structures, the alignment was fit manually for conserved secondary structures.

2.3. Yeast expression constructs

For expression in yeast several vectors were used with the desired genes under control of inducible promoters. Vector pRS416 (*cen*, *URA3*, *MET17* (formerly *MET25*) promoter) [24], YEp52 (2 μ , *LEU2*, *GAL10*) as well as pYES2 (2 μ , *URA3*, *GAL1* promoter) were used. *TbMCA1–5* were cloned into pRS416, YEp52 and pYES2 (only *TbMCA4*). *ScMCA1* was cloned from *Schizosaccharomyces pombe* by RT-PCR using primers CAT GAC AAG CTT CCA TGA GCT ACA ACT CCA ATT CTT ATA AC (sense) and GAC CTG AAG CTT CTA TAA AAC CAT GGC AAG ATT CAT GTC (antisense) into pRS416. *ScMCA1* was cloned from BY4742 wild-type strain genomic DNA using primers GGT CAG AAG CTT CCA TGA AGA TGA GCC TCG AAG TTT ATC TAA AC (sense) and GCA TAC AAG CTT ATT TCT ACA TAA TAA ATT GCA GAT TTA C (antisense) into pRS416, YEp52 and pYES2. *ScMCA1*^{DN-terminus} was made accordingly using sense primer CAG ATA AGC TTA CCA TGG CTA GAA AGG CTT TGA TTA TCG GTA (initiator ATG bold, thus replacing amino acids GR^{155/156} with MA). *WWM1* was amplified with primers CAG TAC AAG CTT ATG GCT CAA AGT AAA AGT CCT CC (sense) and ACA GTC AAG CTT TTA AAA GCT ACT ACC GTC AAA TCC (antisense) and cloned into pRS416. Mouse Bel-X_L (on expression vector, courtesy of Dr. S. Kugler, Göttingen, Germany) was amplified by CAG TCC AAG CTT ATG TCT CAG AGC AAC CGG GAG CTG (sense) and AGC TGC AAG CTT TCA CTT CCG ACT GAA GAG TGA GCC (antisense) into pRS416 and YEp52. Baculovirus p35 (on expression vector, courtesy of Dr. S. Kugler, Göttingen) was amplified by primers GCA ATA ATG TGT GTA ATT TTT CCG GTA GAA ATC (sense) and AGT ACT TAT TTA ATT GTG TTT AAT ATT ACA TTT TTG TTG (antisense) into pRS416 and YEp52.

Green fluorescent protein (GFP) fusion constructs: a GFP gene (ECFP variant, on expression vector, courtesy of Dr. S. Kugler, Göttingen, Germany) was amplified with primers CGC TAA GCT TAT GGT GAG CAA GGG CGA GGA GCT G (sense) and CCG TTA CTT GTA CAG CTC GTC CAT GCC GAG (antisense) into pBluescript KS⁺. Likewise, PCR products of *ScMCA1*, *TbMCA4* and *WWM1* were cloned into pBluescript KS⁺ using sense primers listed above and antisense primers GCT ACA GAA GCT TCA TAA TAA ATT GCA GAT TTA CGT CAA, CAG AAG CTT CAG GAG CGT CCG TTC CTG CG and CAA GCT TAA AGT CAC TAC CGT CAA ATC CAT C, respectively. Subcloned products were cut with *HindIII* (site introduced with primers) and ligated. A second round of PCR with respective sense primers and the GFP antisense primer amplified the GFP fusions that were cloned into pRS416.

Similarly, site-directed mutagenesis was carried out with *ScMCA1* and *TbMCA4*. In a first round of PCR, 5' and 3' overlapping fragments of the respective gene were amplified with the primers listed above and primers introducing nucleotide changes. The mutagenesis primers were complementary to each other. In a second round of PCR, the 5' and 3' overlapping fragments together served as template (1:100) for amplification of full length gene using the primers listed above, carrying the defined nucleotide changes (verified by sequencing). Mutagenesis primers were (complementary antisense primer omitted, changed codon in bold) *ScMCA1*^{C176A}: CAA CTG CGT GGT GCT ATC AAT GAT GCT; *ScMCA1*^{H241Y}: CAT TAT TCT

GGA TAT GGT GGC CAA ACT; *ScMCA1*^{C297A}: TTG TTT GAC TCT GCT CAT TCG GGT ACA; *TbMCA4*^{C98A}: CAG CTC TCT GGT GCT GCC CAT GAC ATT ATG; *TbMCA4*^{H164A}: CAC TAC TCG GGT GCC GGT ACG CGT GCT; and *TbMCA4*^{C218A}: GCT GTG TTC GAC GCC TCG CAC TCC GGT.

2.4. Yeast strains, growth and media

S. cerevisiae strains employed were BY4742 (S288C background) and derivatives thereof carrying a kanamycin resistance cassette replacing and thereby deleting defined ORFs (*ScMCA1*, *WWM1-Δ* strains) (Euroscarf, Frankfurt/M., Germany).

Yeast cells were grown at 28°C with 2% agar to solidify medium on plates or agitated at 170 rpm in case of liquid cultures. Yeast strains were propagated either with YPD or CM medium. YPD contained 1% yeast extract, 2% bacto-peptone and 2% D-glucose. CM contained 0.67% nitrogen base without amino acids and nucleotide bases (Difco, Detroit, MI, USA), 0.5% ammonium sulfate and 0.2% amino acids (all except those needed for marker selection at equal proportion of weight). CM-S was CM medium lacking methionine and cysteine. Yeast transformations were carried out as described [25].

2.5. Clonogenicity assays

Yeast culture harboring expression vectors were grown overnight in non-inducing selective media and then inoculated into inducing selective media (CM-S with or without 2% galactose as carbon source), starting off at OD₆₀₀ of 0.2–0.3. At defined time points samples were withdrawn from the culture. Its OD₆₀₀ was measured (Pharmacia spectrophotometer, in the range of 0.1–0.15) and an aliquot of diluted cell suspension was plated on non-inducing selective plates – 120 μ l of cell suspension 0.1 OD₆₀₀ diluted 1/400 in two steps. This amount was found to correspond to 500 cells as determined by counting using a hemocytometer (Neubauer). After 2–7 days of incubation the final number of colonies was counted. Experiments generally were run in parallel. All series of experiments were repeated at least three times.

2.6. TCC overlay assay

Tetrazolium red (2,3,5-triphenyltetrazolium chloride, Sigma, Germany) was solved (0.5%) in 100 mM Tris-HCl pH 7.0 containing 1% agar; 20 ml were poured on culture plates. This procedure has been described previously [26].

2.7. Microscopy

Cells (200 μ l of OD₆₀₀ 0.5–1) were incubated in water with an appropriate amount of dye (bisbenzimidazole, 1 μ g in 1 μ l stock solution, MitoTracker[®] CMXRos, 1 μ l 20 μ M), washed in water, mounted on slides and viewed using an Olympus BH2-RFCA or a Zeiss Axioplan microscope equipped with Nomarski optics and fluorescence devices.

3. Results

3.1. Sequence analysis of metacaspases

Amino acid sequences of several metacaspases were aligned with respect to putative common secondary structures (as shown as supplementary information on the net: <http://www.pci.chemie.uni-tuebingen.de/midu/miduhome.html>).

This alignment includes the single metacaspases of *Saccharomyces* and *Schizosaccharomyces*, four *Arabidopsis* orthologues and orthologues of *R. sphaeroides* and *G. sulfurreducens*. The two latter α - and δ -proteobacterial sequences are the only bacterial sequences with definite homology to the eukaryotic ones. The genes of the five *T. brucei* metacaspases *TbMCA1–5* were cloned from bloodstream form trypanosomes by RT-PCR, using sequences provided by the TIGR database (www.tigr.org).

The block of highest sequence homology among metacaspases corresponds to the large subunit of caspases [3] carrying the catalytic dyad histidine and cysteine residues, N-terminally flanked by short β -strands as characteristic for Clan CD cysteinyl proteases [1]. As in caspases [3] or gingipain [4] the same

consecutive secondary structures are predicted to occur in all metacaspases alike. In accordance with Clan CD peptidases the catalytic dyad histidine and cysteine residues are well conserved.

Apart from the high homology region, metacaspases show only a limited similarity to each other in the last third of their sequences (see web site as stated above). This region seems to be analogous to the small subunit of caspases [3]. The region that separates both main blocks of homology is variable in length and becomes particularly large in metacaspases of plants ('latex-abundant RNA protein' [27], 'type II metacaspases' [5], which are not included in our alignment). Another common feature of the metacaspases presented here is their proline-rich N-terminus (see Fig. 1); the aforementioned metacaspases lack this N-terminus.

Notably, in *TbMca1p* a tyrosine substitutes for the putative catalytic histidine and serine substitutes for the putative catalytic cysteine. However, a cysteine residue is located immediately adjacent, as is also the case in *TbMca4p*. As in other Clan CD peptidases some conserved glycine residues can be found which may constitute 'oxanion holes', facilitating nucleophilic attack by the catalytic dyad of histidine and cysteine. Residues which may specify the P1 scissile bond amino acid (for which Clan CD members are always highly specific) may be constituted by two conserved aspartate residues. This would imply that metacaspases are specific for a basic P1 amino acid such as arginine or lysine.

An extraordinary feature of metacaspases is the occurrence of a conserved cysteine between the first strand and helix motifs (see web site as stated above). Superposition of these predicted secondary structures upon the known crystal structures of caspase-1 and gingipain [3,4] puts this conserved cysteine in spatial proximity to the catalytic dyad.

3.2. Deletion or overexpression of *ScMCA1* is not detrimental to mitochondrial function

Assuming that metacaspases were introduced into eukaryotes by their mitochondrial progenitor (see Section 4), we expected a physiological function of metacaspases in mitochondrial biogenesis. Since the yeast *S. cerevisiae* is unique in its ability to live with dysfunctional mitochondria, this model eukaryotic organism seems apt to study a putative mitochondrial biogenesis function of metacaspases. However, a yeast strain deleted in its only endogenous metacaspase gene, *ScMCA1* (*mca1Δ* strain), did not show any petite phenotype that is characteristic for compromised essential mitochondrial functions. In addition, no slow growth phenotype, no alteration in clonogenicity rates or impairment of respiratory competence was noted under various regimes tested such as different temperatures, different carbon and nitrogen sources. Likewise, overexpression of *ScMCA1* using different plasmids and promoters (cen *MET17*, 2 μ *GAL1110*) did not lead to any obvious phenotype (data not shown).

3.3. Expression of *T. brucei* metacaspase *TbMCA4* induces a petite phenotype in yeast

Five metacaspase genes of *T. brucei* (*TbMCA1–5*) were heterologously expressed in yeast, as well as the only one of *S. pombe* (*SpMCA1*). Of these, only expression of *TbMCA4* brought about phenotypic consequences. Strains expressing *TbMCA4* were significantly retarded in growth (Fig. 1A). Furthermore, clonogenicity of *TbMCA4* expressing cultures di-

minished gradually but irreversibly over time of expression (Fig. 1C). Clonogenicity loss was preceded by loss of respiratory competence (Fig. 1D,E). Even a short induction of *TbMCA4* was sufficient to abolish respiratory competence completely (Fig. 1D).

The phenotypes induced by *TbMCA4* expression in yeast were completely lost when the putative catalytic dyad residues histidine-164 and cysteine-218 were both independently mutated to alanine. Yeast expressing the respective alleles of *TbMCA4* was able to grow as the control (Fig. 1A). This result clearly demonstrates that the effects of *TbMca4p* depend on the putative catalytic dyad residues. Therefore, the assumption that *TbMca4p* is a cysteine proteinase inferred by sequence homology to Clan CD proteases is confirmed.

In addition, exchange of the other conserved cysteine amongst metacaspases, cysteine-98 in *TbMca4p* (to alanine, the adjacent cysteine was changed to alanine simultaneously), rendered *TbMCA4* inactive as well (Fig. 1A).

To monitor abundance and shape of mitochondria in *TbMCA4* expressing yeast mitochondrial staining was performed. Staining with MitoTracker Green FM dye revealed that there was no significant difference in mitochondrial abundance or shape in *TbMCA4* expressing yeast compared with control cells. However, staining with MitoTracker CMXRos was significantly weaker in *TbMCA4* expressing yeast than in the control cells (see web site as stated above). Accumulation of MitoTracker CMXRos within mitochondria is dependent on free thiols and thus indicative of metabolically functional organelles.

3.4. Metacaspase-GFP fusion constructs localize to the nucleus

To determine intracellular localization of *ScMca1p* and *TbMca4p* in yeast, the gene for GFP was fused to the respective 3' ends of the respective metacaspase genes and expressed in yeast cells upon induction from pRS416.

The *ScMca1*-GFP fusion apparently localized to the nucleus, as did the *TbMca4*-GFP fusion (Fig. 2B,C). Apart from nuclear localization, *ScMca1*-GFP fluorescence could also be monitored throughout the cell (Fig. 2B), this localization was indistinguishable in wild-type and Δ *mca1* yeast (data not shown).

3.5. Metacaspase *ScMCA1* exhibits genetic interactions with *WWM1*

Protease function generally is expected to be controlled tightly. Thus it was not surprising that *ScMCA1* overexpression did not result in a phenotype as seen by heterologous expression of *TbMCA4*. Considering that *ScMca1p* function might be controlled by endogenous activator or inhibitor proteins, we focused on a reported physical interaction of *ScMca1p* with *Wwm1p* (YFL010p) [19].

Interestingly, overexpression of *WWM1* in wild-type yeast (pRS416) caused severe growth retardation and loss of clonogenicity, similar to the effects exerted by *TbMCA4* (Fig. 1B,C). These effects were independent of *ScMCA1*, since a *mca1Δ* strain was also susceptible to *WWM1* overexpression (Fig. 1B, data not shown). A *wwm1Δ* strain was likewise susceptible to *WWM1* pRS416-based expression but otherwise did not reveal any phenotype considering growth, clonogenicity rates and respiratory competence as compared to the respective wild-type strain (data not shown).

Simultaneous overexpression of *WWM1* and *ScMCA1* re-




Fig. 1. Ten-fold dilution series of wild-type yeast spotted on CM/glucose agar plates. A: Vector based expression (pRS416) of *TbMCA4*. Left panel: control (promoter shut off). Right panel: expression of *TbMCA4* and mutant alleles. First row: empty vector (pRS416), second row: *TbMCA4*, third row: *TbMCA4^{C98A}*, fourth row: *TbMCA4^{H1241A}*, fifth row: *TbMCA4^{C218A}*. B: Vector based expression (pRS416) of *WWMI*. Left panel: control (promoter shut off). Right panel: expression of *WWMI*. First row: empty vector in wild-type strain, second row: *WWMI* in wild-type strain, third row: empty vector in *mca1 Δ strain, fourth row: *WWMI* in *mca1 Δ strain. C: Upper panel: growth of wild-type yeast in liquid CM/glucose medium expressing *TbMCA4* (○), *WWMI* (▲) and control (■). Lower panel: clonogenicity (colonies counted) of wild-type yeast cultures expressing *TbMCA4* (○), *WWMI* (▲) and control (■), corresponding to growth shown in upper panel. D: Outgrown colonies of wild-type yeast culture previously having expressed *TbMCA4* (same experiment as depicted in Fig. 2C) divided into respiration competent (white parts of bars) and respiration incompetent (black parts of bars) ones as judged by TCC-overlay assay. E: Example of TCC overlay assay with plate of outgrown colonies of wild-type yeast culture previously expressing *TbMCA4* for 2 h (independent experiment from that shown in Fig. 2C,D). Bigger red staining colonies are respiration competent, smaller white colonies are respiration incompetent.**




Fig. 2. Images of wild-type yeast cells expressing GFP fusion constructs (pRS416). Left panels: GFP fluorescence. Middle panels: Nuclear fluorescence (stained with Hoechst). Right panels: Nomarski optics. A: Control cells (solely GFP). B: *ScMca1*-GFP. C: *TbMca4*-GFP. D: *Wwm1*-GFP.

sulted in suppression of the phenotypes caused by the former alone. Alleles of *ScMCA1* mutated in the conserved cysteine (cysteine-176 and cysteine-297) or histidine (histidine-241) residues were capable to suppress *WWMI* overexpression as effectively as wild-type *ScMCA1*. A truncated form of *ScMCA1*, however, lacking the 5' region encoding the proline-rich N-terminus was not able to suppress overexpressed *WWMI* (Fig. 3A, data on suppression of loss of clonogenicity not shown). Since WW domains like the one found in *Wwm1p* are known to interact with proline-rich protein do-

main [22], suppression of overexpressed *WWMI* by overexpressed *ScMCA1* is most readily explained by titration of *Wwm1p* by the N-terminal domain of *ScMca1p*.

Growth inhibition, loss of clonogenicity and suppression by *ScMCA1* were also achieved with a pRS416-based *WWMI*-GFP fusion construct (data not shown). The resulting *Wwm1*-GFP fluorescence localized to the nucleus (Fig. 2D), but upon suppression by *ScMCA1* was observed in large aggregates distributed throughout the cell, no longer co-localizing with the nucleus (Fig. 3B).




Fig. 3. Suppression of *WWM1*/*Wwm1*-GFP overexpression (pRS416) by overexpression of *ScMCA1* (YEp52) in wild-type yeast. A: Ten-fold dilution series of wild-type yeast carrying *WWM1* on plasmid pRS416 in combination with different YEp52 plasmids. First row: empty YEp52 (control). Second row: YEp52 with *ScMCA1*. Third row: YEp52 with *ScMCA1*^{C176A}. Fourth row: YEp52 with *ScMCA1*^{H241A}. Fifth row: YEp52 with *ScMCA1*^{C297A}. Sixth row: YEp52 with *ScMCA1*^{ΔN-terminus}. Upper left panel: Neither induction of *WWM1* nor expression from YEp52 (2 days incubation). Upper right panel: Induction of *WWM1*, no expression from YEp52 (5 days incubation). Lower left panel: No induction of *WWM1* but expression from YEp52 (3 days incubation). Lower right panel: Induction of *WWM1* and expression from YEp52 (5 days incubation). B: Images of *Wwm1*-GFP expressing wild-type yeast (pRS416) overexpressing *ScMCA1* (YEp52) (7 h in inducing liquid medium – CM-S with galactose). Left panel: *Wwm1*-GFP fluorescence. Middle panel: Nuclear fluorescence (bisbenzimide); Right panel: Nomarski optics.

4. Discussion

Heterologous expression of *TbMCA4* was found to induce respiration deficiency in yeast. This is obviously due to the peculiarity of *S. cerevisiae* of living without fully functional mitochondria, although the organelle itself remains essential nevertheless [28]. Loss of clonogenicity (clonal lethality) occurred as a much later, downstream event of *TbMCA4* expression, so that yeast colonies with irreversibly damaged respiratory competence ('petites') were obtained at high efficiency. Thus it is apparent that the action of *TbMca4p* primarily affects yeast by compromising mitochondrial function. However, no concomitant alteration in mitochondrial abundance, shape or distribution could be observed by Mito-

Tracker Green FM or electron microscopy (data not shown). Only MitoTracker CMXRos, a fluorescent dye specific for free mitochondrial thiols and thus indicative for mitochondrial functions, was not efficiently sequestered by mitochondria of *TbMCA4* expressing yeast, underscoring that abnormal mitochondrial biogenesis takes place besides the loss of respiratory competence.

The site of *TbMca4p* action on yeast seems to be the nucleus as suggested by a *TbMca4p*-GFP fusion reporter. Although this latter construct was non-functional, the nuclear localization of a *ScMca1p*-GFP reporter supports this idea. Furthermore, yeast growth inhibition by *Wwm1p*-GFP also coincided with a nuclear localization of this fusion construct, which was suppressed by *ScMCA1* overexpression. Finally, the effect of *TbMca4p* upon yeast mitochondrial respiration competence and its presumptive nuclear localization suggests that metacaspases are involved in an aspect of transcriptional control affecting mitochondrial biogenesis.

Development of respiration incompetence is very common for mutations in genes necessary or essential for mitochondrial biogenesis (*PET* genes) [29]. For instance, expression of a truncated cytochrome *bc₁* complex assembly protein of which the gene was disrupted resulted in a temperature-dependent petite phenotype [30]. The most striking parallels to the effects of *TbMCA4*, however, were reported from heterologous expression of mammalian Bax in yeast. Bax is an important proapoptotic effector in mammalian PCD which induces growth retardation, loss of clonogenicity and a permanent loss of respiration competence in yeast [17,18,31–34]. These effects of Bax could be suppressed by coexpression of another related Bcl family member, Bcl-x_L [17,32]. A general protection of yeast mitochondria was achieved by expression of another Bcl family member, Bcl-2 [35]. Furthermore, even plant PCD was suppressed by expression of Bcl-x_L [13]. Therefore it appears notable that we were unable to relieve *TbMCA4* expressing yeast by coexpression of mouse Bcl-x_L (data not shown).

Coexpression of antiapoptotic baculovirus p35 also did not affect *TbMCA4* expressing yeast (data not shown), while this caspase inhibitor was able to suppress PCD in plants [15]. If metacaspases are mediators of PCD in plants and unicellular eukaryotes, which appears to be an attractive speculation [5,10], we expected Bcl-x_L and p35 to relieve *TbMCA4* expressing yeast. Our aforementioned experimental results do not support this idea, however.

The physiological relevance of *WWM1* still remains unsolved. It is intriguing to notice that moderate overexpression of this gene was inhibiting cellular proliferation in such a potent manner, considering the high transcript levels that were reported from growing yeast for this gene [36]. In a recent overexpression screen *WWM1* was found to be one of the strongest growth inhibitors, which caused cells to arrest in G1 [37]. Suppression of *WWM1* by the yeast metacaspase *ScMCA1* appears an interesting fact, although the physiological relevance of this interaction remains elusive. Based on our results, we would like to suggest that *Wwm1p* plays a specific role in metacaspase function. Since *Wwm1p*-GFP localization was altered by overexpression of *ScMCA1*, we assume that exclusive nuclear localization of *Wwm1p*-GFP is due to an increased dosage of *Wwm1p*.

Sequence analysis shows that metacaspases constitute a new group of cysteinyl proteases. Accordingly, exchange of the

putative catalytic dyad residues of *TbMca4p* by site-directed mutagenesis resulted in loss of all phenotypes caused by expression of the respective wild-type gene *TbMCA4* in yeast. In addition, exchange of the other conserved cysteine residue within metacaspases also rendered *TbMca4p* inactive. This indicates that metacaspases possess a second catalytic nucleophile, which might be important either for maturation or for the catalytic activity itself. Notably, a conserved cysteine is also found in *Gpi8p* transamidases at a comparable site (our unpublished observations). Exchange of this residue in the essential yeast *ScGpi8p* was possible, however [38]. So far, we cannot exclude that metacaspases are not typical proteases, but rather transamidases, which mediate protein lipidation similar to the related *Gpi8* proteins.

In our experiments, tagging of *TbMca4p* (N- or C-terminally with *c-myc* or FLAG epitopes or with GFP) always resulted in a loss of function (data not shown). These tagged variants might not fold properly and consequently are hindered in their activity or in their (self-)maturation. It appears fortuitous that *TbMca4p* becomes active in yeast at all, since expression of the four other *T. brucei* metacaspases, the single metacaspase of *S. pombe*, or overexpression of *ScMCA1* did not lead to any visible phenotype.

In order to demonstrate proteinase activity of *TbMca4p*, raw extracts of yeast expressing *TbMCA4* or purified recombinant *TbMca4p* from *Escherichia coli* were incubated with a set of synthetic quarterpeptides with either N, D, R or K preceding a fluorogenic group (7-amino,4-methylcoumarin). Protease activity could not be monitored, however (data not shown).

Metacaspase-like sequences with limited homology to metacaspases occur in many bacteria. Since the putative secondary structures of these different sequences appear to be very similar to metacaspases, they appear to be distant homologues of metacaspases [5]. However, only the α -proteobacterial *Rsmca1p* of *R. sphaeroides* (and the δ -proteobacterial *Gsmca1p*) shows a high degree of sequence homology with the eukaryotic metacaspases. Thus, in view of the endosymbiont hypothesis [39], it appears likely that metacaspases were acquired by the eukaryotic cell from its protomitochondrial endosymbiont. Interestingly, analysis of the yeast mitochondrial proteome revealed that only a minority of proteins (50 out of 400) are clearly derived from the α -proteobacterial ancestor, reflecting an unexpectedly high degree of coevolutionary development of symbiont and host [40]. If the metacaspases are indeed of α -proteobacterial origin, they would be quite unique for their non-mitochondrial localization. The only other, also hypothetical, example known so far are aspartic proteases involved in cell cycle control [41].

Acknowledgements: We thank Drs. R.T. Schwarz (Marburg, Germany), K.-U. Fröhlich (Graz, Austria), S. Kugler (Göttingen, Germany) and M. Ligr (Stuttgart, Germany) for generous gifts of plasmids and genes.

References

- [1] Chen, J.M., Rawlings, N.D., Stevens, R.A. and Barrett, A.J. (1998) FEBS Lett. 441, 361–365.
- [2] Uhlmann, F., Wernic, D., Poupert, M.A., Koonin, E.V. and Nasmyth, K. (2000) Cell 103, 375–386.
- [3] Walker, N.P., Talanian, R.V., Brady, K.D., Dang, L.C., Bump, N.J., Ferez, C.R., Franklin, S., Ghayur, T., Hackett, M.C. and Hammill, L.D. (1994) Cell 78, 343–352.
- [4] Eichinger, A., Beisel, H.G., Jacob, U., Huber, R., Medrano, F.J., Banbula, A., Potempa, J., Travis, J. and Bode, W. (1999) EMBO J. 18, 5453–5462.
- [5] Uren, A.G., O'Rourke, K., Aravind, L.A., Pisabarro, M.T., Seshagiri, S., Koonin, E.V. and Dixit, V.M. (2000) Mol. Cell 6, 961–967.
- [6] Skulachev, V.P. (1996) FEBS Lett. 397, 7–10.
- [7] Mignotte, B. and Vayssières, J.L. (1998) Eur. J. Biochem. 252, 1–15.
- [8] Green, D.R. and Reed, J.C. (1998) Science 281, 1309–1312.
- [9] Brenner, C. and Kroemer, G. (2000) Science 289, 1150–1151.
- [10] Lam, E., Kato, N. and Lawton, M. (2001) Nature 411, 848–853.
- [11] Ameisen, J.C. (1996) Science 272, 1278–1279.
- [12] Aravind, L., Dixit, V.M. and Koonin, E.V. (2001) Science 291, 1279–1284.
- [13] Dickman, M.B., Park, Y.K., Oltersdorf, T., Li, W., Clemente, T. and French, R. (2001) Proc. Natl. Acad. Sci. USA 98, 6957–6962.
- [14] Arnoult, D., Tatischeff, I., Estaquier, J., Girard, M., Sureau, F., Tissier, J.P., Grodet, A., Dellinger, M., Traincard, F., Kahn, A., Ameisen, J.C. and Petit, P.X. (2001) Mol. Biol. Cell 12, 3016–3030.
- [15] Hansen, G. (2000) Mol. Plant Microbe Interact. 13, 649–657.
- [16] Fisher, A.J., Cruz, W., Zoog, S.J., Schneider, C.L. and Friesen, P.D. (1999) EMBO J. 18, 2031–2039.
- [17] Gross, A., Pilcher, K., Blachly-Dyson, E., Basso, E., Jockel, J., Bassik, M.C., Korsmeyer, S.J. and Forte, M. (2000) Mol. Cell Biol. 20, 3125–3136.
- [18] Harris, M.H., Vander Heiden, M.G., Kron, S.J. and Thompson, C.B. (2000) Mol. Cell Biol. 20, 3590–3696.
- [19] Uetz, P., Giot, L., Cagney, G., Mansfield, T.A., Judson, R.S., Knight, J.R., Lockshon, D., Narayan, V., Srinivasan, M., Pochar, P., Qureshi-Emili, A., Li, Y., Godwin, B., Conover, D., Kalbfleisch, T., Vijayadamar, G., Yang, M., Johnston, M., Fields, S. and Rothberg, J.M. (2000) Nature 403, 623–627.
- [20] Ito, T., Chiba, T., Ozawa, R., Yoshida, M., Hattori, M. and Sakai, Y. (2001) Proc. Natl. Acad. Sci. USA 98, 4569–4574.
- [21] Garay-Arroyo, A., Colmenero-Flores, J.M., Garciarrubio, A. and Covarrubias, A.A. (2001) J. Biol. Chem. 275, 5668–5674.
- [22] Sudol, M., Sliwa, K. and Russo, T. (2001) FEBS Lett. 490, 190–195.
- [23] Steinborn, K., Szallies, A., Mecke, D. and Duzsenko, M. (2000) Biol. Chem. 381, 1071–1077.
- [24] Sikorski, R.S. and Hieter, P. (1989) Genetics 122, 19–27.
- [25] Schiestl, R.H. and Gietz, R.D. (1989) Curr. Genet. 16, 339–346.
- [26] Ogur, M., John, R.S. and Nagai, S. (1957) Science 125, 928–929.
- [27] Shin, D.H., Kang, H. and Han, K.-H. (1999) Plant Physiol. 121, 1384.
- [28] Yaffe, M.P. (1999) Science 283, 1493–1497.
- [29] Tzagoloff, A. and Dieckmann, C.L. (1990) Microbiol. Rev. 54, 211–225.
- [30] Schmitt, M.E. and Trumppower, B.L. (1991) J. Biol. Chem. 266, 14958–14963.
- [31] Greenhalf, W., Stephan, C. and Chaudhuri, B. (1996) FEBS Lett. 380, 169–175.
- [32] Manon, S., Chaudhuri, B. and Guerin, M. (1997) FEBS Lett. 415, 29–32.
- [33] Priault, M., Camougrand, N., Chaudhuri, B., Schaeffer, J. and Manon, S. (1999) FEBS Lett. 456, 232–238.
- [34] Kissova, I., Polcic, P., Kempna, P., Zeman, I., Sabova, L. and Kolarov, J. (2000) FEBS Lett. 471, 113–118.
- [35] Longo, V.D., Ellerby, L.M., Bredesen, D.E., Valentine, J.S. and Gralla, E.B. (1997) J. Cell Biol. 137, 1581–1588.
- [36] Naitou, M., Hagiwara, H., Hanaoka, F., Eki, T. and Murakami, Y. (1997) Yeast 13, 1275–1290.
- [37] Stevenson, L.F., Kennedy, B.K. and Harlow, E. (2001) Proc. Natl. Acad. Sci. USA 98, 3946–3951.
- [38] Meyer, U., Benghezal, M., Imhof, I. and Conzelmann, A. (2000) Biochemistry 39, 3461–3471.
- [39] Gray, M.W., Burger, G. and Lang, B.F. (1999) Science 283, 1476–1481.
- [40] Kurland, C.G. and Andersson, S.G. (2000) Microbiol. Mol. Biol. Rev. 64, 786–820.
- [41] Krylov, D.M. and Koonin, E.V. (2001) Curr. Biol. 11, R584–587.