



Erratum

Erratum to ‘2015 HRS/EHRA/APHRS/SOLAECE expert consensus statement on optimal implantable cardioverter-defibrillator programming and testing’ [Journal of Arrhythmia 32/1 (2016) 1–28]

Bruce L. Wilkoff, MD, FHRS, CCDS^a, Laurent Fauchier, MD, PhD^{b,1},
 Martin K. Stiles, MBChB, PhD^{c,2}, Carlos A. Morillo, MD, FRCPC, FHRS^{d,3},
 Sana M. Al-Khatib, MD, MHSc, FHRS, CCDS^e, Jesoes Almendral, MD, PhD, FESC^{f,1},
 Luis Aguinaga, MD, PhD, FACC, FESC^{g,3}, Ronald D. Berger, MD, PhD, FHRS^h,
 Alejandro Cuesta, MD, PhD, FESC^{i,3}, James P. Daubert, MD, FHRS^e,
 Sergio Dubner, MD, FACC^{j,ai,3}, Kenneth A. Ellenbogen, MD, FHRS^k, N.A. Mark Estes III, MD^{l,4},
 Guilherme Fenelon^{m,3}, Fermin C. Garcia, MD^{n,3}, Maurizio Gasparini, MD^{o,1},
 David E. Haines, MD, FHRS^p, Jeff S. Healey, MD, MSc, FRCPC, FHRS^d,
 Jodie L. Hurtwitz, MD^{q,5}, Roberto Keegan, MD^{r,3}, Christof Kolb, MD^{s,1},
 Karl-Heinz Kuck, MD, FHRS^{t,1}, Germanas Marinskis, MD, FESC^{u,1},
 Martino Martinelli, MD, PhD^v, Mark McGuire, MBBS, PhD^{w,2}, Luis G. Molina, MD, DSc^{x,3},
 Ken Okumura, MD, PhD^{y,2}, Alessandro Proclemer, MD^{z,1}, Andrea M. Russo, MD, FHRS^{aa},
 Jagmeet P. Singh, MD, DPhil, FHRS^{ab}, Charles D. Swerdlow, MD, FHRS^{ac},
 Wee Siong Teo, MBBS, FHRS^{ad,2}, William Uribe, MD, FHRS^{ae,3}, Sami Viskin, MD^{af,1},
 Chun-Chieh Wang, MD^{ag,2}, Shu Zhang, MD^{ah,2}

^a Cleveland Clinic, Cleveland, OH, United States

^b Centre Hospitalier Universitaire Trousseau, Tours, France

^c Waikato Hospital, Hamilton, New Zealand

^d Department of Medicine, Cardiology Division, McMaster University–Population Health Research Institute, Hamilton, Canada

^e Duke University Medical Center, Durham, NC, United States

^f Grupo HM Hospitales, Universidad CEU San Pablo, Madrid, Spain

^g Centro Privado De Cardiología, Tucuman, Argentina

^h Johns Hopkins University, Baltimore, MD, United States

ⁱ Servicio de Arritmias, Instituto de Cardiología Infantil, Montevideo, Uruguay

^j Clinica y Maternidad Suizo Argentina

^k Virginia Commonwealth University Medical Center, Richmond, VA, United States

^l New England Medical Center, Boston, MA, United States

^m Federal University of Sao Paulo, Sao Paulo, Brazil

ⁿ Hospital of the University of Pennsylvania, Philadelphia, PA, United States

^o Humanitas Research Hospital, Milan, Italy

^p William Beaumont Hospital Division of Cardiology, Royal Oak, MI, United States

^q North Texas Heart Center, Dallas, TX, United States

^r Hospital Privado del Sur, Bahia Blanca, Argentina

^s Deutsches Herzzentrum Munchen, Munich, Germany

^t Allgemeines Krankenhaus St. Georg, Hamburg, Germany

^u Vilnius University, Clinic of Cardiac and Vascular Diseases, Lithuania

^v Instituto do Coração, Universidade de Sao Paulo, Sao Paulo, Brazil

^w Royal Prince Alfred Hospital, Sydney, Australia

^x Mexico's National University, Mexico's General Hospital, Mexico City, Mexico

^y Hirosaki University Graduate School of Medicine, Hirosaki, Aomori, Japan

^z Azienda Ospedaliero Universitaria S. Maria della Misericordia– Udine, Udine, Italy

^{aa} Cooper University Hospital, Camden, NJ, United States

^{ab} Massachusetts General Hospital, Harvard Medical School, Boston, MA, United States

^a*Cedars-Sinai Medical Center, Beverly Hills, CA, United States*

^a*National Heart Centre Singapore, Singapore, Singapore*

^a*CES Cardiología and Centros Especializados San Vicente Fundación, Medellín y Rionegro, Colombia*

^a*Tel Aviv Sourasky Medical Center and Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel*

^a*Chang Gung Memorial Hospital, Taipei, Taiwan*

^a*National Center for Cardiovascular Disease and Beijing Fu Wai Hospital, Peking Union Medical College and China Academy of Medical Sciences, Beijing, China*

^a*De Los Arcos Sanatorio, Buenos Aires, Argentina*

There was an error on the above article. The sentence “Even if a dual-chamber ICD is implanted, dual-chamber discrimination should be programmed only if the atrial lead becomes chronic or if atrial sensing is unreliable” is not correct. It should read: “Even if a

dual-chamber ICD is implanted, dual-chamber discrimination should be programmed only if the atrial lead becomes chronic or if atrial sensing is reliable”.

¹ Representative of the European Heart Rhythm Association (EHRA).

² Representative of the Asia-Pacific Heart Rhythm Society (APHRS).

³ Representative of the American College of Cardiology (ACC).

⁴ Representative of the American Heart Association (AHA).

⁵ Representative of the Sociedad Latinoamericana de Estimulación Cardíaca y Electrofisiología (SOLAECE).