

## EDITORIAL COMMENT

# Can Mold Allergy be Diagnosed with a Skin Test or Specific IgE Antibodies?

Ren-Bin Tang\*

*Department of Pediatrics, Taipei Veterans General Hospital, and  
National Yang-Ming University School of Medicine, Taipei, Taiwan, R.O.C.*

In this issue of the *Journal of the Chinese Medical Association*, Liang et al<sup>1</sup> describe a study that evaluates the skin test (Tori Ltd, Tokyo, Japan) and specific IgE antibodies (ImmunoCAP system; Pharmacia, Uppsala, Sweden), which are designed to identify mold allergy. Five fungal antigens (*Candida*, *Alternaria*, *Aspergillus*, *Cladosporium* and *Penicillium*) were included in this study.

With the steadily increasing prevalence of atopic disorders in the pediatric population over the past several decades, it is more essential than ever to develop a simple, rapid test that identifies inhalant allergens in these atopic individuals.<sup>2</sup> Fungi are well known as sources of allergens that cause allergic rhinitis and allergic asthma in Taiwan.<sup>3-5</sup> *Penicillium citrinum* and *Aspergillus fumigatus* are prevalent indoor airborne fungal species that have been implicated in human respiratory allergic disorders.<sup>6,7</sup> Fungal allergens, on the whole, show significant differences in incidence among different age groups.<sup>5,6</sup>

Allergy skin testing for immediate hypersensitivity is a cornerstone in the evaluation of the patient with allergic disease.<sup>8,9</sup> Skin prick tests are quick, low cost, highly sensitive and safe. This seemingly simple test, however, is subject to multiple variables that can affect the result. Some of them are patient-dependent, such as the patient age, underlying skin condition, or use of medications that can interfere with the test results. Testing-dependent variables include the quality of the extracts used, the testing technique and device used, the location on the body to which the tests are applied, and the distance between individual test sites.<sup>10</sup> Finally, individual physician scoring and interpretation of allergen skin test results may add further variability.

Specific serum IgE is widely used in the diagnosis of IgE-mediated allergic diseases. The Pharmacia ImmunoCAP system (CAP) represents the standard method of diagnosis and is accurate enough to diagnose specific IgE.<sup>11</sup> Previous studies have established a strong agreement of both the multiple allergosorbent chemiluminescent assay (MAST-CLA) and CAP with the skin prick test. Both MAST-CLA and CAP were comparable in their ability for diagnosing allergies to tested allergens. They can be used as a screening test to measure allergens.<sup>12-16</sup> Inhalant allergens of house dust mites and fungi are also detected by the immunoblotting method.<sup>17,18</sup> A good correlation between the immunodot assay and the skin test was confirmed in asthmatic children.<sup>19</sup> The electroblotting technique is fast, convenient, and highly suitable for both allergen composition studies and screening of antibody specificities. As new and more effective immunomodulatory approaches for specific IgE activity become available, it will be more important than ever to diagnose these mold allergens, so that appropriate interventions can be initiated.<sup>6,7,20</sup>

Skin test and specific IgE usually correlate equally well with clinical allergic symptoms. Concordance for results of skin testing and the radioallergosorbent test was high for most allergens except for *Candida* allergens.<sup>21-23</sup> Both tests can be used as grounds for instituting immunotherapy in an efficient manner. When CAP is used as grounds for immunotherapy, a skin test with an initial diluted dose of immunotherapy should be done to evaluate the tolerable dose of the allergenic extract.<sup>24</sup>

Chapman<sup>25</sup> considered that the clinical aspects of fungal sensitivity are essential for assessment of exposure potential and clinical testing. Allergy skin test material

\*Correspondence to: Dr. Ren-Bin Tang, Department of Pediatrics, Taipei Veterans General Hospital, 201, Section 2, Shih-Pai Road, Taipei 112, Taiwan, R.O.C.

E-mail: [rbtang@vghtpe.gov.tw](mailto:rbtang@vghtpe.gov.tw) • Received: September 5, 2005 • Accepted: November 8, 2005

is unavailable for most airborne fungi. Those that are available are not standardized. A major management approach for patients with proven sensitivity to fungal antigens and a clear correlation with clinical illness is avoidance of fungal sources.<sup>25</sup>

The article by Liang et al<sup>1</sup> in this issue illustrates the difficulties that arise in a study attempting to evaluate mold allergy with the skin test and CAP. This article describes no significant correlation between the two tests to *Candida*, *Alternaria*, *Aspergillus*, *Cladosporium* and *Penicillium*. The positive rates of the skin test and CAP were 56% versus 9% for *Candida*; 22% versus 1% for *Alternaria*; 16% versus 9% for *Aspergillus*; 15% versus 1% for *Cladosporium*; and 32% versus 8% for *Penicillium*. There was no attempted correlation with clinical disease, and the possible clinical application of the data to mold allergy was tenuous. As reported by Piette et al,<sup>26</sup> the improper performance of skin prick tests leads to unreliability, as it may result in an unacceptable number of false-positive results and lead to wrong allergy diagnosis and incorrect treatment.

## References

- Liang KL, Su MC, Jiang RS. Comparison of the skin test and ImmunoCAP system in the evaluation of mold allergy. *J Chin Med Assoc* 2006;69:3-6.
- Tang RB. Risk factors associated with developed asthma. *J Chin Med Assoc* 2005;68:199-201.
- Tang RB, Shen HD, Chen SJ, Lee CY. Detection of IgE reactivity to fungus antigens by immunoblotting in allergic diseases in children. *J Chin Med Assoc* 2003;66:453-9.
- Shen HD, Choo KB, Tang RB, Lee CF, Yeh JY, Han SH. Allergenic components of *Candida albicans* identified by immunoblot analysis. *Clin Exp Allergy* 1989;19:191-5.
- Lee CS, Tang RB, Chung RL. The evaluation of allergens and allergic diseases in children. *J Microbiol Immunol Infect* 2000;33:227-32.
- Chou H, Chang CY, Tsai JJ, Tang RB, Lee SS, Wang SR, Shen HD. The prevalence of IgE antibody reactivity against the alkaline serine protease major allergen of *Penicillium chrysogenum* increases with the age of asthmatic patients. *Ann Allergy Asthma Immunol* 2003;90:248-53.
- Lai HY, Tam MF, Tang RB, Chou H, Chang CY, Tsai JJ, Shen HD. cDNA cloning and immunological characterization of a newly identified enolase allergen from *Penicillium citrinum* and *Aspergillus fumigatus*. *Int Arch Allergy Immunol* 2002;127:181-90.
- McCann WA, Ownby DR. The reproducibility of the allergy skin test scoring and interpretation by board-certified/board-eligible allergists. *Ann Allergy Asthma Immunol* 2002;89:368-71.
- Portnoy JM. What do allergy skin tests really mean? *Ann Allergy Asthma Immunol* 2002;89:335-6.
- Wood RA. Skin testing: making the most of every prick. *Ann Allergy Asthma Immunol* 2002;88:347-9.
- Bousquet J, Chané P, Chanal I, Michel FB. Comparison between RAST and Pharmacia CAP system: a new automated specific IgE assay. *J Allergy Clin Immunol* 1990;5:1039-43.
- Tang RB, Chen BS, Chen SJ, Wu KG, Hwang B. Determination of the food specific IgE antibodies: comparison of MAST-CLA and CAP systems. *J Chin Med Assoc* 1996;57:219-23.
- Tang RB, Tsai LC, Hwang HM, Hwang BT, Wu KG, Hung MW. The prevalence of allergic disease and IgE antibodies to house dust mite in schoolchildren in Taiwan. *Clin Exp Allergy* 1990;20:33-8.
- Tang RB, Wu KG, Hwang BT. Comparison between skin testing and *in vitro* testing for diagnosis of allergen in asthmatic children. *J Chin Med Assoc* 1994;54:246-50.
- Tang RB, Chen BS, Wu KG, Hwang B. Comparison of food specific IgE antibody test (RAST) and skin tests in children with atopic dermatitis. *J Chin Med Assoc* 1993;52:161-5.
- Finnerty JP, Summerell S, Holgate ST. Relationship between skin-prick tests, the multiple allergosorbent test and symptoms of allergic disease. *Clin Exp Allergy* 1989;19:51-6.
- Shen HD, Choo KB, Tang RB, Lee CF, Yeh JY, Han SH. Allergenic components of *Candida albicans* identified by Immunoblot analysis. *Clin Exp Allergy* 1989;19:191-6.
- Tang RB, Tsai LC, Hung MW, Hwang BT, Wu KG. Detection of house dust mite allergens and immunoblot analysis in asthmatic children. *J Asthma* 1988;25:83-8.
- Chao PL, Peng HJ, Tang RB, Hung MW, Tsai LC. Serum specific IgE reactivity to recombinant Der fl 1 in asthmatic children. *J Asthma* 2001;38:333-40.
- Chang CY, Chou H, Tam MF, Tang RB, Lai HY, Shen HD. Characterization of enolase allergen from *Rhodotorula mucilaginosa*. *J Biomed Sci* 2002;9:645-55.
- Tang RB, Wu KK. Total serum IgE, allergy skin testing, and the radioallergosorbent test for the diagnosis of allergy in asthmatic children. *Ann Allergy Asthma Immunol* 1989;64:432-5.
- Tsai LC, Tang RB, Hung MW, Chen HM, Hwang LA. Correlation between serum IgE and specific IgE antibody titer to house dust mite in children with asthma. *J Asthma* 1988;25:7-13.
- Tang RB, Chang HN, Lin FM, Chang YF, Chou NS. Serum IgE, skin and radioallergosorbent tests for house dust and mites in asthmatic children. *J Asthma* 1986;23:245-9.
- Kniker WT. Is the choice of allergy skin testing versus *in vitro* determination of specific IgE no longer a scientific issue? *Ann Allergy Asthma Immunol* 1989;62:373-4.
- Chapman JA. Update on airborne mold and mold allergy. *Allergy Asthma Proc* 1999;20:289-92.
- Piette V, Bourret E, Bousquet J, Demoly P. Prick tests to aeroallergens: is it possible simply to wipe the device between tests? *Allergy* 2002;57:940-2.