




Cairo University  
Bulletin of Faculty of Pharmacy, Cairo University

[www.elsevier.com/locate/bfopcu](http://www.elsevier.com/locate/bfopcu)  
[www.sciencedirect.com](http://www.sciencedirect.com)



## ORIGINAL ARTICLE

# Sagitol C, a new cytotoxic pyridoacridine alkaloid from the sponge *Oceanapia* sp.

Sabrin R.M. Ibrahim <sup>a,\*</sup>, Gamal A. Mohamed <sup>b</sup>, Ehab S. Elkhayat <sup>b</sup>,  
Mostafa A. Fouad <sup>c</sup>, Peter Proksch <sup>d</sup>

<sup>a</sup> Department of Pharmacognosy, Faculty of Pharmacy, Assiut University, Assiut 71526, Egypt

<sup>b</sup> Department of Pharmacognosy, Faculty of Pharmacy, Al-Azhar University, Assiut 71524, Egypt

<sup>c</sup> Department of Pharmacognosy, Faculty of Pharmacy, Minia University, Minia 61519, Egypt

<sup>d</sup> Institut für Pharmazeutische Biologie und Biotechnologie, Heinrich-Heine-Universität Düsseldorf, Universitätsstr. 1, Geb.26.23, Düsseldorf D-40225, Germany

Received 24 April 2013; accepted 31 May 2013

Available online 25 July 2013

## KEYWORDS

*Oceanapia*;  
Pyridoacridine alkaloid;  
Sagitol C;  
Cytotoxicity

**Abstract** A new pyridoacridine alkaloid named sagitol C (**2**) together with two known compounds; kuanoniamine C (**1**) and sagitol (**3**) were isolated from the EtOAc fraction of the Indonesian sponge *Oceanapia* sp. Their chemical structures were established on the basis of physical and spectroscopic methods 1D and 2D NMR, in addition to mass spectrometry and comparison with literature data. Sagitol C was found to exhibit cytotoxic activity when tested against different cancer cell lines.

© 2013 Production and hosting by Elsevier B.V. on behalf of Faculty of Pharmacy, Cairo University.  
Open access under [CC BY-NC-ND license](http://creativecommons.org/licenses/by-nc-nd/4.0/).

## 1. Introduction

*Oceanapia* comprises more than 50 nominal species recorded for the Indo-west Pacific region alone, relatively abundant in both soft and hard substrates, and widely distributed.<sup>1</sup> Pyridoacridine alkaloids have been isolated from sponges, ascidians, anemones, and prosobranch.<sup>2</sup> They are characterised by

11*H*-pyrido[4,3,2-*mn*]acridine moiety and have so far only been reported from marine organisms. The colours exhibited by pyridoacridine alkaloids may vary depending on their pH. This physicochemical property is correlated with the presence of at least two basic nitrogens in the aromatic ring systems.<sup>3</sup> They are known to exhibit a wide range of biological activities including: anti-bacterial, anti-fungal, anti-viral, anti-parasitic, insecticidal, anti-tumour, topoisomerase inhibition, and anti-fouling.<sup>4</sup> *Oceanapia* species have yielded different classes of natural products including: alkaloids,<sup>1–5</sup> sphingolipid-like compounds,<sup>6</sup> sphingolipids,<sup>7,8</sup> polyhydroxy sterols,<sup>9</sup> acetylenes,<sup>10</sup> and dithiocyanates.<sup>11</sup> In the course of our investigation on bioactive compounds from the marine sponges, we have isolated a new pyridoacridine alkaloid named sagitol C (**2**), in addition to two known pyridoacridine alkaloids; kuanoniamine C (**1**) and sagitol (**3**) (Fig. 1). We have also investigated the effect of sagitol C (**2**) on the growth of three tumour cell


\* Corresponding author. Tel.: +20 88 2141330; fax: +20 88 2332776.

E-mail address: [sabrinshaur@gmail.com](mailto:sabrinshaur@gmail.com) (S.R.M. Ibrahim).

Peer review under responsibility of Faculty of Pharmacy, Cairo University.



Production and hosting by Elsevier



**Figure 1** Chemical structures of the isolated compounds.

lines: mouse lymphoma (L5178Y), rat brain (PC12), and human cervix (Hela).

## 2. Experimental

### 2.1. General procedures

Mass spectra (ESIMS) were recorded on a Finnigan MAT TSQ-7000 triple stage quadrupole mass spectrometer. HRESIMS were measured on a Finnigan MAT 95 mass spectrometer. UV spectra were recorded in absolute MeOH on a Shimadzu 1601 UV/VIS spectrophotometer. 1D and 2D NMR spectra (chemical shifts in ppm, coupling constants in Hz) were recorded on a Bruker DRX400 NMR spectrometer using standard Bruker software and DMSO- $d_6$  as solvent, with TMS as the internal reference. NMR spectra were referenced to the solvent signals (DMSO- $d_6$ : 2.49 ppm for  $^1\text{H}$  and 39.9 ppm for  $^{13}\text{C}$ ). Solvents were distilled prior to use, and spectral grade solvents were used for spectroscopic measurements. Column chromatographic separation was performed on silica gel 60 (0.04–0.063 mm), RP-18 (0.04–0.063 mm Merck), and Sephadex LH-20 (0.25–0.1 mm, Merck). TLC was performed on precoated TLC plates with silica gel 60 F<sub>254</sub> (layer thickness 0.2 mm, Merck). The chromatograms were developed using the following solvent system:  $\text{CHCl}_3$ :MeOH (90:10, S<sub>1</sub>). The compounds were detected by UV absorption at  $\lambda_{\text{max}}$  255 and 366 nm followed by spraying with Dragendorff's reagent.

### 2.2. Animal material

The sponge was collected by scuba diving at a depth of 7–9 m from Ambon, Indonesia in 2004. Freshly collected sponges were frozen immediately after collection and then freeze-dried. The dull-red sponge has a fistular growth form, consisting of an irregular turnip-shaped main body up to 8 cm high and 4 cm in diameter, from which issue several long hollow fistules 7–8 cm long and up to 1 cm in diameter. The surface is optically smooth and slightly rough when dry. The consistency of the freeze-dried specimen is very fragile, crumbly, and dusty. The skeleton of the periphery is a halichondroid, tangential, multilayered crust of intercrossing single spicules. This is carried by thin subectosomal spicule tracts of 2–3 spicules in cross section, following a meandering course and anastomosing irregularly. The skeleton of the interior of the sponge is scanty, consisting of a loose irregular reticulation of mostly single

spicules bound at the nodes with a little spongin. Spicules are exclusively curved, somewhat flexuous oxeas with variously pointed or blunt apices that are uniform in size 260–330 × 3–6  $\mu\text{m}$ .<sup>3</sup> It was kindly identified by Prof. Rob W.M. van Soest (Zoological Museum, University of Amsterdam). A voucher specimen is kept under registration no. ZMA POR 11007 in the Zoölogisch Museum, Amsterdam.

### 2.3. Extraction and isolation

The freeze-dried sponge (110 g) was extracted several times with a ratio of 1:1  $\text{CH}_3\text{Cl}$ :MeOH (1 L × 4). The total extract was evaporated to dryness (398 mg) and subjected to vacuum liquid chromatography (VLC) over silica gel using *n*-hexane, EtOAc, and MeOH as solvents. The EtOAc fraction (164 mg) was chromatographed over Sephadex LH-20 column (150 g × 50 × 2 cm) using MeOH as an eluent. 50 mL fractions were collected and monitored by TLC to obtain three subfractions OE-1–OE-3. Subfraction OE-1 (39 mg) was subjected to RP-18 column (0.04–0.063 mm; 40 g × 25 × 1 cm) using MeOH:H<sub>2</sub>O gradients to give **1** (7 mg, yellow amorphous powder). Subfraction OE-2 (57 mg) was chromatographed over silica gel column (100 g × 50 × 3 cm) using  $\text{CHCl}_3$ :MeOH gradient to obtain impure compounds **2** and **3**. Separately, the impure **2** and **3** were purified on RP-18 column (0.04–0.063 mm; 40 g × 25 × 1 cm) using MeOH:H<sub>2</sub>O gradients to give **2** (3 mg, orange residue) and **3** (5 mg, orange residue).

**Kuanoniamine C (1)**: It was isolated as yellow amorphous powder (7 mg);  $R_f = 0.78$  (S<sub>1</sub>). UV (MeOH)  $\lambda_{\text{max}}$ : 267 and 358 nm; ESIMS  $m/z$ : 375  $[\text{M} + \text{H}]^+$ .  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ ):  $\delta_{\text{H}}$  11.4 (1H, brs, NH), 9.41 (1H, s, H-11), 8.49 (1H, brs, H-2), 8.28 (1H, dd,  $J = 8.2, 1.8$  Hz, H-4), 7.88 (1H, brs, H-3), 7.71 (1H, dd,  $J = 8.2, 1.8$  Hz, H-6), 7.71 (1H, brd,  $J = 8.2$  Hz, H-7), 7.25 (1H, dd,  $J = 8.2, 1.8$  Hz, H-5), 3.20 (2H, m, H-14), 3.15 (2H, m, H-13), 2.07 (2H, q,  $J = 7.6$  Hz, H-16), 0.97 (3H, t,  $J = 7.6$  Hz, H-17).  $^{13}\text{C}$  NMR (100 MHz, DMSO- $d_6$ ):  $\delta_{\text{C}}$  174.6 (C-15), 153.2 (C-11), 143.5 (C-12b), 143.0 (C-2), 140.7 (C-12a), 140.4 (C-7a), 140.2 (C-9a), 139.4 (C-3a), 135.0 (C-6), 133.5 (C-8a), 125.4 (C-4), 122.8 (C-5), 118.3 (C-12c), 117.6 (C-7), 114.3 (C-3b), 108.2 (C-3), 107.7 (C-9), 36.3 (C-13), 30.9 (C-14), 28.4 (C-16), 9.7 (C-17).

**Sagitol C (2)**: It was isolated as orange residue (3 mg);  $R_f = 0.71$  (S<sub>1</sub>). UV (MeOH)  $\lambda_{\text{max}}$ : 254 and 366 nm. NMR data: see Table 1; HRESIMS  $m/z$ : 405.1464 (calcd for  $\text{C}_{22}\text{H}_{21}\text{N}_4\text{O}_2\text{S}$ ,  $[\text{M} + \text{H}]^+$ , 405.1463).

**Table 1** NMR data of compound **2** (400 and 100 MHz, DMSO-*d*<sub>6</sub>).

No.	$\delta_{\text{H}}$ [mult., <i>J</i> (Hz)]	$\delta_{\text{C}}$ (mult.)	$^1\text{H}$ - $^1\text{H}$ COSY	HMBC
2	9.15 d (5.4)	149.1 (CH)	3	12b
3	8.92 d (5.4)	117.2 (CH)	2	3a, 12c
3a	—	140.5 (C)	—	—
3b	—	122.5 (C)	—	—
4	9.05 dd (8.2, 1.3)	124.0 (CH)	5	6
5	8.06 dt (8.2, 1.3)	131.6 (CH)	4, 6, 7	7
6	8.11 dt (8.2, 1.3)	131.8 (CH)	6, 5	7, 7a
7	8.47 dd (8.2, 1.3)	132.0 (CH)	5, 7	—
7a	—	146.5 (C)	—	—
8a	—	162.7 (C)	—	—
9	—	72.5 (C)	—	—
9a	—	146.3 (C)	—	—
11	9.72 s	162.1 (CH)	—	12a
12a	—	149.0 (C)	—	—
12b	—	152.0 (C)	—	—
12c	—	115.5 (C)	—	—
13	3.20 m	47.2 (CH <sub>2</sub> )	14	9, 8a, 9a
14	2.78 m	41.5 (CH <sub>2</sub> )	13	9, 15
15	—	174.3 (C)	—	—
16	2.61 m	39.9 (CH)	17, 18	—
17	1.25 d (6.9)	26.3 (CH <sub>3</sub> )	16	15, 16
18	1.25 d (6.9)	26.3 (CH <sub>3</sub> )	16	15, 16

Sagitol (**3**): It was isolated as orange residue (5 mg);  $R_f = 0.69$  (S1). UV (MeOH)  $\lambda_{\text{max}}$ : 264 and 340 nm. ESIMS  $m/z$ : 391  $[\text{M} + \text{H}]^+$ .  $^1\text{H}$  NMR (400 MHz, DMSO-*d*<sub>6</sub>):  $\delta_{\text{H}}$  9.29 (1H, s, H-11), 8.87 (1H, d,  $J = 5.8$  Hz, H-2), 8.72 (1H, d,  $J = 7.8$  Hz, H-4), 8.49 (1H, d,  $J = 5.8$  Hz, H-3), 7.97 (1H, t,  $J = 7.8$  Hz, H-6), 8.27 (1H, d,  $J = 7.8$  Hz, H-7), 7.81 (1H, t,  $J = 7.5$  Hz, H-5), 2.93, 2.75 (2H, each m, H-14), 2.90, 2.49 (2H, m, H-13), 1.71 (2H, q,  $J = 7.3$  Hz, H-16), 0.79 (3H, t,  $J = 6.9$  Hz, H-17).  $^{13}\text{C}$  NMR (100 MHz, DMSO-*d*<sub>6</sub>):  $\delta_{\text{C}}$  174.2 (C-15), 162.5 (C-8a), 155.6 (C-11), 151.5 (C-12b), 150.5 (C-2), 146.5 (C-7a), 146.1 (C-9a), 140.1 (C-3a), 132.5 (C-6), 129.8 (C-7), 128.5 (C-5), 123.5 (C-4), 122.3 (C-3b), 116.2 (C-3), 115.1 (C-12c), 72.8 (C-9), 46.5 (C-13), 35.4 (C-14), 28.2 (C-16), 9.5 (C-17).



#### 2.4. Cytotoxicity test

Cytotoxicity tests were carried out by Prof. Dr. Müller (Mainz University, Germany). The cytotoxicity against L5178Y (mouse lymphoma cells), Hela (human cervix carcinoma cells) and PC12 (brain tumour cells of the rats) was determined using the microculture tetrazolium assay (MTT), and compared to that of untreated controls. Of the test samples, stock solutions in ethanol 96% (v/v) were prepared. Exponentially growing cells were harvested, counted and diluted appropriately. Of the cell suspension, 50  $\mu\text{L}$  containing 3750 cells were pipetted into 96-well microtitre plates. Subsequently, 50  $\mu\text{L}$  of a solution of the test samples containing the appropriate concentration was added to each well. The concentration range was 12.3 and 24.6  $\mu\text{M}$ . The small amount of ethanol present in the wells did not affect the experiments. The test plates were incubated at 37  $^{\circ}\text{C}$  with 5%  $\text{CO}_2$  for 72 h. A solution of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) was prepared at 5 mg/mL in phosphate buffered saline (PBS; 1.5 mM  $\text{KH}_2\text{PO}_4$ , 6.5 mM  $\text{Na}_2\text{HPO}_4$ , 137 mM NaCl,

2.7 mM KCl; pH 7.4) and from this solution, 20  $\mu\text{L}$  was pipetted into each well. The yellow MTT penetrates the healthy living cells and in the presence of mitochondrial dehydrogenases, MTT is transformed to its blue formazan complex. After an incubation period of 4 h at 37  $^{\circ}\text{C}$  in a humidified incubator with 5%  $\text{CO}_2$ , the medium was centrifuged (15 min, 20  $^{\circ}\text{C}$ , 210g) with 200  $\mu\text{L}$  DMSO, the cells were lysed to liberate the formed formazan product. After thorough mixing, the absorbance was measured at 520 nm using a scanning microtitre-well spectrophotometer. The colour intensity is correlated with the number of healthy living cells.<sup>12–14</sup>

### 3. Discussion and conclusion

Compound **2** was isolated as an orange residue. Positive mode ESIMS showed pseudo-molecular ion peaks at  $m/z$  405  $[\text{M} + \text{H}]^+$  and 809  $[2\text{M} + \text{H}]^+$ , which were consistent with the molecular formula  $\text{C}_{22}\text{H}_{21}\text{N}_4\text{O}_2\text{S}$  as confirmed by HRESIMS, which was an increase of 14 mass units compared to **3**. The UV spectrum showed absorption maxima at 254 and 366 nm, which are characteristic for a pyridoacridine moiety.<sup>2,3,16</sup> The NMR data of **2** and **3** were quite similar for the pyridoacridine moiety, but the signals associated with the terminal ethyl moiety found in **3**, were not present. Instead, new signals for two methyl groups at  $\delta_{\text{H}}$  1.25 (6H, d,  $J = 6.9$  Hz, H-17,18)/ $\delta_{\text{C}}$  26.3 (C-17,18) and methine at  $\delta_{\text{H}}$  2.61 (1H, m, H-16)/ $\delta_{\text{C}}$  39.9 (C-16), suggest the presence of an isopropyl group. The pyridoacridine moiety was evident from signals at  $\delta_{\text{H}}$  9.72 (1H, s, H-11), 9.15 (1H, d,  $J = 5.4$  Hz, H-2), 9.05 (1H, dd,  $J = 8.2, 1.3$  Hz, H-4), 8.92 (1H, d,  $J = 5.4$  Hz, H-3), 8.47 (1H, dd,  $J = 8.2, 1.3$  Hz, H-7), 8.11 (1H, dt,  $J = 8.2, 1.3$  Hz, H-6), and 8.06 (1H, dt,  $J = 8.2, 1.3$  Hz, H-5), which was supported by the correlations

**Figure 2** Some key COSY and HMBC correlations of **2**.**Figure 3** Possible fragmentation pattern of **2**.

**Table 2** *In vitro* growth inhibitory activity of **2**.

Sample	L5178Y			Hela cell			PC12 cell		
	% Growth inhibition		ED <sub>50</sub> (μM)	% Growth inhibition		ED <sub>50</sub> (μM)	% Growth inhibition		ED <sub>50</sub> (μM)
	12.3 μM	24.6 μM		12.3 μM	24.6 μM		12.3 μM	24.6 μM	
3	81	93	0.7	74	88	0.9	37	76	2.3

observed in COSY and HMBC spectra (Fig. 2) and confirmed by the fragment ion peak at  $m/z$  274  $[M-C_6H_{13}NO_2+H]^+$  (Fig. 3). Moreover, the <sup>1</sup>H NMR spectrum (Table 1) showed two coupled methylene groups at  $\delta_H$  3.20 (2H, m, H-13) and 2.78 (2H, m, H-14), suggesting the presence of ethylene moiety. The connectivity of the ethylene group with the pyridoacridine moiety was confirmed through the HMBC correlations of H-13 and H-14 to C-9 ( $\delta_C$  72.5) and H-13 to C-8a ( $\delta_C$  162.7) and C-9a ( $\delta_C$  146.2). The location of the isopropyl moiety was established by the HMBC of H-17 and H-18 showed cross peaks with C-16 ( $\delta_C$  39.9) and C-15 ( $\delta_C$  174.3). By comparison of the literature data with the data obtained from the <sup>1</sup>H NMR, COSY, HMQC, and HMBC, the structure of **2** was unambiguously elucidated. Thus, **2** was identified as a new natural product, for which we propose the name sagitol C.

The known compounds were identified by an analysis of the spectroscopic data 1D and 2D NMR and comparison of their data with those in the literature to be: kuanoniamine C (**1**)<sup>3,15</sup> and sagitol (**3**).<sup>16</sup> The cytotoxic effect of **2** was tested against L5178Y, PC12, and Hela cell lines. It gave 93%, 88% and 76% growth suppression against the tested cell lines at a concentration of 24.6 μM and 81%, 74% and 37% at a concentration of 12.3 μM with ED<sub>50</sub>(s) of 0.7, 0.9, and 2.3 μM, respectively (Table 2).

#### 4. Conflict of interest

None.

#### References

- Kijjoa A, Wattanadilok R, Campos N, Nascimento MSJ, Pinto M, et al. Anticancer activity evaluation of kuanoniamines A and C isolated from the marine sponge *Oceanapia sagittaria*, collected from the Gulf of Thailand. *Mar Drugs* 2007;**5**:6–22.
- Nilar, Sidebottom PJ, Carte BK, Butler MS. Three new pyridoacridine type alkaloids from a Singaporean Ascidian. *J Nat Prod* 2002;**65**:1198–200.
- Eder C, Schupp P, Proksch P, Wray V, Steube K, et al. Bioactive pyridoacridine alkaloids from the micronesia sponge *Oceanapia* sp.. *J Nat Prod* 1998;**61**:301–5.
- Marshall KM, Barrows LR. Biological activities of pyridoacridines. *Nat Prod Rep* 2004;**21**:731–51.
- Singh KS, Das B, Naik CG. Quinolizidines alkaloids: petrosin and xestospongins from the sponge *Oceanapia* sp. *J Chem Sci* 2011;**123**:601–7.
- Makarievva TN, Guzii AG, Denisenko VA, Dmitrenok PS, Stonik VA. New two headed sphingolipid like compounds from the marine sponge *Oceanapia* sp. *Russian Chem Bull* 2008;**57**:669–73.
- Bensemhoun J, Bombarda I, Aknin M, Faure R, Vacelet J, et al. Marine bifunctional sphingolipids from the sponge *Oceanapia ramsayi*. *Molecules* 2008;**13**:772–8.
- Guzii AG, Makarieva TN, Denisenko VA, Svetashev VI, Rodkina SA, et al. New cerebrosides from the marine sponge *Oceanapia* sp. *Russian Chem Bull* 2006;**55**:928–33.
- Fu X, Ferreira ML, Schmitz FJ, Kelly M. Tamosterone sulfates: a C-14 epimeric pair of polyhydroxylated sterols from a new *Oceanapiid* sponge genus. *J Org Chem* 1999;**64**:6706–9.
- Matsunaga S, Okada Y, Fusetani N, van Soest RW. An antimicrobial C(14) acetylenic acid from a marine sponge *Oceanapia* species. *J Nat Prod* 2000;**63**:690–1.
- Capon RJ, Skene C, Liu EH, Lacey E, Gill JH, et al. The isolation and synthesis of novel nematocidal dithiocyanates from an Australian marine sponge, *Oceanapia* sp. *J Org Chem* 2001;**66**:7765–9.
- Ibrahim SRM, Mohamed GA, Elkhayat ES, Gouda YG, Proksch P. Strepsiamide A–C, new ceramides from the marine sponge *Strepsichordaia lendfeldi*. *Nat Prod Commun* 2008;**3**:205–9.
- Ibrahim SRM, Edrada-Ebel R, Mohamed GA, Youssef DTA, Wray V, et al. Callyaerin G, a new cytotoxic cyclic peptide from the marine sponge *Callyspongia aerizusa*. *ARKIVOC* 2008;**12**:164–71.
- Carmichael J, DeGraff WG, Gazdar AF, Minna JD, Mitchell JB. Evaluation of a tetrazolium-based semi-automated colorimetric assay: assessment of radio sensitivity. *Cancer Res* 1987;**47**:943–6.
- Bishop MJ, Ciufolini MA. Total synthesis of kuanoniamines and dercitins. *J Am Chem Soc* 1992;**114**:10081–2.
- Salomon CE, Faulkner DJ. Sagitol, a pyridoacridine alkaloid from the sponge *Oceanapia sagittaria*. *Tetrahedron Lett* 1996;**37**:9147–8.