



Cyclic mechanical stretch reduces myofibroblast differentiation of primary lung fibroblasts

Marjolein E. Blaauboer^{a,b,*}, Theo H. Smit^c, Roeland Hanemaaijer^b, Reinout Stoop^b, Vincent Everts^a

^a Department of Oral Cell Biology, Academic Centre for Dentistry Amsterdam (ACTA), University of Amsterdam and VU University Amsterdam, Research Institute MOVE, Amsterdam, The Netherlands

^b TNO Quality of Life, Biosciences, Leiden, The Netherlands

^c Department of Orthopaedics, VU Medical Center, Research Institute MOVE, Amsterdam, The Netherlands

ARTICLE INFO

Article history:

Received 5 November 2010

Available online 20 November 2010

Keywords:

Myofibroblast

Differentiation

TGF β 1

Lung fibrosis

Cyclic mechanical loading

ABSTRACT

In lung fibrosis tissue architecture and function is severely hampered by myofibroblasts due to excessive deposition of extracellular matrix and tissue contraction. Myofibroblasts differentiate from fibroblasts under the influence of transforming growth factor (TGF) β ₁ but this process is also controlled mechanically by cytoskeletal tension. In healthy lungs, the cytoskeleton of fibroblasts is mechanically strained during breathing. In stiffer fibrotic lung tissue, this mechanical stimulus is reduced, which may influence fibroblast-to-myofibroblast differentiation. Therefore, we investigated the effect of cyclic mechanical stretch on fibroblast-to-myofibroblast differentiation.

Primary normal human lung fibroblasts were grown on BioFlex culture plates and stimulated to undergo myofibroblast differentiation by 10 ng/ml TGF β ₁. Cells were either or not subjected to cyclic mechanical stretch (sinusoidal pattern, maximum elongation 10%, 0.2 Hz) for a period of 48 h on a Flexercell apparatus. mRNA expression was analyzed by real-time PCR.

Cyclic mechanical loading reduced the mRNA expression of the myofibroblast marker α -smooth muscle actin and the extracellular matrix proteins type-I, type-III, and type-V collagen, and tenascin C. These outcomes indicate that fibroblast-to-myofibroblast differentiation is reduced. Cyclic mechanical loading did not change the expression of the fibronectin ED-A splice variant, but did decrease the paracrine expression of TGF β ₁, thereby suggesting a possible regulation mechanism for the observed effects. The data suggest that cyclic loading experienced by healthy lung cells during breathing may prevent fibroblasts from differentiating towards myofibroblasts.

© 2010 Elsevier Inc. Open access under the [Elsevier OA license](http://creativecommons.org/licenses/by/3.0/).

1. Introduction

Fibrosis is a process of deregulated wound healing after inflammation or other injury, resulting in the formation of excessive extracellular matrix (ECM) and tissue contraction. The disease progresses rapidly and can occur in many different tissues. Due to the destruction of tissue architecture, it results in severe tissue dysfunction and, depending on the tissue, may lead to death. One of the most fatal forms is lung fibrosis, which has an incidence of 5–10 per 100,000 and is often fatal within 3–5 years after diagnosis [1].

The exact cause of lung fibrosis is still mostly unknown but almost certainly multifactorial. So far, there is no effective treatment. Recent investigation of the cellular processes underlying the

formation of fibrotic tissue has shown that, after minor damage to the lung tissue, fibroblasts inside the parenchymal tissue of the lung differentiate into myofibroblasts [2]. These cells are specialized in wound healing and produce large amounts of extracellular matrix proteins, such as collagens, tenascin C and the ED-A splice variant of fibronectin. They also express α -smooth muscle actin (α SMA), which enables the cells to strongly contract the wound during wound healing. After tissue repair and once the balance of remodeling is restored in the repaired tissue the myofibroblasts undergo apoptosis. However, in fibrosis, myofibroblasts stay active after the wound has healed, creating excessive, strongly contracted extracellular matrix.

Fibroblast-to-myofibroblast differentiation is regulated by both chemical and mechanical factors [2]. Chemically, transforming growth factor (TGF) β ₁ is the major molecule stimulating myofibroblast differentiation. It is present in a latent complex in the ECM, from which it can be released to locally induce myofibroblast differentiation. TGF β ₁ can be released either by proteases [3–5] or through mechanical tension in the tissue [6].

* Corresponding author at: Department of Oral Cell Biology, Academic Centre for Dentistry Amsterdam (ACTA), University of Amsterdam and VU University Amsterdam, Research Institute MOVE, Gustav Mahlerlaan 3004, 1081 LA Amsterdam, The Netherlands. Fax: +31 20 5980333.

E-mail address: me.blaauboer@acta.nl (M.E. Blaauboer).

Besides playing a role in latent TGF β ₁ release, mechanical tension also directly influences myofibroblast differentiation: in splinted wounds, the static tension in the tissue results in more ED-A fibronectin and α SMA expression, indicating more myofibroblast differentiation [7]. However, in lung tissue, the breathing cycle will result in external loads on the lung cells of a cyclic nature. Because cells will have trouble adapting to the constantly changing tensional status of external cyclic loading [8], this cyclic loading could prevent the build-up of internal cytoskeletal tension and thus myofibroblast differentiation. Therefore, we aimed to establish whether cyclic mechanical stretch regulates the differentiation of lung fibroblasts into myofibroblasts. To study this, we investigated the effect of cyclic mechanical stretch on the myofibroblast marker α SMA, and extracellular matrix proteins such as collagens, tenascin C and the ED-A splice variant of fibronectin, which has been shown to be essential for fibrosis [9].

2. Materials and methods

2.1. Cell culture

Primary normal human lung fibroblast cells were obtained from Lonza Walkersville, Inc. (Walkersville, MD, USA). Cells were cultured in Dulbecco's minimal essential medium (D-MEM; Invitrogen, Paisley, UK) supplemented with 10% fetal clone serum (FCS; HyClone, South Logan, UT, USA) and 1% antibiotic–antimycotic solution (100 U/ml penicillin, 100 μ g/ml streptomycin, and 250 ng/ml amphotericin B (PSA), Sigma–Aldrich, St. Louis, MO, USA) in an incubator set at 37 °C, 95% humidity, and 5% CO₂. Once grown to confluency, cells were trypsinized using 0.5% trypsin (Sigma–Aldrich) and 0.1% EDTA (Merck, Darmstadt, Germany) in PBS (Invitrogen).

For each experiment, cells from passage 6 to 10 were seeded at a density of 3.000 or 10.000 cells/cm² on BioFlex collagen I coated six-well culture plates (BioFlex, Flexcell International Corp, McKeesport, PA, USA). After allowing the cells to attach for 24 h, medium was replaced by D-MEM with 1% FCS and 1% PSA and 24 h later myofibroblast differentiation was induced in 50% of the samples by replacing the medium with D-MEM supplemented with 1% FCS and 1% PSA containing 2–20 ng/ml recombinant human TGF β ₁ (PeproTech EC, London, UK). Fibroblasts were cultured for 4–72 h in the presence or absence of TGF β ₁.

2.2. Cyclic mechanical loading

To assess the effect of cyclic mechanical loading on the expression of fibrosis-related genes, fibroblasts cells were subjected to 24 or 48 h of cyclic mechanical strain in a Flexercell FX4000 apparatus (Flexcell International Corp) placed in an incubator set at 37 °C, 95% humidity and 5% CO₂. Cyclic mechanical loading was applied in a sinusoidal pattern with a frequency of 0.2 Hz and a maximum elongation of 10%. Control cells not receiving cyclic mechanical loading were placed in the same incubator next to the loading station of the Flexercell apparatus. During mechanical loading, 50% of the samples were stimulated with 10 ng/ml TGF β ₁. After mechanical loading, cell viability was checked with a WST-1 assay (Roche Diagnostics, Mannheim, Germany) and mRNA samples were collected.

2.3. Quantitative real-time PCR

mRNA was isolated using an RNeasy Mini Kit for RNA extraction (Qiagen, Hilden, Germany). The mRNA concentration was measured using a Nanodrop spectrophotometer (NanoDrop Technologies; Thermo-Fischer Scientific, Wilmington, DE, USA). mRNA

was reverse-transcribed to complementary DNA (cDNA) using a High Capacity RNA-to-cDNA Kit (Applied Biosystems, Foster City, CA, USA). mRNA expression of COL1A1, COL1A2, COL3A1, COL5A2, PLOD2, α SMA, tenascin C (TNC), TGF β ₁, TGF β ₂, TGF β ₃, fibronectin, and fibronectin ED-A and the housekeeping gene GAPDH was analyzed by Real-Time PCR performed on a 7500 Fast Real-Time PCR system (Applied Biosystems). GAPDH mRNA expression was determined using TaqMan[®] Rodent GAPDH Control Reagents (Applied Biosystems). Fibronectin and its ED-A splice variant were analyzed using custom designed primers (Applied Biosystems) in combination with SYBR green (Invitrogen). All other genes were analyzed using unique TaqMan[®] Assays-on-Demand™ Gene Expression kits (Table 1; Applied Biosystems) specific for human.

2.4. Statistics

Statistical analyses were performed using SPSS (SPSS Inc., Chicago, IL, USA). Groups were compared using ANOVAs with Bonferroni-adjusted *t*-tests as post hoc tests.

3. Results

3.1. TGF β ₁ increases myofibroblast differentiation

Culturing lung fibroblasts for 24 h with different concentrations of TGF β ₁ resulted in an induction of myofibroblast differentiation at all concentrations studied (0, 2, 5, 10, and 20 ng/ml) as indicated by an increase in mRNA expression of COL1A1 and α SMA (data not shown). Time-line experiments showed that COL1A1 mRNA expression was already increased after 4 h of TGF β ₁ stimulation, whereas the mRNA expression of α SMA was increased from 24 h on (Fig. 1A and B).

3.2. Cyclic mechanical loading reduces myofibroblast differentiation

Cyclic mechanical loading decreased mRNA expression of COL1A1, COL1A2, COL3A1, COL5A2, and tenascin C (TNC), in both TGF β ₁ stimulated and non-stimulated cells (Fig. 2). The effect of cyclic mechanical loading was most pronounced for COL1A1. mRNA expression of the myofibroblast marker α SMA was also lower after cyclic mechanical loading. mRNA expression of PLOD2, an enzyme involved in collagen cross-linking, was not influenced by cyclic mechanical loading. The WST-1 assay showed no cell loss after 48 h of loading (data not shown). Taken together, these data indicate that cyclic mechanical loading reduces TGF β ₁-induced myofibroblast differentiation.

3.3. Cyclic mechanical loading does not influence fibronectin ED-A mRNA expression

Both normal fibronectin (from 4 h on) and the ED-A splice variant (from 24 h on) mRNA levels were higher when cells were stimulated with TGF β ₁ (Fig. 3A and B). However, these two fibronectin forms were not significantly reduced after 48 h of cyclic mechanical loading (Fig. 3C and D).

3.4. Cyclic mechanical loading reduces endogenous TGF β ₁ mRNA expression

Since TGF β ₁-induced myofibroblast differentiation was repressed by cyclic mechanical loading, we examined whether this could be explained by a loading-induced change in autocrine TGF β ₁ stimulation. We first analyzed the mRNA expression of the different isoforms of TGF β and the effect of exogenous TGF β ₁ hereupon. TGF β ₁, TGF β ₂, and TGF β ₃ mRNA expression was measured after 4,

Table 1

Gene product, Gene, Entrez Gene ID, primers, and Assay-on-Demand™ used in this study.

Gene product	Gene	Gene ID	Forward primer	Reverse primer
Fibronectin	FN1	2335	CaggATCACTTACggAgAgACA	gCCAgTgACAgCATAACAgTg
Fibronectin ED-A	FN1	2335	CCCTAAaggACTggCATTCA	CATCCTCAgggCTCgAgTAg
Gene product	Gene	Gene ID	Assay-on-Demand™	
α_1 Chain type-I collagen	COL1A1	NM_000088.3	Hs01076777_m1	
α_2 Chain type-I collagen	COL1A2	NM_000089.3	Hs01028956_m1	
α_1 Chain type-III collagen	COL3A1	NM_000090.3	Hs00164103_m1	
α_2 Chain type-V collagen	COL5A2	NM_000393.3	Hs00169768_m1	
Lysyl hydroxylase 2	PLOD2	5352	Hs01118190_m1	
α -Smooth muscle actin	ACTA2	59	Hs00426835_g1	
Tenascin C	TNC	NM_002160.2	Hs01115665_m1	
TGF β_1	TGFB1	NM_000660.4	Hs99999918_m1	
TGF β_2	TGFB2	7042	Hs00234244_m1	
TGF β_3	TGFB3	NM_003239.2	Hs00234245_m1	




Fig. 1. TGF β_1 increases mRNA expression of collagen type-I α_1 and α -smooth muscle actin. Stimulation of normal human lung fibroblasts with application of 10 ng/ml transforming growth factor (TGF) β_1 (\blacktriangle) in the medium significantly increases the expression of (A) collagen type-I α_1 (COL1A1) and (B) α -smooth muscle actin (α SMA) at different time points (4, 8, 24, 48, and 72 h), compared no TGF β_1 stimulation (\blacksquare). Gene expression is shown relative to mRNA expression of the household gene GAPDH. Data are shown as mean \pm SD ($n = 3$). Significant differences with 0 ng/ml TGF β_1 are indicated. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.




Fig. 2. Cyclic mechanical loading reduces myofibroblast differentiation. Stimulating normal human lung fibroblasts for 48 h with cyclic mechanical loading reduces mRNA expression of COL1A1, COL1A2, COL3A1, COL5A2, α -smooth muscle actin (α SMA) and tenascin C (TNC), independent of 10 ng/ml TGF β_1 stimulation. mRNA expression of PLOD2, encoding for an enzyme important for collagen cross-linking, is not influenced by cyclic mechanical loading. Gene expression is shown relative to mRNA expression of the household gene GAPDH. Data are from one representative experiment out of three and shown as mean \pm SD ($n = 6$). Significant differences are indicated. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.




Fig. 3. TGFβ₁ stimulation but not cyclic mechanical loading influences the expression of fibronectin and its ED-A splice variant in fibroblasts. Stimulation of normal human lung fibroblasts with application of 10 ng/ml transforming growth factor (TGF) β₁ (▲) in the medium significantly increases the expression of (A) fibronectin (FN) and (B) the ED-A splice variant of fibronectin (FN ED-A) at different time points (4, 8, 24, 48, and 72 h), compared to no TGFβ₁ stimulation (■). Cyclic mechanical strain (48 h) does not change (C) fibronectin or (D) fibronectin ED-A expression. Gene expression is shown relative to mRNA expression of the household gene GAPDH. Data are from one representative experiment out of three and shown as mean ± SD (n = 3 or 6). Significant differences between 0 and 10 ng/ml are indicated. *p < 0.05, **p < 0.01, ***p < 0.001.




Fig. 4. TGFβ₁ stimulation and cyclic mechanical loading influence the expression of TGFβs in fibroblasts. Stimulation of normal human lung fibroblasts with application of 10 ng/ml transforming growth factor (TGF) β₁ (▲) in the medium significantly increases the expression of (A) TGFβ₁ and (B) TGFβ₂ and reduces the expression of (C) TGFβ₃ at different time points (4, 8, 24, 48, and 72 h), compared to no TGFβ₁ stimulation (■). Cyclic mechanical strain (48 h) reduces (D) TGFβ₁ expression only in the presence of external TGFβ₁ and only slightly. (E) TGFβ₃ expression is reduced by 48 h of cyclic mechanical strain both in the absence and presence of external TGFβ₁. Gene expression is shown relative to mRNA expression of the household gene GAPDH. Data are from one representative experiment out of three and shown as mean ± SD (n = 3 or 6). Significant differences between 0 and 10 ng/ml are indicated. *p < 0.05, **p < 0.01, ***p < 0.001.

8, 24, 48 or 72 h of culturing in the presence or absence of 10 ng/ml TGF β ₁ (Fig. 4A–C). Both TGF β ₁ and TGF β ₂ mRNA expression were higher after stimulation with TGF β ₁, whereas the expression of TGF β ₃ was lower in the presence of TGF β ₁. Cyclic mechanical stretch lowered the expression of TGF β ₁ when cells were cultured with exogenous TGF β ₁ (Fig. 4D). mRNA expression of TGF β ₃ decreased also under stretch conditions (Fig. 4E).

4. Discussion

In this study, we aimed to establish the effect of cyclic mechanical loading on myofibroblast differentiation in lung fibroblasts. We show that application of 48 h of cyclic mechanical loading to primary human lung fibroblasts reduces the expression of fibrosis-related genes. Our findings thus indicate a reduced myofibroblast differentiation under these conditions. Although cyclic mechanical loading did not affect fibronectin ED-A mRNA expression, it did reduce endogenous expression of TGF β ₁. The reduced myofibroblast differentiation may thus be the consequence of a reduced autocrine TGF β ₁ stimulation.

In our experiments, lung fibroblasts were subjected to a cyclic mechanical loading regime consisting of a sinusoidal wave-form with a frequency of 0.2 Hz and amplitude of 10% elongation. The frequency was chosen to represent a normal breathing cycle that lasts 5 s. The chosen elongation lies within the elongation range of the epithelial basement membrane measured during breathing at 40–100% of the total lung capacity [10]. We therefore believe that this cyclic mechanical loading regime represents the loading that lung fibroblasts undergo in healthy tissue *in vivo*.

To explore a possible mechanism for the intriguing finding that myofibroblast differentiation reduced under stretching conditions, we tested whether fibronectin ED-A mRNA expression was influenced in our experiments. Since fibronectin ED-A has been shown to be essential for myofibroblast differentiation [9], a reduction of fibronectin ED-A caused by cyclic mechanical loading could be a mechanism. However, mRNA expression of fibronectin ED-A proved to be unaffected under cyclic mechanical stretch conditions (see Fig. 3D), thus making it very unlikely to play a role in the reduced differentiation.

A second mechanism that could explain a reduced myofibroblast differentiation after mechanical stretch is related to the level of TGF β in the cell culture system. Next to exogenously added growth factor the cells have the capacity to express endogenous TGF β . Here, it was found that lung fibroblasts indeed express higher levels of TGF β ₁ and TGF β ₂ mRNA in response to exogenous TGF β ₁, suggesting a positive autocrine feedback mechanism. The expression of TGF β ₃ was found to be lower after stimulation by exogenous TGF β ₁, which is in line with the suggestion that TGF β ₃ can act as an antagonist of TGF β ₁ [11]. We then explored the possibility that cyclic mechanical stretch affects this positive autocrine feedback by reducing TGF β ₁ expression. After cyclic mechanical stretch, the mRNA expression of TGF β ₁ was indeed found to be lower (see Fig. 4D). These findings appear to provide an explanation for the reduced myofibroblast differentiation after cyclic mechanical loading. Surprisingly, the mRNA expression of TGF β ₃ is also lower after cyclic mechanical loading. This contradicts the earlier mentioned alleged antagonistic effect of TGF β ₃.

Although we found a consistent reduction in expression of myofibroblast-related genes after cyclic mechanical loading, others observed that expression of collagens [12,13] or tenascin C [14–16] was higher after fibroblasts were subjected to a comparable loading regime. Several differences in experimental set-up could have led to this discrepancy. In our experiments we tried to optimally mimic the *in vivo* conditions for our lung fibroblasts by culturing them on collagen type-I coated membranes, an extracellular

matrix protein abundantly expressed in the lung connective tissue, and applying a loading regime mimicking breathing. The other studies have used fibroblasts from different origin, a shorter loading protocol, and a different coating on the membranes. Such a different coating is recognized by different integrins [17,18]. Since integrins are known to play an important role in mechanotransduction, this could be an important factor influencing the effect of mechanical loading on fibroblastic cells.

Our results show that cyclic mechanical loading resulted in a lower expression of myofibroblast-related genes, such as α SMA, collagens, and tenascin C. During the development of lung fibrosis *in vivo*, tissue stiffens and fibroblasts experience probably less cyclic mechanical loading during the breathing cycle. In light of our findings we hypothesize that this facilitates fibroblast-to-myofibroblast differentiation, resulting in a positive feedback loop with the devastating disease of fibrosis as the end result.

In summary, cyclic mechanical stretch reduces fibroblast-to-myofibroblast differentiation of primary lung fibroblasts. This may result from a decreased autocrine TGF β ₁ stimulation. Our findings suggest that tissue stiffening during fibrosis facilitates fibroblast-to-myofibroblast differentiation.

References

- [1] M. Selman, V.J. Thannickal, A. Pardo, D.A. Zisman, F.J. Martinez, J.P. Lynch III, Idiopathic pulmonary fibrosis: pathogenesis and therapeutic approaches, *Drugs* 64 (2004) 405–430.
- [2] J.J. Tomasek, G. Gabbiani, B. Hinz, C. Chaponnier, R.A. Brown, Myofibroblasts and mechano-regulation of connective tissue remodelling, *Nat. Rev. Mol. Cell Biol.* 3 (2002) 349–363.
- [3] G. Ge, D.S. Greenspan, BMP1 controls TGF β 1 activation via cleavage of latent TGF β -binding protein, *J. Cell Biol.* 175 (2006) 111–120.
- [4] D. Mu, S. Cambier, L. Fjellbirkeland, J.L. Baron, J.S. Munger, H. Kawakatsu, D. Sheppard, V.C. Broaddus, S.L. Nishimura, The integrin α (v) β 8 mediates epithelial homeostasis through MT1-MMP-dependent activation of TGF- β 1, *J. Cell Biol.* 157 (2002) 493–507.
- [5] Q. Yu, I. Stamenkovic, Cell surface-localized matrix metalloproteinase-9 proteolytically activates TGF- β and promotes tumor invasion and angiogenesis, *Genes Dev.* 14 (2000) 163–176.
- [6] P.J. Wipff, D.B. Rifkin, J.J. Meister, B. Hinz, Myofibroblast contraction activates latent TGF- β 1 from the extracellular matrix, *J. Cell Biol.* 179 (2007) 1311–1323.
- [7] B. Hinz, D. Mastrangelo, C.E. Iselin, C. Chaponnier, G. Gabbiani, Mechanical tension controls granulation tissue contractile activity and myofibroblast differentiation, *Am. J. Pathol.* 159 (2001) 1009–1020.
- [8] R.A. Brown, R. Prajapati, D.A. McGrouther, I.V. Yannas, M. Eastwood, Tensional homeostasis in dermal fibroblasts: mechanical responses to mechanical loading in three-dimensional substrates, *J. Cell Physiol.* 175 (1998) 323–332.
- [9] A.F. Muro, F.A. Moretti, B.B. Moore, M. Yan, R.G. Atrasz, C.A. Wilke, K.R. Flaherty, F.J. Martinez, J.L. Tsui, D. Sheppard, F.E. Baralle, G.B. Toews, E.S. White, An essential role for fibronectin extra type III domain A in pulmonary fibrosis, *Am. J. Respir. Crit. Care Med.* 177 (2008) 638–645.
- [10] D.J. Tschumperlin, S.S. Margulies, Alveolar epithelial surface area–volume relationship in isolated rat lungs, *J. Appl. Physiol.* 86 (1999) 2026–2033.
- [11] M. Shah, D.M. Foreman, M.W. Ferguson, Neutralisation of TGF- β 1 and TGF- β 2 or exogenous addition of TGF- β 3 to cutaneous rat wounds reduces scarring, *J. Cell Sci.* 108 (Pt. 3) (1995) 985–1002.
- [12] P.S. Howard, U. Kucich, R. Taliwal, J.M. Korostoff, Mechanical forces alter extracellular matrix synthesis by human periodontal ligament fibroblasts, *J. Periodont. Res.* 33 (1998) 500–508.
- [13] S.G. Kim, T. Akaike, T. Sasagawa, Y. Atomi, H. Kurosawa, Gene expression of type I and type III collagen by mechanical stretch in anterior cruciate ligament cells, *Cell Struct. Funct.* 27 (2002) 139–144.
- [14] M. Chiquet, A. Sarasa-Renedo, V. Tunc-Civelek, Induction of tenascin-C by cyclic tensile strain versus growth factors: distinct contributions by Rho/ROCK and MAPK signaling pathways, *Biochim. Biophys. Acta* 1693 (2004) 193–204.
- [15] R. Lutz, T. Sakai, M. Chiquet, Pericellular fibronectin is required for RhoA-dependent responses to cyclic strain in fibroblasts, *J. Cell Sci.* 123 (2010) 1511–1521.
- [16] A. Sarasa-Renedo, V. Tunc-Civelek, M. Chiquet, Role of RhoA/ROCK-dependent actin contractility in the induction of tenascin-C by cyclic tensile strain, *Exp. Cell Res.* 312 (2006) 1361–1370.
- [17] E.C. Breen, Mechanical strain increases type I collagen expression in pulmonary fibroblasts *in vitro*, *J. Appl. Physiol.* 88 (2000) 203–209.
- [18] A. van der Flier, A. Sonnenberg, Function and interactions of integrins, *Cell Tissue Res.* 305 (2001) 285–298.