

Patient education was beneficial in primary care in improving self-reported health and reducing health service utilisation. No improvements were found for self efficacy. Improvements in pain and social outcomes were found for case management (GP and nurse). Primary outcome measures included quality of life (study 1 and study 3) and self reported disability (study 2). There were a range of different patient reported outcome measures across studies.

Conclusions: A systematic review identified three trials investigating multidisciplinary approaches to multisite or generalised OA. Trials each used core interventions endorsed by OA guidelines. All three trials targeted osteoarthritis of the hand, hip or knee. Despite each of the three studies having an educational intervention, the studies were very heterogeneous. There are currently very few studies that target multidisciplinary approaches to multisite osteoarthritis. A consistent approach to outcome measurement in future studies of multisite OA is needed.

381 DEMOGRAPHIC AND CLINICAL CHARACTERISTICS OF PRIMARY AND SECONDARY CARE PATIENTS WITH OSTEOARTHRITIS OF HIP AND KNEE ARE QUITE SIMILAR

S. Lanson¹, C. Lucas², C. Veenhof¹. ¹NIVEL (Netherlands Inst. for Hlth.Services Res.), Utrecht, Netherlands; ²Dept. of Clinical Epidemiology, Univ. of Amsterdam, Amsterdam, Netherlands

Purpose: As is stated in many guidelines for patients with osteoarthritis, it is advised to treat patients conservatively before patients are referred to secondary care. Therefore, it can be expected that secondary care patients with osteoarthritis differ from primary care patients. The purpose of the current study is to compare demographic and clinical characteristics of patients with osteoarthritis who are referred to primary care physiotherapists and patients who are referred to secondary care.

Methods: In a cross-sectional study secondary analyses were performed on two cohorts of patients with knee or hip osteoarthritis: one primary care dataset (n=200) and one secondary care dataset (n=279). In both studies comparable measurements were performed at baseline in the same time period. Measurements included, among others, gender, age, location and duration of osteoarthritis, number of comorbidities, pain (VAS), physical function (WOMAC), muscle strength, range of motion, social support and quality of life.

Results: Both groups were similar in most characteristics, with some exceptions. The primary care group consisted of less patients with both hip and knee osteoarthritis and of younger patients compared to the secondary care group. Furthermore, secondary care patients reported more pain (4.0 versus 4.8), less muscle strength and more falls in the last six months.

Conclusions: There are differences between primary and secondary care populations of patients with osteoarthritis, but these differences are minimal. Looking at the results of this research, the question arises whether the health care of patients with osteoarthritis can be organised more efficient and less expensive. However, long-term prospective studies are needed to evaluate the cost effectiveness of both primary and secondary care of patients with osteoarthritis.

382 LONGITUDINAL CHANGES IN PRESCRIPTION USE IN THE OSTEOARTHRITIS INITIATIVE (OAI)

F.R. Saunders, J.E. Jeffrey, R.M. Aspden, J.S. Gregory. Univ. of Aberdeen, Aberdeen, United Kingdom

Purpose: To investigate the longitudinal changes in prescription use in the OAI study

Methods: 48 month medication data were added to data collected for previous clinic visits of the OAI study. The data were sorted by drug classes with particular focus on NSAIDs, Statins and Bisphosphonates, chosen for their association with musculoskeletal diseases. Chi-squared statistics were used to compare groups at baseline and 48 months.


Results: There was an overall decrease in the number of prescriptions for both NSAIDs and Bisphosphonates from baseline to 48 months (by 48%

($P<0.001$) and 37 % ($P=0.264$) respectively), however the number of prescriptions for Statins increased by 58 % from baseline to 48 months ($P<0.001$) (Figure 1).

Analysis showed there were significant differences between prescribing patterns by sex. Statin prescriptions for men and women increased significantly between baseline and 48 months ($P<0.001$) but NSAID prescriptions decreased significantly only in women at 48 months compared with baseline for females ($P<0.001$). The decrease in Bisphosphonate prescriptions from baseline to 48 months were not significant in females ($P=0.59$) or males ($P=0.062$).

The differences between cohorts were analysed at each time point and each of the three drug classes showed significant differences ($P<0.001$). Subset analysis, however, found that both NSAIDs and Bisphosphonates were significantly lower in the progression than the incident cohort (both $P0.05$).

Conclusions: Over the duration of the OAI study the number of Statin prescriptions has significantly increased, whilst NSAID and Bisphosphonate prescriptions have decreased. The increase in Statin prescriptions is likely to be due to two factors, the increase in prescribing these drugs to a normal population for prevention of disease and an aging population. The decrease in the numbers of Bisphosphonate prescriptions may be attributable to patients having a drug holiday, although we do not know for how long the patients have been on Bisphosphonates prior to study entry. These temporal changes in prescription drug use should be borne in mind when analysing the OAI dataset.



383 PSYCHOLOGICAL FACTORS ASSOCIATED WITH DAILY STEP COUNT IN KNEE OSTEOARTHRITIS

Y. Ahamed¹, T. Egerton¹, M.A. Hunt², F.J. Keefe³, C. Bryant¹, G. Jull⁴, P. Connellan⁴, K.L. Bennell¹. ¹Univ. of Melbourne, Melbourne, Australia; ²Univ. of British Columbia, Vancouver, BC, Canada; ³Duke Univ., Durham, NC; ⁴Univ. of Queensland, Brisbane, Australia

Purpose: Despite strong evidence of the effectiveness of daily walking for alleviating self-reported pain and improving physical function for people with knee osteoarthritis (OA), daily physical activity levels remain below recommendations for this population. OA leads to psychological as well as physical dysfunction and while psychological factors are known to influence physical activity levels in other populations, little is known about the extent to which psychological factors influence walking levels among people with knee OA. This knowledge is important as greater understanding of modifiable factors that influence daily walking levels may enable interventions to increase walking behaviors to be better planned and targeted. Previous studies exploring associations between psychological factors and physical activity levels among people with knee OA, using a range of self-report and accelerometer-based physical activity measures, have reported that depression, anxiety and negative affect (encompassing depression, tension, fatigue and anger) are not associated with physical activity levels. These studies have not considered many other important psychological variables, which may be important modifiable mediators of physical activity for people with knee OA. In addition, none

focused primarily on walking behavior. Daily step count recording with a pedometer offers a low cost convenient method of quantifying the amount of walking-related physical activity. The purpose of this study was to identify the role of psychological factors in determining average daily step count among people with painful knee OA.

Methods: Our sample consisted of 166 people with painful knee OA: aged ≥ 50 years, visual analogue scale for pain $\geq 40/100$, radiographic Kellgren & Lawrence grade ≥ 2 , and Western Ontario & McMaster Physical Function (WOMAC PF) ≥ 25 . A multiple regression analysis was carried out with average daily step count recorded over 7 days using a pedometer as the dependent variable. Psychological variables were entered as independent variables after entering age, body mass index, pain (WOMAC pain score), and physical function (WOMAC PF score) as covariates. These demographic variables are known to be associated with physical activity in knee OA. Psychological variables included the subscales of the Depression, Anxiety & Stress (DASS) scale (Depression, Anxiety and Stress), Arthritis Self-Efficacy scale (pain, function and other), Pain Catastrophizing Scale (Rumination, Magnification and Helplessness) and Attitudes to Physical Activity Scale (Self-efficacy, barriers and benefits).

Results: DASS Stress subscale accounted for 6% of the variance in step count after accounting for the covariates (15% of variance) and was the only significant psychological contributor ($\beta = -0.25$, $SE = 52.3$, $p < 0.05$). Higher levels of stress were associated with lower daily step counts.

Conclusion: After controlling for demographic and symptomatic variables, only stress was associated with average daily step count. Our findings are consistent with previous studies showing no associations between depression or anxiety and physical activity levels of people with knee OA. The findings indicate that stress may be an important psychological factor that needs to be addressed in order for people with knee OA to increase their amount of daily walking as a means to becoming more physically active.

384 DEVELOPMENTAL ORIGINS OF SELF-REPORTED ARTHRITIS AND OSTEOARTHRITIS: NEW ENGLAND FAMILY STUDY

C.B. Eaton¹, M.B. Roberts², J.B. Driban³, G. Agha⁴, E. Loucks⁴, S. Buka⁴, ¹Warren Alpert Med. Sch. of Brown Univ., Providence, RI, USA; ²Mem. Hosp. of Rhode Island, Pawtucket, RI, USA; ³Tufts Med. Ctr., Boston, MA, USA; ⁴Brown Univ., Providence, RI, USA

Purpose: Low and high birthweights have been linked to adverse health outcomes in adulthood, including body composition, obesity and diabetes. Fetal programming related to maternal stress and under- or over-nutrition during middle to late gestation has been proposed as a possible mechanism for these associations. Animal ecology studies show an association between early life poor nutrition and greater OA prevalence later in life. We hypothesized that osteoarthritis linked to both bone and cartilage metabolism might also be operating under such a developmental origins hypothesis in humans.

Methods: Birth weight (an indicator of fetal growth) and maternal smoking status (a measure of intra-uterine stress) were assessed in 609 offspring in Providence, Rhode Island as part of the Collaborative Perinatal Project (CPP) between 1959 and 1966 and self-reported arthritis was prospectively ascertained on average 42 years later. Small for gestational age (SGA) was defined as ≤ 10 th%ile and LGA as ≥ 90 th%ile for the entire Providence CPP cohort ($n=4,184$). Maternal smoking during each trimester was also assessed via self report. Self-reported arthritis and osteoarthritis was based upon the question derived from the National Health Interview Survey: Have you ever been told by a doctor or health professional that you have arthritis or rheumatism? If yes, what type of arthritis do/did you have? Rheumatoid arthritis (RA), osteoarthritis(OA), another type of arthritis? (please specify). Logistic regression analysis determined odds ratios (OR) for risk of self-reported arthritis and OA adjusting for age, race, gender, and educational attainment for both birthweight adjusted for gestational age (model 1) and maternal smoking (model 2). An additional covariate of maternal SES was added to the maternal smoking analysis. Those reporting RA, both RA and OA, and other arthritis were excluded from the OA analysis.

Results: Of the 609 offspring with self-reported arthritis data, 52 reported a physician diagnosis of OA, 45 RA, 5 reported both and 8 reported other

types of arthritis and 486 reported no arthritis. A modest signal that was not statistically significant was found for associations of birthweight adjusted for gestational age and maternal smoking with self-reported arthritis and for maternal smoking with OA. (see table)

Conclusion: We found modest evidence to support the developmental origins of arthritis hypothesis in this preliminary analysis. The small sample size, young age of our cohort and the lack of x-ray documentation of self-reported arthritis and osteoarthritis limit the inferences that can be drawn from our pilot study. If future larger studies find stable associations between intra-uterine exposures and OA, then epigenetic mechanisms to explain these results should be pursued.

Odds Ratio and 95% CI of self-reported arthritis and OA

Self-reported Arthritis	OR	95% CI	Self-reported OA	OR	95% CI
LGA vs AGA	1.03	0.51-2.09		0.56	0.16-1.92
SGA vs AGA	1.16	0.59-2.29		0.86	0.28-2.59
Model 2					
Maternal Smoking	1.04	0.65-1.65		1.06	0.55-2.05

385 ASSOCIATIONS BETWEEN DIETARY PATTERNS, TEA & COFFEE DRINKING AND OSTEOARTHRITIS

A. Wood, R. Aspden, D.M. Reid, H.M. Macdonald, University of Aberdeen Musculoskeletal Research, Univ. of Aberdeen, Aberdeen, United Kingdom

Purpose: There are few studies that have explored the relationship between diet and prevalence of osteoarthritis (OA). Although it is usual to study individual nutrients, a dietary pattern approach has the advantage of assessing food combinations, and it avoids the errors associated with nutrient databases. We have shown that dietary patterns were associated with markers of bone health: a healthy diet being related to lower bone turnover; and both a processed food diet, and a diet low in milk and tea consumption, being associated with lower hip bone mineral density (BMD). The aim of this study was to test the relationship between dietary patterns and osteoarthritis.

Methods: Data from the 1998-2000 visit of the Aberdeen Prospective Osteoporosis Screening Study (APOSS) were used for this study. Women were aged between 50–62 y. They were asked whether they suffered from OA. This was validated by radiological records in a subset who also stated the site of OA (which included hands, hips, knees, feet, spine, shoulders and elbows). Diet was assessed using food frequency questionnaires (FFQ) ($n=3229$) and dietary patterns were generated by principal components analysis using energy-adjusted food intakes. Logistic regression was used to test the relationship between dietary pattern and prevalence of self reported OA.

Results: A total of 596 women had self reported OA compared to 2633 who reported no OA. Five dietary patterns (accounting for 26% of the total variance in diet) were tested with OA prevalence, using logistic regression. One dietary pattern, which had negative factor loadings for milk, tea and cereal, and positive loadings for crisps/ nuts, sauce/ jam and confectionery, was associated with lower OA prevalence. This relationship remained significant after adjustment for weight, height, age, national deprivation category, smoking, physical activity and also hip BMD. Further exploration of tea drinking showed that whereas self-reported OA was associated with more tea drinking; coffee drinking appeared to be protective with higher consumption in women who did not report OA. Further, although numbers were small both coffee drinking and being heavier were significantly associated with reduced risk of hand OA (reported hand OA, $n=148$; did not report hand OA $n=96$).

Tea and coffee consumption according to self reported osteoarthritis (OA)

Mean [SD]	OA n 596	No OA n 2633	P
Tea intake (cups/week)	18.5 [14.7]	16.3 [14.0]	<0.001
Coffee intake (cups/week)	12.8 [12.1]	14.8 [13.2]	0.007
Tea and coffee intake (cups/week)	31.4 [13.9]	31.1 [13.9]	0.701