



Contents lists available at ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra

On the Hard Lefschetz property of stringy Hodge numbers

Jan Schepers¹

Katholieke Universiteit Leuven, Departement Wiskunde, Celestijnenlaan 200B, 3001 Leuven, Belgium

ARTICLE INFO

Article history:

Received 7 July 2007

Available online 20 November 2008

Communicated by Laurent Moret-Bailly

Keywords:

Algebraic geometry

Stringy invariants

ABSTRACT

For projective varieties with a certain class of 'mild' isolated singularities and for projective threefolds with arbitrary Gorenstein canonical singularities, we show that the stringy Hodge numbers satisfy the Hard Lefschetz property (i.e. $h_{st}^{p,q} \leq h_{st}^{p+1,q+1}$ for $p+q \leq d-2$, where d is the dimension of the variety). This result fits nicely with a 6-dimensional counterexample of Mustață and Payne for the Hard Lefschetz property for stringy Hodge numbers in general. We also give such an example, ours is a hypersurface singularity.

© 2008 Elsevier Inc. All rights reserved.

1. Introduction

1.1. Stringy Hodge numbers of projective varieties with Gorenstein canonical singularities were introduced by Batyrev in [Ba]. They are defined if the *stringy E-function*, which is in general a rational function of two variables u and v , is in fact a polynomial. The idea is that they should be the Hodge numbers of a conjectural 'string cohomology.' Several constructions of such string cohomology spaces were made by Borisov and Mavlyutov in [BM], and they also made a connection to the orbifold cohomology of Chen and Ruan from [CR]. Moreover, Yasuda showed that the stringy Hodge numbers are the Hodge numbers of the orbifold cohomology for varieties with Gorenstein quotient singularities (see [Ya, Remark 1.4(2)]).

1.2. In this paper we want to study under which conditions the Hard Lefschetz property holds for stringy Hodge numbers. By the Hard Lefschetz property we mean the inequalities that are imposed if there would be an analogue of the Hard Lefschetz Theorem for the conjectural string cohomology, i.e.

$$h_{st}^{p,q} \leq h_{st}^{p+1,q+1} \quad \text{for } p+q \leq d-2,$$

E-mail address: janschepers1@gmail.com.

¹ Supported by the Netherlands Organisation for Scientific Research (NWO).

where d is the dimension of the variety. Fernandez gave in [Fe] a criterion for the Hard Lefschetz Theorem to hold for orbifold cohomology. He also shows that this criterion fails in general for generic Calabi–Yau hypersurfaces in the weighted projective space $\mathbb{P}(1, 1, 1, 3, 3)$ [Fe, Example 4.4]. However, it follows for instance from the main theorem below that this cannot be seen from the Hodge or Betti numbers of the orbifold cohomology ($h_{orb}^{p,q} = h_{st}^{p,q} \leq h_{orb}^{p+1,q+1} = h_{st}^{p+1,q+1}$ for $p + q \leq 1$). In fact, the orbifold Hodge numbers where in this case already studied by Poddar (see [Po, Section 5 and especially Corollary 2]).

1.3. Mustařa and Payne first gave an example of a 6-dimensional projective toric variety with stringy Betti number $b_{6,st} = h_{st}^{3,3}$ strictly smaller than $b_{4,st} = h_{st}^{2,2}$ (see [MP, Example 1.1]). This example was used to disprove a conjecture of Hibi on the unimodality of the so-called h -vector or δ -vector of a reflexive polytope. This vector actually gives the stringy Betti numbers of the toric variety defined by the fan over the faces of the polytope ([BD, Theorem 7.2] and [MP, Theorem 1.2]). For many more examples of reflexive polytopes with nonunimodal h -vector we refer to [Pa].

1.4. The main result of this paper is the following. The used notions are explained in Section 2.

Main theorem. *Let Y be either*

- *a projective variety of dimension $d = 3$ with Gorenstein canonical singularities, or*
- *a projective variety of dimension $d \geq 4$ with at most isolated Gorenstein singularities that admits a log resolution with all discrepancy coefficients of exceptional components $> \lfloor \frac{d-4}{2} \rfloor$. (This condition does not depend on the chosen log resolution.)*

Write the stringy E -function of Y as a power series $\sum_{i,j \geq 0} b_{i,j} u^i v^j$. Then for $i + j \leq d - 2$, we have $(-1)^{i+j} b_{i,j} \leq (-1)^{i+j+2} b_{i+1,j+1}$. In particular, if the stringy E -function of Y is a polynomial, then $h_{st}^{p,q}(Y) \leq h_{st}^{p+1,q+1}(Y)$ for $p + q \leq d - 2$.

For the proof of this theorem we refer to Section 3. In Section 4 we compare the above theorem with the example of Mustařa and Payne. We also discuss an explicit example of a 6-dimensional projective variety with an isolated canonical hypersurface singularity that does not satisfy the Hard Lefschetz property.

2. Stringy Hodge numbers

2.1. Let X be an arbitrary complex algebraic set of dimension d . It is well known that the cohomology with compact support $H_c^*(X)$ carries a natural mixed Hodge structure (we always use cohomology with coefficients in the complex numbers). The data of this mixed Hodge structure are put in the Hodge–Deligne polynomial

$$H(X; u, v) := \sum_{p,q=0}^d \left[\sum_{i=0}^{2d} (-1)^i h^{p,q}(H_c^i(X)) \right] u^p v^q,$$

where $h^{p,q}$ denotes the dimension of the (p, q) -component $H^{p,q}(H_c^i(X))$. The Hodge–Deligne polynomial is a generalized Euler characteristic: if Y is a Zariski-closed subset of X , then $H(X; u, v) = H(X \setminus Y; u, v) + H(Y; u, v)$ and for a product of algebraic sets X and X' we have $H(X \times X'; u, v) = H(X; u, v)H(X'; u, v)$.

2.2. Now let Y be a normal irreducible variety. Assume that Y is \mathbb{Q} -Gorenstein; i.e. rK_Y is a Cartier divisor for some $r \in \mathbb{Z}_{>0}$. If K_Y is already Cartier, then Y is called Gorenstein. For example, all hypersurfaces and more generally all complete intersections are Gorenstein. Choose a log resolution

$f : X \rightarrow Y$ of Y . This is a proper birational map f from a nonsingular variety X such that the exceptional locus is a divisor D with only smooth components $D_i, i \in I$, that have normal crossings. We can write

$$rK_X - f^*(rK_Y) = \sum_{i \in I} b_i D_i,$$

with all $b_i \in \mathbb{Z}$. Using \mathbb{Q} -coefficients this becomes $K_X - f^*(K_Y) = \sum a_i D_i$ with $a_i = b_i/r$. The rational number a_i is called the discrepancy coefficient of D_i . We call Y log terminal, canonical or terminal if all $a_i > -1, \geq 0$ or > 0 respectively (this does not depend on the chosen log resolution).

2.3. Definition. (See [Ba, Definition 3.1].) Let Y be log terminal. Choose a log resolution $f : X \rightarrow Y$ with irreducible exceptional components $D_i, i \in I$. Denote the discrepancy coefficient of D_i by a_i . For a subset J of I we use the notations

$$D_J := \bigcap_{j \in J} D_j \quad \text{and} \quad D_J^\circ := D_J \setminus \bigcup_{i \in I \setminus J} D_i.$$

This gives a stratification of X as $\bigsqcup_{J \subset I} D_J^\circ$. The stringy E -function of Y is defined by the formula

$$E_{st}(Y; u, v) := \sum_{J \subset I} H(D_J^\circ; u, v) \prod_{j \in J} \frac{uv - 1}{(uv)^{a_j+1} - 1}.$$

Batyrev used motivic integration to show that this formula does not depend on the chosen log resolution [Ba, Theorem 3.4].

2.4. Remark.

1. If Y is Gorenstein (and thus automatically canonical) then $E_{st}(Y; u, v)$ is a rational function in u and v . It lives in $\mathbb{Z}[[u, v]] \cap \mathbb{Q}(u, v)$.
2. If Y is smooth, then $E_{st}(Y; u, v) = H(Y; u, v)$. If Y admits a crepant resolution (i.e. a log resolution $f : X \rightarrow Y$ such that $K_X = f^*(K_Y)$) then $E_{st}(Y; u, v) = H(X; u, v)$. More generally, for a projective birational morphism $g : Y' \rightarrow Y$ from a normal variety Y' such that $K_{Y'} = g^*(K_Y)$ one has $E_{st}(Y'; u, v) = E_{st}(Y; u, v)$ [Ba, Theorem 3.12].
3. An alternative formula for $E_{st}(Y; u, v)$ is

$$E_{st}(Y; u, v) = \sum_{J \subset I} H(D_J; u, v) \prod_{j \in J} \frac{uv - (uv)^{a_j+1}}{(uv)^{a_j+1} - 1}.$$

2.5. Assume now that Y is in addition projective of dimension d . Batyrev proves the following relation in [Ba, Theorem 3.7]:

$$E_{st}(Y; u, v) = (uv)^d E_{st}(Y; u^{-1}, v^{-1}). \tag{*}$$

If Y is also Gorenstein canonical and if $E_{st}(Y; u, v)$ is a polynomial $\sum_{p,q} a_{p,q} u^p v^q$ then Batyrev defines the stringy Hodge numbers of Y as $h_{st}^{p,q}(Y) := (-1)^{p+q} a_{p,q}$. Remark the following:

- (1) If Y is smooth then the stringy Hodge numbers are equal to the usual Hodge numbers of Y . If Y has a crepant desingularization X , then the stringy Hodge numbers of Y are equal to the Hodge numbers of X .

- (2) $h_{st}^{p,q}(Y) = h_{st}^{q,p}(Y)$.
- (3) From (*) above it follows that $h_{st}^{p,q}(Y)$ can only be nonzero for $0 \leq p, q \leq d$ and that $h_{st}^{p,q}(Y) = h_{st}^{d-p,d-q}(Y)$.
- (4) From Remark 2.4(3) we have $h_{st}^{0,0}(Y) = 1$.

Batyrev also made the following very intriguing conjecture.

Conjecture. (See [Ba, Conjecture 3.10].) *Stringy Hodge numbers are nonnegative.*

2.6. Example. Canonical surface singularities are classified and are precisely the *A–D–E* singularities. It is well known that they admit a crepant resolution and thus the conjecture above is trivially true for them by Remark 2.5(1). Of course, the Hard Lefschetz property is also satisfied in this case.

In [SV1] the conjecture was proved for the same class of varieties that is treated by the main theorem of this paper, and thus also for threefolds in full generality.

3. Proof of the main theorem

3.1. Let us for convenience repeat the statement of the theorem.

Theorem. *Let Y be either*

- *a projective variety of dimension $d = 3$ with Gorenstein canonical singularities, or*
- *a projective variety of dimension $d \geq 4$ with at most isolated Gorenstein singularities that admits a log resolution with all discrepancy coefficients of exceptional components $> \lfloor \frac{d-4}{2} \rfloor$.*

Write the stringy E-function of Y as a power series $\sum_{i,j \geq 0} b_{i,j} u^i v^j$. Then for $i + j \leq d - 2$, we have $(-1)^{i+j} b_{i,j} \leq (-1)^{i+j+2} b_{i+1,j+1}$.

Remark.

- (1) In particular, if the stringy E-function of Y is a polynomial, then $h_{st}^{p,q}(Y) \leq h_{st}^{p+1,q+1}(Y)$ for $p+q \leq d - 2$.
- (2) It is not hard to check that the lower bound on the discrepancies for the second class of varieties from the theorem does not depend on the chosen log resolution. Note that isolated Gorenstein terminal four- and fivefold singularities are included in the theorem.
- (3) Note also that $b_{0,0} = E_{st}(Y; 0, 0) = 1$ by Remark 2.4(3), so $b_{i,i} \geq 1$ for $i \leq d/2$.

Proof of the theorem. Let us first treat the second case. So Y is of dimension $d \geq 4$. Let $f : X \rightarrow Y$ be a log resolution with X projective and such that f is an isomorphism when restricted to the inverse image of the nonsingular part of Y . Denote by D the total exceptional locus of f . In [SV1, Remark 3.5(3)] the following description of the numbers $(-1)^{i+j} b_{i,j}$ for $i + j \leq d$ was given in this case:

$$(-1)^{i+j} b_{i,j} = \dim \ker(H^{d-i,d-j}(H^{2d-i-j}(X)) \rightarrow H^{d-i,d-j}(H^{2d-i-j}(D))) + S_{i,j},$$

where one has to remark that

- (1) the map $H^k(X) \rightarrow H^k(D)$ induced by inclusion is surjective for $k \geq d$ and thus $H^k(D)$ carries then a pure Hodge structure of weight k ,
- (2) $S_{i,j}$ is nonnegative and for even d only nonzero for $i = j = d/2$ and for odd d only nonzero for $i = j = (d - 1)/2$ and $\{i, j\} = \{(d - 1)/2, (d + 1)/2\}$. The term $S_{i,j}$ has to be introduced for contributions of the exceptional components with the lowest allowed discrepancy coefficients. If

one does not put the above lower bound on the discrepancy coefficients, it becomes more often nonzero, much more complicated and can even be negative (see the example in [SV2]).

If we denote $\ker(H^{d-i,d-j}(H^{2d-i-j}(X)) \rightarrow H^{d-i,d-j}(H^{2d-i-j}(D)))$ by $K_{i,j}^{2d-i-j}$ then it suffices to prove that $\dim K_{i,j}^{2d-i-j} \leq \dim K_{i+1,j+1}^{2d-i-j-2}$ for $i + j \leq d - 2$. Denote $\ker(H^k(X) \rightarrow H^k(D))$ by K^k .

We will use the following construction of de Cataldo and Migliorini. Embed Y in a projective space \mathbb{P}^r and take a generic hyperplane section Y_s of Y . So Y_s is nonsingular and does not contain any of the singular points of Y . Let $X_s := f^{-1}(Y_s)$ and denote by η_s the fundamental class of X_s in $H^{1,1}(H^2(X))$. If one dualizes the surjective map $H^k(X) \rightarrow H^k(D)$ for $k \geq d$ and uses Poincaré duality on X , one obtains an injection $H_k(D) \rightarrow H^{2d-k}(X)$. Now the spaces

$$H^0(X), H^1(X), \frac{H^2(X)}{H_{2d-2}(D)}, \dots, \frac{H^{d-1}(X)}{H_{d+1}(D)}, H^d(X), \\ K^{d+1}, \dots, K^{2d-2}, H^{2d-1}(X), H^{2d}(X)$$

satisfy the Hard Lefschetz Theorem with respect to the cup product with η_s . This result is discussed by de Cataldo and Migliorini in Section 2.3 and the beginning of Section 2.4 from [dCM2] for dimensions 3 and 4, but it is not hard to see that their argument works in any dimension. It also follows from their earlier work [dCM1]; see, in particular, Section 2.4 therein. We note that the choice of the space $H^d(X)$ in the middle is somewhat arbitrary, it can be replaced by any subspace containing the image of $\frac{H^{d-2}(X)}{H_{d+2}(D)}$. We can take K^d for that. To prove this, it suffices to show that

$$H^{d-2}(X) \xrightarrow{\cup \eta_s} H^d(X) \rightarrow H^d(D)$$

forms a complex. If we dualize, this means that

$$H_d(D) \rightarrow H_d(X) \rightarrow H_{d-2}(X)$$

should be a complex as well, where $H_d(X) \rightarrow H_{d-2}(X)$ corresponds to intersecting with X_s . And this is clear.

Summarizing, we obtain that the maps $\cup \eta_s : K^{d+k} \rightarrow K^{d+k+2}$ are surjective for $0 \leq k \leq d - 2$. Since $\cup \eta_s$ is a morphism of Hodge structures of type (1,1), we also get that $\cup \eta_s : K_{i+1,j+1}^{2d-i-j-2} \rightarrow K_{i,j}^{2d-i-j}$ is surjective for $i + j \leq d - 2$.

Now let Y be a projective threefold with arbitrary Gorenstein canonical singularities. By the main theorem of [Re] we can find a projective variety Z with terminal singularities and a projective birational crepant morphism $g : Z \rightarrow Y$. So $E_{st}(Z) = E_{st}(Y)$ by Remark 2.4(2). The point is that terminal threefold singularities are automatically isolated (see for instance [Ma, Corollary 4-6-6]) and thus we can apply the above reasoning for Z (the used results of de Cataldo and Migliorini remain valid for Z , as well as the description of the numbers $(-1)^{i+j}b_{i,j}$ from [SV1] for $i + j \leq 3$, now with $S_{i,j}$ always zero). □

4. Examples

4.1. Example. We first compare Example 1.1 of [MP] with the main theorem. Let f be the vector $(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ in \mathbb{R}^6 and N the lattice $\mathbb{Z}^6 + \mathbb{Z} \cdot f$. Denote the standard basis vectors of \mathbb{R}^6 by e_1, \dots, e_6 . Mustață and Payne consider the polytope P with vertices

$$\{e_1, \dots, e_6, e_1 - f, \dots, e_6 - f\}.$$

It is a reflexive polytope and the projective toric variety Y defined by the fan Σ over the faces of the polytope has stringy E -function

$$(uv)^6 + 6(uv)^5 + 8(uv)^4 + 6(uv)^3 + 8(uv)^2 + 6uv + 1.$$

The fan Σ has eight cones of maximal dimension, namely

- a cone σ generated by e_1, \dots, e_6 ,
- τ generated by $e_1 - f, \dots, e_6 - f$,
- and six nonsimplicial cones ρ_i generated by all vectors to vertices except e_i and $e_i - f$.

As sketched in [MP, Example 3.1] one can make a triangulation of the boundary of P by taking the convex hulls of $\{e_1, \dots, \widehat{e_j}, \dots, e_k, e_k - f, \dots, e_6 - f\}$ and $\{e_1, \dots, e_j, e_j - f, \dots, \widehat{e_k - f}, \dots, e_6\}$ for $1 \leq j < k \leq 6$. This triangulation is regular in the sense of [BG, Section 1.F] as can be seen for instance by repeatedly applying Lemma 1.65 from that book. This implies that the toric variety Z given by the fan over this triangulation is projective. Moreover, the toric morphism from Z to Y is crepant, since no new rays and hence no exceptional components of codimension 1 were introduced. Thus $E_{st}(Z) = E_{st}(Y)$. Since the singular locus of Z is given by the union of the orbits of the torus action corresponding to cones in the fan that cannot be generated by a part of the basis of the lattice, we see that Z has exactly two isolated singular points coming from the cones σ and τ . The resolution of singularities of these points is particularly easy: we subdivide the fan by adding the rays generated by f and by $-f$. This does introduce two irreducible exceptional components of codimension 1 and from the theory of toric varieties it is well known that their discrepancy coefficients are 1. So the example of the variety Z shows that the lower bound on the discrepancies in the main theorem is crucial. Even for isolated singularities with a very easy resolution, the theorem cannot be extended.

4.2. We conclude this paper by presenting another 6-dimensional example that was obtained independently of the one of Mustařa and Payne. The resolution of singularities is much more complicated, but it has the advantage of being a hypersurface singularity. We will need the formula for the Hodge–Deligne polynomial of a Fermat hypersurface and for a quasihomogeneous affine hypersurface with an isolated singularity at the origin.

Denote the Fermat hypersurface of dimension d and degree l by $Y_l^{(d)}$. So $Y_l^{(d)}$ is given by

$$\{x_0^l + \dots + x_{d+1}^l = 0\} \subset \mathbb{P}_{\mathbb{C}}^{d+1}.$$

Dais shows in [Da, Lemma 3.3] that the Hodge–Deligne polynomial of $Y_l^{(d)}$ is given by

$$H(Y_l^{(d)}; u, v) := \sum_{p=0}^d u^p (v^p + (-1)^d \mathcal{G}(d+1, p+1 | l-1, p)) v^{d-p},$$

where

$$\mathcal{G}(\kappa, \lambda | \nu, \xi) := \sum_{j=0}^{\lambda} (-1)^j \binom{\kappa+1}{j} \binom{\nu(\lambda-j) + \xi}{\kappa}$$

for $(\kappa, \lambda, \nu, \xi) \in \mathbb{Z}_{\geq 0}^4$ and $\kappa \geq \lambda$ (if $m > n$, the binomial coefficient $\binom{n}{m}$ must be interpreted as 0).

Let $f \in \mathbb{C}[x_1, \dots, x_{r+1}]$ be a quasihomogeneous polynomial of degree d with respect to the weights w_1, \dots, w_{r+1} and assume that the origin is an isolated singularity of $Y := f^{-1}(0)$. According to [Da, Section 2] the Hodge–Deligne polynomial of Y equals

$$(uv)^r + (-1)^{r-1}(uv - 1) \sum_{p=0}^{r-1} \dim_{\mathbb{C}} M(f)_{(p+1)d - (w_1 + \dots + w_{r+1})} u^p v^{r-1-p},$$

where $M(f)_{(p+1)d - (w_1 + \dots + w_{r+1})}$ denotes the piece of degree $(p + 1)d - (w_1 + \dots + w_{r+1})$ of the Milnor algebra

$$M(f) := \frac{\mathbb{C}[x_1, \dots, x_{r+1}]}{\left(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_{r+1}}\right)}.$$

Indeed, this is a graded \mathbb{C} -algebra if we give x_i degree w_i . The needed dimensions can be computed from the Poincaré series

$$P_{M(f)}(t) := \sum_{k \geq 0} (\dim_{\mathbb{C}} M(f)_k) t^k,$$

which in this case simply equals

$$P_{M(f)}(t) = \frac{(1 - t^{d-w_1}) \dots (1 - t^{d-w_{r+1}})}{(1 - t^{w_1}) \dots (1 - t^{w_{r+1}})}.$$

4.3. Example. In order to construct the announced 6-dimensional projective variety Y (which, as we will explain below, does not satisfy the Hard Lefschetz property), we define the auxiliary hypersurface

$$Y' := \{x_1^5 z^3 + x_2^5 z^3 + x_3^8 + x_4^8 + x_5^8 + x_6^8 + x_7^8 = 0\} \subset \mathbb{P}^7,$$

where we consider $z = 0$ as the hyperplane at infinity. At infinity there is a singular \mathbb{P}^1 and the origin of the affine chart $z \neq 0$ is singular as well. Our variety Y will consist of a resolution of the singular \mathbb{P}^1 and will thus have one isolated hypersurface singularity. Let us first describe the resolution process at infinity. Thereby we want to compute the Hodge–Deligne polynomial of the nonsingular part Y_{ns} of Y , since the stringy E -function of Y can be written as $H(Y_{ns}; u, v) +$ contribution of the singular point. So we keep track of the contributions in every step. We blow up in the singular line. The exceptional locus after this first step consists of five disjoint components $D_1^\infty, \dots, D_5^\infty$ (all isomorphic to \mathbb{P}^5) and one other component D_6^∞ , isomorphic to a \mathbb{P}^4 -bundle over \mathbb{P}^1 and singular for the strict transform of Y' . The intersection of D_6^∞ and another D_i^∞ is isomorphic to \mathbb{P}^4 . Since we are only interested in the nonsingular part and since we will blow up in D_6^∞ in the following step, the contribution of the first step to the Hodge–Deligne polynomial of Y_{ns} will be $5(uv)^5$ (the nonsingular exceptional part consists of five disjoint \mathbb{A}^5 's). So in the second step we blow up with D_6^∞ as center. Again there appear five new disjoint components $E_1^\infty, \dots, E_5^\infty$ and one component E_6^∞ intersecting the others transversally and singular for the strict transform of Y' . Here the E_i^∞ are \mathbb{P}^1 -bundles over \mathbb{P}^4 and E_6^∞ is isomorphic to a \mathbb{P}^4 -bundle over \mathbb{P}^1 . The intersection of E_6^∞ with an E_i^∞ is isomorphic to \mathbb{P}^4 and thus the contribution of this step to the Hodge–Deligne polynomial of Y_{ns} is $5(uv)^5 + 5(uv)^4 + 5(uv)^3 + 5(uv)^2 + 5uv$. In the next step we blow up with E_6^∞ as center. There are two new exceptional components. The first, F^∞ , is a \mathbb{P}^4 -bundle over \mathbb{P}^1 that contains five disjoint singular components for the strict transform of Y' , all isomorphic to \mathbb{P}^4 . The second component G^∞ is isomorphic to a \mathbb{P}^1 -bundle over a $Y_8^{(3)}$ -bundle over \mathbb{P}^1 , where $Y_8^{(3)}$ is the Fermat hypersurface of dimension 3 and

degree 8 (notation as in 4.2). The intersection $F^\infty \cap G^\infty$ is isomorphic to a $Y_8^{(3)}$ -bundle over \mathbb{P}^1 . All of this means that the contribution of this step to the Hodge–Deligne polynomial of Y_{ns} equals

$$(uv - 4)((uv)^4 + (uv)^3 + (uv)^2 + uv + 1) + ((uv)^2 + uv)H(Y_8^{(3)}; u, v).$$

In the final step we blow up in the five remaining singular components. This gives five new disjoint exceptional components $H_1^\infty, \dots, H_5^\infty$, whose Hodge–Deligne polynomial equals

$$(uv)^5 + 2(uv)^4 + 2(uv)^3 + 2(uv)^2 + 2uv + 1 + uvH(Y_8^{(3)}; u, v).$$

So the total contribution of the exceptional locus above the singular \mathbb{P}^1 of Y' to the Hodge–Deligne polynomial of Y_{ns} is

$$16(uv)^5 + 12(uv)^4 + 12(uv)^3 + 12(uv)^2 + 12uv + 1 + ((uv)^2 + 6uv)H(Y_8^{(3)}; u, v).$$

To compute the Hodge–Deligne polynomial of Y_{ns} we must add the Hodge–Deligne polynomial of the nonsingular part of Y' . At infinity this is the double projective cone over $Y_8^{(3)}$ minus the singular \mathbb{P}^1 , with contribution

$$(uv)^2H(Y_8^{(3)}; u, v),$$

and the contribution of the nonsingular part of Y' in the affine chart $z \neq 0$ can be computed by the method of Dais from 4.2; it equals

$$(uv)^6 - 1 - (uv - 1)(140u^4v + 140uv^4 + 4060u^2v^3 + 4060u^3v^2).$$

The formula for $H(Y_8^{(3)}; u, v)$ is

$$(uv)^3 + (uv)^2 + uv + 1 - 35u^3 - 35v^3 - 1015u^2v - 1015uv^2,$$

so finally the Hodge–Deligne polynomial of Y_{ns} equals


$$A := (uv)^6 + 18(uv)^5 + 20(uv)^4 + 20(uv)^3 + 20(uv)^2 + 18uv - 210u^5v^2 - 210u^2v^5 - 6090u^4v^3 - 6090u^3v^4 - 70u^4v - 70uv^4 - 2030u^3v^2 - 2030u^2v^3.$$

Next we compute the contribution of the singular point given by the origin of

$$\{x_1^5 + x_2^5 + x_3^8 + x_4^8 + x_5^8 + x_6^8 + x_7^8 = 0\} \subset \mathbb{A}^7$$

to the stringy E -function of Y by a log resolution. We remark that one can also use the combinatorial procedure of [SV2, Section 4]. We first blow up in the singular point itself. This gives five exceptional components D_1, \dots, D_5 , all isomorphic to \mathbb{P}^5 and intersecting in the new singular locus (isomorphic to \mathbb{P}^4). Blowing up in this intersection gives two new exceptional components, but one of them (isomorphic to a \mathbb{P}^1 -bundle over \mathbb{P}^4) is singular for the strict transform of Y . The other one, called E , is isomorphic to a \mathbb{P}^2 -bundle over $Y_8^{(3)}$. Its intersection with the D_i is contained in the new singular locus. Blowing up in this singular locus creates six new exceptional components. Five of them behave similarly. They are isomorphic to a \mathbb{P}^1 -bundle over \mathbb{P}^4 , they are disjoint and each of them has an intersection with one D_i isomorphic to \mathbb{P}^4 (they are now the only components that intersect the D_i ; for simplicity, we use the same name for an exceptional component throughout the resolution process, instead of speaking of strict transforms). We call these components F_1, \dots, F_5 and choose the numbering compatible with those of the D_i . The sixth exceptional component is called G and is also

isomorphic to a \mathbb{P}^1 -bundle over \mathbb{P}^4 . The new singular locus is $E \cap G$ and it is isomorphic to a \mathbb{P}^1 -bundle over $Y_8^{(3)}$. The intersection of an F_i and G is isomorphic to \mathbb{P}^4 and has a $Y_8^{(3)}$ in common with the singular locus (this $Y_8^{(3)}$ is the intersection of F_i with E). In the final step we blow up in the remaining singular locus. There is one new exceptional component, called H . Its Hodge–Deligne polynomial equals $((uv)^2 + 7uv + 1)H(Y_8^{(3)}; u, v)$ and it splits off E from the other components. The intersections of H with the F_i , with G and with E are all isomorphic to a \mathbb{P}^1 -bundle over $Y_8^{(3)}$. This final blow up also adds $uvH(Y_8^{(3)}; u, v)$ to the Hodge–Deligne polynomial of the F_i . And there are threefold intersections $F_i \cap G \cap H$ isomorphic to $Y_8^{(3)}$. The result is a normal crossing divisor with the following intersection diagram:



The discrepancy coefficients are 1 for the D_i and 0 for all the other components. Thus one can compute that

$$B := 8(uv)^5 + 15(uv)^4 + 10(uv)^3 + 15(uv)^2 + 8uv + 1 - 70u^5v^2 - 70u^2v^5 - 2030u^4v^3 - 2030u^3v^4 - 210u^4v - 210uv^4 - 6090u^3v^2 - 6090u^2v^3$$

is the contribution of the singular point to the stringy E -function. Then the stringy E -function of Y is just $A + B$ which equals

$$E_{st}(Y; u, v) = (uv)^6 + 26(uv)^5 + 35(uv)^4 + 30(uv)^3 + 35(uv)^2 + 26uv + 1 - 280u^5v^2 - 280u^2v^5 - 8120u^4v^3 - 8120u^3v^4 - 280u^4v - 280uv^4 - 8120u^3v^2 - 8120u^2v^3,$$

and thus $h_{st}^{2,2}(Y) > h_{st}^{3,3}(Y)$.

Acknowledgments

I want to thank Wim Veys for helpful discussions and for encouraging me to communicate these results. I also want to thank the referee of an earlier paper for bringing references [Fe] and [MP] to my attention.

References

[Ba] V.V. Batyrev, Stringy Hodge numbers of varieties with Gorenstein canonical singularities, in: Integrable Systems and Algebraic Geometry, Kobe/Kyoto 1997, World Sci. Publ., 1999, pp. 1–32.
 [BD] V.V. Batyrev, D. Dais, Strong McKay correspondence, string-theoretic Hodge numbers and mirror symmetry, Topology 35 (1996) 901–929.
 [BG] W. Bruns, J. Gubeladze, Polytopes, Rings and K -theory, Springer-Verlag, in press.
 [BM] L.A. Borisov, A.R. Mavlyutov, String cohomology of Calabi–Yau hypersurfaces via mirror symmetry, Adv. Math. 180 (2003) 355–390.
 [CR] W. Chen, Y. Ruan, A new cohomology theory of orbifold, Comm. Math. Phys. 248 (2004) 1–31.
 [Da] D. Dais, On the string-theoretic Euler number of a class of absolutely isolated singularities, Manuscripta Math. 105 (2001) 143–174.
 [dCM1] M.A. de Cataldo, L. Migliorini, The Hodge theory of algebraic maps, Ann. Sci. École Norm. Sup. (4) 38 (2005) 693–750.

- [dCM2] M.A. de Cataldo, L. Migliorini, Intersection forms, topology of maps and motivic decomposition for resolutions of threefolds, in: Algebraic Cycles and Motives, vol. 1, in: London Math. Soc. Lecture Note Ser., vol. 343, Cambridge Univ. Press, 2007, pp. 102–137.
- [Fe] J. Fernandez, Hodge structures for orbifold cohomology, Proc. Amer. Math. Soc. 134 (2006) 2511–2520.
- [Ma] K. Matsuki, Introduction to the Mori Program, Universitext, Springer-Verlag, 2002.
- [MP] M. Mustață, S. Payne, Ehrhart polynomials and stringy Betti numbers, Math. Ann. 333 (2005) 787–795.
- [Pa] S. Payne, Ehrhart series and lattice triangulations, Discrete Comput. Geom. 40 (2008) 365–376.
- [Po] M. Poddar, Orbifold Hodge numbers of Calabi–Yau hypersurfaces, Pacific J. Math. 208 (2003) 151–167.
- [Re] M. Reid, Minimal models of canonical threefolds, in: Algebraic Varieties and Analytic Varieties, Tokyo, 1981, in: Adv. Stud. Pure Math., vol. 1, North-Holland, 1983, pp. 131–180.
- [SV1] J. Schepers, W. Veys, Stringy Hodge numbers for a class of isolated singularities and for threefolds, Int. Math. Res. Not. 2007 (2007), article ID rnm016, 14 pages.
- [SV2] J. Schepers, W. Veys, Stringy E -functions of hypersurfaces and of Brieskorn singularities, Adv. Geom., in press.
- [Ya] T. Yasuda, Twisted jets, motivic measures and orbifold cohomology, Compos. Math. 240 (2004) 396–422.