

Contents lists available at ScienceDirect

Physics Letters B

www.elsevier.com/locate/physletbComplex structures and zero-curvature equations for σ -modelsDmitri Bykov^{a,b,*}^a Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut, Am Mühlenberg 1, D-14476 Potsdam-Golm, Germany^b Steklov Mathematical Institute of Russ. Acad. Sci., Gubkina str. 8, 119991 Moscow, Russia

ARTICLE INFO

Article history:

Received 9 May 2016

Received in revised form 26 June 2016

Accepted 28 June 2016

Available online 1 July 2016

Editor: M. Cvetič

ABSTRACT

We construct zero-curvature representations for the equations of motion of a class of σ -models with complex homogeneous target spaces, not necessarily symmetric. We show that in the symmetric case the proposed flat connection is gauge-equivalent to the conventional one.

© 2016 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. The models

A σ -model is a field theory describing maps $X: \Sigma \rightarrow \mathcal{M}$ from a worldsheet Σ to a target space \mathcal{M} . In this paper the worldsheet Σ will be a two-dimensional Euclidean manifold, and the target-space \mathcal{M} will be required to have the following properties¹:

- \mathcal{M} is a homogeneous space G/H , G semi-simple and compact
- \mathcal{M} has an integrable G -invariant complex structure \mathbb{I} (1)
- The Killing metric \mathbb{G} on \mathcal{M} is Hermitian w.r.t. \mathbb{I}

We will show that, for a target space \mathcal{M} with these properties, one can define a σ -model, whose equations of motion may be rewritten as the flatness condition for a one-parameter family of connections A_u , $u \in \mathbb{C}^*$. This is an extension to a broader class of target spaces of a property that is encountered in σ -models with symmetric target spaces [1,2]. In the latter case it is an important sign of integrability of the model: it may be used to find Bäcklund transformations [3,4], and it is a starting point for the construction of classical solutions of the models [5].

Complex simply-connected homogeneous manifolds G/H with G semi-simple were classified long ago [6]. They are given by the following theorem: any such manifold G/H corresponds to a subgroup H , whose semi-simple part coincides with the semi-simple part of the centralizer of a toric subgroup of G .

For the case of $G = SU(N)$, for example, invariant complex structures exist on those of the manifolds

$$\mathcal{M}_{n_1, \dots, n_m | N} = \frac{SU(N)}{S(U(n_1) \times \dots \times U(n_m))},$$

$$m \geq 0, \quad n_i > 0, \quad \sum_{i=1}^m n_i \leq N, \quad (2)$$

that are even-dimensional. If $\sum_{i=1}^m n_i = N$, the manifold in (2) is a flag manifold (for a review of flag manifolds see [7,8]). Otherwise, it is a toric bundle over a flag manifold. The fiber $U(1)^{2s}$ ($2s = N - \sum_{i=1}^m n_i$) is even-dimensional, since the flag manifold itself is even-dimensional.

If one relaxes the requirements of simple connectedness of G/H and semi-simplicity of G in the above theorem, there are more examples, such as $S^1 \times S^3 = U(2)$, which is even hyper-complex, i.e. it has a continuous family of invariant complex structures (such manifolds were considered and classified in [9] and [10]). In this case, in place of the Killing metric one can take the metric $\hat{\mathbb{G}}_{ij} := \text{tr}(T_i T_j)$, where T_i are the generators of \mathfrak{g} taken in a representation suitably chosen in order to make sure that $\hat{\mathbb{G}}$ is non-degenerate and Hermitian with respect to \mathbb{I} .

The models, which will be of interest for us in the present paper, are defined by the following action:

$$S[\mathbb{G}, \mathbb{I}] := \int_{\Sigma} d^2z \|\partial X\|_{\mathbb{G}}^2 + \int_{\Sigma} X^* \omega, \quad (3)$$

where ω is the Kähler form corresponding to the pair (\mathbb{G}, \mathbb{I}) , defined as

$$\omega = \mathbb{G} \circ \mathbb{I}. \quad (4)$$

In general, the Killing metric \mathbb{G} is not Kähler, i.e. the Kähler form is not closed: $d\omega \neq 0$. This might be the case, even if the manifold \mathcal{M} admits a Kähler metric – it is in general different from \mathbb{G} . As an example of such phenomenon one can consider the flag

* Correspondence to: Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut, Am Mühlenberg 1, D-14476 Potsdam-Golm, Germany.

E-mail addresses: dmitri.bykov@aei.mpg.de, dbykov@mi.ras.ru.

¹ Generalizations to non-semi-simple groups G are possible, but will be elaborated elsewhere.

manifold $\frac{SU(3)}{S(U(1)^3)}$. The σ -model (3) for the flag manifold was investigated in detail in [11,12]. Other examples of models of the class (1) are provided by Hermitian symmetric spaces – symmetric spaces with complex structure. These manifolds are Kähler, and the invariant metric is essentially unique (up to scale), thus leading to the closedness of ω : $d\omega = 0$. We will discuss this special case in Section 2.1.

2. Structure of the target space

The Lie algebra \mathfrak{g} of the Lie group G may be decomposed as follows: $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$, where \mathfrak{h} is the Lie algebra of H and \mathfrak{m} is the orthogonal complement to \mathfrak{h} with respect to the Killing metric \mathbb{G} . We will assume that the quotient space G/H possesses an almost complex structure \mathbb{I} . We are not postulating that \mathbb{I} be integrable – this will rather follow from the requirement of the existence of a Lax connection. The almost complex structure acts on \mathfrak{m} and may be diagonalized, its eigenvalues being $\pm i$. We denote the $\pm i$ -eigenspaces by \mathfrak{m}_{\pm} :

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}_+ \oplus \mathfrak{m}_-, \quad \mathbb{I} \circ \mathfrak{m}_{\pm} = \pm i \mathfrak{m}_{\pm}. \quad (5)$$

G -invariance of the almost complex structure implies that $[\mathfrak{h}, \mathfrak{m}_{\pm}] \subset \mathfrak{m}_{\pm}$. We introduce the current

$$J = g^{-1}dg = J_0 + J_+ + J_-, \quad J_0 \in \mathfrak{h}, \quad J_{\pm} \in \mathfrak{m}_{\pm}. \quad (6)$$

It takes values in the Lie algebra \mathfrak{g} , therefore we have decomposed it according to the decomposition (5) of the Lie algebra. In these terms the action (3) may be rewritten as follows:

$$\mathcal{S}[\mathbb{G}, \mathbb{I}] := \int_{\Sigma} d^2z \langle (J_+)_z, (J_-)_{\bar{z}} \rangle_{\mathbb{G}}. \quad (7)$$

Henceforth we will be using bracket notation for the scalar product of two elements $\alpha, \beta \in \mathfrak{g}$: $\langle \alpha, \beta \rangle_{\mathbb{G}} := \mathbb{G}(\alpha, \beta)$. The Noether current, constructed using the above action, will be denoted by K . It is derived by taking an infinitesimal (z, \bar{z}) -dependent variation $g \rightarrow e^{\epsilon(z, \bar{z})} \circ g$ in the above action, which leads to

$$\begin{aligned} \delta \mathcal{S}[\mathbb{G}, \mathbb{I}] &= \int_{\Sigma} d^2z \left[\left\langle (g^{-1} \partial_z \epsilon)_+, (J_-)_{\bar{z}} \right\rangle_{\mathbb{G}} + \left\langle (J_+)_z, (g^{-1} \partial_{\bar{z}} \epsilon)_- \right\rangle_{\mathbb{G}} \right] \\ &= \int_{\Sigma} d^2z \left[\left\langle g^{-1} \partial_z \epsilon, (J_-)_{\bar{z}} \right\rangle_{\mathbb{G}} + \left\langle (J_+)_z, g^{-1} \partial_{\bar{z}} \epsilon \right\rangle_{\mathbb{G}} \right] \quad (8) \end{aligned}$$

To pass to the second line we have used the following properties of the metric: $\langle \mathfrak{h}, \mathfrak{m}_{\pm} \rangle_{\mathbb{G}} = 0$, $\langle \mathfrak{m}_{\pm}, \mathfrak{m}_{\pm} \rangle_{\mathbb{G}} = 0$. The latter is a consequence of the Hermiticity of \mathbb{G} . Indeed, by definition of Hermiticity, $\langle \mathbb{I} \circ u, \mathbb{I} \circ v \rangle_{\mathbb{G}} = \langle u, v \rangle_{\mathbb{G}}$ for $u, v \in \mathfrak{m}$. Then, $\langle \mathfrak{m}_+, \mathfrak{m}_+ \rangle_{\mathbb{G}} = 0$, since $\mathbb{I} \circ \mathfrak{m}_+ = i \mathfrak{m}_+$. In other words, \mathfrak{m}_+ (and \mathfrak{m}_-) is an isotropic subspace of \mathfrak{m} . Using the invariance of the Killing metric \mathbb{G} under the adjoint action of G , $\langle gag^{-1}, gbg^{-1} \rangle_{\mathbb{G}} = \langle a, b \rangle_{\mathbb{G}}$, we obtain

$$\delta \mathcal{S}[\mathbb{G}, \mathbb{I}] = \int_{\Sigma} d^2z \left[\left\langle \partial_z \epsilon, g(J_-)_{\bar{z}} g^{-1} \right\rangle_{\mathbb{G}} + \left\langle \partial_{\bar{z}} \epsilon, g(J_+)_z g^{-1} \right\rangle_{\mathbb{G}} \right]. \quad (9)$$

Using the non-degeneracy of \mathbb{G} , we deduce the conservation (on the e.o.m.) of the Noether current, defined as follows:

$$K = g \cdot \underbrace{2((J_+)_z dz + (J_-)_{\bar{z}} d\bar{z})}_{:=S} \cdot g^{-1} = g S g^{-1} \quad (10)$$

Since the target space $\mathcal{M} = G/H$ is homogeneous, the equations of motion of the model are equivalent to its conservation:

$$d * K = 0. \quad (11)$$

In order to be able to build a family of flat connections we require that K be flat (this will be used in (20)–(21) below):

$$dK - K \wedge K = 0. \quad (12)$$

We have to show, of course, that it is possible to satisfy this relation. Equations (11)–(12) may be rewritten in terms of S (introduced in (10)) as follows:

$$d * S + \{J, * S\} = 0 \quad (13)$$

$$dS + \{J - \frac{1}{2}S, S\} = 0 \quad (14)$$

One checks directly that $J - \frac{1}{2}S = (J_-)_z dz + (J_+)_{\bar{z}} d\bar{z} + J_0$. Therefore

$$\begin{aligned} d * S + \{J, * S\} &= -2idz \wedge d\bar{z} \left(\mathcal{D}_{\bar{z}}(J_+)_{\bar{z}} - [(J_+)_{\bar{z}}, (J_+)_{\bar{z}}] + \mathcal{D}_z(J_-)_{\bar{z}} \right. \\ &\quad \left. + [(J_-)_z, (J_-)_{\bar{z}}] \right) \quad (15) \end{aligned}$$

$$\begin{aligned} dS + \{J - \frac{1}{2}S, S\} &= -2dz \wedge d\bar{z} \left(\mathcal{D}_z(J_+)_{\bar{z}} - [(J_+)_{\bar{z}}, (J_+)_{\bar{z}}] - \mathcal{D}_{\bar{z}}(J_-)_{\bar{z}} \right. \\ &\quad \left. - [(J_-)_z, (J_-)_{\bar{z}}] \right) \quad (16) \end{aligned}$$

\mathcal{D} is the covariant derivative for the gauge group H : $\mathcal{D}_j M_k := \partial_j M_k + [(J_0)_j, M_k]$ ($j, k = z, \bar{z}$). We have used the definition of the Hodge star $*dz = idz$, $*d\bar{z} = -id\bar{z}$.

The conditions (15)–(16) are equivalent, if

$$[\mathfrak{m}_+, \mathfrak{m}_+] \subset \mathfrak{m}_+, \quad [\mathfrak{m}_-, \mathfrak{m}_-] \subset \mathfrak{m}_- \quad (17)$$

(The commutators $[(J_+)_{\bar{z}}, (J_+)_{\bar{z}}]$ and $[(J_-)_z, (J_-)_z]$ then lie in the subspaces \mathfrak{m}_+ , \mathfrak{m}_- respectively.) In this case the equations take the form

$$\mathcal{D}_{\bar{z}}(J_+)_{\bar{z}} - [(J_+)_{\bar{z}}, (J_+)_{\bar{z}}] = 0 \quad (18)$$

$$\mathcal{D}_z(J_-)_{\bar{z}} + [(J_-)_z, (J_-)_{\bar{z}}] = 0 \quad (19)$$

If the metric \mathbb{G} is Hermitian w.r.t. the chosen almost complex structure \mathbb{I} , the conditions (17) are equivalent to the integrability of \mathbb{I} . Indeed, integrability of the almost complex structure means, in general, that $[\mathfrak{m}_+, \mathfrak{m}_+] \subset \mathfrak{m}_+ \oplus \mathfrak{h}$. To see this, note that the almost complex structure \mathbb{I} has been defined by $\mathbb{I} \circ J_{\pm} = \pm i J_{\pm}$. Integrability of \mathbb{I} is equivalent to the statement that (anti)-holomorphic forms generate a differential ideal, i.e. that $d(J_-)_a \sim \sum_b R_{ab} \wedge (J_-)_b$

for some one-forms R_{ab} . Since $dJ + J \wedge J = 0$, we have

$$dJ_- = \left[-J_0 \wedge J_0 + (\text{terms involving } J_-) - J_+ \wedge J_+ \right]_{\mathfrak{m}_-}.$$

Therefore integrability of \mathbb{I} requires that $[J_+ \wedge J_+]_{\mathfrak{m}_-} = 0$, i.e. $[\mathfrak{m}_+, \mathfrak{m}_+] \subset \mathfrak{m}_+ \oplus \mathfrak{h}$. We see that the conditions (17) define an integrable complex structure. Conversely, suppose we have an integrable complex structure on G/H , and \mathfrak{m}_{\pm} are its holomorphic and anti-holomorphic subspaces. Then, $[a, b] = c + \gamma$, where $a, b, c \in \mathfrak{m}_+$ and $\gamma \in \mathfrak{h}$. Since $\langle \mathfrak{m}_+, \mathfrak{h} \rangle_{\mathbb{G}} = 0$, taking the scalar product with an arbitrary element $\gamma' \in \mathfrak{h}$, one obtains $\langle \gamma', [a, b] \rangle_{\mathbb{G}} = \langle \gamma', \gamma \rangle_{\mathbb{G}}$. Using $\langle [a, \gamma'], b \rangle_{\mathbb{G}} + \langle \gamma', [a, b] \rangle_{\mathbb{G}} = 0$, we obtain $\langle \gamma', \gamma \rangle_{\mathbb{G}} = -\langle [a, \gamma'], b \rangle_{\mathbb{G}} = \langle a', b \rangle_{\mathbb{G}}$, where $a' = [\gamma', a] \in \mathfrak{m}_+$. As discussed earlier, \mathfrak{m}_+ is isotropic if the metric \mathbb{G} is Hermitian, therefore $\langle \gamma', \gamma \rangle_{\mathbb{G}} = 0$ for all $\gamma' \in \mathfrak{h}$, leading to $\gamma = 0$ due to non-degeneracy of \mathbb{G} . We conclude that $[\mathfrak{m}_+, \mathfrak{m}_+] \subset \mathfrak{m}_+$.

Consider now the following family of connections A_u , indexed by a parameter $u \in \mathbb{C}^*$:

$$A_u = \frac{1-u}{2} K_z dz + \frac{1-u^{-1}}{2} K_{\bar{z}} d\bar{z}. \quad (20)$$

Conservation and flatness of the Noether current K , eqs. (11)–(12), imply that A_u is flat for all u :

$$dA_u - A_u \wedge A_u = 0 \quad \text{for all } u \in \mathbb{C}^*. \quad (21)$$

2.1. Symmetric spaces

As discussed in Section 1, Hermitian symmetric spaces satisfy the requirements (1) and hence fall in our class. This case is characterized by the relation $[m_{\pm}, m_{\pm}] = 0$. Indeed, this follows from the symmetric space property $[m, m] \subset \mathfrak{h}$ and the existence of integrable complex structure (17). Conversely, if $[m_{\pm}, m_{\pm}] = 0$, one shows, using ad -invariance of the Killing metric on \mathfrak{g} , that $[m_+, m_-]$ is orthogonal to m_{\pm} , and hence $[m_+, m_-] \subset \mathfrak{h}$. Sigma-models with symmetric target spaces are known to be integrable, however the canonical Lax connection in this case is apparently different from the one in (20). Indeed, the connection usually employed in the analysis of σ -models with symmetric target spaces has the form

$$\tilde{A}_{\lambda} = \frac{1-\lambda}{2} \tilde{K}_z dz + \frac{1-\lambda^{-1}}{2} \tilde{K}_{\bar{z}} d\bar{z}, \quad (22)$$

where \tilde{K} is the Noether current derived using the canonical action

$$S[\mathbb{G}] = \int_{\Sigma} d^2z \|\partial X\|_{\mathbb{G}}^2. \quad (23)$$

In the case of a Hermitian symmetric target space the difference between the two actions, (3) and (23), is a topological term:

$$S[\mathbb{G}, \mathbb{I}] - S[\mathbb{G}] = \int_{\Sigma} X^* \omega, \quad (24)$$

where $d\omega = 0$ if \mathcal{M} is symmetric.

Therefore the two actions lead to the same equations of motion. Nevertheless, the Noether currents K and \tilde{K} are different. The canonical Noether current \tilde{K} is defined as follows:

$$\tilde{K} = 2g \cdot [g^{-1} dg]_{\mathfrak{m}} \cdot g^{-1} \quad (25)$$

The notation $[\dots]_{\mathfrak{m}}$ means that one should take the projection on the linear subspace $\mathfrak{m} \subset \mathfrak{g}$. This is certainly different from (10), since it follows from (25) that \tilde{K} may be written as (taking into account that $\mathfrak{m} = \mathfrak{m}_+ \oplus \mathfrak{m}_-$)

$$\begin{aligned} \tilde{K} &= 2g \cdot (J_+ + J_-) \cdot g^{-1} \\ &= 2g \cdot (((J_+)_z + (J_-)_z) dz + ((J_+)_{\bar{z}} + (J_-)_{\bar{z}}) d\bar{z}) \cdot g^{-1} \end{aligned} \quad (26)$$

On the other hand, \tilde{K} is also flat: $d\tilde{K} - \tilde{K} \wedge \tilde{K} = 0$. Checking this property does not, in fact, require using the equations of motion – it is purely a consequence of the structure of the Lie algebra of the symmetric space (in particular, the fact that $[m, m] \subset \mathfrak{h}$). Moreover, the flatness condition may be solved, in this case, in a local fashion²:

$$\tilde{K} = -\widehat{g}^{-1} d\widehat{g}, \quad \text{where } \widehat{g} = \sigma(g)g^{-1}, \quad (27)$$

σ being Cartan’s involution on the Lie group G . By definition, σ is a group homomorphism, $\sigma(g_1 g_2) = \sigma(g_1)\sigma(g_2)$, and $\sigma(h) = h$ for

$h \in H$. The formula $\widehat{g} = \sigma(g)g^{-1}$, viewed as a map $g \in G/H \rightarrow \widehat{g} \in G$, describes the Cartan embedding

$$G/H \hookrightarrow G. \quad (28)$$

Flatness and conservation of the current \tilde{K} lead to the flatness of the family \tilde{A}_{λ} . A question naturally arises of what the relation between A_u and \tilde{A}_{λ} is. The answer is that the connections A_u and \tilde{A}_{λ} are gauge-equivalent, if one makes the following identification of spectral parameters:

$$\lambda = u^{1/2}. \quad (29)$$

It follows that, whenever $|u| = 1$, one has $|\lambda| = 1$, therefore if one connection is unitary, the other is as well. Relation (29) is also consistent with the analysis in [5] of the limiting behavior of the holonomies of the connection around the cycles of Σ as $u \rightarrow 0$ (or $u \rightarrow \infty$). The gauge transformation Ω relating A_u and \tilde{A}_{λ} ,

$$\tilde{A}_{\lambda} = \Omega A_u \Omega^{-1} - \Omega d\Omega^{-1}, \quad (30)$$

may be constructed explicitly. The following formula holds for the case when the target-space is the Grassmannian $G_{n_1, n_1+n_2} := \frac{SU(n_1+n_2)}{S(U(n_1) \times U(n_2))}$ and the complex structure is chosen so that it splits \mathfrak{m} as $\mathfrak{m} = \begin{pmatrix} 0_{n_1 \times n_1} & \mathfrak{m}_+ \\ \mathfrak{m}_- & 0_{n_2 \times n_2} \end{pmatrix}$:

$$\begin{aligned} \Omega &= g \Lambda g^{-1}, \\ \text{where } \Lambda &= \lambda^{\frac{1}{2} \frac{n_1-n_2}{n_1+n_2}} \text{diag}(\underbrace{\lambda^{-1/2}, \dots, \lambda^{-1/2}}_{n_1}, \underbrace{\lambda^{1/2}, \dots, \lambda^{1/2}}_{n_2}) \end{aligned} \quad (31)$$

In terms of the ‘dynamical’ projectors Π_{n_1}, Π_{n_2} on the subspaces of $\mathbb{C}^{n_1+n_2}$ of the corresponding dimensions, Ω can also be written as

$$\Omega = \lambda^{\frac{1}{2} \frac{n_1-n_2}{n_1+n_2}} \left(\lambda^{-1/2} \Pi_{n_1} + \lambda^{1/2} \Pi_{n_2} \right) \quad (32)$$

In particular, at the point $\lambda = -1$ we obtain

$$\Omega|_{\lambda=-1} = e^{-i\pi \frac{n_2}{n_1+n_2}} (\Pi_{n_1} - \Pi_{n_2}), \quad (33)$$

which is the explicit representation of Cartan’s embedding for the case

$$G_{n_1, n_1+n_2} \hookrightarrow SU(n_1 + n_2). \quad (34)$$

At this point $u = 1$. Since $A_{u=1} \equiv 0$, one has $\tilde{A}_{\lambda=-1} = \tilde{K} = -\Omega d\Omega^{-1}$, which coincides with representation (27), provided $\Omega = \widehat{g}^{-1}$.

3. Discussion

In this letter we showed that the σ -models with complex homogeneous target spaces, defined by action (3), have the following property: their equations of motion may be rewritten as the flatness condition for a one-parameter family of connections. This representation is ubiquitously encountered in the realm of integrable models in two dimensions, and is often believed to be the characteristic, cornerstone property of these models [3–5]. As we saw in Section 2.1, in the case of symmetric target spaces the proposed flat connection is gauge-equivalent to the conventional one, upon redefinition of the spectral parameter (29).

In recent years there has been renewed interest in the classical solutions of σ -models, due to advances in the understanding of their relation to the divergences of the perturbative expansion in quantum field theory [13]. When the target-space is the flag manifold $\mathcal{M} = \frac{SU(3)}{S(U(1)^3)}$ and the worldsheet is a sphere, $\Sigma = \mathbb{C}P^1$,

² ‘Local’ means here that \widehat{g} is a local function of the fields of the model.

earlier [12] we have been able to construct all solutions to the e.o.m. of the model (3). The zero-curvature representation introduced in this letter might allow to construct explicit solutions of the model (3), in general.

Interestingly, the action (3) coincides in form with the bosonic part of the action introduced in [14]. In that paper the main focus was on a particular nilpotent supersymmetry of the full action, generated by a supercharge Q , $Q^2 = 0$. It was shown that the correlation functions of Q -closed observables in those models could be computed. They are related to the moduli spaces of \mathbb{I} -holomorphic curves in the target space. (The holomorphic curves are absolute minima of the action (3); the action vanishes on them.) Moreover, the correlation functions of Q -closed observables are ‘topological’, meaning that they do not change under smooth deformations of the metric \mathbb{G} and of the complex structure \mathbb{I} . On the other hand, the zero-curvature representation, which we have constructed in the present paper, might allow to find other solutions of these models, not just the holomorphic curves. This zero-curvature representation only holds, however, if we choose the Killing metric and a compatible complex structure on the target space \mathcal{M} of the σ -model.

If the models described in the present letter can be consistently quantized, an interesting question is whether the integrable properties persist in the quantized version. It is known that for some σ -models this is not the case. One such example is the $\mathbb{C}\mathbb{P}^N$ model with $N \geq 2$ [15,16]. On the other hand, in the supersymmetric $\mathbb{C}\mathbb{P}^N$ model this problem is resolved [17]. Therefore it is important to determine, what type of supersymmetry the proposed models allow for. Conditions for (2, 2) supersymmetry in the presence of a B -field were found in [18]. It was shown in [19] that, when the target space is a flag manifold, these conditions are not satisfied, unless the B -field is a closed two-form. We leave a full clarification of these issues for future work.

Acknowledgements

I would like to thank C. Devchand and V. Pestun for discussions and the referee of Physics Letters B for useful comments. I am indebted to Prof. A.A. Slavnov and to my parents for support and

encouragement. My work was supported in part by the grant RFBR 14-01-00695-a.

References

- [1] K. Pohlmeyer, Integrable hamiltonian systems and interactions through quadratic constraints, *Commun. Math. Phys.* 46 (3) (1976) 207–221.
- [2] H. Eichenherr, M. Forger, On the dual symmetry of the nonlinear sigma models, *Nucl. Phys. B* 155 (1979) 381.
- [3] K. Uhlenbeck, Harmonic maps into Lie groups: classical solutions of the chiral model, *J. Differ. Geom.* 30 (1) (1989) 1–50.
- [4] C. Devchand, J. Schiff, Hidden symmetries of the principal chiral model unveiled, *Commun. Math. Phys.* 190 (1998) 675–695.
- [5] N.J. Hitchin, Harmonic maps from a 2-torus to the 3-sphere, *J. Differ. Geom.* 31 (3) (1990) 627–710.
- [6] H.-C. Wang, Closed manifolds with homogeneous complex structure, *Am. J. Math.* 76 (1954) 1–32.
- [7] D.V. Alekseevsky, Flag manifolds, *Zb. Rad. (Beogr.)* 6 (14) (1997) 3–35, 11th Yugoslav Geometrical Seminar (Divčibare, 1996).
- [8] A. Arvanitoyeorgos, An Introduction to Lie Groups and the Geometry of Homogeneous Spaces, Student Mathematical Library, vol. 22, American Mathematical Society, Providence, RI, 2003.
- [9] P. Spindel, A. Sevrin, W. Troost, A. Van Proeyen, Extended supersymmetric sigma models on group manifolds. 1. The complex structures, *Nucl. Phys. B* 308 (1988) 662.
- [10] D. Joyce, Compact hypercomplex and quaternionic manifolds, *J. Differ. Geom.* 35 (3) (1992) 743–761.
- [11] D. Bykov, Integrable properties of sigma-models with non-symmetric target spaces, *Nucl. Phys. B* 894 (2015) 254–267.
- [12] D. Bykov, Classical solutions of a flag manifold σ -model, *Nucl. Phys. B* 902 (2016) 292–301.
- [13] G.V. Dunne, M. Unsal, Resurgence and trans-series in quantum field theory: the $\mathbb{C}\mathbb{P}^{N-1}$ model, *J. High Energy Phys.* 11 (2012) 170.
- [14] E. Witten, Topological sigma models, *Commun. Math. Phys.* 118 (1988) 411.
- [15] Y.Y. Goldschmidt, E. Witten, Conservation laws in some two-dimensional models, *Phys. Lett. B* 91 (1980) 392–396.
- [16] E. Abdalla, M. Forger, M. Gomes, On the origin of anomalies in the quantum nonlocal charge for the generalized nonlinear σ models, *Nucl. Phys. B* 210 (1982) 181–192.
- [17] M. Gomes, E. Abdalla, M.C.B. Abdalla, On the nonlocal charge of the $\mathbb{C}\mathbb{P}^{(N-1)}$ model and its supersymmetric extension to all orders, *Phys. Rev. D* 27 (1983) 825.
- [18] S.J. Gates Jr., C.M. Hull, M. Rocek, Twisted multiplets and new supersymmetric nonlinear sigma models, *Nucl. Phys. B* 248 (1984) 157–186.
- [19] D.V. Alekseevsky, L. David, A note about invariant SKT structures and generalized Kähler structures on flag manifolds, *Proc. Edinb. Math. Soc. (2)* 55 (3) (2012) 543–549.