

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.ScienceDirect.com)

## Physics Letters B

[www.elsevier.com/locate/physletb](http://www.elsevier.com/locate/physletb)

## Phenomenological high precision neutron–proton delta-shell potential



R. Navarro-Pérez, J.E. Amaro\*, E. Ruiz-Arriola

Departamento de Física Atómica, Molecular y Nuclear and Instituto Carlos I de Física Teórica y Computacional Universidad de Granada, E-18071 Granada, Spain

## ARTICLE INFO

## Article history:

Received 15 March 2013

Received in revised form 27 May 2013

Accepted 29 May 2013

Available online 1 June 2013

Editor: W. Haxton

## Keywords:

Potential scattering

Pn interaction

One pion exchange

## ABSTRACT

We provide a successful fit for proton–neutron scattering below pion production threshold up to LAB energies of 350 MeV. We use seven high-quality fits based on potentials with different forms as a measure of the systematic uncertainty. We represent the interaction as a sum of delta shells in configuration space below the 3 fm and a charge-dependent one-pion exchange potential above 3 fm together with magnetic effects. Special attention is paid to estimate the errors of the phenomenological interaction.

© 2013 Elsevier B.V. Open access under [CC BY license](http://creativecommons.org/licenses/by/3.0/).

The study of the NN interaction has been a central and recurrent topic in Nuclear Physics for many years (see e.g. [1,2] and references therein). The standard approach to constrain the interaction is to analyze NN scattering data below pion production threshold and to undertake a partial wave analysis (PWA), the quality of the fit being given by the  $\chi^2/\text{d.o.f.}$  value. Only by the mid 90s was it possible to fit about 4000 selected NN scattering data after discarding about further 1000 of  $3\sigma$  inconsistent data with a  $\chi^2/\text{d.o.f.} \lesssim 1$  and incorporating charge dependence (CD) for the One Pion Exchange (OPE) potential as well as magnetic and vacuum polarization effects [3]. This benchmark partial wave analysis (PWA) was carried out using an energy-dependent potential for the short range part for which nuclear structure calculations become hard to formulate. Thus, energy-independent high-quality potentials were subsequently produced with almost identical  $\chi^2/\text{d.o.f.} \sim 1$  for a gradually increasing database [4–7]. While any of these potentials provides individually satisfactory fits to the available data there are no published error estimates of the potential parameters. Moreover it should also be noticed that the existing high-quality potentials are different in their specific form; they range from local to non-local in different versions of non-locality. Thus, scattering phase shifts and observable amplitudes are not identical and in fact the existing set of high-quality potentials as a whole provides a distribution of scattering observables accounting for systematic uncertainties in addition to the statistical uncertainties obtained from the fitted data for each in-

dividual potential. Given the fact that these interactions are just constrained to the elastic scattering data (and eventually to the deuteron) which go up to the pion production threshold, one is physically probing the interaction with a resolution not finer than the shortest de Broglie wavelength  $\Delta\lambda = \hbar/\sqrt{M_N m_\pi} \sim 0.5$  fm. Thus, for practical purposes it may be advantageous to consider coarse grained interactions. This is actually the physics underlying the so-called  $V_{\text{low}k}$  approach [8] in which an effective interaction in a restricted model space is built. By starting from different high-quality potentials with a common charge-dependent OPE interaction the CM-momenta above  $\Lambda \sim \sqrt{M_N m_\pi}$  are eliminated by a suitable transformation and a remarkable universal interaction is obtained for  $p \leq \Lambda$ . Many of the applications of such an appealing interaction have recently been reviewed [9].

On the other hand, when switching from the NN problem to the many body nuclear problem the features and the form of the interaction are relevant in terms of computational cost and feasibility (see e.g. [10] and references therein). The lack of knowledge of a precise potential form with finer resolution than  $\Delta r \sim 0.5$  fm suggests to search for a description of scattering data directly in terms of a coarse grained potential sampled at some sensible “thick points”. Any sampling procedure necessarily redistributes the interaction strength and smooths the potential as compared to the zero resolution limit  $\Delta r \rightarrow 0$  implicit in most potential approaches and generating the troublesome short distance cores. This requires short distance correlations in the wave function to ensure the finiteness of the energy [10]. A desirable way to sample the interaction is to provide an acceptable  $\chi^2$ -fit with the minimal number of sampling points [11]; by implementing this minimal sampling we just try to avoid statistical dependence between the strengths of the potential at the chosen sampling points.

\* Corresponding author.

E-mail addresses: [rnavarrop@ugr.es](mailto:rnavarrop@ugr.es) (R. Navarro-Pérez), [amaro@ugr.es](mailto:amaro@ugr.es) (J.E. Amaro), [earriola@ugr.es](mailto:earriola@ugr.es) (E. Ruiz-Arriola).

Our motivation to proceed in this fashion is to make error propagation in nuclear structure calculations more direct since in the absence of statistical correlations errors may just be added in quadrature [12].

In the present work we report a first-stage form of a high-quality potential accommodating these desirable features, namely complying with existing high-quality fits and at the same time accounting for the systematic uncertainties. Our “coarse grained” potential is simple enough to undertake an analysis of the errors in its parameters as well as in the resulting phase shifts from a global fit to the experimental data. Our research is conveniently done in two stages. First we compile the phase shifts provided by the partial wave analysis (PWA) and by the high-quality potentials of Refs. [3–7] for a discrete set of energies  $E_i$ . By averaging those phase shifts for each energy and partial wave, we generate a set of “pseudodata” and “pseudo-errors” that we will use to perform a fit of our potential parameters for each partial wave, with their uncertainties. Since the pseudodata have been generated from high-quality potentials, our first-stage model, after fitting, will be a faithful representation of the experimental knowledge. A second stage of our research [14] will be to use the first-stage parameters as a starting point for a global partial wave analysis to pp and np scattering data below pion production threshold up to LAB energies of 350 MeV.

In this Letter we present the results of the first-stage fit which is interesting in its own right, since the existing systematic errors relevant for Nuclear Physics can already be extracted. Actually, the second-stage fit is currently being done and will be presented in a forthcoming publication [14] where the specific statistical errors will be considered. Note that the first-stage parameters are not checked against the experimental database, but to the pseudodata. Therefore calling it already “high-quality” could seem not justified a priori. At this point we should anticipate that, as presented in Refs. [13,14], a  $\chi^2/\text{d.o.f.}$  value can be computed with the experimental data and the first-stage optimal parameters. The optimal value turns out to be very close to one ( $\chi^2/\text{d.o.f.} = 1.12$ ), when about 4-digits are taken into account, although errors deduced directly from experimental data turn out to be much smaller than the systematic errors addressed here. The reason is that here we are fitting pseudodata constructed from a family of high-quality potentials, and the resulting potential inherits some of its characteristics. This fact makes the first-stage fit to pseudodata highly interesting and useful despite its simplicity. Our procedure allows to undertake an analysis of the systematic errors in the first-stage parameters. We do this by assuming that the systematic error inherent to any specific choice of the potential form corresponds to individual uncorrelated pseudodata.<sup>1</sup> Hence we may invoke the central limit theorem to undertake the traditional statistical treatment to the mean average and the corresponding standard deviation of Refs. [3–7] without any further ado. We will use this compilation as our pseudo-database.

A convenient representation to sample the short distance component of the NN interaction was already suggested by Aviles [15] almost 40 years ago in terms of delta shells which for any partial wave  $^{2S}(l', l)_J$  we take as

$$V_{l,l'}^{JS}(r) = \frac{1}{2\mu_{np}} \sum_{n=1}^N (\lambda_n)_{l,l'}^{JS} \delta(r - r_n) \quad r \leq r_c \quad (1)$$

<sup>1</sup> We have actually checked that, within the corresponding statistical uncertainty, the absence of correlations among the different partial waves of the six high-quality potentials by a direct evaluation of the correlation coefficient (see Ref. [11] for a definition) holds for every single LAB energy below 350 MeV.

with  $\mu_{np}$  the reduced np mass and  $r_c = 3$  fm. In the spirit of Refs. [3–7] for  $r \geq r_c$  we use the well-known long-distance tail of the NN potential

$$V(\vec{r}) = V_{EM}(\vec{r}) + V_{OPE}(\vec{r}), \quad r > r_c, \quad (2)$$


where  $V_{EM}$  is the electromagnetic potential of Ref. [5], and  $V_{OPE}$  is the one-pion exchange potential.

The solution of the corresponding Schrödinger equation in coupled channels is straightforward; for any  $r_n < r < r_{n+1}$  with  $r_N < r_c$  we have free solutions and log-derivatives are discontinuous at the  $r_n$ -points so that one generates an accumulated S-matrix at any sampling point providing a discrete version of Calogero’s variable phase equation [16]. Although this potential is formally local, the fact that we are coarse graining the interaction enables to encode efficiently nonlocalities operating below the finest resolution  $\Delta r$ . Of course, once we admit that the interaction below  $r_c$  is unknown there is no advantage in prolonging the well-known charge-dependent OPE tail and other electromagnetic effects for  $r < r_c$ . The low energy expansion of the discrete variable phase equations was used in Ref. [17] to determine threshold parameters in all partial waves. The relation to the well-known Nyquist theorem of sampling a signal with a given bandwidth has been discussed in Ref. [18]. Some of the advantages of using this simple potential for nuclear structure calculations as well as the connection to the  $V_{\text{lowk}}$  approach have been spelled out already [19].

To generate the pseudodata we use the standard LAB energy values usually listed in the high-quality potentials, namely  $E_{\text{LAB}} = 1, 5, 10, 25, 50, 100, 150, 200, 250, 300, 350$  MeV and fit to the mean phase-shift values at those energies with an error equal to the standard deviation. In this way we account for the systematic errors due to the different representations of the potentials [3–7]. We find that they are generally larger than those quoted by the original PWA where only statistical uncertainties were explicitly discussed for a fixed potential form [3]. With these pseudodata sets and the given energies we undertake a phase-shift fit and determine errors using the standard covariance matrix.

As expected from Nyquist theorem, we need at most  $N = 5$  sampling points which for simplicity are taken to be equidistant with  $\Delta r = 0.6$  fm between the origin and  $r_c = 3$  fm. This is the minimal number which provides an acceptable fit to the pseudodata compiled from Refs. [3–7]. Our results for the np phase shifts for all partial waves with total angular momentum  $J \leq 5$  are depicted in Fig. 1. The fitting parameters  $(\lambda_n)_{l,l'}^{JS}$  entering the delta-shell potentials, Eq. (1), are listed in Table 1 with their deduced uncertainties. Of course, a definitive assessment on systematic errors would require testing all possible potential forms. Thus, the errors will generally be larger than those estimated here. We find that correlations among the different  $(\lambda_n)_{l,l'}^{JS}$  values within a given partial wave channel are unimportant, and hence these parameters are essentially independent from each other. This is a direct consequence of our strategy to minimize the number of sampling points. We find that introducing more points or equivalently reducing  $\Delta r$  generates unnecessary correlations. Also, lowering the value of  $r_c$  below 3 fm, requires overlapping the short-distance potential, Eq. (1), with the OPE plus em corrections. Note that in order to encompass all different forms of potentials, 48 parameters are needed in contrast to the about 40 independent parameters found in the original 1990 and subsequent models.

We determine the deuteron properties by solving the bound state problem in the  $^3S_1$ – $^3D_1$  channel using the corresponding parameters listed in Table 1. The predictions are presented in Table 2 where our quoted errors are obtained from propagating Table 1.




**Fig. 1.** np phase shifts in degrees for  $J \leq 5$  as a function of the LAB energy. The curves have been calculated with the fitted potential. The points with error bars represent the pseudodata obtained as the mean value and standard deviation of the phase shifts provided by the PWA [3] and the six high-quality potentials [4–7]. We fit the energies  $E_{\text{LAB}} = 1, 5, 10, 25, 50, 100, 150, 200, 250, 300, 350$  MeV.

**Table 1**  
Fitting delta-shell parameters  $(\lambda_n)_{l,J}^S$  (in  $\text{fm}^{-1}$ ) with their errors for all states in the  $JS$  channel and the corresponding  $\chi^2$ -value for  $J \leq 5$  in np scattering. We take  $N = 5$  equidistant points with  $\Delta r = 0.6$  fm. A dash symbol “–” indicates that the corresponding parameter has been fixed to  $(\lambda_n)_{l,J}^S = 0$ .

| Wave         | $\lambda_1$ | $\lambda_2$ | $\lambda_3$ | $\lambda_4$ | $\lambda_5$ | $\chi^2/\text{D.o.f.}$ |
|--------------|-------------|-------------|-------------|-------------|-------------|------------------------|
| $^1S_0$      | 2.12(7)     | -0.987(7)   | –           | -0.087(2)   | –           | 0.3476                 |
| $^3P_0$      | –           | 1.26(4)     | -0.43(1)    | –           | -0.037(2)   | 0.6589                 |
| $^1P_1$      | –           | 1.23(2)     | –           | 0.079(4)    | –           | 0.0088                 |
| $^3P_1$      | –           | 1.33(2)     | –           | 0.053(2)    | –           | 0.4323                 |
| $^1D_2$      | –           | –           | -0.252(3)   | –           | -0.0163(9)  | 0.6946                 |
| $^3D_2$      | –           | –           | -0.596(8)   | -0.08(1)    | -0.050(4)   | 0.6144                 |
| $^1F_3$      | –           | –           | 0.34(1)     | –           | 0.010(2)    | 0.3812                 |
| $^3F_3$      | –           | –           | –           | 0.060(2)    | –           | 0.4177                 |
| $^1G_4$      | –           | –           | -0.22(2)    | –           | -0.0137(9)  | 0.8090                 |
| $^3G_4$      | –           | –           | –           | -0.267(3)   | –           | 1.8670                 |
| $^1H_5$      | –           | –           | –           | 0.071(8)    | –           | 0.6577                 |
| $^3H_5$      | –           | –           | –           | 0.04(1)     | –           | 0.4193                 |
| $^3S_1$      | 1.57(4)     | -0.40(1)    | –           | -0.064(3)   | –           | –                      |
| $\epsilon_1$ | –           | -1.69(1)    | -0.379(4)   | -0.216(5)   | -0.027(3)   | –                      |
| $^3D_1$      | –           | –           | 0.52(2)     | –           | 0.041(3)    | 0.4313                 |
| $^3P_2$      | –           | -0.415(6)   | –           | -0.0384(9)  | –           | –                      |
| $\epsilon_2$ | –           | 0.65(1)     | –           | 0.106(2)    | –           | –                      |
| $^3F_2$      | –           | –           | 0.14(3)     | -0.076(6)   | –           | 0.3881                 |
| $^3D_3$      | –           | –           | –           | –           | –           | –                      |
| $\epsilon_3$ | –           | –           | -0.47(3)    | -0.24(1)    | -0.020(4)   | –                      |
| $^3G_3$      | –           | –           | –           | 0.101(6)    | –           | 0.6806                 |
| $^3F_4$      | –           | –           | -0.163(4)   | –           | -0.0101(4)  | –                      |
| $\epsilon_4$ | –           | –           | –           | 0.108(3)    | –           | –                      |
| $^3H_4$      | –           | –           | –           | –           | -0.010(1)   | 0.2659                 |
| $^3G_5$      | –           | –           | –           | 0.025(4)    | –           | –                      |
| $\epsilon_5$ | –           | –           | –           | -0.35(1)    | –           | –                      |
| $^3I_5$      | –           | –           | –           | –           | –           | 0.5354                 |

**Table 2**  
Deuteron static properties compared with empirical values and high-quality potentials calculations.

|  | Delta shell | Empirical [20–25] | Nijm I [4] | Nijm II [4] | Reid93 [4] | AV18 [5] | CD-Bonn [6] |
|--|-------------|-------------------|------------|-------------|------------|----------|-------------|
| $E_d$ (MeV)                                  | 2.2(2)      | 2.224575(9)       | Input      | Input       | Input      | Input    | Input       |
| $\eta$                                       | 0.025(2)    | 0.0256(5)         | 0.02534    | 0.02521     | 0.02514    | 0.0250   | 0.0256      |
| $A_S$ (fm <sup>1/2</sup> )                   | 0.88(3)     | 0.8781(44)        | 0.8841     | 0.8845      | 0.8853     | 0.8850   | 0.8846      |
| $r_m$ (fm)                                   | 1.97(8)     | 1.953(3)          | 1.9666     | 1.9675      | 1.9686     | 1.967    | 1.966       |
| $Q_D$ (fm <sup>2</sup> )                     | 0.272(9)    | 0.2859(3)         | 0.2719     | 0.2707      | 0.2703     | 0.270    | 0.270       |
| $P_D$  | 5.7(2)      | 5.67(4)           | 5.664      | 5.635       | 5.699      | 5.76     | 4.85        |
| $\langle r^{-1} \rangle$ (fm <sup>-1</sup> ) | 0.45(1)     |                   |            | 0.4502      | 0.4515     |          |             |




**Fig. 2.** Some np scattering observables for several energies in the laboratory system as a function of the CM angle. The short-dashed line denotes the predictions by our delta-shell model. The band represents the compilation the six high-quality potentials [3–7] which provided a  $\chi^2/\text{d.o.f.} \lesssim 1$ . For references to the experimental data see [29–31]. We show the following set of observables:  $I_0$  (differential cross section),  $P$  (polarization),  $D$  (depolarization),  $R$  (rotation parameter),  $A_t$ ,  $D_t$ , and  $R_t$  (polarization transfer parameters), and  $C_{nn}$  (spin correlation parameter). For notation and further explanations see Refs. [26,27].

The comparison with experimental values or high-quality potentials where the binding energies are used as an input is satisfactory. This is partly due to the fact that theoretical errors are about 10%. Of course, one may improve on this by using the deuteron binding energy as an input as in Refs. [3–7].

Fitting phase shifts to some accuracy do not necessarily provides angle-dependent scattering amplitudes to the same accuracy

because errors are finite and the relation between phase shifts and observables is non-linear. This is often the case when the form of the potential is kept fixed, so that the channel by channel fit is usually taken as a first step which is afterwards refined by a full fledged analysis of differential cross section and polarization data [3–7]. To check our strategy of fitting first pseudodata (phase shifts) and determining observables afterwards we proceed as fol-



**Fig. 3.** np Wolfenstein parameters for different energies in the laboratory system as a function of the CM angle. Upper panel:  $E_{\text{LAB}} = 50$  MeV. Lower panel:  $E_{\text{LAB}} = 200$  MeV. The band represents the compilation of the PWA and six high-quality potentials [3–7] which provided a  $\chi^2/\text{d.o.f.} \lesssim 1$ .

lows. The complete on-shell np scattering amplitude contains five independent complex quantities, which we choose for definiteness as the Wolfenstein parameters

$$\begin{aligned}
 M(\mathbf{k}_f, \mathbf{k}_i) = & a + m(\boldsymbol{\sigma}_1, \mathbf{n})(\boldsymbol{\sigma}_2, \mathbf{n}) \\
 & + (g - h)(\boldsymbol{\sigma}_1, \mathbf{m})(\boldsymbol{\sigma}_2, \mathbf{m}) \\
 & + (g + h)(\boldsymbol{\sigma}_1, \mathbf{l})(\boldsymbol{\sigma}_2, \mathbf{l}) \\
 & + c(\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2, \mathbf{n}),
 \end{aligned} \quad (3)$$

where  $a, m, g, h, c$  depend on energy and angle, and could in principle be determined directly from experiment as shown in Ref. [26,27] (see also [28] for an exact analytical inversion). We follow an alternative procedure by arranging the parameters as a 10-dimensional real vector  $(a_1, \dots, a_{10})$ , and defining a new set of pseudodata out of the high-quality analyzes, given by the corresponding mean value of the Wolfenstein parameters,  $\bar{a}_i(E_{\text{LAB}}, \theta)$ , with their corresponding standard deviations,  $\Delta a_i(E_{\text{LAB}}, \theta)$ . For any given LAB energy and scattering angle  $\theta$  we then compute

$$\chi^2(E_{\text{LAB}}, \theta) = \sum_{i=1}^{10} \left[ \frac{\bar{a}_i(E_{\text{LAB}}, \theta) - a_i(E_{\text{LAB}}, \theta)}{\Delta a_i(E_{\text{LAB}}, \theta)} \right]^2. \quad (4)$$

The total  $\chi^2$  value is obtained as an average over the chosen reference energies,  $E_{\text{LAB}} = 1, 5, 10, 25, 50, 100, 150, 200, 250, 300$ ,

350 MeV, and a dense sampling of  $\theta$ -values in one degree steps.<sup>2</sup> The result is

$$\chi^2/\text{d.o.f.} = 0.78,$$

which is equivalent to carry a complete  $\chi^2$ -fit to the mean average scattering amplitude. In Fig. 2 we illustrate the situation for a set of observables as a function of the CM angle and for representative energies. One sees that our model (1) describes the data, and (2) it differs from the six high-quality potentials [3–7] mainly when no data are available.

Similar results are obtained for the Wolfenstein parameters themselves [32] as can be seen by direct inspection of Fig. 3 for the sampling energies of  $E_{\text{LAB}} = 50$  MeV and  $E_{\text{LAB}} = 200$  MeV.

To summarize, we have determined a high-quality neutron–proton interaction which is based on a few delta shells for the lowest partial waves, in addition to charge-dependent electromagnetic interactions and one pion exchange, and provides a good starting point for Nuclear Physics applications. We provide error estimates on our fitting parameters accounting both for systematic and statistical uncertainties from pseudodata provided by present day high-quality analyzes of neutron–proton scattering data. Deuteron properties are compatible with experimental or recommended values. Our method allows to coarse grain long-range

<sup>2</sup> We have tried finer energy grids and the  $\chi^2/\text{d.o.f.}$  remains stable and below unity.

Coulomb interactions in a rather natural way and hence to discuss proton–proton scattering data.

### Acknowledgements

We want to thank D.R. Entem and M. Pavón Valderrama for a critical reading of the manuscript. This work is supported by Spanish DGI (grant FIS2011-24149) and Junta de Andalucía (grant FQM225). R.N.P. is supported by a Mexican CONACYT grant.

### References

- [1] R. Machleidt, *Adv. Nucl. Phys.* 19 (1989) 189.
- [2] R. Machleidt, D. Entem, *Phys. Rept.* 503 (2011) 1.
- [3] V.G.J. Stoks, R.A.M. Kompl, M.C.M. Rentmeester, J.J. de Swart, *Phys. Rev. C* 48 (1993) 792.
- [4] V.G.J. Stoks, R.A.M. Klomp, C.P.F. Terheggen, J.J. de Swart, *Phys. Rev. C* 49 (1994) 2950.
- [5] R.B. Wiringa, V.G.J. Stoks, R. Schiavilla, *Phys. Rev. C* 51 (1995) 38.
- [6] R. Machleidt, *Phys. Rev. C* 63 (2001) 024001.
- [7] F. Gross, A. Stadler, *Phys. Rev. C* 78 (2008) 014005.
- [8] S.K. Bogner, T.T.S. Kuo, A. Schwenk, *Phys. Rept.* 386 (2003) 1.
- [9] S.K. Bogner, R.J. Furnstahl, A. Schwenk, *Prog. Part. Nucl. Phys.* 65 (2010) 94.
- [10] S.C. Pieper, R.B. Wiringa, *Ann. Rev. Nucl. Part. Sci.* 51 (2001) 53.
- [11] J.R. Taylor, *An Introduction to Error Analysis: The Study of Uncertainties in Physical Measurements*, 2nd ed., University Science Books, 1997.
- [12] R. Navarro-Perez, J.E. Amaro, E. Ruiz-Arriola, arXiv:1202.6624 [nucl-th], 2012.
- [13] R. Navarro-Perez, J.E. Amaro, E. Ruiz Arriola, Few-body systems, <http://dx.doi.org/10.1007/s00601-012-0537-5>, arXiv:1209.6269, 2013.
- [14] R. Navarro-Perez, J.E. Amaro, E. Ruiz Arriola, arXiv:1304.0895 [nucl-th].
- [15] J.B. Aviles, *Phys. Rev. C* 6 (1972) 1467.
- [16] F. Calogero, *Variable Phase Approach to Potential Scattering*, Academic Press, New York, 1967.
- [17] M. Pavon-Valderrama, E. Ruiz-Arriola, *Phys. Rev. C* 72 (2005) 044007.
- [18] D.R. Entem, E. Ruiz-Arriola, M. Pavon-Valderrama, R. Machleidt, *Phys. Rev. C* 77 (2008) 044006.
- [19] R. Navarro-Perez, J.E. Amaro, E. Ruiz-Arriola, *Prog. Part. Nucl. Phys.* 67 (2012) 359.
- [20] C. Van Der Leun, C. Alderliesten, *Nucl. Phys. A* 380 (1982) 261.
- [21] I. Borbély, W. Grüebler, V. König, P.A. Schmelzbach, A.M. Mukhamedzhanov, *Phys. Lett. B* 106 (1985) 17.
- [22] N.L. Rodning, L.D. Knutson, *Phys. Rev. C* 41 (1990) 898.
- [23] S. Klarsfeld, J. Martorell, J.A. Oteo, M. Nishimura, D.W.L. Sprung, *Nucl. Phys. A* 456 (1986) 373.
- [24] D.M. Bishop, L.M. Cheung, *Phys. Rev. A* 20 (1979) 381.
- [25] J.J. de Swart, C.P.F. Terheggen, V.G.J. Stoks, arXiv:nucl-th/9509032, 1995.
- [26] J. Bystricky, F. Lehar, P. Winternitz, *J. Phys. (Paris)* 39 (1978) 1.
- [27] P. La France, P. Winternitz, *J. Phys. (Paris)* 41 (1980) 1391.
- [28] H. Kamada, W. Glöckle, H. Witala, J. Golak, R. Skibiński, *Few-Body Syst.* 50 (2011) 231.
- [29] <http://nn-online.org>.
- [30] <http://gwdac.phys.gwu.edu>.
- [31] <http://play.google.com/store/apps/details?id=es.ugr.amaro.nnprovider>.
- [32] R. Navarro-Perez, J.E. Amaro, E. Ruiz Ariola, in: *Proc. 7th International Workshop on Chiral Dynamics*, JLab, August 2012, arXiv:1301.6949.