

SIRT7 Represses Myc Activity to Suppress ER Stress and Prevent Fatty Liver Disease

Jiyung Shin,^{1,13} Ming He,^{1,2,13} Yufei Liu,^{3,13} Silvana Paredes,^{4,5,13} Lidia Villanova,^{4,5,6} Katharine Brown,¹ Xiaolei Qiu,¹ Noushin Nabavi,¹ Mary Mohrin,¹ Kathleen Wojnoonski,⁷ Patrick Li,⁸ Hwei-Ling Cheng,^{9,10} Andrew J. Murphy,¹¹ David M. Valenzuela,¹¹ Hanzhi Luo,¹ Pankaj Kapahi,⁸ Ronald Krauss,⁷ Raul Mostoslavsky,¹² George D. Yancopoulos,¹¹ Frederick W. Alt,^{9,10,*} Katrin F. Chua,^{4,5,*} and Danica Chen^{1,*}

¹Program in Metabolic Biology, Nutritional Sciences, & Toxicology, University of California, Berkeley, CA 94720, USA

²Key Laboratory of Cell Differentiation and Apoptosis of Ministry of Education, Department of Pathophysiology, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China

³Department of Molecular & Cell Biology, University of California, Berkeley, CA 94720, USA

⁴Division of Endocrinology, Gerontology, and Metabolism, Department of Medicine, Stanford University School of Medicine, Stanford, CA 94305, USA

⁵Geriatric Research, Education, and Clinical Center, VA Palo Alto Health Care System, Palo Alto, CA 94304, USA

⁶Department of Experimental Medicine, Sapienza University, Rome 00161, Italy

⁷Department of Atherosclerosis Research, Children's Hospital Oakland Research Institute, Oakland, CA 94609, USA

⁸Buck Institute for Research on Aging, 8001 Redwood Boulevard, Novato, CA 94945, USA

⁹Howard Hughes Medical Institute, The Children's Hospital, CBR Institute for Biomedical Research, Harvard University Medical School, Boston, MA 02115, USA

¹⁰Department of Genetics, Harvard Medical School, Boston, MA 02115, USA

¹¹Regeneron Pharmaceuticals, 777 Old Saw Mill River Road, Tarrytown, NY 10591, USA

¹²The Massachusetts General Hospital Cancer Center, Harvard Medical School, Boston, MA 02114, USA

¹³These authors contributed equally to this work

*Correspondence: alt@enders.tch.harvard.edu (F.W.A.), kfchua@stanford.edu (K.F.C.), danicac@berkeley.edu (D.C.)

<http://dx.doi.org/10.1016/j.celrep.2013.10.007>

This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-No Derivative Works License, which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

SUMMARY

Nonalcoholic fatty liver disease is the most common chronic liver disorder in developed countries. Its pathogenesis is poorly understood, and therapeutic options are limited. Here, we show that SIRT7, an NAD⁺-dependent H3K18Ac deacetylase, functions at chromatin to suppress ER stress and prevent the development of fatty liver disease. SIRT7 is induced upon ER stress and is stabilized at the promoters of ribosomal proteins through its interaction with the transcription factor Myc to silence gene expression and to relieve ER stress. SIRT7-deficient mice develop chronic hepatosteatosis resembling human fatty liver disease. Myc inactivation or pharmacological suppression of ER stress alleviates fatty liver caused by SIRT7 deficiency. Importantly, SIRT7 suppresses ER stress and reverts the fatty liver disease in diet-induced obese mice. Our study identifies SIRT7 as a cofactor of Myc for transcriptional repression and delineates a druggable regulatory branch of the ER stress response that prevents and reverts fatty liver disease.

INTRODUCTION

Nonalcoholic fatty liver disease (NAFLD) affects one-third of adults and an increasing number of children in developed countries and is strongly associated with obesity and insulin resistance (Browning and Horton, 2004; Browning et al., 2004; Cohen et al., 2011). NAFLD begins with aberrant accumulation of triglyceride in the liver (steatosis). Hepatic steatosis can proceed to nonalcoholic steatohepatitis (NASH), a condition associated with hepatocyte injury, inflammation, and fibrosis. Steatohepatitis can further progress to cirrhosis and liver cancer (Argo and Caldwell, 2009; Starley et al., 2010).

The endoplasmic reticulum (ER) stress response (also known as the unfolded protein response [UPR^{ER}]), a signal transduction pathway that is activated in response to the accumulation of unfolded proteins in the ER, has emerged as a critical regulator of lipid homeostasis in the liver (Basseri and Austin, 2012; Cnop et al., 2012; Fu et al., 2012; Hotamisligil, 2010; Ozcan and Tabas, 2012). The initial phase of the UPR^{ER} is suppression of protein translation and increased production of molecular chaperones to promote protein folding, allowing the cells to cope with an increased protein-folding demand and restore protein homeostasis (Hetz, 2012; Walter and Ron, 2011). Prolonged ER stress has been implicated in the development of numerous diseases, including fatty liver disease (Hotamisligil, 2010; Ozcan and Tabas, 2012). Identification of crucial UPR regulators with

precise ER-stress-relieving properties that are amenable for therapeutic targeting represents attractive opportunities for pharmacological intervention of fatty liver and a wide spectrum of human diseases.

The sirtuin family of nicotinamide adenine dinucleotide (NAD⁺)-dependent deacetylases is profoundly implicated in metabolic regulation (Bellet et al., 2011; Finkel et al., 2009; Gillum et al., 2010; Hirschey et al., 2011; Houtkooper et al. 2012; Imai and Guarente, 2010). Sirtuins are well-sought-after drug targets for metabolic disorders, as their enzymatic activities are amenable for regulation (Baur et al., 2012). SIRT7, a histone H3 lysine 18 (H3K18) deacetylase that binds to the promoters of a specific set of gene targets for transcriptional repression (Barber et al., 2012), is the only mammalian sirtuin whose function in metabolic regulation remains unknown.

We set out to fill this gap in knowledge by asking whether SIRT7 governs metabolic homeostasis under physiological conditions. Here, we show that SIRT7 has a physiological function in metabolic regulation that occurs through a chromatin-dependent signaling pathway that maintains metabolic homeostasis. Finally, we show that SIRT7 can be targeted to restore metabolic homeostasis in animals with metabolic disorders.

RESULTS

SIRT7-Deficient Mice Develop Steatosis Resembling Human Fatty Liver Disease

Among the metabolic tissues, SIRT7 is the most highly expressed in the liver (Ford et al., 2006). To gain insight into SIRT7 function, we generated SIRT7 knockout (KO) mice (Figures 1A and 1B; Figure S1A). Livers of SIRT7 KO mice fed a chow diet appeared paler and slightly larger than the wild-type (WT) controls with 100% penetrance (Figures 1C and S1B). Hematoxylin and eosin (H&E) staining showed that SIRT7 KO hepatocytes were markedly vacuolated, with the accumulated material staining positive for fat with the Oil Red O stain (Figure 1C). Quantification of the triglyceride extracted from the livers by a colorimetric assay showed that SIRT7 KO livers had a 2.5-fold increase in triglyceride content (Figure 1D). Compared to the WT controls, the livers of SIRT7 KO mice had increased expression of inflammatory markers (Figure 1E), and immunohistochemistry showed increased staining for F4/80 (Figure 1C), a marker for tissue macrophages, indicating the progression to steatohepatitis. Notably, although fatty liver disease is often associated with obesity (Cohen et al., 2011; Ozcan et al., 2006), SIRT7 KO mice were leaner than littermate controls (Figures S1C and S1D). Thus, SIRT7 deficiency results in fatty liver without obesity.

Next, we investigated the lipid metabolic pathways in SIRT7 KO livers. The expression of lipogenic genes was increased in SIRT7 KO livers compared to the WT controls, whereas the expression of genes in the fatty acid oxidation pathway was unchanged (Figure 1E), suggesting that increased lipogenesis may be a contributing factor for hepatosteatosis. Despite increased lipid content in the livers of SIRT7 KO mice (Figure 1D), the levels of plasma triglyceride were 4-fold lower in SIRT7 KO mice compared to WT controls (Figure S1E). Reduced plasma triglycerides in SIRT7 KO mice is not due to reduced food intake or malabsorption of lipid (Figures S1F and S1G), suggesting that

SIRT7 KO mice may have reduced very-low-density lipoprotein (VLDL) secretion, the lipoprotein responsible for hepatic lipid export. In a well-established VLDL-TG secretion assay, SIRT7 KO mice had a 50% reduction in VLDL-TG secretion (Figure 1F). Quantification of the VLDL particles (sizes ranging from 250–550 Å) using a gas-phase differential electrical mobility analyzer showed greatly reduced VLDL concentration in the blood of SIRT7 KO mice compared to the WT controls (Figure 1G). Together, these data indicate that SIRT7 KO mice develop fatty liver because of increased lipogenesis and reduced VLDL secretion.

Hepatic SIRT7 Autonomously Prevents the Development of Fatty Liver

To investigate whether the fatty liver phenotype of SIRT7 KO mice is due to SIRT7 deficiency in the liver or systemic effects of SIRT7 deletion, we reintroduced SIRT7 specifically in the livers of SIRT7 KO mice via adenoassociated virus 8 (AAV8)-mediated gene transfer (Figure 2A). Strikingly, liver-specific reconstitution of SIRT7 in SIRT7 KO mice reversed the fatty liver phenotype (Figures 2B–2D), suppressed hepatic inflammation and lipogenesis (Figure 2E), and rescued the VLDL-TG secretion defect (Figure 2F). Importantly, this reversal of hepatosteatosis was not due to unphysiological levels of SIRT7 overexpression, because the reconstituted expression of SIRT7 was comparable to endogenous levels (Figure 2A). Together, these data indicate that hepatic SIRT7 autonomously prevents the development of fatty liver.

SIRT7 Activation Is a Critical Event of the UPR to Alleviate ER Stress

Activation of the UPR pathways induces inflammation (Hotamisligil, 2010), perturbs hepatic lipid metabolism by modulating lipogenesis and lipoprotein metabolism (Kammoun et al., 2009; Lee et al., 2008; Ota et al., 2008; Rutkowski et al., 2008; So et al., 2012; Wang et al., 2012; Zhang et al., 2011), and results in the development of fatty liver, reminiscent of essential aspects of SIRT7 KO phenotype in the liver (Figure 1). We therefore hypothesized that SIRT7 might be an essential regulator of the UPR. ER stress triggers finely regulated signaling events and transcriptional activation of ER stress response target genes with well-defined *cis*-elements (Hetz, 2012; Walter and Ron, 2011). Analysis of the SIRT7 promoter using MATInspector identified potential binding elements for XBP1, a key regulator of the UPR, that preferentially binds to sequences containing an ACGT core (Figure S2A) (Acosta-Alvear et al., 2007; Sha et al., 2009). Thus, SIRT7 may be transcriptionally upregulated upon ER stress. Indeed, treatment with chemical inducers of ER stress, tunicamycin and thapsigargin, increased the expression of SIRT7 at both the mRNA and protein levels in various cell types (Figures 3A and 3B; Figures S2B–S2D). However, ER stress did not induce SIRT7 expression in XBP1 KO mouse embryonic fibroblasts (MEFs) (Figures 3A and 3B). Furthermore, ectopic expression of the spliced XBP1 (XBP1s), the active form of XBP1, induced SIRT7 expression to the same degree as it induced Erdj4, a known XBP1s target (Figures S2E and S2F). Additionally, in a luciferase assay, XBP1s induced transcription driven by the SIRT7 promoter, but not the SIRT3 promoter (Figure S2G). SIRT7 activity is dependent on NAD⁺, but




Figure 1. SIRT7 Prevents the Development of Fatty Liver Disease

(A) Schematic representation of the SIRT7 locus and the SIRT7 KO-targeting vector.

(B) Southern blot confirming the generation of SIRT7 KO mice.

(C) Morphology, H&E staining, Oil Red O staining, and F4/80 staining showing increased lipid accumulation and inflammation in the livers of SIRT7 KO mice compared to WT controls.

(D) Quantification of triglyceride extracted from livers in a colorimetric assay showing increased triglyceride content in SIRT7 KO livers. $n = 6$.

(E) Gene expression analysis by quantitative PCR showing increased inflammation and lipogenesis, but not fatty acid oxidation in SIRT7 KO livers. $n = 4$.

(F) A VLDL-TG secretion assay showing defective VLDL-TG secretion for SIRT7 KO mice. $n = 6$.

(G) Quantification of VLDL particle concentration in the blood using a gas-phase differential electrical mobility analyzer showing reduced VLDL particle concentration in the blood of SIRT7 KO mice. $n = 6$.

Error bars represent SEM. * $p < 0.05$. ** $p < 0.01$. *** $p < 0.001$. ns: $p > 0.05$. See also [Figure S1](#).

tunicamycin treatment did not change the cellular NAD⁺ levels ([Figure S2H](#)). Thus, SIRT7 is induced transcriptionally by XBP1 upon ER stress.

Next, we tested the effects of altered SIRT7 levels on ER stress. Overexpression of SIRT7 substantially reduced ER stress in response to tunicamycin, as evidenced by the reduced eIF2 α




Figure 2. Hepatic SIRT7 Prevents the Development of Fatty Liver Autonomously

(A–F) Data shown are the comparison for the livers of WT, SIRT7 KO, and SIRT7 KO mice expressing SIRT7 in livers via AAV8-mediated gene transfer. (A) Western analysis for SIRT7 expression. (B) Liver morphology. (C) Liver triglyceride quantification. (D) H&E and Oil Red O staining. (E) Gene expression by quantitative PCR showing that SIRT7 expression in liver suppresses hepatic inflammation and lipogenesis of SIRT7 KO mice. (F) In a VLDL-TG secretion assay, SIRT7 expression in liver rescued VLDL-TG secretion defects of SIRT7 KO mice. $n = 4$. Error bars represent SEM. * $p < 0.05$. ** $p < 0.01$. *** $p < 0.001$. ns: $p > 0.05$.

Together, these data suggest that SIRT7 activation is a critical event of the UPR to alleviate ER stress.

Myc Recruits SIRT7 to Repress the Expression of Ribosomal Proteins and to Suppress ER Stress

How does SIRT7 regulate the UPR? The initial phase of the UPR is suppression of protein translation and increased production of molecular chaperones to re-establish homeostasis (Hetz, 2012; Walter and Ron, 2011). SIRT7 deacetylates H3K18Ac at specific gene promoters to repress transcription, and a major class of SIRT7-target genes are involved in protein translation and ribosome biogenesis (Barber et al., 2012). Thus, SIRT7 may alleviate ER stress by suppressing ribosome biogenesis and protein translation. Although SIRT7 lacks known DNA binding motifs, previous

work has shown that it is recruited to a subset of its target promoters via interaction with the transcription factor ELK4 (Barber et al., 2012). However, ELK4 was dispensable for SIRT7 promoter occupancy at other target promoters, including the promoters of ribosomal protein genes, leaving open the question of how SIRT7 is recruited to such promoters. Recently, Myc has been shown to coordinate the transcriptional control of ribosomal components and serve as a master regulator of ribosome biogenesis (Kim et al., 2000; van Riggelen et al., 2010). Chromatin remodeling is believed to be central to Myc function in modulating the expression of its target genes (Knoepfler et al., 2006; van Riggelen et al., 2010). Myc binding on target chromatin is associated with the H3K18Ac mark, a substrate of SIRT7 (Martinato et al., 2008). We therefore hypothesized that Myc stabilizes SIRT7 at the promoters of ribosomal proteins to mediate chromatin remodeling and gene repression. To probe a potential connection between Myc and SIRT7, we tested whether SIRT7 physically interacts with Myc in vivo. Western blot analysis of the immunoprecipitates from

phosphorylation and expression of ER stress response genes (Figures 3C–3F; Figures S2I and S2J). A catalytically inactive SIRT7 mutant (H187Y) did not suppress ER stress (Figures 3C–3F), linking the catalytic activity of SIRT7 to ER stress management. To determine whether endogenous SIRT7 prevents ER stress, we stably knocked down SIRT7 expression using two independent short hairpin RNAs that specifically target SIRT7. SIRT7 depletion led to constitutive induction of ER stress (Figures 3G and S2K). Similarly, ER stress response genes were also upregulated in SIRT7 KO MEFs (Figure 3H). ER stress induces a reduction in the polysome-to-monosome ratio, indicative of translational initiation blockade (Kawai et al., 2004). Ribosomal profiling via a sucrose gradient showed that in contrast to WT MEFs, which had a high polysome-to-monosome ratio, SIRT7 KO MEFs had a reduction in polysomes and an increase in low-molecular-weight monosomes, consistent with increased ER stress in SIRT7 KO MEFs (Figure 3I). SIRT7 also suppressed ER stress-induced cell death (Figures 3J, 3K, S2L, and S2M), but not general apoptosis (Figures S2N and S2O).

work has shown that it is recruited to a subset of its target promoters via interaction with the transcription factor ELK4 (Barber et al., 2012). However, ELK4 was dispensable for SIRT7 promoter occupancy at other target promoters, including the promoters of ribosomal protein genes, leaving open the question of how SIRT7 is recruited to such promoters.

Recently, Myc has been shown to coordinate the transcriptional control of ribosomal components and serve as a master regulator of ribosome biogenesis (Kim et al., 2000; van Riggelen et al., 2010). Chromatin remodeling is believed to be central to Myc function in modulating the expression of its target genes (Knoepfler et al., 2006; van Riggelen et al., 2010). Myc binding on target chromatin is associated with the H3K18Ac mark, a substrate of SIRT7 (Martinato et al., 2008). We therefore hypothesized that Myc stabilizes SIRT7 at the promoters of ribosomal proteins to mediate chromatin remodeling and gene repression. To probe a potential connection between Myc and SIRT7, we tested whether SIRT7 physically interacts with Myc in vivo. Western blot analysis of the immunoprecipitates from




Figure 3. SIRT7 Suppresses ER Stress

(A and B) Gene expression analysis by quantitative PCR (A) and western blotting (B) showing increased SIRT7 expression upon treatment of ER stress inducer thapsigargin (TG) in WT, but not XBP1 KO, MEFs.

(C–F) Western blots (C) and quantitative PCR (D–F) showing reduced ER stress in tunicamycin-treated stable HepG2 cells overexpressing (OE) WT, but not catalytically inactive (H187Y), SIRT7, as indicated by eIF2 α phosphorylation levels and ER stress response gene expression.

(G) Western blots showing increased ER stress in SIRT7 knockdown (KD) stable HepG2 cells.

(legend continued on next page)

Flag-tagged SIRT7-transfected cells revealed that Myc was associated with Flag-SIRT7 (Figure 4A). Western blot analysis of endogenous SIRT7 immunoprecipitates also revealed a specific interaction with Myc (Figure 4B). Thus, SIRT7 physically interacts with Myc in cells.

To investigate whether SIRT7 is stabilized at the promoters of ribosomal proteins through its interaction with Myc, we assessed SIRT7 and Myc co-occupancy at these promoter regions. Chromatin immunoprecipitation (ChIP) was performed with SIRT7 and Myc antibodies to assess the genomic occupancy of endogenous SIRT7 and Myc, compared to immunoglobulin G (IgG)-negative background control. SIRT7 bound to the core promoters of ribosomal proteins (Barber et al., 2012). Myc was also detected at the same regions as SIRT7 on the core promoters of ribosomal proteins, but not at the genomic region 6 kb upstream of the transcription start site (Figures 4C–4E; Figures S3A and S3B), consistent with the notion that Myc generally occupies the core promoter regions of actively transcribed genes (Lin et al., 2012). In contrast, Myc was not detected at the promoters of NME1 and COPS2, where SIRT7 binding is mediated through ELK4 (Figures 4F and S3C) (Barber et al., 2012). Myc depletion using small interfering RNA (siRNA) led to a significant reduction in SIRT7 occupancy at the promoters of ribosomal proteins (Figures 4G, 4H, S3D, and S3E). However, Myc depletion had no effect on SIRT7 occupancy at the promoters of NME1 and COPS2 (Figures 4I and S3F). These data suggest that Myc targets SIRT7 specifically to the promoters of ribosomal proteins.

The specific association of SIRT7 and Myc at the promoters of ribosomal proteins, but not at ELK4 target genes, suggests that SIRT7 might specifically influence the expression of ribosomal proteins via Myc. Therefore, we examined the effects of Myc inhibition on SIRT7-mediated gene expression. As shown previously, SIRT7 KD led to increased expression of ribosomal proteins (Figures 4J and 4K) (Barber et al., 2012). Tunicamycin treatment induced SIRT7 expression (Figures 3A and 3B; Figures S2B–S2D) and suppressed the expression of ribosomal proteins in control cells, but not in SIRT7 KD cells (Figures 4J and 4K), indicating that ER stress triggers the transcriptional silencing of ribosomal proteins and that this effect requires SIRT7. Strikingly, Myc inactivation via siRNA or by a specific inhibitor, 10058-F4, abrogated the elevated expression of ribosomal proteins, but not NME1 in SIRT7 KD cells (Figures 4J, 4K, S3G, and S2H), demonstrating that the observations on ribosomal proteins are due to Myc-dependent effects of SIRT7 on gene expression. Together, these data suggest that Myc targets SIRT7 specifically to the promoters of ribosomal proteins for transcriptional silencing.

The observation that ER stress represses ribosomal proteins in a SIRT7-dependent manner suggests that upon ER stress, the induction of SIRT7 might function to suppress ribosomal protein expression and protein translation, in order to relieve ER

stress. To test this possibility, we assessed the effects of suppressing Myc-mediated expression of ribosomal proteins on SIRT7-associated ER stress management. Myc inactivation by siRNA or 10058-F4 abrogated the increased ER stress in SIRT7 KD cells (Figures 4K and S3I). Myc inactivation also blunted the effects of SIRT7 on ER stress resistance (Figures S3J and S3K). Together, these results indicate that SIRT7 is targeted to the promoters of ribosomal proteins by interacting with Myc and alleviates ER stress by countering Myc-dependent expression of these genes.

ER Stress Underlies the Development of Fatty Liver Caused by SIRT7 Deficiency

We next determined whether ER stress underlies the development of fatty liver in SIRT7-deficient mice. SIRT7 KO livers showed increased ER stress, as evidenced by the induction of e-IF2 α phosphorylation and ER stress response genes, as well as a reduction in the polysome-to-monosome ratio (Figures 5A–5C). Reintroduction of SIRT7 in the livers of SIRT7 KO mice via AAV8-mediated gene transfer reduced ER stress (Figure 5D) and reverted the fatty liver phenotype (Figures 2A–2E). Moreover, treatment of SIRT7 KO mice with TUDCA, a small molecule chaperone that has been shown to alleviate ER stress in vivo (Ozcan et al., 2006), partially rescued the fatty liver phenotype (Figures 5E and 5F). Finally, liver-specific Myc knockdown in SIRT7 KO mice via AAV8-mediated gene transfer reduced ER stress (Figures 6A and 6B), suppressed hepatic inflammation and lipogenesis (Figures 6C and 6D), and reversed the fatty liver phenotype (Figures 6D and 6E). Thus, SIRT7 prevents the development of fatty liver by suppressing ER stress.

SIRT7 Reverts Fatty Liver Associated with Diet-Induced Obesity

A high-fat, high-calorie diet is associated with increased ER stress and the development of fatty liver disease (Oyadomari et al., 2008). We next asked whether SIRT7 could be targeted to alleviate high-fat diet-induced ER stress and the development of fatty liver disease. We overexpressed SIRT7 specifically in the livers of mice fed a high-fat diet via AAV8-mediated gene transfer. Consistent with previous reports (Oyadomari et al., 2008), high-fat diet feeding led to an increase in ER stress markers (Figures 7A and 7B), accumulation of TG in the liver (Figures 7C–7E), possibly because of decreased VLDL secretion and increased lipogenesis (Figures 7A and 7F), and increased inflammation (Figures 7A and 7D). Strikingly, SIRT7 overexpression in the livers of high-fat-diet-fed mice suppressed ER stress (Figures 7A and 7B), VLDL secretion (Figure 7F), lipogenesis (Figure 7A), and inflammation (Figures 7A and 7D) and rescued the fatty liver phenotype (Figures 7C–7E). Thus, SIRT7 activation represents an attractive approach to revert ER stress-mediated fatty liver.

(H and I) Increased ER stress in SIRT7 KO MEFs. Quantitative PCR showing increased expression of ER stress-induced genes in SIRT7 KO MEFs (H). Polysome profiling showing a reduced polysome-to-monosome ratio, an ER stress marker indicative of translational initiation blockade (I).

(J and K) SIRT7 prevents ER stress-induced cell death. Stable SIRT7 overexpression (OE) or knockdown (KD1 or KD2) cells used in Figures 3C and 3G were treated with tunicamycin (2 μ g/ml for J and 1 μ g/ml for K) for 24 hr. Apoptosis was scored with Annexin V staining.

Error bars represent SEM. * $p < 0.05$. ** $p < 0.01$. *** $p < 0.001$. See also Figure S2.




Figure 4. Myc Recruits SIRT7 to Repress the Expression of Ribosomal Proteins and to Suppress ER Stress

(A) Western analysis showing colIP of Flag-tagged SIRT7 and endogenous Myc in 293T cells.
 (B) Western blots showing colIP of endogenous SIRT7 and Myc in Hep G2 cells.
 (C–F) ChIP-qPCR (mean \pm SEM) showing Myc occupancy at the RPS20 proximal promoter, but not 6 kb upstream, compared to IgG-negative control samples. Myc occupancy at the γ -tubulin and NME1 promoters was used as negative controls. All samples were normalized to input DNA.
 (G) Western blots showing knockdown of Myc with siRNA in cells used in (H and I).
 (H and I) Reduction of SIRT7 occupancy at the RPS20, but not the NME1, promoter in Myc knockdown cells determined by ChIP (mean \pm SEM).
 (legend continued on next page)




Figure 5. SIRT7 Prevents the Development of Fatty Liver by Suppressing ER Stress

(A–C) Western analysis (A), quantitative PCR (B), and polysome profiling (C) showing increased ER stress in SIRT7 KO livers.

(D) Quantitative PCR showing decreased expression of ER stress response genes in SIRT7 KO livers by reintroduction of SIRT7 via AAV8-mediated gene transfer.

(E and F) H&E, Oil Red O staining, and F4/80 staining of liver sections (E), and quantification of liver triglyceride (F) showing a small molecule chaperone TUDCA partially reverses fatty liver phenotype of SIRT7 KO mice. $n = 6$.

Error bars represent SEM. * $p < 0.05$. ** $p < 0.01$. *** $p < 0.001$.

(phosphorylation of eIF2 α , blockade of polysome assembly and translational initiation, and induction of ER stress response genes), apoptosis, and inflammation and, specifically in the liver, increased lipogenesis and reduced VLDL secretion (Figure S4B). Importantly, SIRT7 upregulation alleviates ER stress and restores hepatic metabolic homeostasis in diet-induced obese animals, providing an avenue to treat fatty liver disease and likely other ER stress-associated pathologies.

Our observation that SIRT7 opposes Myc-dependent gene regulation is intriguing. In addition to its role in fatty liver pathology, the dynamic balance between SIRT7 and Myc activities could play a pivotal role in the context of tumorigenesis. SIRT7 may keep the Myc oncogene in check to prevent tumorigenesis. However, under certain conditions, Myc can also trigger apoptosis as an evolved tumor defense mechanism (Evan et al., 1992; Soucek and Evan, 2010).

Our study suggests that unopposed Myc activation in the absence of SIRT7 results in ER stress. This may be an underlying mechanism for Myc-induced apoptosis in cancer cells that are particularly sensitive to ER stress (Ozcan and Tabas, 2012). Indeed, we find that SIRT7 prevents ER stress-induced cell death in a Myc-dependent manner (Figures S3J and S3K). These observations are consistent with previous findings that SIRT7 can promote cancer cell survival and maintain oncogenic transformation (Barber et al., 2012).

Our findings regarding the role of SIRT7 in ER stress management might also be important for aging biology. Indeed, SIRT7 has several potential links to aging. SIRT7 KO mice have shortened life spans and exhibit phenotypes linked to

DISCUSSION

Our work uncovered a physiological role of SIRT7 in maintaining hepatic metabolic homeostasis, demonstrated the feasibility of targeting SIRT7 to restore metabolic homeostasis in animals with metabolic disorders, and revealed a regulatory branch of the UPR that is amenable for therapeutic targeting. The canonical UPR leads to the phosphorylation of eIF2 α and the suppression of translation to alleviate ER stress (Ozcan and Tabas, 2012; Walter and Ron, 2011). We show that SIRT7 is induced by XBP1 upon ER stress and is recruited to the promoters of ribosomal proteins via Myc to repress gene expression and to alleviate ER stress (Figure S4A). In SIRT7-deficient cells, failure to engage SIRT7-mediated ER stress management results in constitutive ER stress that induces activation of the canonical UPR

(J) Western analysis showing Myc inactivation by a small molecule inhibitor abrogates increased expression of ribosomal proteins, but not NME1, in stable SIRT7 KO cells used in Figure 3G.

(K) Western analysis showing Myc inactivation via siRNA abrogates ER stress and increased expression of ribosomal proteins, but not NME1, in stable SIRT7 KO cells used in Figure 3G.

Error bars represent SEM. * $p < 0.05$. *** $p < 0.001$. ns: $p > 0.05$. See also Figure S3.




Figure 6. SIRT7 Suppresses ER Stress and Prevents the Development of Fatty Liver by Repressing Myc

(A–E) Data shown are the comparison of WT, SIRT7 KO, and SIRT7 KO mice with Myc knock-down in livers via AAV8-mediated gene transfer. (A) Myc expression by quantitative PCR. (B) ER stress in the livers by western analysis. (C) Quantitative PCR analysis for the expression of inflammation, lipogenesis, and fatty acid oxidation genes. (D) H&E, Oil Red O staining, and F4/80 staining of liver sections. (E) Liver triglyceride quantification. n = 5.

Error bars represent SEM. * p < 0.05. ** p < 0.01. *** p < 0.001. ns: p > 0.05. See also Figure S4.

with Kpn1 and the membranes were probed with a 500 bp fragment covering exons 1–3. Chimeric mice were generated by injection of targeted ES clones into C57BL6/J blastocysts. Male chimeras were mated with 129Sv females to generate F1 heterozygous mice, which were interbred to generate homozygous KO mice, which were screened by Southern blotting as described above (Figure S1A).

All mice were housed on a 12:12 hr light:dark cycle at 25°C. Experiments were performed using 4- to 6-month-old male littermates. Mice were randomized into each of the groups. Samples were processed blindly during the experiments, and the outcome was assessment. All animal procedures were in accordance with the animal care committee at the University of California, Berkeley. The high-fat diet was provided by OpenSource Diets (D12079B). For TUDCA treatment, mice received intraperitoneal injections of 500 mg/kg/day TUDCA or PBS twice daily for 20 days as described previously (Cnop et al., 2012). VLDL-TG secretion assay was performed as described previously (Haeusler et al., 2010). Blood was collected from the mouse tail vein and kept on ice until centrifugation (1,500 g, 15 min at 4°C). VLDL concentration was quantified as described using a gas-phase differential electrical mobility analyzer (Caulfield et al., 2008). Plasma triglyceride was measured in accordance with the manufacturer's instruction (Wako Diagnostics). Triglycerides were extracted from liver tissues as described and were extracted from feces as described (Zadravec et al., 2010). Extracted triglyceride was quantified in accordance with the manufacturer's instruction (Wako Diagnostics). Liver tissues were processed for H&E staining, Oil Red O staining, and F4/80 staining as described previously (Sun et al., 2012).

For AAV-mediated gene transfer to the mouse liver, SIRT7 or Myc knock-down target sequence was cloned into dsAAV-RSVeGFP-U6 vector. dsAAV8 was produced by triple-transfection and CsCl purification, and virus titer was determined as previously described (Gao et al., 2006). Each mouse was injected with 3×10^{11} genome copies of virus via tail vein.

Cell Culture, RNAi, and Viral Transduction

293T, Hepa 1-6, and Hep G2 cells were acquired from the American Type Culture Collection. WT and SIRT7 KO MEFs were generated in accordance with standard procedures (Greber et al., 2007). XBP-1 KO MEFs were acquired from L. Glimcher (Lee et al., 2008). Cells were cultured in advanced Dulbecco's modified Eagle's medium (Invitrogen) supplemented with 1%

aging (Vakhrusheva et al., 2008). Furthermore, SIRT7 expression decreases in some aging tissues (Vakhrusheva et al., 2008). Suppression of ribosomal proteins and ER stress leads to life span extension in yeast (Steffen et al., 2012). Thus, the interplay between SIRT7, ER stress, and protein translation may represent an evolutionarily conserved phenomenon in aging.

EXPERIMENTAL PROCEDURES

Mice

Sirt7^{-/-} mice were produced by the VelociGene method (Valenzuela et al., 2003) (VelociGene allele identification number: VG321). The SIRT7 KO-targeting vector was constructed by replacing exons 4 to 11 with a LacZ gene inserted in-frame after the first few base pairs of exon 4 (Figures 1A and 1B). An Srf1-linearized targeting vector (30 μ g) was electroporated into CJ7, 129SJ ES cells as described previously (Valenzuela et al., 2003). Positives clones were identified by Taqman analysis (Valenzuela et al., 2003) and confirmed by Southern blot analysis in which genomic DNA was digested




Figure 7. SIRT7 Rescues High-Fat Diet-Induced Fatty Liver

(A–F) Data shown are a comparison of livers of mice fed a chow diet, a high-fat diet, and a high-fat diet with SIRT7 reintroduced specifically in the liver via AAV8-mediated gene transfer. (A) Quantitative PCR analyses for gene expression of ER stress, inflammation, and lipogenesis. (B) Western blotting for ER stress markers. (C) Morphology. (D) H&E staining, Oil Red O staining, and F4/80 staining. (E) Quantification of liver triglyceride. (F) A VLDL-TG secretion assay. $n = 6$. Error bars represent SEM. ** $p < 0.01$. *** $p < 0.001$.

penicillin-streptomycin (Invitrogen) and 10% FBS (Invitrogen). For ER stress induction, cells were treated with tunicamycin (Sigma, 1 $\mu\text{g/ml}$) or thapsigargin (Sigma, 0.1 μM) for 24 hr for biochemical analysis or 48 hr for cell survival analysis. Cells were treated with 0.5 μM staurosporin for 48 hr for cell survival anal-

ysis. Cell survival was scored with trypan blue staining or Annexin V staining (Biolegend). For Myc inactivation, cells were treated with 10058-F4 (Sigma, 2 μM) for 24 hr before analysis. Cellular NAD⁺ concentration was quantified using EnzyChrom NAD/NADH Assay Kit (Bioassay System).

SIRT7 knockdown target sequences are as follows, as previously described (Barber et al., 2012):

S7KD1, 5'-CACCTTTCTGTGAGAACGGAA-3',
S7KD2, 5'-TAGCCATTTGTCCTTGAGGAA-3'.

Myc knockdown target sequences are as follows, as previously described (Caulfield et al., 2008):

Myc KD (human), 5'-GGACTATCCTGCTGCCAAG-3',
Myc KD (mouse), 5'-CCCAAGGTAGTGATCCTCAAA-3'.

Double-stranded siRNAs were purchased from Thermo Scientific and were transfected into cells via RNAiMax (Invitrogen) in accordance with the manufacturer's instruction.

For lentiviral packaging, 293T cells were cotransfected with packaging vectors (pCMV-dR8.2 dvpr and pCMV-VSV-G) and the pSiCoR-SIRT7 knockdown or empty construct. For retroviral packaging, 293T cells were cotransfected with packaging vectors (pVPack-VSV-G, pVPack-GP) and pBABE-SIRT7, pBABE-SIRT7 H187Y, or empty construct. Viral supernatant was harvested after 48 hr. For transduction, cells were incubated with virus-containing supernatant in the presence of 8 μ g/ml polybrene. After 48 hr, infected cells were selected with puromycin (1 μ g/ml).

Coimmunoprecipitations

Coimmunoprecipitations (coIPs) were performed as previously described (Qiu et al., 2010) with Flag-resin (Sigma) or Protein A/G beads (Santa Cruz) for SIRT7 IP. Elution was performed with either Flag peptide (Sigma) or 100 mM Glycine solution (pH 3) for SIRT7 IP. Antibodies are provided in Table S1.

ChIP and mRNA Analysis

Cells were prepared for ChIP as previously described (Dahl and Collas, 2007), with the exception that DNA was washed and eluted using a PCR purification kit (QIAGEN) rather than by phenol-chloroform extraction. RNA was isolated from cells or tissue using Trizol reagent (Invitrogen) and purified using the RNeasy Mini Kit (QIAGEN). cDNA was generated using the qScript cDNA SuperMix (Quanta Biosciences). Gene expression was determined by real-time PCR using Eva quantitative PCR (qPCR) SuperMix Kit (BioChain Institute) on an ABI StepOnePlus system. All data were normalized to ActB or GAPDH expression. Antibodies and PCR primer details are provided in Tables S1, S2, and S3.

Polysomal Profiling

MEFs or liver tissues were collected for polysomal profiling as described (Zid et al., 2009). Briefly, ten million MEF cells or 0.1 g of liver were harvested and homogenized on ice in 400 μ l of solubilization buffer (300 mM NaCl, 50 mM Tris-HCl [pH 8.0], 10 mM MgCl₂, 1 mM EGTA, 200 μ g/ml heparin, 1 mM DTT, 400 U/ml RNasin plus (Promega), 1X complete, Mini Protease Inhibitor Cocktail (Roche), 0.2 mg/ml cycloheximide, 1% Triton X-100, and 0.1% sodium deoxycholate). Additional solubilization buffer was added for a total of 1 ml. Cell lysate was placed back on ice for 10 min before centrifuging at 16,000 g for 15 min at 4°C. The supernatant was applied to the top of a 10%–50% continuous sucrose gradient in high salt resolving buffer (140 mM NaCl, 25 mM Tris-HCl [pH 8.0], and 10 mM MgCl₂) and centrifuged in a Beckman SW41Ti rotor (Beckman Coulter) at 180,000 g for 90 min at 4°C. Gradients were fractionated with continuous monitoring of absorbance at 254 nm.

Statistical Analysis

The number of mice chosen for each experiment is based on the principle that the minimal number of mice is used to have sufficient statistical power and is comparable to published literature for the same assays performed. No animals were excluded from the analyses. Student's t test was used for data analysis. Error bars represent SE. In all corresponding figures, * represents $p < 0.05$. ** represents $p < 0.01$. and *** represents $p < 0.001$. ns represents $p > 0.05$.

SUPPLEMENTAL INFORMATION

Supplemental Information includes four figures and three tables and can be found with this article online at <http://dx.doi.org/10.1016/j.celrep.2013.10.007>.

ACKNOWLEDGMENTS

We thank the Kay lab for reagents. This work was supported by grants from the National Institutes of Health (R01AG040990 [to D.C.], R01AG028867 [to K.F.C.], training grant 3T32DK007217-36S1 [to S.P.], training grant T32 AG000266 [to M.M.]), the Searle Scholars Program (to D.C.), the Ellison Medical Foundation (to D.C. and K.F.C.), The Glenn Foundation (to D.C.), the American Heart Association (to D.C.), UCOP TRDRP (to D.C.), the Hellman Family Faculty Fund (D.C.), the National Science Foundation (to J.S.), the Siebel Stem Cell Institute (to D.C., M.M., N.N., and X.Q.), the Department of Veterans Affairs (to K.F.C., Merit Award), and the Italian Foundation for Cancer Research (to L.V.). F.A. is an Investigator of the Howard Hughes Medical Institute. A.J.M., D.M.V., and G.D.Y. are employees and shareholders of Regeneron Pharmaceuticals.

Received: September 4, 2013

Revised: September 30, 2013

Accepted: October 3, 2013

Published: November 7, 2013

REFERENCES

- Acosta-Alvarez, D., Zhou, Y., Blais, A., Tsikitis, M., Lents, N.H., Arias, C., Lennon, C.J., Kluger, Y., and Dynlacht, B.D. (2007). XBP1 controls diverse cell type- and condition-specific transcriptional regulatory networks. *Mol. Cell* 27, 53–66.
- Argo, C.K., and Caldwell, S.H. (2009). Epidemiology and natural history of non-alcoholic steatohepatitis. *Clin. Liver Dis.* 13, 511–531.
- Barber, M.F., Michishita-Kioi, E., Xi, Y., Tasselli, L., Kioi, M., Moqtaderi, Z., Tennen, R.I., Paredes, S., Young, N.L., Chen, K., et al. (2012). SIRT7 links H3K18 deacetylation to maintenance of oncogenic transformation. *Nature* 487, 114–118.
- Basseri, S., and Austin, R.C. (2012). Endoplasmic reticulum stress and lipid metabolism: mechanisms and therapeutic potential. *Biochem. Res. Int.* 2012, 841362.
- Baur, J.A., Ungvari, Z., Minor, R.K., Le Couteur, D.G., and de Cabo, R. (2012). Are sirtuins viable targets for improving healthspan and lifespan? *Nat. Rev. Drug Discov.* 11, 443–461.
- Bellet, M.M., Orozco-Solis, R., Sahar, S., Eckel-Mahan, K., and Sassone-Corsi, P. (2011). The time of metabolism: NAD⁺, SIRT1, and the circadian clock. *Cold Spring Harb. Symp. Quant. Biol.* 76, 31–38.
- Browning, J.D., and Horton, J.D. (2004). Molecular mediators of hepatic steatosis and liver injury. *J. Clin. Invest.* 114, 147–152.
- Browning, J.D., Szczepaniak, L.S., Dobbins, R., Nuremberg, P., Horton, J.D., Cohen, J.C., Grundy, S.M., and Hobbs, H.H. (2004). Prevalence of hepatic steatosis in an urban population in the United States: impact of ethnicity. *Hepatology* 40, 1387–1395.
- Caulfield, M.P., Li, S., Lee, G., Blanche, P.J., Salameh, W.A., Benner, W.H., Reitz, R.E., and Krauss, R.M. (2008). Direct determination of lipoprotein particle sizes and concentrations by ion mobility analysis. *Clin. Chem.* 54, 1307–1316.
- Cnop, M., Foufelle, F., and Velloso, L.A. (2012). Endoplasmic reticulum stress, obesity and diabetes. *Trends Mol. Med.* 18, 59–68.
- Cohen, J.C., Horton, J.D., and Hobbs, H.H. (2011). Human fatty liver disease: old questions and new insights. *Science* 332, 1519–1523.
- Dahl, J.A., and Collas, P. (2007). A quick and quantitative chromatin immunoprecipitation assay for small cell samples. *Front. Biosci.* 12, 4925–4931.
- Evan, G.I., Wyllie, A.H., Gilbert, C.S., Littlewood, T.D., Land, H., Brooks, M., Waters, C.M., Penn, L.Z., and Hancock, D.C. (1992). Induction of apoptosis in fibroblasts by c-myc protein. *Cell* 69, 119–128.

- Finkel, T., Deng, C.X., and Mostoslavsky, R. (2009). Recent progress in the biology and physiology of sirtuins. *Nature* 460, 587–591.
- Ford, E., Voit, R., Liszt, G., Magin, C., Grummt, I., and Guarente, L. (2006). Mammalian Sir2 homolog SIRT7 is an activator of RNA polymerase I transcription. *Genes Dev.* 20, 1075–1080.
- Fu, S., Watkins, S.M., and Hotamisligil, G.S. (2012). The role of endoplasmic reticulum in hepatic lipid homeostasis and stress signaling. *Cell Metab.* 15, 623–634.
- Gao, G.P., Lu, Y., Sun, X., Johnston, J., Calcedo, R., Grant, R., and Wilson, J.M. (2006). High-level transgene expression in nonhuman primate liver with novel adeno-associated virus serotypes containing self-complementary genomes. *J. Virol.* 80, 6192–6194.
- Gillum, M.P., Erion, D.M., and Shulman, G.I. (2010). Sirtuin-1 regulation of mammalian metabolism. *Trends Mol. Med.* 17, 8–13.
- Greber, B., Lehrach, H., and Adjaye, J. (2007). Fibroblast growth factor 2 modulates transforming growth factor beta signaling in mouse embryonic fibroblasts and human ESCs (hESCs) to support hESC self-renewal. *Stem Cells* 25, 455–464.
- Haeusler, R.A., Han, S., and Accili, D. (2010). Hepatic FoxO1 ablation exacerbates lipid abnormalities during hyperglycemia. *J. Biol. Chem.* 285, 26861–26868.
- Hetz, C. (2012). The unfolded protein response: controlling cell fate decisions under ER stress and beyond. *Nat. Rev. Mol. Cell Biol.* 13, 89–102.
- Hirschey, M.D., Shimazu, T., Huang, J.Y., Schwer, B., and Verdin, E. (2011). SIRT3 regulates mitochondrial protein acetylation and intermediary metabolism. *Cold Spring Harb. Symp. Quant. Biol.* 76, 267–277.
- Hotamisligil, G.S. (2010). Endoplasmic reticulum stress and the inflammatory basis of metabolic disease. *Cell* 140, 900–917.
- Houtkooper, R.H., Pirinen, E., and Auwerx, J. (2012). Sirtuins as regulators of metabolism and healthspan. *Nat. Rev. Mol. Cell Biol.* 13, 225–238.
- Imai, S., and Guarente, L. (2010). Ten years of NAD-dependent SIR2 family deacetylases: implications for metabolic diseases. *Trends Pharmacol. Sci.* 31, 212–220.
- Kammoun, H.L., Chabanon, H., Hainault, I., Luquet, S., Magnan, C., Koike, T., Ferré, P., and Foufelle, F. (2009). GRP78 expression inhibits insulin and ER stress-induced SREBP-1c activation and reduces hepatic steatosis in mice. *J. Clin. Invest.* 119, 1201–1215.
- Kawai, T., Fan, J., Mazan-Mamczarz, K., and Gorospe, M. (2004). Global mRNA stabilization preferentially linked to translational repression during the endoplasmic reticulum stress response. *Mol. Cell Biol.* 24, 6773–6787.
- Kim, S., Li, Q., Dang, C.V., and Lee, L.A. (2000). Induction of ribosomal genes and hepatocyte hypertrophy by adenovirus-mediated expression of c-Myc in vivo. *Proc. Natl. Acad. Sci. USA* 97, 11198–11202.
- Knoepfler, P.S., Zhang, X.Y., Cheng, P.F., Gafken, P.R., McMahon, S.B., and Eisenman, R.N. (2006). Myc influences global chromatin structure. *EMBO J.* 25, 2723–2734.
- Lee, A.H., Scapa, E.F., Cohen, D.E., and Glimcher, L.H. (2008). Regulation of hepatic lipogenesis by the transcription factor XBP1. *Science* 320, 1492–1496.
- Lin, C.Y., Lovén, J., Rahl, P.B., Paranal, R.M., Burge, C.B., Bradner, J.E., Lee, T.I., and Young, R.A. (2012). Transcriptional amplification in tumor cells with elevated c-Myc. *Cell* 151, 56–67.
- Martinato, F., Cesaroni, M., Amati, B., and Guccione, E. (2008). Analysis of Myc-induced histone modifications on target chromatin. *PLoS ONE* 3, e3650.
- Ota, T., Gayet, C., and Ginsberg, H.N. (2008). Inhibition of apolipoprotein B100 secretion by lipid-induced hepatic endoplasmic reticulum stress in rodents. *J. Clin. Invest.* 118, 316–332.
- Oyadomari, S., Harding, H.P., Zhang, Y., Oyadomari, M., and Ron, D. (2008). Dephosphorylation of translation initiation factor 2alpha enhances glucose tolerance and attenuates hepatosteatosis in mice. *Cell Metab.* 7, 520–532.
- Ozcan, L., and Tabas, I. (2012). Role of endoplasmic reticulum stress in metabolic disease and other disorders. *Annu. Rev. Med.* 63, 317–328.
- Ozcan, U., Yilmaz, E., Ozcan, L., Furuhashi, M., Vaillancourt, E., Smith, R.O., Görgün, C.Z., and Hotamisligil, G.S. (2006). Chemical chaperones reduce ER stress and restore glucose homeostasis in a mouse model of type 2 diabetes. *Science* 313, 1137–1140.
- Qiu, X., Brown, K., Hirschey, M.D., Verdin, E., and Chen, D. (2010). Calorie restriction reduces oxidative stress by SIRT3-mediated SOD2 activation. *Cell Metab.* 12, 662–667.
- Rutkowski, D.T., Wu, J., Back, S.H., Callaghan, M.U., Ferris, S.P., Iqbal, J., Clark, R., Miao, H., Hassler, J.R., Fornek, J., et al. (2008). UPR pathways combine to prevent hepatic steatosis caused by ER stress-mediated suppression of transcriptional master regulators. *Dev. Cell* 15, 829–840.
- Sha, H., He, Y., Chen, H., Wang, C., Zenno, A., Shi, H., Yang, X., Zhang, X., and Qi, L. (2009). The IRE1alpha-XBP1 pathway of the unfolded protein response is required for adipogenesis. *Cell Metab.* 9, 556–564.
- So, J.S., Hur, K.Y., Tarrio, M., Ruda, V., Frank-Kamenetsky, M., Fitzgerald, K., Kotlianski, V., Lichtman, A.H., Iwawaki, T., Glimcher, L.H., and Lee, A.H. (2012). Silencing of lipid metabolism genes through IRE1alpha-mediated mRNA decay lowers plasma lipids in mice. *Cell Metab.* 16, 487–499.
- Soucek, L., and Evan, G.I. (2010). The ups and downs of Myc biology. *Curr. Opin. Genet. Dev.* 20, 91–95.
- Starley, B.Q., Calcagno, C.J., and Harrison, S.A. (2010). Nonalcoholic fatty liver disease and hepatocellular carcinoma: a weighty connection. *Hepatology* 51, 1820–1832.
- Steffen, K.K., McCormick, M.A., Pham, K.M., MacKay, V.L., Delaney, J.R., Murakami, C.J., Kaeberlein, M., and Kennedy, B.K. (2012). Ribosome deficiency protects against ER stress in *Saccharomyces cerevisiae*. *Genetics* 191, 107–118.
- Sun, Z., Miller, R.A., Patel, R.T., Chen, J., Dhir, R., Wang, H., Zhang, D., Graham, M.J., Unterman, T.G., Shulman, G.I., et al. (2012). Hepatic Hdac3 promotes gluconeogenesis by repressing lipid synthesis and sequestration. *Nat. Med.* 18, 934–942.
- Vakhrusheva, O., Smolka, C., Gajawada, P., Kostin, S., Boettger, T., Kubin, T., Braun, T., and Bober, E. (2008). Sirt7 increases stress resistance of cardiomyocytes and prevents apoptosis and inflammatory cardiomyopathy in mice. *Circ. Res.* 102, 703–710.
- Valenzuela, D.M., Murphy, A.J., Frendewey, D., Gale, N.W., Economides, A.N., Auerbach, W., Poueymirou, W.T., Adams, N.C., Rojas, J., Yasenachak, J., et al. (2003). High-throughput engineering of the mouse genome coupled with high-resolution expression analysis. *Nat. Biotechnol.* 21, 652–659.
- van Riggelen, J., Yetil, A., and Felsher, D.W. (2010). MYC as a regulator of ribosome biogenesis and protein synthesis. *Nat. Rev. Cancer* 10, 301–309.
- Walter, P., and Ron, D. (2011). The unfolded protein response: from stress pathway to homeostatic regulation. *Science* 334, 1081–1086.
- Wang, S., Chen, Z., Lam, V., Han, J., Hassler, J., Finck, B.N., Davidson, N.O., and Kaufman, R.J. (2012). IRE1alpha-XBP1s induces PDI expression to increase MTP activity for hepatic VLDL assembly and lipid homeostasis. *Cell Metab.* 16, 473–486.
- Zdravcov, D., Brolinson, A., Fisher, R.M., Carneheim, C., Csikas, R.I., Bertrand-Michel, J., Borén, J., Guillou, H., Rudling, M., and Jacobsson, A. (2010). Ablation of the very-long-chain fatty acid elongase ELOVL3 in mice leads to constrained lipid storage and resistance to diet-induced obesity. *FASEB J.* 24, 4366–4377.
- Zhang, K., Wang, S., Malhotra, J., Hassler, J.R., Back, S.H., Wang, G., Chang, L., Xu, W., Miao, H., Leonardi, R., et al. (2011). The unfolded protein response transducer IRE1alpha prevents ER stress-induced hepatic steatosis. *EMBO J.* 30, 1357–1375.
- Zid, B.M., Rogers, A.N., Katewa, S.D., Vargas, M.A., Kolipinski, M.C., Lu, T.A., Benzer, S., and Kapahi, P. (2009). 4E-BP extends lifespan upon dietary restriction by enhancing mitochondrial activity in *Drosophila*. *Cell* 139, 149–160.