

Available online at www.sciencedirect.com

J. Math. Pures Appl. 95 (2011) 99–135

JOURNAL
DE
MATHÉMATIQUES
PURES ET APPLIQUÉESwww.elsevier.com/locate/matpur

Weak KAM aspects of convex Hamilton–Jacobi equations with Neumann type boundary conditions

Hitoshi Ishii

Department of Mathematics, Faculty of Education and Integrated Arts and Sciences, Waseda University, 1-6-1 Nishi-Waseda, Shinjuku, Tokyo 169-8050, Japan

Received 10 October 2009

Available online 16 October 2010

Dedicated to Professor L.C. Evans on the occasion of his 60th birthday

Abstract

We study convex Hamilton–Jacobi equations $H(x, Du) = 0$ and $u_t + H(x, Du) = 0$ in a bounded domain Ω of \mathbb{R}^n with the Neumann type boundary condition $D_\gamma u = g$ in the viewpoint of weak KAM theory, where γ is a vector field on the boundary $\partial\Omega$ pointing a direction oblique to $\partial\Omega$. We establish the stability under the formations of infimum and of convex combinations of subsolutions of convex Hamilton–Jacobi equations, some comparison and existence results for convex and coercive Hamilton–Jacobi equations with the Neumann type boundary condition as well as existence results for the Skorokhod problem. We define the Aubry set associated with the Neumann type boundary problem and establish some properties of the Aubry set including the existence results for the “calibrated” extremals for the corresponding action functional (or variational problem).

© 2010 Elsevier Masson SAS. All rights reserved.

Résumé

On étudie du point de vue de la théorie KAM-faible, des équations de Hamilton–Jacobi de la forme $H(x, Du) = 0$ et $u_t + H(x, Du) = 0$, pour un hamiltonien convexe H , dans un domaine borné Ω de \mathbb{R}^n , avec une condition au bord de type Neumann oblique : $D_\gamma u = g$, γ étant un champ de vecteurs sur le bord $\partial\Omega$, pointant dans une direction oblique à $\partial\Omega$. On montre la stabilité par infimum et par combinaison convexe des sous-solutions de ces équations lorsque le hamiltonien H est convexe, on établit des théorèmes de comparaison et d’existence lorsque le hamiltonien est convexe et coercif et avec condition de type Neumann au bord, ainsi que des résultats d’existence pour le problème de Skorokhod. On définit l’ensemble de Aubry associé aux conditions de type Neumann et on établit des propriétés de cet ensemble, notamment des résultats d’existence d’extrémales « calibrées » pour la fonctionnelle d’action (ou problème variationnel) correspondante.

© 2010 Elsevier Masson SAS. All rights reserved.

MSC: 35B20; 70H20; 37J50; 49L20

Keywords: Hamilton–Jacobi equations; Neumann type boundary conditions; Weak KAM theory; Aubry–Mather theory; Viscosity solutions

E-mail address: hitoshi.ishii@waseda.jp.

0021-7824/\$ – see front matter © 2010 Elsevier Masson SAS. All rights reserved.

doi:10.1016/j.matpur.2010.10.006

1. Introduction

Let Ω be a bounded, open connected subset of \mathbb{R}^n with C^1 boundary. We denote by Γ its boundary $\partial\Omega$. We consider the Hamilton–Jacobi (HJ for short) equation with the Neumann type (or, in other words, oblique) boundary condition:

$$H(x, Du(x)) = a \quad \text{in } \Omega, \quad (1)$$

$$D_\gamma u(x) = g(x) \quad \text{on } \Gamma. \quad (2)$$

Here a is a constant, H is a given continuous function on $\overline{\Omega} \times \mathbb{R}^n$, called a Hamiltonian, u represents the unknown function on $\overline{\Omega}$, Du denotes the gradient $(u_{x_1}, \dots, u_{x_n})$, $D_\gamma u = D_\gamma u(x)$ denotes the directional derivative $\gamma(x) \cdot Du(x)$ at x , γ is a continuous vector field: $\Gamma \rightarrow \mathbb{R}^n$, and g is a given continuous function on Γ .

In addition to the continuity assumption on H , g , γ , we make the following standing assumptions.

(A1) H is a convex Hamiltonian, i.e., for each $x \in \overline{\Omega}$ the function $H(x, \cdot)$ is convex on \mathbb{R}^n .

(A2) H is coercive. That is, $\lim_{|p| \rightarrow \infty} H(x, p) = \infty$ for all $x \in \overline{\Omega}$.

(A3) γ is oblique to Γ . That is, for any $x \in \Gamma$, if $\nu(x)$ denotes the outer unit normal vector at x , then $\nu(x) \cdot \gamma(x) > 0$.

We consider the initial-value problem with the Neumann type (oblique) boundary condition:

$$u_t(x, t) + H(x, Du(x, t)) = 0 \quad \text{for } (x, t) \in \Omega \times (0, T), \quad (3)$$

$$D_\gamma u(x, t) = g(x) \quad \text{for } (x, t) \in \Gamma \times (0, T), \quad (4)$$

$$u(x, 0) = u_0(x) \quad \text{for } x \in \overline{\Omega}, \quad (5)$$

where $0 < T \leq \infty$ is a given constant, u represents the unknown function on $\overline{\Omega} \times [0, T]$, Du denotes the spatial gradient of u , $D_\gamma u = \gamma \cdot Du$, and u_0 is a given continuous function on $\overline{\Omega}$.

We call (1) and (3) *convex Hamilton–Jacobi equations* if H is a convex Hamiltonian.

The study of weak solutions (i.e., viscosity solutions) of problems (1), (2) and (3)–(5) goes back to Lions [1], and the theory of existence and uniqueness of viscosity solutions of such boundary or initial-boundary value problems including the case of second-order elliptic or parabolic equations has been well-developed. We refer for the developments to [1–6] and references therein. However, if problem (1), (2) has a solution, then it admits clearly multiple solutions and therefore the problem is a bit out of the scope of such developments. Indeed, problem (1), (2) has a solution only if a is assigned a specific value.

The problem of finding a pair $(a, u) \in \mathbb{R} \times C(\overline{\Omega})$ for which u is a solution of (1), (2) is called an ergodic problem in terms of optimal control or an additive eigenvalue problem, and it is also part of weak KAM theory. See [7] for a classical fundamental work on the ergodic problem for (1) in the periodic setting and also [8,9].

Weak KAM theory concerns the link between the HJ equation (1) in a domain Ω , with an appropriate boundary condition on its boundary $\partial\Omega$, and the Lagrangian flow generated by the Lagrangian L given by:

$$L(x, \xi) = \sup_{p \in \mathbb{R}^n} (\xi \cdot p - H(x, p)), \quad (6)$$

(or the extremals (minimizers) to the action functional associated with L). We refer [10,11,8,12] for pioneering work and further developments. We refer to [13] for some results in this direction on HJ equations with the state-constraint boundary condition.

A typical application of weak KAM theory to the evolution equation (3) is in the study of the long-time behavior of solutions of (3) with appropriate initial and boundary conditions. For these applications we refer to [14–20].

Our purpose in this paper is to establish some theorems concerning weak KAM theory for convex HJ equations. Indeed, we define the critical value (or the additive eigenvalue) and the Aubry set associated with (1), (2) and establish some of basic properties of the Aubry set, representation formulas for solutions of (1), (2) and the existence of extremals (or minimizers) for variational formulas of certain types of solutions of (1), (2). Our approach is relatively close to that of [21,22] in view of weak KAM theory. The paper [23] by O.-S. Serea deals with HJ equations on a convex domain with homogeneous Neumann condition in view of weak KAM theory. The requirements on the Lagrangian in [23] (see the conditions (7)–(10)) seem very restrictive. On the other hand, no regularity on the domain

other than the convexity is posed in [23]. In some special cases, the state-constraint problem for (1) is equivalent to the Neumann type problem (1), (2), and thus some results in [13] are related to those obtained here. For this equivalence, we refer for instance to [24].

As is somehow well known in optimal control, the solutions of (3)–(5) or (1), (2) should be given by the value function of the corresponding optimal control problem, whose dynamics is described by the Skorokhod problem. We present here a treatment of the Skorokhod problem suitable to our formulas for the solutions of (3)–(5) or of (1), (2), which is slightly different from usual treatments of the Skorokhod problems for applications to stochastic differential equations. The formulas thus obtained allows us to discuss the dynamical properties like “extremal curves” associated with (1), (2) or (3)–(5).

One of main difficulties in our approach was how to recover the convexity of problems (1), (2) or (3)–(5) in the viscosity formulation. In problems (1), (2) or (3)–(5) the Hamiltonian $H(x, p)$ is convex in p and the boundary condition, $D_\gamma u = g$, is convex in $p = Du$, but, for instance, in the definition of viscosity subsolution of (1), (2), it is formally required that

$$\min\{H(x, Du(x)), D_\gamma u(x) - g(x)\} \leq 0 \quad \text{on } \partial\Omega,$$

where the left-hand side is no more convex in $p = Du(x)$. This difficulty will be indeed solved in Theorems 2.1 and 2.2 below.

This paper is organized as follows. In the next section, we establish the stability under the formations of infimum and of convex combinations of subsolutions of (1), (2) and of (3)–(5). In Section 3 we establish comparison results for sub and supersolutions of (1), (2) and of (3)–(5). Section 4 is devoted to the Skorokhod problem in $\bar{\Omega}$ with reflection direction γ , which is essential to formulate variational representations for solutions of (1), (2) and of (3)–(5), and we establish results concerning existence and stability of solutions of the Skorokhod problem. In Section 5, we prove the existence of a solution of the initial-boundary value problem (3)–(5) as well as a variational formula for the solution. In Section 6, we introduce the critical value and the Aubry set associated with (1), (2), study basic properties of the Aubry set and establish representation formulas, based on the Aubry set, for solutions of (1), (2). In Section 7 we establish the existence of “calibrated” extremals for the variational problem associated with (1), (2).

Notation. Let e_i , with $i = 1, 2, \dots, n$, denote the unit vector of \mathbb{R}^n having unity as its i th coordinate. We $a \wedge b$ and $a \vee b$ for $\min\{a, b\}$ and $\max\{a, b\}$, respectively. For $A \subset \mathbb{R}^n$, $\text{Lip}(A, \mathbb{R}^m)$ (resp., $\text{BUC}(A, \mathbb{R}^m)$ and $\text{UC}(A, \mathbb{R}^m)$) denotes the space of Lipschitz continuous (resp., bounded uniformly continuous and uniformly continuous) functions on A with values in \mathbb{R}^m . For brevity, we may write $\text{Lip}(A)$, $\text{BUC}(A)$ and $\text{UC}(A)$ for $\text{Lip}(A, \mathbb{R}^m)$, $\text{BUC}(A, \mathbb{R}^m)$ and $\text{UC}(A, \mathbb{R}^m)$, respectively. We write A^c to denote the complement of A . For given function g on A with values in \mathbb{R}^m , we write $\|g\|_\infty = \sup_{x \in A} |g(x)|$. For an interval I , we denote by $\text{AC}(I)$ or $\text{AC}(I, \mathbb{R}^n)$ the space of absolutely continuous functions on I with values in \mathbb{R}^n . For a given function $w : A \rightarrow \mathbb{R}$, w^* and w_* denote respectively the upper and lower semicontinuous envelopes of w defined on \bar{Q} . Regarding the definition of (viscosity) solutions, we adopt the following convention: for instance, we consider (1), (2). A function $u : \bar{\Omega} \rightarrow \mathbb{R}$ is a subsolution (resp., a supersolution) provided that u is bounded above (resp., bounded below) and whenever $(x, \phi) \in \bar{\Omega} \times C^1(\bar{\Omega})$ and $u^* - \phi$ (resp., $u_* - \phi$) attains a maximum (resp., a minimum) at x , $H(x, D\phi(x)) \leq a$ (resp., $\geq a$) if $x \in \Omega$ and either $H(x, D\phi(x)) \leq a$ (resp., $\geq a$) or $D_\gamma \phi(x) \leq g(x)$ (resp., $\geq g(x)$) if $x \in \Gamma$. A bounded function $u : \bar{\Omega} \rightarrow \mathbb{R}$ is a solution if it is both a subsolution and a supersolution. In a more general situation where a candidate of solutions, u , is defined on a set which is not necessarily compact, the requirement on u regarding the boundedness to be a solution (resp., subsolution or supersolution) is that it is locally bounded (resp., locally bounded above or locally bounded below).

2. Basic propositions on convex HJ equations

In this section we establish the stability of the operations of infimum and of convex combinations subsolutions of convex HJ equations. We remark that these stability properties, without boundary condition, is the main technical observations in the theory of lower semicontinuous viscosity solutions due to Barron and Jensen [25].

To localize problems (1), (2), or (3)–(5), let U be an open subset of \mathbb{R}^n and set $\Omega_U = U \cap \Omega$, $\Gamma_U = U \cap \Gamma$ and $\Sigma := \Omega_U \cup \Gamma_U = U \cap \bar{\Omega}$.

2.1. Propositions without the coercivity assumption

In this subsection we *do not assume* the coercivity of H . That is, in this subsection we assume only (A1) and (A3). Let $f \in C(\Sigma)$. We consider the HJ equation:

$$\begin{cases} H(x, Du) = f(x) & \text{in } \Omega_U, \\ D_\gamma u(x) = g(x) & \text{on } \Gamma_U, \end{cases} \quad (7)$$

and establish the following theorems.

Theorem 2.1. *Let $\mathcal{S} \subset \text{Lip}(\Sigma)$ be a nonempty family of subsolutions of (7). Set $u(x) = \inf\{v(x) \mid v \in \mathcal{S}\}$ for $x \in \Sigma$ and assume that $u \in C(\Sigma)$. Then u is a subsolution of (7).*

Theorem 2.2. *For $k \in \mathbb{N}$ let $f_k \in C(\Sigma)$ and let $u_k \in \text{Lip}(\Sigma)$ be a subsolution of (7), with f_k in place of f , and $\{\lambda_k\}_{k \in \mathbb{N}}$ a sequence of nonnegative numbers such that $\sum_{k \in \mathbb{N}} \lambda_k = 1$. Assume that the sequences $\{u_k\}_{k \in \mathbb{N}}$ and $\{f_k\}_{k \in \mathbb{N}}$ are uniformly bounded on compact subsets of Σ . Set $u(x) = \sum_{k \in \mathbb{N}} \lambda_k u_k(x)$ and $f(x) = \sum_{k \in \mathbb{N}} \lambda_k f_k(x)$ for $x \in \Sigma$. Then u is a subsolution of (7).*

Before going into the proof of the above two theorems, we give two remarks. (i) If V is an open subset of \mathbb{R}^n satisfying $V \cap \bar{\Omega} \subset U$ and u is a subsolution (resp., a supersolution) of (7), then u is a subsolution (resp., a supersolution) of (7), with V in place of U . (ii) If U_α are open subsets of \mathbb{R}^n for $\alpha \in \Lambda$, where Λ is an index set, and the inclusion,

$$\bar{\Omega} \subset \bigcup_{\alpha \in \Lambda} U_\alpha,$$

holds and $u : \bar{\Omega} \rightarrow \mathbb{R}$ is a subsolution of (7), with $U := U_\alpha$, for any $\alpha \in \Lambda$, then u is a subsolution (resp., a supersolution) of (7), with Ω and Γ in place of Ω_U and Γ_U .

In the rest of this subsection we are devoted to proving Theorems 2.1 and 2.2. It is well known (see for instance [25,21]) that, if $\Gamma_U = \emptyset$, the assertions of Theorems 2.1 and 2.2 are valid. Thus, in order to prove the above two theorems, because of their local property together with the C^1 regularity of Ω , we may assume by use of a C^1 change of variables that for some constant $r > 0$,

$$\begin{aligned} U &= \text{int } B(0, r), & \Omega_U &= \{(x', x_n) \in U \mid x_n < 0\}, \\ \Gamma_U &= \{x = (x', x_n) \in U \mid x_n = 0\}. \end{aligned} \quad (8)$$

Here and later, for $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, we put $x' = (x_1, \dots, x_{n-1})$ and $x = (x', x_n)$.

We set $\mathbb{R}_+^n = \mathbb{R}^{n-1} \times (0, \infty)$ and define the function $\zeta \in C^\infty(\mathbb{R}_+^n \times \mathbb{R}^n)$ by:

$$\zeta(y, z) = \frac{1}{2} \left| z - \frac{z \cdot e_n}{y \cdot e_n} y \right|^2 + \frac{1}{2} (z \cdot e_n)^2.$$

We write $D_{z'}$ for the gradient operator with respect to the variables $z' = (z_1, \dots, z_{n-1})$. For instance, we write $D_{z'} \zeta = (\zeta_{z_1}, \dots, \zeta_{z_{n-1}})$.

Lemma 2.3. *The function $\zeta \in C^\infty(\mathbb{R}_+^n \times \mathbb{R}^n)$ has the properties:*

$$\begin{cases} \zeta(\xi, tz) = t^2 \zeta(\xi, z) & \text{for } (\xi, z, t) \in \mathbb{R}_+^n \times \mathbb{R}^n \times \mathbb{R}, \\ \zeta(\xi, z) > 0 & \text{for } (\xi, z) \in \mathbb{R}_+^n \times (\mathbb{R}^n \setminus \{0\}), \\ \xi \cdot D_z \zeta(\xi, z) = \xi_n z_n & \text{for } (\xi, z) \in \mathbb{R}_+^n \times \mathbb{R}^n. \end{cases}$$

Proof. We observe that

$$D_z \zeta(\xi, z) = z - \frac{z_n}{\xi_n} \xi - \frac{z \cdot \xi}{\xi_n} e_n + \frac{|\xi|^2 z_n}{\xi_n^2} e_n + z_n e_n.$$

and

$$\xi \cdot D_z \zeta(\xi, z) = \xi_n z_n.$$

It is now obvious that the function ζ has all the required properties. \square

We note by the homogeneity of the functions $\zeta(\xi, \cdot)$ that

$$\begin{aligned} C_0^{-1}|z|^2 &\leq \zeta(\xi, z) \leq C_0|z|^2, \\ |D_\xi \zeta(\xi, z)| &\leq C_0|z|^2 \quad \text{and} \quad |D_z \zeta(\xi, z)| \leq C_0|z| \end{aligned} \tag{9}$$

for all $(\xi, z) \in \mathbb{R}_+^n \times \mathbb{R}^n$ and for some constant $1 < C_0 < \infty$.

By assumption (A3) and (8), we have $\inf_{x \in \Gamma_U} \gamma(x) \cdot e_n > 0$. We restrict the domain of definition of γ to Γ_U and then extend that of the resulting vector field to \mathbb{R}^n so that $\gamma \in \text{BUC}(\mathbb{R}^n, \mathbb{R}^n)$ and $\gamma_0^{-1} \leq \gamma \cdot e_n \leq |\gamma| \leq \gamma_0$ on \mathbb{R}^n for some constant $\gamma_0 > 1$. Let ω be the modulus of continuity of γ .

By mollification, we may choose a family $\{\gamma^\delta\}_{\delta \in (0,1)}$ of C^∞ vector fields on \mathbb{R}^n so that $|\gamma(x) - \gamma^\delta(x)| \leq \omega(\delta)$, $|\gamma^\delta(x) - \gamma^\delta(y)| \leq \omega(|x - y|)$ and $|D\gamma^\delta(x)| \leq C_1\omega(\delta)/\delta$ for all $x, y \in \mathbb{R}^n$ and $\delta \in (0, 1)$ and for some constant $C_1 > 1$. Here $|A| := \max\{|A\xi| \mid \xi \in \mathbb{R}^n, |\xi| \leq 1\}$ for $n \times n$ real matrix A . We may also assume that $\gamma_0^{-1} \leq \gamma^\delta \cdot e_n \leq |\gamma^\delta| \leq \gamma_0$ on \mathbb{R}^n .

For $\delta \in (0, 1)$ we set $\psi^\delta(x, y) = \zeta(\gamma^\delta(x), x - y)$ and note that

$$\begin{aligned} D_x \psi^\delta(x, y) &= (D\gamma^\delta(x))^T D_\xi \zeta(\gamma^\delta(x), x - y) + D_z \zeta(\gamma^\delta(x), x - y), \\ D_y \psi^\delta(x, y) &= -D_z \zeta(\gamma^\delta(x), x - y), \end{aligned}$$

where A^T denotes the transposed matrix of the matrix A . From these we get:

$$\begin{aligned} |D_x \psi^\delta(x, y) + D_y \psi^\delta(x, y)| &= |(D\gamma^\delta(x))^T D_\xi \zeta(\gamma^\delta(x), x - y)| \\ &\leq \frac{C_0 C_1 \omega(\delta) |x - y|^2}{\delta}. \end{aligned} \tag{10}$$

Given a bounded function u on $\bar{\Sigma}$, for $\delta > 0$ let $u^\delta \in C(\mathbb{R}^n)$ denote the sup-convolution of u with kernel function $\delta^{-1}\psi^\delta$, i.e.,

$$u^\delta(x) = \sup_{y \in \bar{\Sigma}} \left(u(y) - \frac{1}{\delta} \psi^\delta(x, y) \right).$$

For $s \in (0, r]$ we set:

$$\begin{cases} \Omega_s = \{x = (x_1, \dots, x_n) \in \text{int } B(0, s) \mid x_n < 0\}, \\ \Gamma_s = \{x = (x_1, \dots, x_n) \in \text{int } B(0, s) \mid x_n = 0\}. \end{cases} \tag{11}$$

In particular, we have $\Omega_U = \Omega_r$, $\Gamma_U = \Gamma_r$, $\Sigma = \Omega_r \cup \Gamma_r$ and $\bar{\Sigma} = \bar{\Omega}_r$.

Lemma 2.4. *Let $\mu > 0$ and $0 < \varepsilon < r$. Let $u \in \text{Lip}(\Sigma)$ be a viscosity subsolution of (7), with $f := 0$ and $g := -\mu$. Then there is a constant $\delta_0 > 0$, independent of u , such that if $0 < \delta < \delta_0$, then $v := u^\delta$ is a viscosity subsolution of,*

$$H(x, Dv(x)) = \varepsilon \quad \text{in } \Omega_{r-\varepsilon}. \tag{12}$$

Moreover, if $0 < \delta < \delta_0$, then

$$D_\gamma^+ u^\delta(x) \leq \varepsilon \quad \text{for } x \in \Gamma_{r-\varepsilon}, \tag{13}$$

where

$$D_\gamma^+ v(x) := \limsup_{t \rightarrow 0^+} \frac{v(x) - v(x - t\gamma(x))}{t}.$$

Proof. Let $0 < \delta < 1$. Let $R > 0$ be a Lipschitz constant of u . We may assume by extending by continuity that $u \in \text{Lip}(\bar{\Sigma})$, so that for each $x \in \mathbb{R}^n$ there is a point $y \in \bar{\Sigma}$ such that

$$u^\delta(x) = u(y) - \frac{1}{\delta} \psi^\delta(x, y). \tag{14}$$

Fix $x \in \Omega_{r-\varepsilon} \cup \Gamma_{r-\varepsilon}$ and $y \in \bar{\Sigma}$ so that (14) holds. We collect here some basic estimates. As is standard, we have $u^\delta(x) \geq u(x)$, and

$$\frac{1}{\delta} \psi^\delta(x, y) = u(y) - u^\delta(x) \leq u(y) - u(x) \leq R|x - y|.$$

Noting by (9) that $\psi^\delta(x, y) \geq C_0^{-1}|x - y|^2$, we get:

$$|x - y| \leq C_2\delta, \tag{15}$$

where $C_2 := C_0R$. It follows from (10) that

$$|D_x \psi^\delta(x, y) + D_y \psi^\delta(x, y)| \leq C_3\omega(\delta)\delta, \tag{16}$$

where $C_3 := C_0C_1C_2^2$. By Lemma 2.3, we get:

$$\gamma^\delta(x) \cdot D_y \psi^\delta(x, y) = -\gamma_n^\delta(x)(x_n - y_n). \tag{17}$$

Also, we get:

$$|D_y \psi^\delta(x, y)| \leq C_0|x - y| \leq C_4\delta, \tag{18}$$

$$|D_x \psi^\delta(x, y)| \leq |D_y \psi^\delta(x, y)| + |D_x \psi^\delta(x, y) + D_y \psi^\delta(x, y)| \leq C_4\delta, \tag{19}$$

where $C_4 := C_0C_2 + C_3\omega(1)$.

We now show that u^δ is a subsolution of (12) if $\delta > 0$ is sufficiently small. Let $\phi \in C^1(\bar{\Omega}_{r-\varepsilon})$ and $x \in \Omega_{r-\varepsilon}$. We assume that $u^\delta - \phi$ attains a strict maximum at x , and choose a point $y \in \bar{\Sigma} = \bar{\Omega}_r$ so that (14) holds. We choose a constant $\delta_1 \in (0, 1)$ so that $C_2\delta_1 < \varepsilon$ and assume in what follows that $0 < \delta < \delta_1$. By (15), we have $|x - y| < \varepsilon$. Hence, we have $\partial\Omega_r \setminus \Gamma_r$. Since $y \in \bar{\Omega}_r$, we have two possibilities: $y \in \Omega_r$ or $y \in \Gamma_r$.

Now we treat the case where $y \in \Omega_r$. Then we have:

$$D\phi(x) \in D^+u^\delta(x),$$

$$D\phi(x) + \frac{1}{\delta}D_x\psi^\delta(x, y) = 0 \quad \text{and} \quad \frac{1}{\delta}D_y\psi^\delta(x, y) \in D^+u(y),$$

where $D^+h(x)$ denotes the superdifferential of the function h at x . Using this last inclusion, we get $H(y, D_y\psi^\delta(x, y)/\delta) \leq 0$. According to (18) and (19), we have $|D_y\psi^\delta(x, y)|/\delta \leq C_4$ and $|D\phi(x)| = |D_x\psi^\delta(x, y)|/\delta \leq C_4$. Let ω_H denote the modulus of continuity of the function H restricted to $\bar{\Omega} \times B(0, C_4)$. Using (16) and (15), we obtain:

$$0 \geq H\left(y, \frac{1}{\delta}D_y\psi^\delta(x, y)\right) \geq H(x, D\phi(x)) - \omega_H(|x - y|) - \omega_H(C_3\omega(\delta))$$

$$\geq H(x, D\phi(x)) - \omega_H(C_2\delta) - \omega_H(C_3\omega(\delta)).$$

We choose a $\delta_2 > 0$ so that

$$\omega_H(C_2\delta_2) + \omega_H(C_3\omega(\delta_2)) \leq \varepsilon.$$

Thus, if $y \in \Omega_r$ and $0 < \delta < \delta_1 \wedge \delta_2$, then we have,

$$H(x, D\phi(x)) \leq \varepsilon. \tag{20}$$

Next, we turn to the case where $y \in \Gamma_r$. Then we have,

$$D\phi(x) = -\frac{1}{\delta}D_x\psi^\delta(x, y) \in D^+u^\delta(x) \quad \text{and} \quad \frac{1}{\delta}D_y\psi^\delta(x, y) \in D_\Sigma^+u(y),$$

where $D_\Sigma^+u(y)$ denotes the set of those $p \in \mathbb{R}^n$ for which

$$u(y + \xi) \leq u(y) + p \cdot \xi + o(|\xi|) \quad \text{as } y + \xi \in \Sigma \quad \text{and} \quad \xi \rightarrow 0.$$

By (17), we get:

$$\gamma^\delta(x) \cdot D_y \psi^\delta(x, y) = -\gamma_n(x)(x_n - y_n) = -\gamma_n(x)x_n > 0.$$

Since $|D_y \psi^\delta(x, y)|/\delta \leq C_4$ by (18), we get:

$$\begin{aligned} \gamma(y) \cdot \frac{1}{\delta} D_y \psi^\delta(x, y) &= \gamma^\delta(x) \cdot \frac{1}{\delta} D_y \psi^\delta(x, y) + (\gamma(y) - \gamma^\delta(x)) \cdot \frac{1}{\delta} D_y \psi^\delta(x, y) \\ &> -C_4(\omega(|x - y|) + \omega(\delta)) \geq -C_4(\omega(C_2\delta) + \omega(\delta)). \end{aligned}$$

We select a $\delta_3 > 0$ so that $C_4(\omega(C_2\delta_3) + \omega(\delta_3)) < \mu$, and assume in the following that $0 < \delta < \delta_1 \wedge \delta_3$. Accordingly, we have $(1/\delta)\gamma(y) \cdot D_y \psi^\delta(x, y) > -\mu$. Since u is a viscosity subsolution of (7), with $f := 0$ and $g := -\mu$, we get $H(y, D_y \psi^\delta(x, y)/\delta) \leq 0$. Now, as in the previous case, we obtain:

$$0 \geq H(x, D\phi(x)) - \omega_H(C_2\delta) - \omega_H(C_3\delta).$$

Consequently, if $y \in \partial\Omega_r$ and $0 < \delta < \delta_1 \wedge \delta_2 \wedge \delta_3$, then we have (20). Thus we see that if $0 < \delta < \delta_1 \wedge \delta_2 \wedge \delta_3$, then u^δ is a subsolution of (12).

We now prove that (13) is valid if δ is sufficiently small. Let $x \in \Gamma_{r-\varepsilon}$, and choose a $y \in \bar{\Sigma}$ so that (14) holds. Then, for $t > 0$ sufficiently small, we have:

$$u^\delta(x) - u^\delta(x - t\gamma(x)) \leq -\frac{1}{\delta}(\psi^\delta(x, y) - \psi^\delta(x - t\gamma(x), y)).$$

Hence,

$$D_\gamma^+ u^\delta(x) \leq -\gamma(x) \cdot \frac{1}{\delta} D_x \psi^\delta(x, y). \tag{21}$$

Using (18), (16) and (17), we compute that

$$\begin{aligned} -\gamma(x) \cdot \frac{1}{\delta} D_x \psi^\delta(x, y) &\leq -\gamma^\delta(x) \cdot \frac{1}{\delta} D_x \psi^\delta(x, y) + C_4\omega(\delta) \\ &\leq \gamma^\delta(x) \cdot \frac{1}{\delta} D_y \psi^\delta(x, y) + \frac{\gamma_0}{\delta} |D_x \psi^\delta(x, y) + D_y \psi^\delta(x, y)| + C_4\omega(\delta) \\ &\leq \gamma_0 C_3\omega(\delta) + C_4\omega(\delta). \end{aligned} \tag{22}$$

We select a $\delta_4 > 0$ so that $(\gamma_0 C_3 + C_4)\omega(\delta_4) < \varepsilon$. From (21) and (22), we find that if $0 < \delta < \delta_4$, then (13) holds.

Finally, setting $\delta_0 = \delta_1 \wedge \delta_2 \wedge \delta_3 \wedge \delta_4$, we conclude that if $0 < \delta < \delta_0$, then u^δ is a subsolution of (12) and satisfies (13). \square

Lemma 2.5. *Let $\mu > 0$. Let $u, v \in \text{Lip}(\Sigma)$ be subsolutions of (7), with $f := 0$ and $g := -\mu$. Then $u \wedge v$ is a subsolution of (7), with $f = g = 0$.*

Proof. Fix any $\varepsilon \in (0, r)$. In view of Lemma 2.4, there is a constant $\delta_0 > 0$ such that if $0 < \delta < \delta_0$, then $u := u^\delta, v^\delta$ are solutions of $H(x, Du) \leq \varepsilon$ in the viscosity sense in $\Omega_{r-\varepsilon}$ and satisfy $D_\gamma^+ u \leq \varepsilon$ on $\Gamma_{r-\varepsilon}$. As is well known, since $H(x, \cdot)$ is convex, the function $z^\delta := u^\delta \wedge v^\delta$ is a subsolution of $H(x, Dz^\delta) \leq \varepsilon$ in $\Omega_{r-\varepsilon}$. Also, it is easy to see that $D_\gamma^+ z^\delta(x) \leq \varepsilon$ for $x \in \Gamma_{r-\varepsilon}$. It is then easily checked that z^δ is a subsolution of (7), with $\Omega_U := \Omega_{r-\varepsilon}, \Gamma_U := \Gamma_{r-\varepsilon}, f(x) := \varepsilon$ and $g(x) := \varepsilon$. Sending $\delta \rightarrow 0$ and setting $z := u \wedge v$, we see by the stability of the viscosity property under uniform convergence that z is a viscosity subsolution of (7), with $\Omega_U := \Omega_{r-\varepsilon}, \Gamma_U := \Gamma_{r-\varepsilon}, f(x) := \varepsilon$ and $g(x) := \varepsilon$. But, since $\varepsilon \in (0, r)$ is arbitrary, the function z is a viscosity subsolution of (7), with $f := 0$ and $g := 0$. \square

Noting that for any $u, v \in C(\Sigma), 0 < \lambda < 1$ and $x \in \Gamma_U$,

$$D_\gamma^+(\lambda u + (1 - \lambda)v)(x) \leq \lambda D_\gamma^+ u(x) + (1 - \lambda) D_\gamma^+ v(x),$$

we deduce that the argument of the above proof yields also the following lemma.

Lemma 2.6. Let $\mu > 0$ and $f_1, f_2 \in C(\bar{\Sigma})$. For $i = 1, 2$ let $u_i \in C(\bar{\Sigma})$ be a subsolution of (7), with $f := f_i$ and $g := -\mu$. Let $0 < \lambda < 1$ and set $u = \lambda u_1 + (1 - \lambda)u_2$ and $f = \lambda f_1 + (1 - \lambda)f_2$. Then u is a subsolution of (7), with $g := 0$.

Proof of Theorem 2.1. By the continuity of the function u , we may assume that \mathcal{S} is a sequence $\{u_k\}_{k \in \mathbb{N}}$. Indeed, we can choose a sequence $\{K_m\}_{m \in \mathbb{N}}$ of compact subsets of Σ such that $\Sigma = \bigcup_{m \in \mathbb{N}} K_m$. By a compactness argument, we can choose for each $m \in \mathbb{N}$ a sequence $\{v_{m,j}\}_{j \in \mathbb{N}} \subset \mathcal{S}$ such that $u(x) = \inf\{v_{m,j}(x) \mid j \in \mathbb{N}\}$ for $x \in K_m$. Then we have $u(x) = \inf\{v_{m,j}(x) \mid j, m \in \mathbb{N}\}$ for $x \in \Sigma$. Relabeling $\{v_{m,j}\}$ appropriately, we find a sequence $\{u_k\}$ which replaces \mathcal{S} in the following argument.

Next, we fix any $\mu > 0$. According to the C^1 regularity of Ω and the continuity of g , we may select $\psi_\mu \in C^1(\bar{\Omega})$ so that $g(x) + \mu \leq D_\gamma \psi_\mu(x) \leq g(x) + 2\mu$ for $x \in \Gamma$. Set $v_k(x) = u_k(x) - \psi_\mu(x)$ and $v(x) = u(x) - \psi_\mu(x)$ for $x \in \Sigma$ and observe that $w := v_k$ is a solution of,

$$\begin{cases} \tilde{H}(x, Dw) \leq 0 & \text{in } \Omega_U, \\ D_\gamma w(x) \leq -\mu & \text{on } \Gamma_U, \end{cases} \tag{23}$$

where \tilde{H} is the continuous function on $\bar{\Omega} \times \mathbb{R}^n$ given by $\tilde{H}(x, p) = H(x, p + D\psi_\mu(x)) - f(x)$. By Lemma 2.5, we see that $w_k := v_1 \wedge \dots \wedge v_k$ is a solution of (23), with μ replaced by 0. Since $w_k(x) \rightarrow v(x)$ locally uniformly on Σ as $k \rightarrow \infty$, by the stability of the viscosity property under uniform convergence, we see that v is a solution of (23), with $\mu := 0$. This means that u is a subsolution of (7), with $g(x)$ replaced by $g(x) + 2\mu$. Since $\mu > 0$ is arbitrary, we conclude that u is a subsolution of (7). \square

Proof of Theorem 2.2. Since the property to be shown is local, by replacing U by a smaller one, we may assume that the sequences $\{u_k\}$ and $\{f_k\}$ are uniformly bounded on Σ . We set for $x \in \Sigma$,

$$v_k(x) = \frac{1}{\sum_{j=1}^k \lambda_j} \sum_{j=1}^k \lambda_j u_j(x) \quad \text{and} \quad F_k(x) = \frac{1}{\sum_{j=1}^k \lambda_j} \sum_{j=1}^k \lambda_j f_j(x).$$

Assume that k is sufficiently large, so that $\sum_{j=1}^k \lambda_j > 0$, $v_k \in \text{Lip}(\Sigma)$ and $F_k \in C(\Sigma)$. Moreover, using Lemma 2.6 and arguing as in the previous proof that v_k is a subsolution of (7), with f replaced by F_k . In view of the uniform boundedness of the sequences $\{u_k\}$ and $\{f_k\}$, we see that $v_k(x) \rightarrow u(x)$ and $F_k(x) \rightarrow f(x)$ uniformly on Σ as $k \rightarrow \infty$. By the stability of the viscosity property, we conclude that u is a subsolution of (7). \square

2.2. Propositions under the coercivity assumption

In this subsection, we always assume that (A1)–(A3) hold, and reformulate Theorems 2.1 and 2.2.

Theorem 2.7. Let $\mathcal{S} \subset C(\Sigma)$ be a nonempty subset of subsolutions of (7). Assume that $\inf\{v(x) \mid v \in \mathcal{S}\} > -\infty$ for some $x \in \Sigma$. Then the function,

$$u(x) := \inf\{v(x) \mid v \in \mathcal{S}\}, \tag{24}$$

on Σ is a subsolution of (7).

A consequence of the above theorem is stated as follows. If $\mathcal{S} \subset C(\Sigma)$ is a nonempty subset of solutions of (7) and formula (24) defines a real-valued function u , then u is a solution of (7). Indeed, as is well known, the supersolution property is stable under taking infimums, and therefore u is a supersolution of (7) as well.

Proof. Because of the local nature of our assertion, by replacing U by a smaller one, we may assume that f is bounded on Σ . Then, by the coercivity assumption (A2), we can choose a constant $C > 0$ so that for $(x, p) \in \bar{\Omega} \times \mathbb{R}^n$, if $H(x, p) \leq f(x)$, then $|p| \leq C$. This together with the boundedness and C^1 regularity of Ω implies that \mathcal{S} is equi-Lipschitz continuous on Σ . Consequently, we have $u \in \text{Lip}(\Sigma)$. Applying Theorem 2.1, we find that u is a subsolution of (7). \square

We consider next the evolution equation with the Neumann type boundary condition:

$$\begin{cases} u_t + H(x, Du) = f(x, t) & \text{in } \Omega_U \times (0, T), \\ D_\gamma u = g(x) & \text{on } \Gamma_U \times (0, T), \end{cases} \tag{25}$$

where $f \in C(\Sigma \times (0, T))$.

Theorem 2.8. *Let $\mathcal{S} \subset C(\Sigma \times (0, T))$ be a nonempty subset of subsolutions of (25). Assume that \mathcal{S} is uniformly bounded on compact subsets of $\Sigma \times (0, T)$. Then the function,*

$$u(x, t) := \inf\{v(x, t) \mid v \in \mathcal{S}\}, \tag{26}$$

on $\Sigma \times (0, T)$ is a subsolution of (25).

A remark parallel to the remark after Theorem 2.7 is valid here. Indeed, if $\mathcal{S} \subset C(\Sigma \times (0, T))$ is a nonempty subset of solutions of (25) and it is uniformly bounded on compact subsets of $\Sigma \times (0, T)$, then the function u given by (26) is a solution of (25).

Proof. Because the viscosity property is local, we may assume, by replacing U and the interval $(0, T)$ by smaller ones and by translation in the t -direction if needed, that \mathcal{S} are uniformly bounded on $\Sigma \times (0, T)$. We may also assume that $f \in \text{BUC}(\Sigma)$. Let $C > 0$ be a constant such that $|v(x, t)| \leq C$ for $(x, t) \in \Sigma \times (0, T)$ and $v \in \mathcal{S}$.

Let $\varepsilon > 0$ and introduce the sup-convolution of $v \in \mathcal{S}$ with respect to the t -variable:

$$v^\varepsilon(x, t) = \inf_{0 < s < T} \left(v(x, s) - \frac{1}{2\varepsilon}(t - s)^2 \right) \quad \text{for } (x, t) \in \Sigma \times \mathbb{R}.$$

Setting $\delta = 2\sqrt{\varepsilon C}$, we observe that for $(x, t) \in \Sigma \times (\delta, T - \delta)$,

$$v^\varepsilon(x, t) = \max_{|s-t| \leq \delta} \left(u(x, s) - \frac{1}{2\varepsilon}(t - s)^2 \right),$$

from which we deduce as usual in viscosity solutions theory that v^ε is a subsolution of,

$$\begin{cases} v_t^\varepsilon + H(x, Dv^\varepsilon) = f + \omega(\delta) & \text{in } \Omega_U \times (\delta, T - \delta), \\ D_\gamma v^\varepsilon = g & \text{on } \Gamma_U \times (\delta, T - \delta), \end{cases} \tag{27}$$

where ω is the modulus of continuity of f .

Now, the family of functions $v^\varepsilon(x, \cdot)$, with $x \in \Sigma$ and $v \in \mathcal{S}$, is equi-Lipschitz continuous on $(\delta, T - \delta)$. From this and (27), we see that $H(x, Dv^\varepsilon) \leq C_\varepsilon$ in the viscosity sense in $\Omega_U \times (\delta, T - \delta)$ for all $v \in \mathcal{S}$ and for some constant $C_\varepsilon > 0$. Observe then that for $(x, t) \in \Sigma \times \mathbb{R}$,

$$u^\varepsilon(x, t) := \inf_{0 < s < T} \left(u(x, s) - \frac{1}{2\varepsilon}(t - s)^2 \right) = \inf\{v^\varepsilon(x, t) \mid v \in \mathcal{S}\}.$$

We apply Theorem 2.1, to see that u^ε is a subsolution of (27). Indeed, in order to apply Theorem 2.1, we set $\tilde{\Omega} = \Omega \times (0, T)$, $\tilde{U} = U \times (0, T)$, $\tilde{H}(x, t, p, q) = H(x, p) + q$ and $\tilde{\gamma}(x, t) = (\gamma(x), 0)$, and regard problem (25) as problem (7), with $\tilde{\Omega}$, \tilde{U} , \tilde{H} and $\tilde{\gamma}$ in place of Ω , U , H and γ , respectively.

Next, we observe that for $(x, t) \in \Sigma \times (0, T)$, the family $\{u^\varepsilon(x, t)\}$ converges monotonically to $u(x, t)$ as $\varepsilon \rightarrow 0$, which implies, together with the continuity of u^ε , that $u(x, t)$ is identical to the upper relaxed limit of $u^\varepsilon(x, t)$ as $\varepsilon \rightarrow 0$. Because of the stability of the subsolution property under such a limiting process, we see that u is a subsolution of (25). \square

Theorem 2.9. *For $k \in \mathbb{N}$ let $f_k \in C(\Sigma \times (0, T))$ and $u_k \in \text{USC}(\Sigma \times (0, T))$ be a subsolution of (25), with f_k in place of f . Let $\{\lambda_k\}_{k \in \mathbb{N}}$ be a sequence of nonnegative numbers such that $\sum_{k \in \mathbb{N}} \lambda_k = 1$. Assume that the sequences $\{u_k\}_{k \in \mathbb{N}}$ and $\{f_k\}_{k \in \mathbb{N}}$ are uniformly bounded on compact subsets of $\Sigma \times (0, T)$. Set $u(x, t) = \sum_{k \in \mathbb{N}} \lambda_k u_k(x, t)$ and $f(x, t) = \sum_{k \in \mathbb{N}} \lambda_k f_k(x, t)$ for $(x, t) \in \Sigma \times (0, T)$. Then u is a subsolution of (25).*

Proof. Arguing as in the proof of Theorem 2.8, with use of Theorem 2.1 instead of Theorem 2.2, we conclude that Theorem 2.9 is valid. \square

3. Comparison results

The comparison results presented in this section are more or less well known (see for instance [1,3,4]). The only new feature of our results may be in the point that they are formulated in a localized fashion.

Theorem 3.1. *Let $f_1, f_2 \in C(\Sigma)$ and let $u \in \text{USC}(\overline{\Sigma})$ (resp., $v \in \text{LSC}(\overline{\Sigma})$) be a subsolution (resp., a supersolution) of (7), with f replaced by f_1 (resp., f_2). Assume that $f_1(x) < f_2(x)$ for $x \in \Sigma$. Then $\sup_{\Sigma}(u - v) \leq \sup_{\partial U \cap \overline{\Omega}}(u - v)$.*

We remark that if $\partial U \cap \overline{\Omega} = \emptyset$ in the above theorem, then the right-hand side of the above inequality equals $-\infty$ by definition. In particular, if $\overline{\Omega} \subset U$ in the above theorem, then the theorem asserts that $\sup_{\Omega}(u - v) = -\infty$.

Corollary 3.2. *If $a < b$ and problem (1), (2) has a subsolution, then problem (1), (2), with b in place of a , does not have a supersolution. In particular, if problem (1), (2) has a solution for some $a \in \mathbb{R}$, then problem (1), (2), with a replaced by $b \neq a$, has no solution.*

Proof. Let $a < b$, and assume that there are a subsolution u of (1), (2) and a supersolution of (1), (2), with b in place of a . Note that, for any $c \in \mathbb{R}$, the function $u + c$ is also a subsolution of (1), (2). By Theorem 3.1, we have $u^* + c \leq v_*$ on $\overline{\Omega}$ for $c \in \mathbb{R}$, which is a contradiction. This proves our claim. \square

Lemma 3.3. *Assume that f is bounded on Σ . Then there is a constant $C > 0$, depending only H, f and Ω , such that for any subsolution $u \in \text{USC}(\Sigma)$ of (7) and $x, y \in \Sigma$, $|u(x) - u(y)| \leq C|x - y|$.*

Proof. Let $u \in \text{USC}(\Sigma)$ be a subsolution of (7). By the coercivity assumption (A2) and the boundedness of f , there is a constant $C_0 > 0$ such that for $(x, p) \in \Omega_U$, if $|p| \geq C_0$, then $H(x, p) \geq f(x) + 1$. It follows from (7) that u is a subsolution of $|Du| \leq C_0$ in Ω_U , which implies together with the C^1 regularity of Ω that u is Lipschitz continuous on Ω_U with a Lipschitz constant $C > 0$ depending only on C_0 and Ω .

We next show that $u \in C(\Sigma)$, which guarantees that u is Lipschitz continuous on Σ with the same Lipschitz constant C . To this end, we need only to show that for any fixed $z \in \Gamma_U$, u is continuous at z . By translation, we may assume that $z = 0$. By rotation and localization, we may furthermore assume that U, Ω_U and Γ_U are given by (8). Since $u \in \text{USC}(\Sigma)$ and $u \in \text{Lip}(\Omega_U)$, it is enough to show that

$$u(0) \leq \sup_{\Omega_s} u \quad \text{for } s \in (0, r). \tag{28}$$

Here and later we use the notation Ω_s and Γ_s as defined in (11).


We may assume by replacing $r > 0$ by a smaller one that $\gamma_0 := \inf_{x \in \Gamma_r} \gamma(x) \cdot e_n > 0$. (Recall that e_n denotes the unit vector $(0, \dots, 0, 1) \in \mathbb{R}^n$.) We select a closed convex cone K with vertex at the origin so that $K \setminus \{0\} \subset -\mathbb{R}_+^n$ and $-\gamma(x) + B(0, \delta) \in K$ for all $x \in \Gamma_r$ and for some $\delta > 0$. We denote by N_K the normal cone to K at the origin. That is, we set $N_K = \{\xi \in \mathbb{R}^n \mid \xi \cdot p \leq 0 \text{ for } p \in K\}$. It follows that $\xi \cdot (-\gamma(x)) \leq -\delta|\xi|$ for all $\xi \in N_K$ and $x \in \Gamma_r$. Let d_K denote the distance function from the set K , i.e., $d_K(x) = \text{dist}(x, K)$. As is well known, the function d_K is convex on \mathbb{R}^n , $d_K \in C(\mathbb{R}^n) \cap C^1(\mathbb{R}^n \setminus K)$, $d_K(x) \geq 0$ for $x \in \mathbb{R}^n$ and $Dd_K(x) \in N_K \cap \partial B(0, 1)$ for $x \in \mathbb{R}^n \setminus K$.

Fix any $s \in (0, r)$ and set $\rho = \text{dist}(K, \partial B(0, s) \cap \{x_n = 0\})$. Here and later we use the notation: $\{x_n = 0\} := \{(x', x_n) \in \mathbb{R}^n \mid x_n = 0\}$ and similarly $\{x_n < 0\} := \{(x', x_n) \in \mathbb{R}^n \mid x_n < 0\}$. Note that $0 < \rho \leq s$ and fix any $\varepsilon \in (0, \rho)$. We may assume by replacing $r > 0$ by a smaller one that u is bounded above on $\overline{\Omega}_r$. We choose a constant $C_1 > 0$ so that $\sup_{\overline{\Omega}_r} u \leq C_1$, $\sup_{\Omega_r} |u| \leq C_1$ and $\sup_{\Gamma_r} g \leq C_1$. We select a function $\zeta_\varepsilon \in C^1(\mathbb{R})$ so that $\zeta'_\varepsilon(r) \geq 1$ for $r \in \mathbb{R}$, $\zeta_\varepsilon(r) = r$ for $r \leq \varepsilon$ and $\zeta_\varepsilon(\rho) \geq 2C_1$. We set $A = \max\{1, C_0, (C_1 + 1)/\delta\}$, and define the function $v \in C(\mathbb{R}^n)$ by

$$v(x) := A\zeta_\varepsilon(d_K(x + \varepsilon e_n)) + \sup_{\Omega_s} u = A\zeta_\varepsilon(\text{dist}(x, K - \varepsilon e_n)) + \sup_{\Omega_s} u.$$

Let $V = \Omega_s \setminus (K - \varepsilon e_n)$. We intend to show that $u \leq v$ on the set \overline{V} . To do this, we suppose by contradiction that $\max_{\overline{V}}(u - v) > 0$. Note that

$$\overline{V} \subset \overline{\Omega}_s = (\overline{\Omega}_s \cap \{x_n < 0\}) \cup (\partial B(0, s) \cap \{x_n = 0\}) \cup \Gamma_s.$$



Since $u \in C(\Omega_r)$, it is clear that $u \leq \sup_{\Omega_s} u \leq v$ on $\bar{\Omega}_s \cap \{x_n < 0\}$. For any $x \in \partial B(0, s) \cap \{x_n = 0\}$, we have $\text{dist}(x, K - \epsilon e_n) \geq \text{dist}(x, K) \geq \rho > s$, and hence

$$v(x) \geq \zeta_\epsilon (d_K(x + \epsilon e_n)) - C_1 \geq \zeta_\epsilon(\rho) - C_1 \geq C_1 \geq u(x).$$

Consequently, we have $u(x) \leq v(x)$ for $\bar{V} \cap \Gamma_s$ and therefore there is a point $y \in \Gamma_s$ such that $(u - v)(y) = \max_{\bar{V}}(u - v)$. Since u is a subsolution of (7), with V in place of U , we have either $H(y, Dv(y)) \leq f(y)$ or $D_\gamma v(y) \leq g(y)$. Since $y \in \Gamma_s$ and $\Gamma_s \cap (K - \epsilon e_n) = \emptyset$, we have:

$$Dv(y) = A\zeta'_\epsilon(d_K(y + \epsilon e_n))Dd_K(y + \epsilon e_n).$$

Hence, we get $|Dv(y)| \geq A \geq C_0$ and, by the choice of C_0 , $H(y, Dv(y)) > f(y)$. Also, we get:

$$D_\gamma v(y) = A\zeta'_\epsilon(d_K(y + \epsilon e_n))\gamma(y) \cdot Dd_K(y + \epsilon e_n) \geq A\delta \geq C_1 + 1 > g(y).$$

We are in a contradiction, and thus we conclude that (28) holds. \square

Proof of Theorem 3.1. We first deal with the case where $\bar{\Omega} \cap \partial U \neq \emptyset$. We suppose by contradiction that

$$\max_{\bar{\Sigma}}(u - v) > \max_{\partial U \cap \bar{\Omega}}(u - v). \tag{29}$$

By replacing U by a smaller one (for instance, the set $\{x \in U \mid \text{dist}(x, \partial U) > \epsilon\}$ with sufficiently small $\epsilon > 0$) if needed, we may assume that f_1, f_2 are continuous on $\bar{\Sigma}$ and $\sup_{\bar{\Sigma}}(f_1 - f_2) < 0$. We note by Lemma 3.3 that the function u is Lipschitz continuous on Σ .

We now intend to replace H by a uniformly continuous Hamiltonian, which is not coercive nor convex any more. For this, we define the function $\tilde{H} \in \text{UC}(\Sigma \times \mathbb{R}^n)$ by $\tilde{H}(x, p) = \min\{H(x, p) - f_1(x), 1\}$. Set $\tilde{f}_1(x) = 0$ and $\tilde{f}_2(x) = \min\{f_2(x) - f_1(x), 1\}$ for $x \in \Sigma$. Now, the function u (resp., v) is a subsolution (resp., a supersolution) of (7), with \tilde{H} and \tilde{f}_1 (resp., \tilde{f}_2) in place of H and f . Thus, replacing H, f_1 and f_2 by \tilde{H}, \tilde{f}_1 and \tilde{f}_2 , respectively, we may assume in what follows that $H \in \text{UC}(\Sigma \times \mathbb{R}^n)$.

We select a function $\psi \in C^1(\bar{\Omega})$ so that $D_\gamma \psi(x) > 0$ for all $x \in \Gamma$. Let $\delta > 0$ and set $u_\delta(x) = u(x) - \delta\psi(x)$ and $v_\delta(x) = v(x) + \delta\psi(x)$ for $x \in \bar{\Sigma}$. In view of the uniform continuity of H , selecting $\delta > 0$ small enough, replacing f_1, f_2 by a new ones if necessary, we may assume that u_δ (resp., v_δ) is a subsolution (resp., a supersolution) of (7), with g and f replaced respectively by $g - \epsilon$ (resp., $g + \epsilon$), where ϵ is a positive constant and by f_1 (resp., f_2). We may also assume that (29) holds with u_δ and v_δ in place of u and v , respectively. Henceforth we replace u and v by u_δ and v_δ in our notation, respectively.

If $\sup_{\Gamma_U}(u - v) < \max_{\bar{\Sigma}}(u - v)$, then we have $\max_{\partial \Sigma}(u - v) < \max_{\bar{\Sigma}}(u - v)$ and get a contradiction by arguing as in the standard proof (in the case of the Dirichlet boundary condition) of comparison results where the Lipschitz continuity of u is available.

Thus we assume henceforth that $\sup_{\Gamma_U}(u - v) = \max_{\bar{\Sigma}}(u - v)$. Then the function $u - v$ attains a maximum at a point $z \in \Gamma_U$. By replacing U by an open ball $\text{int } B(z, r)$, with $r > 0$ sufficiently small, and by translation, we may assume that $z = 0$, $\Omega_U = \Omega_r$ and $\Gamma_U = \Gamma_r$, where Ω_r and Γ_r are the sets given by (11). We set $\tilde{\gamma} = \gamma(0)/|\gamma(0)|^2$, $\tilde{u}(x) = u(x) - g(0)\tilde{\gamma} \cdot x - |x|^2$ and $\tilde{v}(x) = v(x) - g(0)\tilde{\gamma} \cdot x$ for $\bar{\Sigma}$. Note that $\tilde{u} - \tilde{v}$ attains a strict maximum at the origin and that $w := \tilde{u}$ is a solution of,

$$\begin{cases} H(x, Dw(x) + g(0)\tilde{\gamma} + 2x) \leq f_1(x) & \text{in } \Omega_r, \\ D_\gamma Dw(x) \leq g(x) - g(0)\gamma(x) \cdot \tilde{\gamma} - 2\gamma(x) \cdot x - \epsilon & \text{on } \Gamma_r, \end{cases}$$

and $w := \tilde{v}$ is a solution of,

$$\begin{cases} H(x, Dw(x) + g(0)\tilde{\gamma}) \geq f_2(x) & \text{in } \Omega_r, \\ D_\gamma Dw(x) \geq g(x) - g(0)\gamma(x) \cdot \tilde{\gamma} + \varepsilon & \text{on } \Gamma_r. \end{cases}$$

Replacing $r > 0$ by a smaller positive number, we may assume that $w := \tilde{u}$ is a solution of,

$$\begin{cases} H(x, Dw(x) + g(0)\tilde{\gamma}) \leq f_1(x) + \varepsilon & \text{in } \Omega_r, \\ D_\gamma Dw(x) \leq -\frac{\varepsilon}{2} & \text{on } \Gamma_r, \end{cases}$$

and $w := \tilde{v}$ is a solution of,

$$\begin{cases} H(x, Dw(x) + g(0)\tilde{\gamma}) \geq f_2(x) & \text{in } \Omega_r, \\ D_\gamma Dw(x) \geq \frac{\varepsilon}{2} & \text{on } \Gamma_r. \end{cases}$$

Reselecting $\varepsilon > 0$ small enough if necessary, we may assume that $\max_{\overline{\Omega}_r} (f_1 + \varepsilon - f_2) < 0$. In the argument which follows, we write u, v, f_1 and H for the functions $\tilde{u}, \tilde{v}, f_1 + \varepsilon$ and $H(x, p + g(0)\tilde{\gamma})$, respectively.

Let $\zeta \in C^\infty(\mathbb{R}_+^n \times \mathbb{R}^n)$ be the function from Lemma 2.3. Set $\phi(x, y) = \zeta(\gamma(0), x - y)$. For $\alpha > 1$ we consider the function $\Phi(x, y) := u(x) - v(y) - \alpha\phi(x, y)$ on $\overline{\Sigma} \times \overline{\Sigma}$. Let $(x_\alpha, y_\alpha) \in \overline{\Sigma}^2$ be a maximum point of Φ . Since $u - v$ attains a strict maximum at the origin, we deduce easily that $x_\alpha, y_\alpha \rightarrow 0$ as $\alpha \rightarrow \infty$. Let $C_1 > 0$ be the Lipschitz constant of the function u . Then, since $\Phi(y_\alpha, y_\alpha) \leq \Phi(x_\alpha, y_\alpha)$, we find that $\alpha\phi(x_\alpha, y_\alpha) \leq C_1|x_\alpha - y_\alpha|$, from which we get $\alpha|x_\alpha - y_\alpha| \leq C_2$, where $C_2 > 0$ is a constant independent of α . If $x_\alpha, y_\alpha \in \Omega_r$, then we have:

$$H(x_\alpha, D_x\phi(x_\alpha, y_\alpha)) \leq f_1(x_\alpha) \quad \text{and} \quad H(y_\alpha, -D_y\phi(x_\alpha, y_\alpha)) \geq f_2(y_\alpha).$$

Here, noting that $D_x\phi(x, y) + D_y\phi(x, y) = 0$, we find that

$$H(x_\alpha, D_x\phi(x_\alpha, y_\alpha)) \leq f_1(x_\alpha) \quad \text{and} \quad H(y_\alpha, D_x\phi(x_\alpha, y_\alpha)) \geq f_2(y_\alpha). \tag{30}$$

Assume instead that $x_\alpha \in \Gamma_r$. By the viscosity property of u , we have either,

$$H(x_\alpha, D_x\phi(x_\alpha, y_\alpha)) \leq f_1(x_\alpha) \quad \text{or} \quad \gamma(x_\alpha) \cdot D_x\phi(x_\alpha, y_\alpha) \leq -\frac{\varepsilon}{2}.$$

Compute that

$$\begin{aligned} \gamma(x_\alpha) \cdot D_x\phi(x_\alpha, y_\alpha) &= \gamma(x_\alpha) \cdot D_z\zeta(\gamma(0), x_\alpha - y_\alpha) \\ &\geq \gamma_n(0) \cdot (-y_{\alpha n}) - C_2C_3\omega_\gamma(|x_\alpha|), \end{aligned}$$

where $C_3 > 0$ is a constant, independent of α , such that $|D_z\zeta(\gamma(0), z)| \leq C_3|z|$ for $z \in \mathbb{R}_+^n \times \mathbb{R}^n$, ω_γ is the modulus of continuity of γ on Γ and $y_{\alpha n} := e_n \cdot y$. Accordingly, if α is large enough, then we have $\gamma(x_\alpha) \cdot D_x\phi(x_\alpha, y_\alpha) > -\varepsilon/2$. Thus, we have $H(x_\alpha, D_x\phi(x_\alpha, y_\alpha)) \leq f_1(x_\alpha)$ if α is large enough. Similarly, in the case where $y_\alpha \in \Gamma_r$, we have $H(y_\alpha, D_x\phi(x_\alpha, y_\alpha)) \geq f_2(y_\alpha)$ if α is large enough. Now, assuming α is large enough, we always have (30), from which get a contradiction, $f_1(0) \geq f_2(0)$, by taking the limit as $\alpha \rightarrow \infty$.

We next turn to the case where $\partial U \cap \overline{\Omega} = \emptyset$. We have:

$$\Omega = (\Omega \cap U) \cup (\Omega \cap U^c) = (\Omega \cap U) \cup (\Omega \cap \text{int}(U^c)).$$

Since Ω is connected and $\Omega \cap U = \Sigma \neq \emptyset$, we see that $\Omega \cap \text{int}(U^c) = \emptyset$ and $\overline{\Omega} \subset U$. We thus need to show that $\sup_{\overline{\Omega}}(u - v) = -\infty$. Indeed, if $\max_{\overline{\Omega}}(u - v) \in \mathbb{R}$, then the argument in the previous case yields a contradiction. The proof is now complete. \square

Theorem 3.4. *Let $u \in \text{USC}(\overline{\Sigma} \times [0, T])$ and $v \in \text{LSC}(\overline{\Sigma} \times [0, T])$ be respectively a subsolution and a supersolution of (25). Assume that $u \leq v$ on $\overline{\Sigma} \times \{0\} \cup (\partial U \cap \overline{\Omega}) \times (0, T)$. Then $u \leq v$ in $\overline{\Sigma} \times [0, T)$.*

Lemma 3.5. *Assume that $f \in C(\Sigma \times (0, T))$ is bounded on $\Sigma \times (0, T)$. Then for any $R > 0$ there is a constant $C_R > 0$, depending only on R, H, f and Ω , for which if $u \in \text{USC}(\Sigma \times (0, T))$ is a subsolution of (25) and if the family $\{u(x, \cdot) \mid x \in \Sigma\}$ is equi-Lipschitz continuous on $(0, T)$ with Lipschitz constant R , then the function u is Lipschitz continuous on $\overline{\Sigma} \times (0, T)$ with Lipschitz constant C_R .*

Proof. Fix any $R > 0$. As in the proof of Lemma 3.3, there is a constant $M_R > 0$, depending only on R, H and f , such that for $(x, p) \in \Sigma \times \mathbb{R}^n$, if $H(x, p) \leq f(x) + R$, then $|p| \leq M_R$. Let $u \in \text{USC}(\Sigma \times (0, T))$ be a subsolution of (25), and assume that the family $\{u(x, \cdot) \mid x \in \Sigma\}$ is equi-Lipschitz continuous on $(0, T)$ with Lipschitz constant R . Then, it is easily seen that for each $t \in (0, T)$, the function $u(\cdot, t)$ is a subsolution of (7), with $H(x, p)$ and $f(x)$ replaced by $|p|$ and C_0 , respectively. By Lemma 3.3, there is a constant $C_R \geq R$, depending only on M_R and Ω , such that the family $\{u(\cdot, t) \mid 0 < t < T\}$ is equi-Lipschitz continuous on Σ , with Lipschitz constant C_R . Then we have $|u(x, t) - u(y, s)| \leq C_R(|x - y| + |t - s|)$ for all $(x, t), (y, s) \in \Sigma \times (0, T)$ and finish the proof. \square

Proof of Theorem 3.4. We follow the line of the proof of Theorem 3.1. For $S < T$ we write:

$$\partial'_p(\Sigma \times (S, T)) = \bar{\Sigma} \times \{S\} \cup (\partial U \cap \bar{\Omega}) \times (S, T).$$

It is enough to show that

$$\sup_{Q_T}(u - v) \leq \sup_{\partial'_p Q_T}(u - v), \tag{31}$$

where $Q_T = \Sigma \times (0, T)$.

To prove (31), we suppose, on the contrary, that

$$\sup_{Q_T}(u - v) > \sup_{\partial'_p Q_T}(u - v). \tag{32}$$

Let $\delta > 0$ and set $\tilde{u}(x, t) = u(x, t) - \delta/(T - t)$ for $(x, t) \in Q_T$. By replacing u by \tilde{u} , we may assume that u is a subsolution of (25) with $f(x)$ replaced by $f(x) - \varepsilon$, where $\varepsilon > 0$ is a constant, and that $\lim_{t \rightarrow T-} \sup_{x \in \bar{\Sigma}}(u - v)(x, t) = -\infty$. By taking the sup-convolution of u in the t -variable, replacing U and the interval $(0, T)$ by smaller (in the sense of inclusion) ones, and translating the smaller interval, we may assume that f is uniformly continuous on Q_T and the family $\{u(x, \cdot) \mid x \in \bar{\Sigma}\}$ is equi-Lipschitz continuous on $(0, T)$. According to Lemma 3.5, the function u is Lipschitz continuous on Q_T . Next, we may replace H by a uniformly continuous function on $\Sigma \times \mathbb{R}^n$. By perturbing u (resp., v) as in the proof of Theorem 3.1 and replacing $\varepsilon > 0$ by a smaller positive number, we may assume that u (resp., v) is a subsolution (resp., a supersolution) of (25), with $f(x, t)$ and $g(x)$ replaced by $f(x, t) - \varepsilon$ and $-\varepsilon$ (resp., $f(x, t)$ and ε). Moreover, we may assume that $u - v$ attains a strict maximum at a point $(z, \tau) \in \Gamma_U \times (0, T)$. Furthermore, we may assume that $z = 0, U = \text{int } B(0, r), \Omega_U = \Omega_r$ and $\Gamma_U = \Gamma_r$, where $r > 0$ and Ω_r, Γ_r are the sets given by (11).

Now we consider the function,

$$\Phi(x, t, y, s) = u(x, t) - v(y, s) - \alpha\phi(x, y) - \alpha(t - s)^2,$$

on the set $\bar{Q}_T \times \bar{Q}_T$, where $\alpha > 1$ is a constant and ϕ is the function used in the proof of Theorem 3.1. Let $(x_\alpha, y_\alpha) \in \bar{Q}_T \times \bar{Q}_T$ be a maximum point of Φ . Arguing as in the proof of Theorem 3.1, we see that if α is sufficiently large, then we always have:

$$\begin{cases} 2\alpha(t_\alpha - s_\alpha) + H(x_\alpha, \alpha D_x \phi(x_\alpha, y_\alpha)) \leq f(x_\alpha) - \varepsilon, \\ 2\alpha(t_\alpha - s_\alpha) + H(y_\alpha, \alpha D_x \phi(x_\alpha, y_\alpha)) \geq f(y_\alpha). \end{cases} \tag{33}$$

Also, using the Lipschitz continuity of u , we find that $\alpha|t_\alpha - s_\alpha| + \alpha|x_\alpha - y_\alpha| \leq C$ for some constant $C > 0$, independent of α . Sending $\alpha \rightarrow \infty$ in (33) yields a contradiction. \square

4. Skorokhod problem

In this section we are concerned with the Skorokhod problem. We recall that $\mathbb{R}_+ = (0, \infty)$ and hence $\bar{\mathbb{R}}_+ = [0, \infty)$. We denote by $L^1_{\text{loc}}(\bar{\mathbb{R}}_+, \mathbb{R}^k)$ (resp., $\text{AC}_{\text{loc}}(\bar{\mathbb{R}}_+, \mathbb{R}^k)$) the space of functions $v : \bar{\mathbb{R}}_+ \rightarrow \mathbb{R}^k$ which are integrable (resp., absolutely continuous) on any bounded interval $I \subset \bar{\mathbb{R}}_+$.

Given $x \in \bar{\Omega}$ and $v \in L^1_{\text{loc}}(\bar{\mathbb{R}}_+, \mathbb{R}^n)$, the Skorokhod problem is to seek for a pair of functions, $(\eta, l) \in \text{AC}_{\text{loc}}(\bar{\mathbb{R}}_+, \mathbb{R}^n) \times L^1_{\text{loc}}(\bar{\mathbb{R}}_+, \mathbb{R})$, such that

$$\begin{cases} \eta(0) = x, & \eta(t) \in \bar{\Omega} & \text{for } t \in \bar{\mathbb{R}}_+, \\ \dot{\eta}(t) + l(t)\gamma(\eta(t)) = v(t) & & \text{for a.e. } t \in \bar{\mathbb{R}}_+, \\ l(t) \geq 0, & l(t) = 0 \text{ if } \eta(t) \in \Omega & \text{for a.e. } t \in \bar{\mathbb{R}}_+. \end{cases} \tag{34}$$

Regarding the solvability of the Skorokhod problem, our main result is the following:

Theorem 4.1. *Let $v \in L^1_{\text{loc}}(\bar{\mathbb{R}}_+, \mathbb{R}^n)$ and $x \in \bar{\Omega}$. Then there exists a pair $(\eta, l) \in AC_{\text{loc}}(\bar{\mathbb{R}}_+, \mathbb{R}^n) \times L^1_{\text{loc}}(\bar{\mathbb{R}}_+, \mathbb{R})$ such that (34) holds.*

We are interested in “regular” solutions in the above theorem. See [26] and references therein for more general viewpoints on the Skorokhod problem. The advantage of the above result is in that it applies to domains with C^1 boundary.

A natural question is the uniqueness of the solution (η, l) in the above theorem. But we do not know if the uniqueness holds or not.

We first establish the following result.

Theorem 4.2. *Let $v \in L^\infty(\bar{\mathbb{R}}_+, \mathbb{R}^n)$ and $x \in \bar{\Omega}$. Then there exists a pair $(\eta, l) \in \text{Lip}(\bar{\mathbb{R}}_+, \mathbb{R}^n) \times L^\infty(\bar{\mathbb{R}}_+, \mathbb{R})$ such that (34) holds.*

We borrow some ideas from [26] in the following proof.

Proof. We may assume that γ is defined on \mathbb{R}^n . Let $\psi \in C^1(\mathbb{R}^n)$ be such that $\psi(x) < 0$ in Ω , $|D\psi(x)| > 1$ for $x \in \Gamma$, $\psi(x) > 0$ for $x \in \mathbb{R}^n \setminus \bar{\Omega}$ and $\liminf_{|x| \rightarrow \infty} \psi(x) > 0$. We can select a constant $\delta > 0$ so that for $x \in \mathbb{R}^n$,

$$\gamma(x) \cdot D\psi(x) \geq \delta |D\psi(x)| \quad \text{and} \quad |D\psi(x)| \geq 1 \quad \text{if } 0 \leq \psi(x) \leq \delta.$$

We set $q(x) = (\psi(x) \vee 0) \wedge \delta$ for $x \in \mathbb{R}^n$. Note that $q(x) = 0$ for $x \in \bar{\Omega}$, $q(x) > 0$ for $x \in \mathbb{R}^n \setminus \bar{\Omega}$, and $\gamma(x) \cdot Dq(x) \geq \delta |Dq(x)|$ for a.e. $x \in \mathbb{R}^n$.

Fix $\varepsilon > 0$ and $x \in \bar{\Omega}$. We consider the initial-value problem for the ODE

$$\dot{\xi}(t) + \frac{1}{\varepsilon} q(\xi(t)) \gamma(\xi(t)) = v(t) \quad \text{for a.e. } t \in \bar{\mathbb{R}}_+, \quad \xi(0) = x. \tag{35}$$

Here ξ represents the unknown function. By the standard ODE theory, there is a unique solution $\xi \in C^1(\bar{\mathbb{R}}_+)$ of (35).

Let $m \geq 2$. We multiply the ODE of (35) by $m q(\xi(t))^{m-1} Dq(\xi(t))$, to get:

$$\frac{d}{dt} q(\xi(t))^m + \frac{m}{\varepsilon} q(\xi(t))^m Dq(\xi(t)) \cdot \gamma(\xi(t)) = m q(\xi(t))^{m-1} Dq(\xi(t)) \cdot v(t) \quad \text{a.e.}$$

Fix any $T \in \mathbb{R}_+$. Integrating over $[0, T]$, we get:

$$\begin{aligned} q(\xi(T))^m - q(\xi(0))^m + \frac{m}{\varepsilon} \int_0^T q(\xi(s))^m \gamma(\xi(s)) \cdot Dq(\xi(s)) \, ds \\ = m \int_0^T q(\xi(s))^{m-1} Dq(\xi(s)) \cdot v(s) \, ds. \end{aligned}$$

Here we have:

$$\int_0^T q(\xi(s))^m \gamma(\xi(s)) \cdot Dq(\xi(s)) \, ds \geq \delta \int_0^T q(\xi(s))^m |Dq(\xi(s))| \, ds,$$

and

$$\begin{aligned} & \int_0^T q(\xi(s))^{m-1} Dq(\xi(s)) \cdot v(s) \, ds \\ & \leq \left(\int_0^T q(\xi(s))^m |Dq(\xi(s))| \, ds \right)^{1-\frac{1}{m}} \left(\int_0^T |v(s)|^m |Dq(\xi(s))| \, ds \right)^{\frac{1}{m}}. \end{aligned}$$

Combining these, we get:

$$\begin{aligned} & q(\xi(T))^m + \frac{m\delta}{\varepsilon} \int_0^T q(\xi(s))^m |Dq(\xi(s))| \, ds \\ & \leq m \left(\int_0^T q(\xi(s))^m |Dq(\xi(s))| \, ds \right)^{1-\frac{1}{m}} \left(\int_0^T |v(s)|^m |Dq(\xi(s))| \, ds \right)^{\frac{1}{m}}. \end{aligned} \tag{36}$$

From this we obtain

$$\frac{\delta}{\varepsilon} \left(\int_0^T q(\xi(s))^m |Dq(\xi(s))| \, ds \right)^{\frac{1}{m}} \leq \left(\int_0^T |v(s)|^m |Dq(\xi(s))| \, ds \right)^{\frac{1}{m}}, \tag{37}$$

and

$$q(\xi(T))^m \leq \left(\frac{\varepsilon}{\delta} \right)^{m-1} m \int_0^T |v(s)|^m |Dq(\xi(s))| \, ds.$$

Hence, setting $C_0 = \|Dq\|_{L^\infty(\mathbb{R}^n)}$, we deduce that

$$q(\xi(t))^m \leq \left(\frac{\varepsilon}{\delta} \right)^{1-\frac{1}{m}} m C_0 T \|v\|_{L^\infty(0,T)}^m \quad \text{for } t \in [0, T]. \tag{38}$$

Henceforth we write ξ_ε for ξ , to indicate the dependence on ε of ξ . We see from (38) that for any $T > 0$,

$$\lim_{\varepsilon \rightarrow 0^+} \max_{t \in [0, T]} \text{dist}(\xi_\varepsilon(t), \Omega) = 0. \tag{39}$$

Also, (38) ensures that for each $T > 0$ there is an $\varepsilon_T > 0$ such that $q(\xi_\varepsilon(t)) < \delta$ for $t \in [0, T]$ and $0 < \varepsilon < \varepsilon_T$.

Now let $T > 0$ and $0 < \varepsilon < \varepsilon_T$. We have $q(\xi_\varepsilon(s)) = \psi(\xi_\varepsilon(s)) \vee 0$ for all $t \in [0, T]$ and hence $q(\xi_\varepsilon(t))^m \times |Dq(\xi_\varepsilon(t))| = q(\xi_\varepsilon(t))^m$ for a.e. $t \in (0, T)$. Accordingly, (37) yields,

$$\frac{\delta}{\varepsilon} \left(\int_0^T q(\xi_\varepsilon(s))^m \, ds \right)^{\frac{1}{m}} \leq (C_0 T)^{\frac{1}{m}} \|v\|_{L^\infty(0,T)}.$$

Sending $m \rightarrow \infty$, we find that $(\delta/\varepsilon)\|q \circ \xi_\varepsilon\|_{L^\infty(0,T)} \leq \|v\|_{L^\infty(0,T)}$, and moreover

$$\frac{\delta}{\varepsilon} \|q \circ \xi_\varepsilon\|_{L^\infty(\mathbb{R}_+)} \leq \|v\|_{L^\infty(\mathbb{R}_+)}. \tag{40}$$

We set $l_\varepsilon = (1/\varepsilon)q \circ \xi_\varepsilon$. Due to (40), we may choose a sequence $\varepsilon_j \rightarrow 0^+$ so that $l_{\varepsilon_j} \rightarrow l$ weakly-star in $L^\infty(\mathbb{R}_+)$ as $j \rightarrow \infty$ for a function $l \in L^\infty(\mathbb{R}_+)$. It is clear that $l(s) \geq 0$ for a.e. $s \in \mathbb{R}_+$.

ODE (35) together with (40) guarantees that $\{\xi_\varepsilon\}_{\varepsilon>0}$ is bounded in $L^\infty(\overline{\mathbb{R}_+})$. Hence, we may assume as well that ξ_{ε_j} converges locally uniformly on $\overline{\mathbb{R}_+}$ to a function $\eta \in \text{Lip}(\overline{\mathbb{R}_+})$ as $j \rightarrow \infty$. It is then obvious that $\eta(0) = x$ and the pair (η, l) satisfies:

$$\eta(t) + \int_0^t (l(s)\gamma(\xi(s)) - v(s)) ds = 0 \quad \text{for } t > 0,$$

from which we get $\dot{\eta}(t) + l(t)\gamma(\eta(t)) = v(t)$ for a.e. $t \in \overline{\mathbb{R}}_+$. It follows from (39) that $\eta(t) \in \overline{\Omega}$ for $t \geq 0$.

In order to show that the pair (η, l) is a solution of (34), we need only to prove that for a.e. $t \in \overline{\mathbb{R}}_+$, $l(t) = 0$ if $\eta(t) \in \Omega$. Set $A = \{t \geq 0 \mid \eta(t) \in \Omega\}$. It is clear that A is an open subset of $[0, \infty)$. We can choose a sequence $\{I_k\}_{k \in \mathbb{N}}$ of closed intervals of A such that $A = \bigcup_{k \in \mathbb{N}} I_k$. Note that for each $k \in \mathbb{N}$, the set $\eta(I_k)$ is a compact subset of Ω and the convergence of $\{\xi_{\varepsilon_j}\}$ to η is uniform on I_k . Hence, for any fixed $k \in \mathbb{N}$, we may choose $J \in \mathbb{N}$ so that $\xi_{\varepsilon_j}(t) \in \Omega$ for all $t \in I_k$ and $j \geq J$. From this, we have $q(\xi_{\varepsilon_j}(t)) = 0$ for $t \in I_k$ and $j \geq J$. Moreover, in view of the weak-star convergence of $\{l_{\varepsilon_j}\}$, we find that for any $k \in \mathbb{N}$,

$$\int_{I_k} l(t) dt = \lim_{j \rightarrow \infty} \int_{I_k} \frac{1}{\varepsilon_j} q(\xi_j(t))^m dt = 0,$$

which yields $l(t) = 0$ for a.e. $t \in I_k$. Since $A = \bigcup_{k \in \mathbb{N}} I_k$, we see that $l(t) = 0$ a.e. in A . The proof is complete. \square

For $x \in \overline{\Omega}$, let $\text{SP}(x)$ denote the set of all triples,

$$(\eta, v, l) \in \text{AC}_{\text{loc}}(\overline{\mathbb{R}}_+, \mathbb{R}^n) \times L^1_{\text{loc}}(\overline{\mathbb{R}}_+, \mathbb{R}^n) \times L^1_{\text{loc}}(\overline{\mathbb{R}}_+),$$

which satisfies (34). We set $\text{SP} = \bigcup_{x \in \overline{\Omega}} \text{SP}(x)$.

We remark that for any $x, y \in \overline{\Omega}$ and $0 < T < \infty$, there exists a triple $(\eta, v, l) \in \text{SP}(x)$ such that $\eta(T) = y$. Indeed, given $x, y \in \overline{\Omega}$ and $0 < T < \infty$, we choose a curve $\eta \in \text{Lip}([0, T], \overline{\Omega})$ so that $\eta(0) = x$ and $\eta(T) = y$. The existence of such a curve is guaranteed since Ω is a domain and has the C^1 regularity. We extend the domain of definition of η to $\overline{\mathbb{R}}_+$ by setting $\eta(t) = y$ for $t > T$. Now, if we set $v(t) = \dot{\eta}(t)$ and $l(t) = 0$ for $t \geq 0$, we have $(\eta, v, l) \in \text{SP}(x)$, which has the property, $\eta(T) = y$. Here and henceforth, for interval I , we denote by $\text{Lip}(I, \overline{\Omega})$ the set of those $\eta \in \text{Lip}(I, \mathbb{R}^n)$ such that $\eta(t) \in \overline{\Omega}$ for $t \in I$. We use such notation for other spaces of functions having values in $\overline{\Omega} \subset \mathbb{R}^n$ as well.

We note also that problem (34) has the following *semi-group* property: for any $(x, t) \in \overline{\Omega} \times \mathbb{R}_+$ and $(\eta_1, v_1, l_1), (\eta_2, v_2, l_2) \in \text{SP}$, if $\eta_1(0) = x$ and $\eta_2(0) = \eta_1(t)$ hold and if (η, v, l) is defined on $\overline{\mathbb{R}}_+$ by:

$$(\eta(s), v(s), l(s)) = \begin{cases} (\eta_1(s), v_1(s), l_1(s)) & \text{for } s \in [0, t), \\ (\eta_2(s-t), v_2(s-t), l_2(s-t)) & \text{for } s \in [t, \infty), \end{cases}$$

then $(\eta, v, l) \in \text{SP}(x)$.

Proposition 4.3. *There is a constant $C > 0$, depending only on Ω and γ , such that for $(\eta, v, l) \in \text{SP}$, we have $|\dot{\eta}(s)| \vee l(s) \leq C|v(s)|$ for a.e. $s \geq 0$.*

An immediate consequence of the above proposition is that for $(\eta, v, l) \in \text{SP}$, if $v \in L^p(\mathbb{R}_+, \mathbb{R}^n)$ (resp., $v \in L^p_{\text{loc}}(\mathbb{R}_+, \mathbb{R}^n)$), with $1 \leq p \leq \infty$, then $(\eta, l) \in L^p(\mathbb{R}_+, \mathbb{R}^{n+1})$ (resp., $(\eta, l) \in L^p_{\text{loc}}(\mathbb{R}_+, \mathbb{R}^{n+1})$).

Proof. Thanks to hypothesis (A3), there is a constant $\delta_0 > 0$ such that $v(x) \cdot \gamma(x) \geq \delta_0$ for $x \in \Gamma$. Let $(\eta, v, l) \in \text{SP}$. According to the C^1 regularity of Ω , there is a function $\psi \in C^1(\mathbb{R}^n)$ such that $\Omega = \{x \in \mathbb{R}^n \mid \psi(x) < 0\}$ and $D\psi(x) \neq 0$ for $x \in \Gamma$. Noting that $\psi(\eta(s)) \leq 0$ for all $s \geq 0$, we find that for any $s > 0$, if $\eta(s) \in \Gamma$ and η is differentiable at s , then

$$0 = \frac{d}{ds} \psi(\eta(s)) = D\psi(\eta(s)) \cdot \dot{\eta}(s).$$

Hence, noting that $D\psi(\eta(s))$ is parallel to $v(\eta(s))$, we see that $v(\eta(s)) \cdot \dot{\eta}(s) = 0$.

Let $s > 0$ be such that $\eta(s) \in \Gamma$, $\dot{\eta}(s)$ exists, $\dot{\eta}(s) + l(s)\gamma(\eta(s)) = v(s)$, $l(s) \geq 0$ and $v(\eta(s)) \cdot \dot{\eta}(s) = 0$. We see immediately that $l(s)\gamma(\eta(s)) \cdot v(\eta(s)) = v(s) \cdot v(\eta(s))$. Hence, we get $\delta_0 l(s) \leq v(s) \cdot v(\eta(s)) \leq |v(s)|$ and $l(s) \leq \delta_0^{-1}|v(s)|$ for a.e. $s \geq 0$. We also have:

$$|\dot{\eta}(s)| \leq |v(s)| + \|\gamma\|_{\infty} |l(s)| \leq \left(1 + \frac{\|\gamma\|_{\infty}}{\delta_0}\right) |v(s)| \quad \text{for a.e. } s \geq 0. \quad \square$$

Let \mathcal{F} be a subset of $L^1(I, \mathbb{R}^m)$, where $I \subset \mathbb{R}$ is an interval. We recall that \mathcal{F} is said to be *uniformly integrable* if for any $\varepsilon > 0$ there is a $\delta > 0$ such that for any $f \in \mathcal{F}$,

$$\left| \int_B f(s) \, ds \right| < \varepsilon \quad \text{whenever } B \subset I \text{ is measurable and } |B| < \delta.$$

Here $|B|$ denotes the Lebesgue measure of $B \subset \mathbb{R}$.

Proposition 4.4. *Let $\{(\eta_k, v_k, l_k)\}_{k \in \mathbb{N}} \subset \text{SP}$. Assume that $\{|v_k|\}$ is uniformly integrable on every intervals $[0, T]$, with $0 < T < \infty$. Then there exist a subsequence $\{\eta_{k_j}, v_{k_j}, l_{k_j}\}_{j \in \mathbb{N}}$ of $\{\eta_k, v_k, l_k\}$ and an $(\eta, v, l) \in \text{SP}$ such that*

$$\begin{aligned} \eta_{k_j}(t) &\rightarrow \eta(t) && \text{uniformly on } [0, T], \\ \dot{\eta}_{k_j} \, dt &\rightarrow \dot{\eta} \, dt && \text{weakly-star in } C([0, T], \mathbb{R}^n)^*, \\ v_{k_j} \, dt &\rightarrow v \, dt && \text{weakly-star in } C([0, T], \mathbb{R}^n)^*, \\ l_{k_j} \, dt &\rightarrow l \, dt && \text{weakly-star in } C([0, T])^*, \end{aligned}$$

for every $T > 0$.

In the above proposition, we denote by X^* the dual space of the Banach space X . Regarding notation in the above proposition, we remark that the weak-star convergence in $C([0, T])^*$ or $C([0, T], \mathbb{R}^n)^*$ is usually stated as the weak convergence of measures.

Proof. By Proposition 4.3, there is a constant $C_0 > 0$ such that for $k \in \mathbb{N}$,

$$|\dot{\eta}_k(s)| \vee l_k(s) \leq C_0 |v_k(s)| \quad \text{for a.e. } s \geq 0. \tag{41}$$

It follows from this that the sequences $\{|\dot{\eta}_k|\}$ and $\{l_k\}$ are uniformly integrable on the intervals $[0, T]$, $0 < T < \infty$. If we set:

$$V_k(t) = \int_0^t v_k(s) \, ds \quad \text{and} \quad L_k(t) = \int_0^t l_k(s) \, ds \quad \text{for } t \geq 0,$$

then the sequences $\{\eta_k\}$, $\{V_k\}$ and $\{L_k\}$ are equi-continuous and uniformly bounded on the intervals $[0, T]$, $0 < T < \infty$. We may therefore choose an increasing sequence $\{k_j\} \subset \mathbb{N}$ so that the sequences $\{\eta_{k_j}\}$, $\{V_{k_j}\}$ and $\{L_{k_j}\}$ converge, as $j \rightarrow \infty$, uniformly on every finite interval $[0, T]$, $0 < T < \infty$, to some functions $\eta \in C(\overline{\mathbb{R}}_+, \overline{\mathbb{R}})$, $V \in C(\overline{\mathbb{R}}_+, \mathbb{R}^n)$ and $L \in C(\overline{\mathbb{R}}_+)$. The uniform integrability of the sequences $\{|\dot{\eta}_k|\}$, $\{|v_k|\}$ and $\{l_k\}$ implies that the functions η , V and L are absolutely continuous on every finite interval $[0, T]$, $0 < T < \infty$.

Fix any $0 < T < \infty$. The uniform integrability of the sequences $\{|\dot{\eta}_k|\}$, $\{|v_k|\}$ and $\{l_k\}$ guarantees that the sequences $\{\dot{\eta}_k \, ds\}$, $\{v_k \, ds\}$ and $\{l_k \, ds\}$ of measures on $[0, T]$ are bounded. That is, we have:

$$\sup_{k \in \mathbb{N}} \int_0^T (|\dot{\eta}_k(s)| + |v_k(s)| + l_k(s)) \, ds < \infty.$$

Hence we may assume without loss of generality that as $j \rightarrow \infty$,

$$\begin{aligned} \dot{\eta}_{k_j} \, ds &\rightarrow \mu_1 && \text{weakly-star in } C([0, T], \mathbb{R}^n)^*, \\ v_{k_j} \, dt &\rightarrow \mu_2 && \text{weakly-star in } C([0, T], \mathbb{R}^n)^*, \\ l_{k_j} \, dt &\rightarrow \mu_3 && \text{weakly-star in } C([0, T])^*, \end{aligned}$$

for some regular Borel measures μ_1 , μ_2 and μ_3 of bounded variations on $[0, T]$. Then, for any $\phi \in C^1([0, T], \mathbb{R}^n)$, using integration by parts twice, we get:

$$\begin{aligned} \int_0^T \phi(s) \mu_1(ds) &= \lim_{j \rightarrow \infty} \int_0^T \phi(s) \dot{\eta}_{k_j}(s) ds \\ &= \lim_{j \rightarrow \infty} \left([\phi \eta_{k_j}]_0^T - \int_0^T \phi'(s) \eta_{k_j}(s) ds \right) \\ &= [\phi \eta]_0^T - \int_0^T \phi'(s) \eta(s) ds = \int_0^T \phi(s) \dot{\eta}(s) ds. \end{aligned}$$

By the density of $C^1([0, T], \mathbb{R}^n)$ in $C([0, T], \mathbb{R}^n)$, we find that

$$\int_0^T \phi(s) \mu_1(ds) = \int_0^T \phi(s) \dot{\eta}(s) ds,$$

which shows that $\mu_1 = \dot{\eta} ds$ on $[0, T]$. Similarly we see that $\mu_2 = \dot{V} ds$ and $\mu_2 = \dot{L} ds$. Thus, setting $v = \dot{V}$ and $l = \dot{L}$, we have as $j \rightarrow \infty$,

$$\begin{aligned} \dot{\eta}_{k_j} ds &\rightarrow \dot{\eta} ds \quad \text{weakly-star in } C([0, T], \mathbb{R}^n)^*, \\ v_{k_j} dt &\rightarrow v ds \quad \text{weakly-star in } C([0, T], \mathbb{R}^n)^*, \\ l_{k_j} dt &\rightarrow l ds \quad \text{weakly-star in } C([0, T])^*. \end{aligned}$$

Note here that the above weak-star convergence is valid for every $0 < T < \infty$.

Since $\dot{\eta}_k(s) + l_k(s)\gamma(\eta_k(s)) = v_k(s)$ for a.e. $s \geq 0$, integrating this over $[0, t]$, with $0 < t < \infty$, and sending $k = k_j$ as $j \rightarrow \infty$, we get:

$$\eta(t) - \eta(0) + \int_0^t l(s)\gamma(\eta(s)) ds = \int_0^t v(s) ds \quad \text{for } t > 0,$$

which ensures that $\dot{\eta}(s) + l(s)\gamma(\eta(s)) = v(s)$ for a.e. $s \geq 0$. It is obvious that $\eta(s) \in \overline{\Omega}$ for $s \geq 0$. Finally, we argue as in the last part of the proof of Theorem 4.2, to find that for a.e. $s \in \mathbb{R}_+$, $l(s) = 0$ if $\eta(s) \in \Omega$. The proof is complete. \square

Proof of Theorem 4.1. Fix any $x \in \overline{\Omega}$ and $v \in L^1_{\text{loc}}(\overline{\mathbb{R}}_+, \mathbb{R}^n)$. In view of the semi-group property of problem (34), we may assume that $v(s) = 0$ for $s \geq 1$, so that $v \in L^1(\overline{\mathbb{R}}_+, \mathbb{R}^n)$. We define the sequence $\{v_k\}_{k \in \mathbb{N}} \subset L^\infty(\overline{\mathbb{R}}_+, \mathbb{R}^n)$ by:

$$v_k(s) = \begin{cases} v(s) & \text{if } |v(s)| \leq k, \\ 0 & \text{otherwise.} \end{cases}$$

Since $|v_k(s)| \leq |v(s)|$ for $s \geq 0$, we see that the sequence $\{|v_k|\}$ is uniformly integrable on $\overline{\mathbb{R}}_+$. According to Theorem 4.2, there is a sequence $\{(\eta_k, l_k)\} \subset \text{Lip}(\overline{\mathbb{R}}_+, \mathbb{R}^n) \times L^\infty(\overline{\mathbb{R}}_+, \overline{\mathbb{R}}_+)$ such that $(\eta_k, v_k, l_k) \in \text{SP}(x)$ for all $k \in \mathbb{N}$. Then applying Proposition 4.4, we deduce that there is an $(\eta, l) \in \text{AC}_{\text{loc}}(\overline{\mathbb{R}}_+, \mathbb{R}^n) \times L^1_{\text{loc}}(\overline{\mathbb{R}}_+, \overline{\mathbb{R}}_+)$ such that $(\eta, v, l) \in \text{SP}(x)$. \square

5. Cauchy problem with the Neumann type boundary condition

In this section we introduce the value function of an optimal control problem associated with the initial-boundary value problem (3)–(5), and show that it is a (unique) solution of problem (3)–(5).

Let L denote the function $L \in \text{LSC}(\overline{\Omega} \times \mathbb{R}^n, \mathbb{R} \cup \{\infty\})$, called the Lagrangian of H , given by (6). The value function w of the optimal control with the dynamics given by (34), the running cost (L, g) and the pay-off u_0 is given by:

$$w(x, t) = \inf \left\{ \int_0^t (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds + u_0(\eta(t)) \mid (\eta, v, l) \in \text{SP}(x) \right\}$$

for $(x, t) \in \overline{\Omega} \times \mathbb{R}_+$. (42)

Under our hypotheses, the Lagrangian L may take the value ∞ and, on the other hand, there is a constant $C_0 > 0$ such that $L(x, \xi) \geq -C_0$ for $(x, \xi) \in \overline{\Omega} \times \mathbb{R}^n$. Thus, it is reasonable to interpret,

$$\int_0^t L(\eta(s), -v(s)) \, ds = \infty,$$

if the function: $s \mapsto L(\eta(s), -v(s))$ is not integrable, which we adopt here.

It is well known (and also easily seen) that the value function w satisfies the dynamic programming principle,

$$w(x, s + t) = \inf \left\{ \int_0^t (L(\eta(\tau), -v(\tau)) + g(\eta(\tau))l(\tau)) \, d\tau + w(\eta(t), s) \mid (\eta, v, l) \in \text{SP}(x) \right\}$$

for $x \in \overline{\Omega}$ and $t, s \in \mathbb{R}_+$.

Theorem 5.1. *The value function w is continuous on $\overline{\Omega} \times \mathbb{R}_+$ and it is a solution of (3)–(4). Moreover, w satisfies (5) in the sense that $\lim_{t \rightarrow 0^+} w(x, t) = u_0(x)$ uniformly for $x \in \overline{\Omega}$.*

The above theorem clearly ensures the existence of a solution of (3)–(5). This together with Theorem 3.4, with $U := \mathbb{R}^n$, establishes the unique existence of a solution of (3)–(5). For the solvability of stationary and evolution problem for HJ equations, we refer to [1–4,6,5].

Another aspect of the theorem above is that it gives a variational formula for the unique solution of (3)–(5). This is a classical observation on the value functions in optimal control, and, in this regard, we refer for instance to [1,2].

The variational formula (42) is sometimes called the Lax–Oleinik formula.

For the proof of Theorem 5.1, we need the following three lemmas. We set $Q = \overline{\Omega} \times \mathbb{R}_+$.

Lemma 5.2. *Let $\psi \in C^1(\overline{Q})$ be a classical subsolution of (3)–(4). Assume that $\psi(x, 0) \leq u_0(x)$ for $x \in \overline{\Omega}$. Then $w \geq \psi$ on \overline{Q} .*

Proof. Let $(x, t) \in Q$ and $(\eta, v, l) \in \text{SP}(x)$. We have:

$$\begin{aligned} \psi(\eta(t), 0) - \psi(\eta(0), t) &= \int_0^t \frac{d}{ds} \psi(\eta(s), t - s) \, ds \\ &= \int_0^t (D\psi(\eta(s), t - s) \cdot \dot{\eta}(s) - \psi_t(\eta(s), t - s)) \, ds \\ &= \int_0^t (D\psi(\eta(s), t - s) \cdot (v(s) - l(s)\gamma(\eta(s))) - \psi_t(\eta(s), t - s)) \, ds. \end{aligned}$$

Now, using the subsolution property of ψ and the inequality $\psi(\cdot, 0) \leq u_0$, we get:

$$\begin{aligned} \psi(x, t) - u_0(\eta(t)) &\leq \int_0^t (-D\psi(\eta(s), t - s) \cdot v(s) + l(s)D\psi(\eta(s)) \cdot \gamma(\eta(s)) + \psi_t(\eta(s), t - s)) \, ds \end{aligned}$$

$$\begin{aligned} &\leq \int_0^t (H(\eta(s), D\psi(\eta(s), t-s)) + L(\eta(s), -v(s)) + l(s)D\psi(\eta(s)) \cdot \gamma(\eta(s)) \\ &\quad + \psi_t(\eta(s), t-s)) \, ds \\ &\leq \int_0^t (L(\eta(s), -v(s)) + l(s)g(\eta(s))) \, ds. \end{aligned}$$

Thus we conclude that $\psi(x, t) \leq w(x, t)$. \square

Lemma 5.3. *For any $\varepsilon > 0$ there is a constant $C_\varepsilon > 0$ such that $w(x, t) \geq u_0(x) - \varepsilon - C_\varepsilon t$ for $(x, t) \in Q$.*

Proof. We fix any $\varepsilon > 0$ and choose a function $u_0^\varepsilon \in C^1(\overline{\Omega})$ so that $|u_0(x) - u_0^\varepsilon(x)| \leq \varepsilon$ for $x \in \overline{\Omega}$. We choose a function $\psi_0 \in C^1(\mathbb{R}^n)$ so that $\Omega = \{x \in \mathbb{R}^n \mid \psi_0(x) < 0\}$ and $D\psi_0(x) \neq 0$ for $x \in \Gamma$. By multiplying ψ_0 by a positive constant, we may find a function $\psi^\varepsilon \in C^1(\overline{\Omega})$ so that $\gamma(x) \cdot D(u_0^\varepsilon + \psi^\varepsilon)(x) \geq g(x)$ for $x \in \Gamma$. Next, approximating the function: $r \mapsto (-\varepsilon) \vee (\varepsilon \wedge r)$ on \mathbb{R} by a smooth function, we build a function $\zeta_\varepsilon \in C^1(\mathbb{R})$ so that $|\zeta_\varepsilon(r)| \leq \varepsilon$ for $r \in \mathbb{R}$ and $\zeta'_\varepsilon(0) = 1$. Note that $D(\zeta_\varepsilon \circ \psi^\varepsilon)(x) = D\psi^\varepsilon(x)$ for $x \in \Gamma$ and $|u_0(x) - u_0^\varepsilon(x) - \zeta_\varepsilon \circ \psi^\varepsilon(x)| \leq 2\varepsilon$ for $x \in \overline{\Omega}$. We choose a constant $C_\varepsilon > 0$ so that $H(x, D(u_0^\varepsilon + \zeta_\varepsilon \circ \psi^\varepsilon)(x)) \leq C_\varepsilon$ for $x \in \overline{\Omega}$. Finally we define the function $\phi^\varepsilon \in C^1(\overline{Q})$ by $\phi^\varepsilon(x, t) = -2\varepsilon + u_0^\varepsilon(x) + \zeta_\varepsilon \circ \psi^\varepsilon(x) - C_\varepsilon t$, and observe that ϕ^ε is a classical subsolution of (3), (4) and that $\phi^\varepsilon(x, 0) \leq u_0(x)$ for $x \in \overline{\Omega}$. By Lemma 5.2, we get $\phi^\varepsilon(x, t) \leq w(x, t)$ for $(x, t) \in Q$. Hence, we obtain $w(x, t) \geq u_0(x) - 4\varepsilon - C_\varepsilon t$ for all $(x, t) \in Q$. \square

Lemma 5.4. *There is a constant $C > 0$ such that $w(x, t) \leq u_0(x) + Ct$ for $(x, t) \in Q$.*

Proof. Let $(x, t) \in Q$. Set $\eta(s) = x$, $v(s) = 0$ and $l(s) = 0$ for $s \geq 0$. Then $(\eta, v, l) \in \text{SP}(x)$. Hence, we have:

$$w(x, t) \leq u_0(x) + \int_0^t L(x, 0) \, ds = u_0(x) + tL(x, 0) \leq u_0(x) - t \min_{p \in \mathbb{R}^n} H(x, p).$$

Setting $C = -\min_{\overline{\Omega} \times \mathbb{R}^n} H$, we get $w(x, t) \leq u_0(x) + Ct$. \square

Lemma 5.5. *Let $t > 0$, $x \in \overline{\Omega}$, $\phi \in C^1(\overline{\Omega} \times [0, t])$ and $\varepsilon > 0$. Then there is a triple $(\eta, v, l) \in \text{SP}(x)$ such that for a.e. $s \in (0, t)$,*

$$H(\eta(s), D\phi(\eta(s), t-s)) + L(\eta(s), -v(s)) \leq \varepsilon - v(s) \cdot D\phi(\eta(s), t-s).$$

We postpone the proof of the above lemma and give now the proof of Theorem 5.1.

Proof of Theorem 5.1. By Lemmas 5.3 and 5.4, there is a constant $C > 0$ and for each $\varepsilon > 0$ a constant $C_\varepsilon > 0$ such that $-\varepsilon - C_\varepsilon t \leq w(x, t) - u_0(x) \leq Ct$ for all $(x, t) \in Q$. This shows that w is a real-valued function on Q and that

$$\lim_{t \rightarrow 0^+} w(x, t) = u_0(x) \quad \text{uniformly for } x \in \overline{\Omega}. \tag{43}$$

We next prove that w is a subsolution of (3), (4). Let $(\hat{x}, \hat{t}) \in Q$ and $\phi \in C^1(\overline{Q})$. Assume that $w^* - \phi$ attains a strict maximum at (\hat{x}, \hat{t}) . We need to show that if $\hat{x} \in \Omega$, then $\phi_t(\hat{x}, \hat{t}) + H(\hat{x}, D\phi(\hat{x}, \hat{t})) \leq 0$, and if $\hat{x} \in \Gamma$, then either

$$\phi_t(\hat{x}, \hat{t}) + H(\hat{x}, D\phi(\hat{x}, \hat{t})) \leq 0 \quad \text{or} \quad \gamma(\hat{x}) \cdot D\phi(\hat{x}, \hat{t}) \leq g(\hat{x}). \tag{44}$$

We are here concerned only with the case where $\hat{x} \in \Gamma$. The other case can be treated similarly. To prove (44), we argue by contradiction. Thus we suppose that (44) were false. We may choose an $\varepsilon \in (0, 1)$ so that $\hat{t} - 2\varepsilon > 0$ and for $(x, t) \in (\overline{\Omega} \cap B(\hat{x}, 2\varepsilon)) \times [\hat{t} - 2\varepsilon, \hat{t} + 2\varepsilon]$,

$$\phi_t(x, t) + H(x, D\phi(x, t)) \geq 2\varepsilon \quad \text{and} \quad \gamma(x) \cdot D\phi(x, t) - g(x) \geq 2\varepsilon, \tag{45}$$

where γ and g are assumed to be defined and continuous on $\overline{\Omega}$. We may assume that $(w^* - \phi)(\hat{x}, \hat{t}) = 0$. Set

$$B = (\partial B(\hat{x}, 2\varepsilon) \times [\hat{t} - 2\varepsilon, \hat{t} + 2\varepsilon] \cup B(\hat{x}, 2\varepsilon) \times \{\hat{t} - 2\varepsilon\}) \cap \overline{Q},$$

and $m = -\max_B(w^* - \phi)$. Note that $m > 0$ and $w(x, t) \leq \phi(x, t) - m$ for $(x, t) \in B$. We choose a point $(\bar{x}, \bar{t}) \in \overline{\Omega} \cap B(\hat{x}, \varepsilon) \times [\hat{t} - \varepsilon, \hat{t} + \varepsilon]$ so that $(w - \phi)(\bar{x}, \bar{t}) > -\varepsilon^2 \wedge m$. We apply Lemma 5.5, to find a triple $(\eta, v, l) \in \text{SP}(\bar{x})$ such that for a.e. $s \geq 0$,

$$H(\eta(s), D\phi(\eta(s), \bar{t} - s)) + L(\eta(s), -v(s)) \leq \varepsilon - v(s) \cdot D\phi(\eta(s), \bar{t} - s). \tag{46}$$

Note that $\sigma := \bar{t} - (\hat{t} - 2\varepsilon) \geq \varepsilon$ and $\text{dist}(\bar{x}, \partial B(\hat{x}, 2\varepsilon)) \geq \varepsilon$. Set

$$S = \{s \in [0, \sigma] \mid \eta(s) \in \partial B(\hat{x}, 2\varepsilon)\} \quad \text{and} \quad \tau = \inf S.$$

We consider first the case where $\tau = \infty$, i.e., the case $S = \emptyset$. By the dynamic programming principle, we have:

$$\begin{aligned} \phi(\bar{x}, \bar{t}) &< w(\bar{x}, \bar{t}) + \varepsilon^2 \\ &\leq \int_0^\sigma (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds + w(\eta(\sigma), \bar{t} - \sigma) + \varepsilon^2 \\ &\leq \int_0^\sigma (L(\eta(s), -v(s)) + g(\eta(s))l(s) + \varepsilon) \, ds + \phi(\eta(\sigma), \bar{t} - \sigma). \end{aligned}$$

Hence, we obtain:

$$\begin{aligned} 0 &< \int_0^\sigma \left(L(\eta(s), -v(s)) + g(\eta(s))l(s) + \varepsilon + \frac{d}{ds} \phi(\eta(s), \bar{t} - s) \right) ds \\ &\leq \int_0^\sigma (L(\eta(s), -v(s)) + g(\eta(s))l(s) + \varepsilon \\ &\quad + D\phi(\eta(s), \bar{t} - s) \cdot \dot{\eta}(s) - \phi_t(\eta(s), \bar{t} - s)) \, ds \\ &\leq \int_0^\sigma (L(\eta(s), -v(s)) + g(\eta(s))l(s) + \varepsilon \\ &\quad + D\phi(\eta(s), \bar{t} - s) \cdot (v(s) - l(s)\gamma(\eta(s)) - \phi_t(\eta(s), \bar{t} - s))) \, ds. \end{aligned}$$

Now, using (46) and (45), we get:

$$\begin{aligned} 0 &< \int_0^\sigma (2\varepsilon - H(\eta(s), D\phi(\eta(s), \bar{t} - s)) + g(\eta(s))l(s) \\ &\quad - l(s)D\phi(\eta(s), \bar{t} - s) \cdot \gamma(\eta(s)) - \phi_t(\eta(s), \bar{t} - s)) \, ds \\ &\leq \int_0^\sigma l(s)(g(\eta(s)) - \gamma(\eta(s)) \cdot D\phi(\eta(s), \bar{t} - s)) \, ds \leq 0, \end{aligned}$$

which is a contradiction.

Next we consider the case where $\tau < \infty$. Observe that $\tau > 0$, and

$$\begin{aligned} \phi(\bar{x}, \bar{t}) &< w(\bar{x}, \bar{t}) + m \\ &\leq \int_0^\tau (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds + w(\eta(\tau), \bar{t} - \tau) + m \\ &\leq \int_0^\tau (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds + \phi(\eta(\tau), \bar{t} - \tau). \end{aligned}$$

Using (46) and (45) as before, we compute that

$$\begin{aligned} 0 &< \int_0^\tau (L(\eta(s), -v(s)) + g(\eta(s))l(s) - \phi_t(\eta(s), \bar{t} - s) \\ &\quad + D\phi(\eta(s), \bar{t} - s) \cdot v(s) - l(s)\gamma(\eta(s)) \cdot D\phi(\eta(s), \bar{t} - s)) \, ds \\ &\leq \int_0^\tau \{ \varepsilon - H(\eta(s), D\phi(\eta(s), \bar{t} - s)) - \phi_t(\eta(s), \bar{t} - s) \\ &\quad + l(s)[g(\eta(s)) - \gamma(\eta(s)) \cdot D\phi(\eta(s), \bar{t} - s)] \} \, ds < 0, \end{aligned}$$

which is again a contradiction. Thus, we conclude that w is a subsolution of (3), (4).

Now, we turn to the proof of the supersolution property of w . Let $\phi \in C^1(\bar{Q})$ and $(\hat{x}, \hat{t}) \in \bar{\Omega} \times (0, \infty)$. Assume that $w_* - \phi$ attains a strict minimum at (\hat{x}, \hat{t}) . We show that if $\hat{x} \in \Omega$, then $\phi_t(\hat{x}, \hat{t}) + H(\hat{x}, D\phi(\hat{x}, \hat{t})) \geq 0$, and if $\hat{x} \in \Gamma$, then

$$\phi_t(\hat{x}, \hat{t}) + H(\hat{x}, D\phi(\hat{x}, \hat{t})) \geq 0 \quad \text{or} \quad \gamma(\hat{x}) \cdot D\phi(\hat{x}, \hat{t}) \geq g(\hat{x}). \tag{47}$$

We only consider the case where $\hat{x} \in \Gamma$, and leave it to the reader to check the details in the other case. To show (47), we suppose by contradiction that (47) were false. That is, we have $\phi_t(\hat{x}, \hat{t}) + H(\hat{x}, D\phi(\hat{x}, \hat{t})) < 0$ and $\gamma(\hat{x}) \cdot D\phi(\hat{x}, \hat{t}) - g(\hat{x}) < 0$. There is an $\varepsilon > 0$ such that if we set $R := B(\hat{x}, 2\varepsilon) \times [\hat{t} - 2\varepsilon, \hat{t} + 2\varepsilon]$, then, for $(x, t) \in R \cap \bar{Q}$, we have:

$$\phi_t(x, t) + H(x, D\phi(x, t)) < 0 \quad \text{and} \quad \gamma(x) \cdot D\phi(x, t) - g(x) < 0.$$

Here we may assume that $\hat{t} - 2\varepsilon > 0$ and $(u_* - \phi)(\hat{x}, \hat{t}) = 0$. Set $m := \min_{\bar{Q} \cap \partial R} (u_* - \phi) (> 0)$. We may choose a point $(\bar{x}, \bar{t}) \in \bar{Q}$ so that $(u_* - \phi)(\bar{x}, \bar{t}) < m$, $|\bar{x} - \hat{x}| < \varepsilon$ and $|\bar{t} - \hat{t}| < \varepsilon$. We select a triple $(\eta, v, l) \in \text{SP}(\bar{x})$ so that

$$u(\bar{x}, \bar{t}) + m > \int_0^{\bar{t}} (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds + u_0(\eta(\bar{t})).$$

We set $\tau = \min\{s \geq 0 \mid (\eta(s), \bar{t} - s) \in \partial R\}$. It is clear that $\tau > 0$, $\eta(s) \in R \cap \bar{Q}$ for $s \in [0, \tau]$ and, if $|\eta(\tau) - \hat{x}| < 2\varepsilon$, then $\tau = \bar{t} - (\hat{t} - 2\varepsilon) > \varepsilon$. Accordingly, we have:

$$\begin{aligned} \phi(\bar{x}, \bar{t}) + m &> \int_0^\tau (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds + u(\eta(\tau), \bar{t} - \tau) \\ &\geq \int_0^\tau (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds + \phi(\eta(\tau), \bar{t} - \tau) + m. \end{aligned}$$

Hence, we get:

$$\begin{aligned}
 0 &> \int_0^\tau (L(\eta(s), -v(s)) + g(\eta(s)l(s) + D\phi(\eta(s), \bar{t} - s)) \cdot \dot{\eta}(s) \\
 &\quad - \phi_t(\eta(s), \bar{t} - s)) \, ds \\
 &\geq \int_0^\tau (-v(s) \cdot D\phi(\eta(s), \bar{t} - s) - H(\eta(s), D\phi(\eta(s), \bar{t} - s)) - g(\eta(s))l(s) \\
 &\quad + D\phi(\eta(s), \bar{t} - s) \cdot \dot{\eta}(s) - \phi_t(\eta(s), \bar{t} - s)) \, ds > 0,
 \end{aligned}$$

which is a contradiction.

It remains to show that w is continuous on Q . According to (43), we have $w^*(\cdot, 0) = w_*(\cdot, 0) = u_0$ on $\bar{\Omega}$. Thus, applying the comparison theorem (Theorem 3.4 with $U := \mathbb{R}^n$), we see that $w^* \leq w_*$ on \bar{Q} , which guarantees that $w \in C(Q)$. This completes the proof. \square

For the proof of Lemma 5.5, we need the following basic lemma.

Lemma 5.6. *Let $R > 0$. There is a constant $C > 0$, depending only on R and H , such that for any $(x, p, v) \in \bar{\Omega} \times B(0, R) \times \mathbb{R}^n$, if $H(x, p) + L(x, -v) \leq 1 - v \cdot p$, then we have $|v| \leq C$.*

Proof. We may choose a constant $C_1 > 0$ so that $C_1 \geq \max_{\bar{\Omega} \times B(0, 2R)} |H|$. Observe that

$$L(x, -v) \geq \max_{p \in B(0, 2R)} (-v \cdot p) - C_1 = 2R|v| - C_1 \quad \text{for } (x, v) \in \bar{\Omega} \times \mathbb{R}^n.$$

Let $(x, p, v) \in \bar{\Omega} \times B(0, R) \times \mathbb{R}^n$ be a point such that $H(x, p) + L(x, -v) \leq 1 - v \cdot p$. Then we have $-C_1 + 2R|v| - C_1 \leq 1 + |v||p| \leq 1 + R|v|$. Consequently, we get $R|v| \leq 2C_1 + 1$. \square

For $i \in \mathbb{N}$ we introduce the function $L_i \in C(\bar{\Omega} \times \mathbb{R}^n)$ by setting:

$$L_i(x, \xi) = \max_{p \in B(0, i)} (\xi \cdot p - H(x, p)).$$

Observe that $L_i(x, \xi) \leq L(x, \xi)$ and $\lim_{i \rightarrow \infty} L_i(x, \xi) = L(x, \xi)$ for $(x, \xi) \in \bar{\Omega} \times \mathbb{R}^n$ and that every L_i is uniformly continuous on bounded subsets of $\bar{\Omega} \times \mathbb{R}^n$.

Proof of Lemma 5.5. Fix $k \in \mathbb{N}$. Set $\delta = t/k$ and $s_j = (j - 1)\delta$ for $j = 1, 2, \dots, k$. We define inductively a sequence $\{(x_j, \eta_j, v_j, l_j)\}_{j=1}^k \subset \bar{\Omega} \times \text{SP}$. We set $x_1 = x$ and choose a $\xi_1 \in \mathbb{R}^n$ so that

$$H(x_1, D\phi(x_1, t)) + L(x_1, -\xi_1) \leq \varepsilon - \xi_1 \cdot D\phi(x_1, t).$$

Set $v_1(s) = \xi_1$ for $s \geq 0$ and choose a pair $(\eta_1, l_1) \in \text{Lip}(\bar{\mathbb{R}}_+, \bar{\Omega}) \times L^\infty(\bar{\mathbb{R}}_+, \bar{\mathbb{R}}_+)$ so that $(\eta_1, v_1, l_1) \in \text{SP}(x_1)$. According to Theorem 4.2, such a pair always exists.

Suppose now that we are given (x_i, η_i, v_i, l_i) for all $i = 1, 2, \dots, j - 1$ and for some $j \leq k$. Then set $x_j = \eta_{j-1}(\delta)$, choose a $\xi_j \in \mathbb{R}^n$ so that

$$H(x_j, D\phi(x_j, t - s_j)) + L(x_j, -\xi_j) \leq \varepsilon - \xi_j \cdot D\phi(x_j, t - s_j), \tag{48}$$

set $v_j(s) = \xi_j$ for $s \geq 0$, and select a pair $(\eta_j, l_j) \in \text{Lip}(\bar{\mathbb{R}}_+, \bar{\Omega}) \times L^\infty(\bar{\mathbb{R}}_+, \mathbb{R}^n)$ so that $(\eta_j, v_j, l_j) \in \text{SP}(x_j)$. Thus, by induction, we have chosen a sequence $\{(x_j, \eta_j, v_j, l_j)\}_{j=1}^k \subset \bar{\Omega} \times \text{SP}$ such that $x_1 = \eta_1(0)$, $x_j = \eta_{j-1}(\delta) = \eta_j(0)$ for $j = 2, \dots, k$ and for each $j = 1, 2, \dots, k$, (48) holds with $\xi_j = v_j(s)$ for all $s \geq 0$. Notice that the choice of x_j, η_j, v_j, l_j , with $j = 1, \dots, k$, depends on k , which is not explicit in our notation.

Next, we define a triple $(\bar{\eta}_k, \bar{v}_k, \bar{l}_k) \in \text{SP}(x)$ by setting:

$$(\bar{\eta}_k(s), \bar{v}_k(s), \bar{l}_k(s)) = (\eta_j(s - s_j), v_j(s - s_j), l_j(s - s_j))$$

for $s_j \leq s < s_{j+1}$ and $j = 1, 2, \dots, k - 1$, and

$$(\bar{\eta}_k(s), \bar{v}_k(s), \bar{l}_k(s)) = (\eta_k(s - s_k), v_k(s - s_k), l_k(s - s_k)),$$

for $s \geq s_k$. We may assume that $\varepsilon < 1$ and, by Lemma 5.6, we find a constant $C_1 > 0$, independent of k , such that $\max_{s \geq 0} |\bar{v}_k(s)| = \max_{1 \leq j \leq k} |\xi_j| \leq C_1$. By Proposition 4.3, we find a constant $C_2 > 0$, independent of k , such that $\|d\bar{\eta}_k/ds\|_{L^\infty(\mathbb{R}_+)} \vee \|\bar{l}_k\|_{L^\infty(\mathbb{R}_+)} \leq C_2$. Now, we define the step function χ_k on \mathbb{R}_+ by setting $\chi_k(s) = s_j$ for $s_j \leq s < s_{j+1}$ and $j = 1, 2, \dots, k$ and $\chi_k(s) = s_k$ for $s \geq s_k$, and observe that (48), with $1 \leq j \leq k$, can be rewritten as

$$\begin{aligned} & H(\bar{\eta}_k(\chi_k(s)), D\phi(\bar{\eta}_k(\chi_k(s)), t - \chi_k(s))) + L(\bar{\eta}_k(\chi_k(s)), -\bar{v}_k(s)) \\ & \leq \varepsilon - \bar{v}_k(s) \cdot D\phi(\bar{\eta}_k(\chi_k(s)), t - \chi_k(s)) \quad \text{for } 0 \leq s \leq t. \end{aligned} \tag{49}$$

We may invoke Proposition 4.4, to find a triple $(\eta, v, l) \in \text{SP}(x)$ and a subsequence of $\{(\bar{\eta}_k, \bar{v}_k, \bar{l}_k)\}_{k \in \mathbb{N}}$, which will be denoted again by the same symbol, so that for every $0 < T < \infty$, as $k \rightarrow \infty$, $\bar{\eta}_k \rightarrow \eta$ uniformly on $[0, T]$, $\bar{v}_k ds \rightarrow v ds$ weakly-star in $C([0, T], \mathbb{R}^n)^*$ and $\bar{l}_k ds \rightarrow l ds$ weakly-star in $C([0, T])^*$. We may moreover assume that $\bar{v}_k \rightarrow v$ weakly-star in $L^\infty(\mathbb{R}_+, \mathbb{R}^n)$ and $\bar{l}_k \rightarrow l$ weakly-star in $L^\infty(\mathbb{R}_+)$ as $k \rightarrow \infty$.

Since $\bar{v}_k \rightarrow v$ weakly in $L^2(0, t)$, we may choose a sequence $\{\lambda_k\}_{k \in \mathbb{N}}$ of finite sequences $\lambda_k = (\lambda_{k,1}, \lambda_{k,2}, \dots, \lambda_{k,N_k})$ of nonnegative numbers such that

$$\sum_{j=1}^{N_k} \lambda_{k,j} = 1 \quad \text{and} \quad \hat{v}_k := \sum_{j=1}^{N_k} \lambda_{k,j} \bar{v}_{k+j} \quad \text{converge to } v \text{ in } L^2(0, t).$$

Here we may moreover assume by selecting a subsequence of $\{(\bar{\eta}_k, \bar{v}_k, \bar{l}_k)\}$ that as $k \rightarrow \infty$, $\hat{v}_k(s) \rightarrow v(s)$ for a.e. $s \in (0, t)$.

Fix any $i \in \mathbb{N}$ and $\theta > 1$. In view of the uniform continuity of the functions H and L_i on bounded subsets of $\bar{\Omega} \times \mathbb{R}^n$ and the uniform convergence of $\{\bar{\eta}_k\}$ to η on $[0, t]$, from (49), we get:

$$\begin{aligned} & H(\eta(s), D\phi(\eta(s), t - s)) + L_i(\eta(s), -\bar{v}_k(s)) \\ & \leq \theta\varepsilon - \bar{v}_k(s) \cdot D\phi(\eta(s), t - s) \quad \text{for } s \in (0, t), \end{aligned}$$

for sufficiently large k , say, for $k \geq k_\theta$, and hence, by taking the convex combination,

$$\begin{aligned} & H(\eta(s), D\phi(\eta(s), t - s)) + L_i(\eta(s), -\hat{v}_k(s)) \\ & \leq \theta\varepsilon - \hat{v}_k(s) \cdot D\phi(\eta(s), t - s) \quad \text{for } s \in (0, t), \end{aligned}$$

for $k \geq k_\theta$. Sending $k \rightarrow \infty$, we get for a.e. $s \in (0, t)$,

$$H(\eta(s), D\phi(\eta(s), t - s)) + L_i(\eta(s), -v(s)) \leq \theta\varepsilon - v(s) \cdot D\phi(\eta(s), t - s),$$

and, because of the arbitrariness of i and $\theta > 1$, we obtain for a.e. $s \in (0, t)$,

$$H(\eta(s), D\phi(\eta(s), t - s)) + L(\eta(s), -v(s)) \leq \varepsilon - v(s) \cdot D\phi(\eta(s), t - s). \quad \square$$

6. Aubry sets and formulas for solutions of (1), (2)

In this section we define the Aubry set associated with (1), (2). Our argument here is very close to that of [21,22].

By the C^1 regularity of Ω and assumption (A3), there is a function $\psi \in C^1(\bar{\Omega})$ such that $D_\gamma \psi(x) > 0$ for $x \in \Gamma$. By multiplying ψ by a positive constant, we may assume that $D_\gamma \psi(x) \geq |g(x)|$ for $x \in \Gamma$. Selecting a constant $C_- \in \mathbb{R}$ small enough, we may have $H(x, D\psi(x)) \geq C_-$ for $x \in \Omega$. It is easy to check that the function ψ is a supersolution of (1), (2), with C_- in place of a . Similarly, if we choose a constant $C_+ \in \mathbb{R}$ large enough, then the function $-\psi$ is a subsolution of (1), (2), with C_+ in place of a .

We define the *critical value* (or additive eigenvalue) c by:

$$c = \inf\{a \in \mathbb{R} \mid \text{there is a subsolution of (1), (2)}\}.$$

Obviously we have $c \leq C_+$. By Corollary 3.2, we see as well that $c \geq C_-$. In particular, we have $c \in \mathbb{R}$. For any decreasing sequence $\{a_k\}$ converging to c , there is a sequence $\{u_k\} \subset \text{USC}(\bar{\Omega})$ such that for every $k \in \mathbb{N}$, u_k is a

subsolution of (1), (2), with a_k in place of a . By Lemma 3.3, with $U = \mathbb{R}^n$, we find that $\{u_k\}$ is equi-Lipschitz continuous on $\overline{\Omega}$. By adding a constant to u_k , we may assume that $\{u_k\}$ is uniformly bounded on $\overline{\Omega}$. By choosing a subsequence, we may thus assume that the sequence $\{u_k\}$ converges to a function $u \in \text{Lip}(\overline{\Omega})$ as $k \rightarrow \infty$. By the stability of the viscosity property under uniform convergence, we see that u is a subsolution of (1), (2), with c in place of a .

Henceforth in this section, we normalize $c = 0$ by replacing H by $H - c$, and we are concerned only with problem (1), (2), with $a = 0$, that is, the problem:

$$\begin{cases} H(x, Du(x)) = 0 & \text{in } \Omega, \\ D_\gamma u(x) = g(x) & \text{on } \Gamma. \end{cases} \tag{50}$$

We introduce the function d on $\overline{\Omega} \times \overline{\Omega}$ by:

$$d(x, y) = \sup\{v(x) - v(y) \mid v \text{ is a subsolution of (50)}\}. \tag{51}$$

According to Lemma 3.3, the family of functions $d(\cdot, y)$, with $y \in \overline{\Omega}$, is equi-Lipschitz continuous on $\overline{\Omega}$. By the stability of the viscosity property, we see that for any $y \in \overline{\Omega}$, the function $d(\cdot, y)$ is a subsolution of (50). It is easily seen that

$$d(x, y) \leq d(x, z) + d(z, y) \quad \text{for } x, y, z \in \overline{\Omega}.$$

Also, in view of the Perron method, we find that for every $y \in \overline{\Omega}$, the function $d(\cdot, y)$ is a solution of,

$$\begin{cases} H(x, Du(x)) = 0 & \text{in } \Omega \setminus \{y\}, \\ D_\gamma u(x) = g(x) & \text{on } \Gamma \setminus \{y\}, \end{cases} \tag{52}$$

which is just problem (7), with $f := 0$ and $U := \mathbb{R}^n \setminus \{y\}$.

We define the Aubry set \mathcal{A} associated with (50) (or (1), (2) with generic a) by:

$$\mathcal{A} = \{y \in \overline{\Omega} \mid d(\cdot, y) \text{ is a solution of (50)}\}.$$

Theorem 6.1. *The Aubry set \mathcal{A} is a nonempty and compact.*

Remark 1. If we define the function d_a on $\overline{\Omega} \times \overline{\Omega}$ by:

$$d_a(x, y) = \sup\{v(x) - v(y) \mid v \text{ is a subsolution of (1), (2)}\},$$

then $d_a(x, y) = \sup \emptyset = -\infty$ for $a < 0$. Moreover, if we define the Aubry set \mathcal{A}_a for $a > 0$ by:

$$\mathcal{A}_a = \{y \in \overline{\Omega} \mid d_a(\cdot, y) \text{ is a solution of (1), (2)}\},$$

then $\mathcal{A}_a = \emptyset$.

The nonemptiness of \mathcal{A} will be proved based on the following observation.

Lemma 6.2. *Let $y \in \overline{\Omega} \setminus \mathcal{A}$. Then there are functions $v \in \text{Lip}(\overline{\Omega})$ and $f \in C(\overline{\Omega})$ such that $f(y) < 0$, $f(x) \leq 0$ for $x \in \overline{\Omega}$ and v is a subsolution of (7), with $U = \mathbb{R}^n$.*

Proof. Fix any $y \in \overline{\Omega} \setminus \mathcal{A}$ and set $u(x) = d(x, y)$ for $x \in \overline{\Omega}$. For definiteness, we consider the case where $y \in \Gamma$. We leave it to the reader to check the other case. Since u is not a supersolution of (50) while it is a solution of (52), we find a C^1 function ϕ on $\overline{\Omega}$ such that $u - \phi$ attains a strict minimum at y , $H(y, D\phi(y)) < 0$ and $D_\gamma \phi(y) < g(y)$. By continuity, there is an open neighborhood V of y such that

$$H(x, D\phi(x)) < 0 \quad \text{for } x \in \Omega_V \quad \text{and} \quad D_\gamma \phi(x) < g(x) \quad \text{for } x \in \Gamma_V. \tag{53}$$

We may assume by adding a constant to ϕ that $u(y) = \phi(y)$. Note that $\min_{\overline{\Omega} \setminus V} (u - \phi) > 0$, and select a constant $\varepsilon > 0$ small enough so that $(u - \phi)(x) > \varepsilon$ for $x \in \overline{\Omega} \setminus V$. We may choose an open neighborhood W of V^c such that $(u - \phi)(x) > \varepsilon$ for $x \in \overline{\Omega} \cap W$. We set $v(x) = u(x) \vee (\phi(x) + \varepsilon)$ for $x \in \overline{\Omega}$.

Observe that $v(x) = u(x)$ for $x \in W \cap \overline{\Omega}$, which ensures that v is a subsolution of (7), with $f(x) := 0$ and $U := W$. On the other hand, there is an open neighborhood $Y \subset V$ of y such that $\phi(x) + \varepsilon > u(x)$ for $x \in Y \cap \overline{\Omega}$. It is clear that $\overline{\Omega} \cap Y \cap W = \emptyset$. In view of (53), we may choose a function $f \in C(\overline{\Omega})$ so that $f(y) < 0$, $f(x) \leq 0$ for $x \in Y$, $f(x) = 0$ for $x \in \overline{\Omega} \setminus Y$ and

$$H(x, D\phi(x)) \leq f(x) \quad \text{for } x \in \Omega_V \quad \text{and} \quad D_\gamma \phi(x) \leq g(x) \quad \text{for } x \in \Gamma_V.$$

It is easily seen that v is a subsolution of (7), with $U := V$. Finally, we note that v is a subsolution of (7), with $U := \mathbb{R}^n$, and finish the proof. \square

Proof of Theorem 6.1. The compactness of \mathcal{A} follows directly from the stability of the viscosity property under uniform convergence.

To see that $\mathcal{A} \neq \emptyset$, we suppose by contradiction that $\mathcal{A} = \emptyset$. By Lemma 6.2, for each $y \in \overline{\Omega}$ there are functions $v_y \in \text{Lip}(\overline{\Omega})$ and $f_y \in C(\overline{\Omega})$ such that $f_y(y) < 0$, $f_y(x) \leq 0$ for $x \in \overline{\Omega}$ and v_y is a subsolution of (7), with $f := f_y$ and $U := \mathbb{R}^n$. By the compactness of $\overline{\Omega}$, we may choose a finite sequence $\{y_j\}_{j=1}^J \subset \overline{\Omega}$ so that $\sum_{j=1}^J f_{y_j}(x) < 0$ for $x \in \overline{\Omega}$. Theorem 2.2, with $U := \mathbb{R}^n$, guarantees that the function $v(x) = (1/J) \sum_{j=1}^J v_{y_j}(x)$ on $\overline{\Omega}$ is a subsolution of (7), with $U := \mathbb{R}^n$ and $f(x) := (1/J) \sum_{j=1}^J f_{y_j}(x)$. We choose a constant $a < 0$ so that $f(x) \leq a$ for $x \in \overline{\Omega}$ and observe that v is a subsolution of (7), with $f := a$ and $U := \mathbb{R}^n$. This contradicts the fact that $c = 0$. The proof is complete. \square

Proposition 6.3. *The function d can be represented as*

$$d(x, y) = \inf \left\{ \int_0^t (L(\eta(s), -v(s)) + g(\gamma(s))l(s)) \, ds \mid t > 0, (\eta, v, l) \in \text{SP}(x), \eta(t) = y \right\}. \tag{54}$$

Proof. Fix any $y \in \overline{\Omega}$. We denote by $w(x)$ the right-hand side of (54). According to Theorem 5.1, the function,

$$u(x, t) := \inf \left\{ \int_0^t L(\eta(s), -v(s)) + g(\eta(s))l(s) \, ds + d(\eta(t), y) \mid (\eta, v, l) \in \text{SP}(x) \right\},$$

is a solution of (3)–(5), with $u_0 := d(\cdot, y)$. Noting that the function $d(x, y)$, as a function of $(x, t) \in \overline{\Omega} \times \overline{\mathbb{R}}_+$, is a subsolution of (3)–(5) with $u_0 := d(\cdot, y)$, by applying the comparison theorem (Theorem 3.4, with $U = \mathbb{R}^n$), we see that $d(x, y) \leq u(x, t)$ for $(x, t) \in \overline{\Omega} \times \overline{\mathbb{R}}_+$. Since $d(y, y) = 0$, we have $\inf_{t>0} u(x, t) \leq w(x)$ for $x \in \overline{\Omega}$. Consequently, we have $d(x, y) \leq w(x)$ for $x \in \overline{\Omega}$.

By the C^1 regularity of Ω , for each $x \in \overline{\Omega}$ we may choose a Lipschitz continuous curve η on $[0, t]$ connecting x to y in $\overline{\Omega}$, with a Lipschitz constant independent of x . Here $t > 0$ is an appropriate constant, and moreover we may assume that $t \leq C_1|x - y|$ for some constant $C_1 > 0$ independent of x . As is well known and easily shown, $L(x, \xi)$ is bounded on $\overline{\Omega} \times B(0, \delta)$, if $\delta > 0$ is chosen sufficiently small. Fix such a constant $\delta > 0$ and choose a constant $C_2 > 0$ so that $L(x, \xi) \leq C_2$ for $(x, \xi) \in \overline{\Omega} \times B(0, \delta)$. By scaling, we may assume that $|\dot{\eta}(s)| \leq \delta$ for a.e. $s \in [0, t]$. Noting that $(\eta, \dot{\eta}, 0) \in \text{SP}(x)$, we get:

$$w(x) \leq \int_0^t L(\eta(s), -\dot{\eta}(s)) \, ds \leq C_2 t \leq C_1 C_2 |x - y|.$$

In particular, we may conclude that w is continuous at y and $w(y) = 0$.

To complete the proof, it is enough to show that w is a subsolution of (50). Indeed, once this is done, by the definition of d , we get:

$$w(x) = w(x) - w(y) \leq d(x, y) \quad \text{for } x \in \overline{\Omega},$$

which guarantees that $d(x, y) = w(x)$ for $x \in \overline{\Omega}$.

To prove the subsolution property of w , we just need to follow the argument of the proof of Theorem 5.1. Let $\hat{x} \in \overline{\Omega}$ and $\phi \in C^1(\overline{\Omega})$. Assume that $w^* - \phi$ attains a strict maximum at \hat{x} . We need to show that if $\hat{x} \in \Omega$, then $H(\hat{x}, D\phi(\hat{x})) \leq 0$, and if $\hat{x} \in \Gamma$, then either

$$H(\hat{x}, D\phi(\hat{x})) \leq 0 \quad \text{or} \quad \gamma(\hat{x}) \cdot D\phi(\hat{x}) \leq g(\hat{x}). \tag{55}$$

We are here concerned only with the case where $\hat{x} \in \Gamma$, and leave the proof in the other case to the reader. To show (55), we suppose by contradiction that (55) were false. Then we may choose an $\varepsilon \in (0, 1)$ so that for $x \in \overline{\Omega} \cap B(\hat{x}, 2\varepsilon)$,

$$H(x, D\phi(x)) \geq 2\varepsilon \quad \text{and} \quad \gamma(x) \cdot D\phi(x) - g(x) \geq 2\varepsilon, \tag{56}$$

where γ and g are, as usual, assumed to be defined and continuous on $\overline{\Omega}$. We may also assume that $(w^* - \phi)(\hat{x}) = 0$. Set $B = \partial B(\hat{x}, 2\varepsilon) \cap \overline{\Omega}$, and $m = -\max_B(w^* - \phi)$. Obviously, we have $m > 0$ and $w(x) \leq \phi(x) - m$ for $x \in B$. We choose a point $\bar{x} \in \overline{\Omega} \cap B(\hat{x}, \varepsilon)$ so that $(w - \phi)(\bar{x}) > -\varepsilon^2 \wedge m$. We apply Lemma 5.5, to obtain a triple $(\eta, v, l) \in \text{SP}(\bar{x})$ such that for a.e. $s \geq 0$,

$$H(\eta(s), D\phi(\eta(s))) + L(\eta(s), -v(s)) \leq \varepsilon - v(s) \cdot D\phi(\eta(s)). \tag{57}$$

Note that $\text{dist}(\bar{x}, \partial B(\hat{x}, 2\varepsilon)) \geq \varepsilon$, and set $\tau = \inf\{s > 0 \mid \eta(s) \in \partial B(\hat{x}, 2\varepsilon)\}$. Consider first the case where $\tau = \infty$, which means that $\eta(s) \in \text{int } B(\hat{x}, 2\varepsilon)$ for all $s \geq 0$. By the dynamic programming principle, we have:

$$\phi(\bar{x}) < w(\bar{x}) + \varepsilon^2 \leq \int_0^\varepsilon (L(\eta(s), -v(s)) + g(\eta(s))l(s) + \varepsilon) ds + \phi(\eta(\varepsilon)).$$

Hence, we obtain:

$$\begin{aligned} 0 &< \int_0^\varepsilon (L(\eta(s), -v(s)) + g(\eta(s))l(s) + \varepsilon + D\phi(\eta(s)) \cdot \dot{\eta}(s)) ds \\ &\leq \int_0^\varepsilon \{L(\eta(s), -v(s)) + g(\eta(s))l(s) + \varepsilon + D\phi(\eta(s)) \cdot (v(s) - l(s)\gamma(\eta(s)))\} ds. \end{aligned}$$

Now, using (57) and (56), we get:

$$0 < \int_0^\varepsilon \{2\varepsilon - H(\eta(s), D\phi(\eta(s))) + l(s)[g(\eta(s)) - D\phi(\eta(s)) \cdot \gamma(\eta(s))]\} ds \leq 0,$$

which is a contradiction.

Consider next the case where $\tau < \infty$. Note that

$$\begin{aligned} \phi(\bar{x}) < w(\bar{x}) + m &\leq \int_0^\tau (L(\eta(s), -v(s)) + g(\eta(s))l(s)) ds + w(\eta(\tau)) + m \\ &\leq \int_0^\tau (L(\eta(s), -v(s)) + g(\eta(s))l(s)) ds + \phi(\eta(\tau)). \end{aligned}$$

Using (57) and (56) as before, we obtain

$$0 < \int_0^\tau \{\varepsilon - H(\eta(s), D\phi(\eta(s))) + l(s)[g(\eta(s)) - \gamma(\eta(s)) \cdot D\phi(\eta(s))]\} ds < 0.$$

This is again a contradiction, and we conclude that w is a subsolution of (50). \square

We give another characterization of the Aubry set associated with (50).

Theorem 6.4. Let $\tau > 0$ and $y \in \overline{\Omega}$. Then we have $y \in \mathcal{A}$ if and only if,

$$\inf \left\{ \int_0^t (L(\eta(s), -v(s)) + g(\eta(s))l(s)) ds \mid t > \tau, (\eta, v, l) \in \text{SP}, \eta(0) = \eta(t) = y \right\} = 0. \tag{58}$$

Lemma 6.5. Let $u_0 \in C(\overline{\Omega})$ and let $u \in C(\overline{\Omega} \times \overline{\mathbb{R}}_+)$ be the solution of (3)–(5). Set $u^-(x) = \liminf_{t \rightarrow \infty} u(x, t)$ for $x \in \overline{\Omega}$. Then $u^- \in \text{Lip}(\overline{\Omega})$ and it is a solution of (50).

Proof. Thanks to Theorem 6.1, there is a solution $\phi \in \text{Lip}(\overline{\Omega})$ of (50). By adding a constant to ϕ if needed, we may assume that $\phi(x) \leq u_0(x)$ for $x \in \overline{\Omega}$. Let $C > 0$ be a constant such that $u_0(x) \leq \phi(x) + C$ for $x \in \overline{\Omega}$. By comparison, we get $\phi(x) \leq u(x, t) \leq \phi(x) + C$ for $x \in \overline{\Omega}$.

Setting $v(x, t) = \inf_{s > t} u(x, s)$ for $(x, t) \in \overline{\Omega} \times \mathbb{R}_+$, we note that $u^-(x) = \sup_{t > 0} v(x, t)$ for $x \in \overline{\Omega}$. Applying Theorem 2.8 (and the remark after it) to the family $\{u(\cdot, \cdot + s)\}_{s > 0}$ of solutions of (3), (4), we see that v is a solution of (3), (4). Observe also that $v \in \text{USC}(\overline{\Omega} \times \mathbb{R}_+)$ and the functions $v(x, \cdot)$, with $x \in \overline{\Omega}$, are nondecreasing on \mathbb{R}_+ . This monotonicity of v guarantees that the functions $v(\cdot, t)$, with $t > 0$, are subsolutions of (50), which implies that the family $\{v(\cdot, t)\}_{t > 0}$ is equi-Lipschitz continuous on $\overline{\Omega}$. Accordingly, we have $u^- \in \text{Lip}(\overline{\Omega})$. By the Dini lemma, we see that $u^-(x) = \lim_{t \rightarrow \infty} v(x, t)$ uniformly for $x \in \overline{\Omega}$. By the stability of the viscosity property under uniform convergence, we conclude that u^- is a solution of (50). \square

Proof of Theorem 6.4. Fix any $\tau > 0$ and $y \in \overline{\Omega}$. By Proposition 6.3, we have:

$$\inf \left\{ \int_0^t (L(\eta(s), -v(s)) + g(\eta(s))l(s)) ds \mid (\eta, v, l) \in \text{SP}, \eta(0) = \eta(t) = y \right\} \geq d(y, y) = 0 \quad \text{for } t > 0. \tag{59}$$

We assume that $y \in \mathcal{A}$ and show that (58) holds. Note that the function $u(x, t) = d(x, y)$ on $\overline{\Omega} \times \mathbb{R}$ is the unique solution of the initial-boundary value problem (3)–(5), with $u_0 := d(\cdot, y)$. By Theorem 5.1, we get:

$$0 = d(y, y) = \inf \left\{ \int_0^\tau (L(\eta(s), -v(s)) + g(\eta(s))l(s)) ds + d(\eta(\tau), y) \mid (\eta, v, l) \in \text{SP}(y) \right\}.$$

Fix any $\varepsilon > 0$ and choose a triple $(\eta, v, l) \in \text{SP}(y)$ so that

$$\varepsilon > \int_0^\tau (L(\eta(s), -v(s)) + g(\eta(s))l(s)) ds + d(\eta(\tau), y).$$

In view of Proposition 6.3, by modifying (η, v, l) on the set (τ, ∞) if necessary, we may assume that for some $t > \tau$,

$$d(\eta(\tau), y) + \varepsilon > \int_\tau^t (L(\eta(s), -v(s)) + g(\eta(s))l(s)) ds \quad \text{and} \quad \eta(t) = y.$$

Thus, we obtain,

$$2\varepsilon > \int_0^t (L(\eta(s), -v(s)) + g(\eta(s))l(s)) ds \quad \text{and} \quad \eta(0) = \eta(t) = y,$$

which ensures together with (59) that (58) holds.

Next we assume that (58) holds and show that $y \in \mathcal{A}$. Let u be the unique solution of problem (3)–(5), with initial data $d(\cdot, y)$. Since $d(\cdot, y)$, regarded as a function on $\overline{\Omega} \times \overline{\mathbb{R}}_+$, is a subsolution of (3), (4), by comparison, we see that $d(x, y) \leq u(x, t)$ for $(x, t) \in \overline{\Omega} \times [0, \infty)$. As in Lemma 6.5, we set $u^-(x) = \liminf_{t \rightarrow \infty} u(x, t)$ for $x \in \overline{\Omega}$, to find that $u^- \in \text{Lip}(\overline{\Omega})$ and u^- is a solution of (50). It follows that $d(x, y) \leq u^-(x)$ for $x \in \overline{\Omega}$. It is easily seen from (58) that for each $k \in \mathbb{N}$,

$$\inf \left\{ \int_0^t (L(\eta(s), v(s)) + g(\eta(s))l(s)) ds \mid t > k\tau, (\eta, v, l) \in \text{SP}, \eta(0) = \eta(t) = y \right\} = 0.$$

On the other hand, we have:

$$\inf_{t > k\tau} u(y, t) \leq \inf \left\{ \int_0^t (L(\eta(s), v(s)) + g(\eta(s))l(s)) ds \mid t > k\tau, (\eta, v, l) \in \text{SP}, \eta(0) = \eta(t) = y \right\}.$$

These together ensure that $u^-(y) \leq 0$ and hence $d(x, y) \geq u^-(x)$ for $x \in \overline{\Omega}$. Thus we find that $d(x, y) = u^-(x)$ and conclude that $y \in \mathcal{A}$. \square

Theorem 6.6. *Let $u \in \text{USC}(\overline{\Omega})$ and $v \in \text{LSC}(\overline{\Omega})$ be respectively a subsolution and a supersolution of (50). Assume that $u(x) \leq v(x)$ for $x \in \mathcal{A}$. Then $u(x) \leq v(x)$ for $x \in \overline{\Omega}$.*

Lemma 6.7. *There exist functions $\psi \in \text{Lip}(\overline{\Omega})$ and $f \in C(\overline{\Omega})$ such that $f(x) \leq 0$ for $x \in \overline{\Omega}$, $f(x) < 0$ for $x \in \overline{\Omega} \setminus \mathcal{A}$ and ψ is a subsolution of (7), with $U := \mathbb{R}^n$.*

Proof. By Lemma 6.2, for each $y \in \overline{\Omega} \setminus \mathcal{A}$ there are functions $f_y \in C(\overline{\Omega})$ and $\psi_y \in C(\overline{\Omega})$ such that $f_y(y) < 0$, $f_y(x) \leq 0$ for $x \in \overline{\Omega}$ and ψ_y is a subsolution of (7), with $U := \mathbb{R}^n$ and $f := f_y$. Since $\{\psi_y\}_{y \in \overline{\Omega} \setminus \mathcal{A}}$ is equi-Lipschitz continuous on $\overline{\Omega}$, we may assume by adding to ψ_y an appropriate constant $C_y \in \mathbb{R}$ if necessary that $\{\psi_y\}_{y \in \overline{\Omega} \setminus \mathcal{A}}$ is uniformly bounded on $\overline{\Omega}$. Also, we may assume without any loss of generality that $\{f_y\}_{y \in \overline{\Omega} \setminus \mathcal{A}}$ is uniformly bounded on $\overline{\Omega}$. We may choose a sequence $\{y_j\}_{j \in \mathbb{N}} \subset \overline{\Omega} \setminus \mathcal{A}$ so that $\inf_{j \in \mathbb{N}} f_{y_j}(x) < 0$ for $x \in \overline{\Omega} \setminus \mathcal{A}$. Now we set $\psi(x) = \sum_{j \in \mathbb{N}} 2^{-j} \psi_{y_j}(x)$ for $x \in \overline{\Omega}$, and observe in view of Theorem 2.2 that ψ is a subsolution of (7), with $U := \mathbb{R}^n$ and f given by $f(x) = \sum_{j \in \mathbb{N}} 2^{-j} f_{y_j}(x)$ for $x \in \overline{\Omega}$. Finally, we note that $f(x) \leq 0$ for $x \in \overline{\Omega}$, $f(x) < 0$ for $x \in \overline{\Omega} \setminus \mathcal{A}$ and $\psi \in \text{Lip}(\overline{\Omega})$. The proof is complete. \square

Proof of Theorem 6.6. Due to Lemma 6.7, there are functions $f \in C(\overline{\Omega})$ and $\psi \in \text{Lip}(\overline{\Omega})$ such that $f(x) \leq 0$ for $x \in \overline{\Omega}$, $f(x) < 0$ for $x \in \overline{\Omega} \setminus \mathcal{A}$ and ψ is a subsolution of (7), with $U := \mathbb{R}^n$. Fix any $0 < \varepsilon < 1$ and set $u_\varepsilon(x) = (1 - \varepsilon)u(x) + \varepsilon\psi(x)$ for $x \in \overline{\Omega}$. Then the function u_ε is a subsolution of (7), with $U := \mathbb{R}^n$ and f replaced by εf . We apply Theorem 3.1, with $U := \mathbb{R}^n \setminus \mathcal{A}$, to obtain $u_\varepsilon \leq v$ on $\overline{\Omega}$, which implies that $u \leq v$ on $\overline{\Omega}$. \square

Theorem 6.8. *Let $u \in C(\overline{\Omega})$ be a solution of (50). Then*

$$u(x) = \min\{u(y) + d(x, y) \mid y \in \mathcal{A}\} \quad \text{for } x \in \overline{\Omega}. \tag{60}$$

Proof. We denote by $w(x)$ the right-hand side of (60). We note first by the remark after Theorem 2.7 that w is a solution of (50). Next, by the definition of d , we have $u(x) - u(y) \leq d(x, y)$ for $x, y \in \overline{\Omega}$. Hence we get $u(x) \leq w(x)$ for $x \in \overline{\Omega}$. Also, by the definition of w , we have $w(x) \leq u(x)$ for $x \in \mathcal{A}$. Thus we have $u(x) = w(x)$ for $x \in \mathcal{A}$. By Theorem 6.6, we conclude that $u = w$ on $\overline{\Omega}$. \square

Corollary 6.9. *If $u \in C(\overline{\Omega})$ is a solution of (50), then*

$$u(x) = \inf \left\{ \int_0^t (L(\eta(s), -v(s)) + g(\eta(s))l(s)) ds + u(\eta(t)) \mid t > 0, (\eta, v, l) \in \text{SP}(x), \eta(t) \in \mathcal{A} \right\} \quad \text{for } x \in \overline{\Omega}.$$

Theorem 6.8 and Proposition 6.3 yield the above assertion.

7. Calibrated extremals

As in the previous section, we assume throughout this section that the critical value c is equal to zero.

Lemma 7.1. *Let $0 < T < \infty$ and $\{(\eta_k, v_k, l_k)\}_{k \in \mathbb{N}} \subset \text{SP}$. Assume that there is a constant $C > 0$, independent of $k \in \mathbb{N}$, such that*

$$\int_0^T (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s)) \, ds \leq C \quad \text{for } k \in \mathbb{N}.$$

Then there exists a triple $(\eta, v, l) \in \text{SP}$ such that

$$\begin{aligned} & \int_0^T (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds \\ & \leq \liminf_{k \rightarrow \infty} \int_0^T (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s)) \, ds. \end{aligned}$$

Moreover, for the triple (η, v, l) , there is a subsequence $\{(\eta_{k_j}, v_{k_j}, l_{k_j})\}$ of $\{(\eta_k, v_k, l_k)\}$ such that as $j \rightarrow \infty$,

$$\eta_{k_j}(0) \rightarrow \eta(0), \tag{61}$$

$$\dot{\eta}_{k_j}(t) \, dt \rightarrow \dot{\eta}(t) \, dt \quad \text{weakly-star in } C([0, T], \mathbb{R}^n)^*, \tag{62}$$

$$v_{k_j}(t) \, dt \rightarrow v(t) \, dt \quad \text{weakly-star in } C([0, T], \mathbb{R}^n)^*, \tag{63}$$

$$l_{k_j}(t) \, dt \rightarrow l(t) \, dt \quad \text{weakly-star in } C([0, T])^*. \tag{64}$$

Of course, under the hypotheses of the above theorem, the functions,

$$\eta_{k_j}(t) = \eta_{k_j}(0) + \int_0^t \dot{\eta}_{k_j}(s) \, ds,$$

converge to $\eta(t)$ uniformly on $[0, T]$ as $j \rightarrow \infty$.

Proof. We may assume without loss of generality that $\eta_k(t) = \eta_k(T)$, $v_k(t) = 0$ and $l_k(t) = 0$ for $t \geq T$ and $k \in \mathbb{N}$.

According to Proposition 4.3, there is a constant $C_0 > 0$ such that for any $(\eta, v, l) \in \text{SP}$, $|\dot{\eta}(t)| \vee |l(t)| \leq C_0|v(t)|$ for a.e. $t \geq 0$. Note that for each $A > 0$ there is a constant $C_A > 0$ such that $L(x, \xi) \geq A|\xi| - C_A$ for $(x, \xi) \in \overline{\Omega} \times \mathbb{R}^n$. From this lower bound of L , it is obvious that for $(x, \xi, r) \in \overline{\Omega} \times \mathbb{R}^n \times \overline{\mathbb{R}}_+$, if $r \leq C_0|\xi|$, then

$$L(x, \xi) + g(x)r \geq A|\xi| - C_A - C_0\|g\|_\infty|\xi|, \tag{65}$$

which ensures that there is a constant $C_1 > 0$ such that for $(\eta, v, l) \in \text{SP}$,

$$L(\eta(s), -v(s)) + g(\eta(s))l(s) + C_1 \geq 0 \quad \text{for a.e. } s \geq 0. \tag{66}$$

Using (66), we obtain for any measurable $B \subset [0, T]$,

$$\begin{aligned} & \int_B (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s) + C_1) \, ds \\ & \leq \int_0^T (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s) + C_1) \, ds \leq C + C_1T. \end{aligned}$$

This together with (65) yields,

$$(A - C_0\|g\|_\infty) \int_B |v_k(s)| \, ds \leq C_A|B| + C + C_1T \quad \text{for } A > 0. \tag{67}$$

This shows that the sequence $\{|v_k|\}$ is uniformly integrable on \mathbb{R}_+ .

We choose an increasing sequence $\{k_j\} \subset \mathbb{N}$ so that

$$\begin{aligned} & \liminf_{k \rightarrow \infty} \int_0^T (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s)) \, ds \\ &= \lim_{j \rightarrow \infty} \int_0^T (L(\eta_{k_j}(s), -v_{k_j}(s)) + g(\eta_{k_j}(s))l_{k_j}(s)) \, ds. \end{aligned}$$

Thanks to estimate (67), in view of Proposition 4.4, we may assume by replacing $\{k_j\}$ by a subsequence if needed that there is a triple $(\eta, v, l) \in \text{SP}$ such that the convergences (61)–(64) hold. Here we may assume that $(\eta(t), v(t), l(t)) = (\eta(T), 0, 0)$ for $t \geq T$.

In what follows, we write (η_j, v_j, l_j) for $(\eta_{k_j}, v_{k_j}, l_{k_j})$ for notational simplicity. It remains to show that

$$\begin{aligned} & \int_0^T (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds \\ & \leq \lim_{j \rightarrow \infty} \int_0^T (L(\eta_j(s), -v_j(s)) + g(\eta_j(s))l_j(s)) \, ds. \end{aligned}$$

In view of the monotone convergence theorem, we need to show that for each $m \in \mathbb{N}$,

$$\begin{aligned} & \int_0^T (L_m(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds \\ & \leq \lim_{j \rightarrow \infty} \int_0^T (L(\eta_j(s), -v_j(s)) + g(\eta_j(s))l_j(s)) \, ds, \end{aligned}$$

where $L_m(x, \xi) := \max_{p \in B(0, m)} (\xi \cdot p - H(x, p))$ for $(x, \xi) \in \overline{\Omega} \times \mathbb{R}^n$. Note that

$$\begin{aligned} & L_m(x, \xi) \leq L_{m+1}(x, \xi) \quad \text{for } (x, \xi) \in \overline{\Omega} \times \mathbb{R}^n \text{ and } m \in \mathbb{N}, \\ & \lim_{m \rightarrow \infty} L_m(x, \xi) = L(x, \xi) \quad \text{for } (x, \xi) \in \overline{\Omega} \times \mathbb{R}^n, \end{aligned}$$

and the functions L_m are uniformly continuous on bounded subsets of $\overline{\Omega} \times \mathbb{R}^n$.

We fix any $m \in \mathbb{N}$. In view of the selection theorem of Kuratowski and Ryll–Nardzewski, we may choose a Borel function $P_m : \overline{\Omega} \times \mathbb{R}^n \rightarrow B(0, m)$, so that

$$L_m(x, \xi) = \xi \cdot P_m(x, \xi) - H(x, P_m(x, \xi)) \quad \text{for } (x, \xi) \in \overline{\Omega} \times \mathbb{R}^n. \tag{68}$$

Indeed, if we define the multifunction: $\overline{\Omega} \times \mathbb{R}^n \rightarrow 2^{\mathbb{R}^n}$ by,

$$F(x, \xi) = \{p \in B(0, m) \mid L_m(x, \xi) = \xi \cdot p - H(x, p)\},$$

then (i) $F(x, \xi)$ is a nonempty closed set for every $(x, \xi) \in \overline{\Omega} \times \mathbb{R}^n$ and (ii) $F^{-1}(K)$ is a closed set whenever $K \subset \mathbb{R}^n$ is closed. From (ii), we see easily that $F^{-1}(U)$ is a F_σ -set (and hence a Borel set) whenever $U \subset \mathbb{R}^n$ is open. Hence, as claimed above, by the theorem of Kuratowski and Ryll–Nardzewski (see, for instance, [27, Theorem 1.5]), there exists a Borel function: $P_m : \overline{\Omega} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $P_m(x, \xi) \in F(x, \xi)$ for all $(x, \xi) \in \overline{\Omega} \times \mathbb{R}^n$.

Set $p(t) = P_m(\eta(t), -v(t))$ for $t \geq 0$. Let ρ_ε , with $\varepsilon > 0$, be a mollification kernel on \mathbb{R} whose support is contained in $[-\varepsilon, 0]$ and set $p_\varepsilon(t) = \rho_\varepsilon * p(t)$ for $t \geq 0$.

We fix any $\varepsilon > 0$, and observe by the definition of L that

$$\begin{aligned} I &:= \int_0^T (L(\eta_j(s), -v_j(s)) + g(\eta_j(s))l_j(s)) \, ds \\ &\geq \int_0^T (-v_j(s) \cdot p_\varepsilon(s) - H(\eta_j(s), p_\varepsilon(s)) + g(\eta_j(s))l_j(s)) \, ds. \end{aligned}$$

From this, in view of (61)–(64), we find that

$$I \geq \int_0^T (-v(s) \cdot p_\varepsilon(s) - H(\eta(s), p_\varepsilon(s)) + g(\eta(s))l(s)) \, ds. \tag{69}$$

Note here that $|p_\varepsilon(s)| \leq m$ for $s \geq 0$ and $p_\varepsilon \rightarrow p$ in $L^1(0, T)$ as $\varepsilon \rightarrow 0$. In particular, for some sequence $\varepsilon_k \rightarrow +0$, we have $p_{\varepsilon_k}(t) \rightarrow p(t)$ for a.e. $t \in [0, T]$ as $k \rightarrow \infty$. Sending $\varepsilon \rightarrow 0$ along the sequence $\varepsilon = \varepsilon_k$ and using (68), from (69) we obtain:

$$\begin{aligned} I &\geq \int_0^T (-v(s) \cdot p(s) - H(\eta(s), p(s)) + g(\eta(s))l(s)) \, ds \\ &= \int_0^T (L_m(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds, \end{aligned}$$

which completes the proof. \square

Theorem 7.2. *Let $u_0 \in C(\overline{\mathcal{D}})$ and let $u \in C(\overline{\mathcal{D}} \times \overline{\mathbb{R}}_+)$ be the unique solution of (3)–(5). Let $(x, t) \in \overline{\mathcal{D}} \times \mathbb{R}_+$. Then there exists a triple $(\eta, v, l) \in \text{SP}(x)$ such that*

$$u(x, t) = \int_0^t (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds + u_0(\eta(t)).$$

If, in addition, $u \in \text{Lip}(\overline{\mathcal{D}} \times (\alpha, \beta))$, with $0 \leq \alpha < \beta \leq \infty$, then the triple (η, v, l) , restricted to (α, β) , belongs to $\text{Lip}(\alpha, \beta) \times L^\infty(\alpha, \beta) \times L^\infty(\alpha, \beta)$.

Here we should note that the infimum on the right-hand side of formula (42) is always attained, which is a consequence of the above theorem and Theorem 5.1.

Proof. Fix $(x, t) \in \overline{\mathcal{D}}$. By Theorem 5.1, we can choose a sequence $\{(\eta_k, v_k, l_k)\} \subset \text{SP}(x)$ such that

$$u(x, t) = \lim_{k \rightarrow \infty} \left\{ \int_0^t (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s)) \, ds + u_0(\eta_k(t)) \right\}.$$

By virtue of Lemma 7.1, there are an increasing sequence $\{k_j\} \subset \mathbb{N}$ and an $(\eta, v, l) \in \text{SP}$ such that $\eta_{k_j}(s) \rightarrow \eta(s)$ uniformly on $[0, t]$, and

$$\begin{aligned} &\int_0^t (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds \\ &\leq \liminf_{k \rightarrow \infty} \int_0^t (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s)) \, ds. \end{aligned}$$

Now it is easy to see that

$$u(x, t) \geq \int_0^t (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds + u_0(\eta(t)),$$

but we have already the opposite inequality by Theorem 5.1.

Now, assume in addition that $u \in \text{Lip}(\overline{\Omega} \times (\alpha, \beta))$, where $0 \leq \alpha < \beta \leq \infty$. Let $C > 0$ be a Lipschitz constant of the function u on the set $\overline{\Omega} \times [\alpha, \beta]$. Let $C_0 > 0$ be the constant from Proposition 4.3, so that $|\dot{\eta}(s)| \vee l(s) \leq C_0|v(s)|$ for a.e. $s \geq 0$. As in the proof of Proposition 4.4, for each $A > 0$ we choose a constant $C_A > 0$ so that $L(y, \xi) \geq A|\xi| - C_A$ for $(y, \xi) \in \overline{\Omega} \times \mathbb{R}^n$. Fix any finite interval $[a, b] \subset (\alpha, \beta)$. Then, with help of the dynamic programming principle, we get:

$$\begin{aligned} \int_a^b (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds &= u(\eta(b), b) - u(\eta(a), a) \\ &\leq C(|\eta(b) - \eta(a)| + |b - a|) \leq \int_a^b (C|\dot{\eta}(s)| + C) \, ds \\ &\leq \int_a^b (CC_0|v(s)| + C) \, ds. \end{aligned}$$

On the other hand, for any $A > 0$, we have:

$$\int_a^b (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds \geq \int_a^b ((A - C_0\|g\|_\infty)|v(s)| - C_A) \, ds.$$

Combining these, we obtain:

$$\int_a^b ((A - C_0\|g\|_\infty - CC_0)|v(s)| - C_A - C) \, ds \leq 0.$$

We fix $A > 0$ so that $A \geq C_0\|g\|_\infty + CC_0 + 1$ and get:

$$\int_a^b (|v(s)| - C_A - C) \, ds \leq 0.$$

Since a, b are arbitrary as far as $\alpha < a < b < \beta$, we conclude from the above that $|v(s)| \leq C_A + C$ for a.e. $s \in (\alpha, \beta)$. By Proposition 4.3, we see that $(\eta, v, l) \in \text{Lip}(\alpha, \beta) \times L^\infty(\alpha, \beta) \times L^\infty(\alpha, \beta)$. \square

Theorem 7.3. *Let $\phi \in \text{Lip}(\overline{\Omega})$ be a solution of (1), (2), with $a := 0$. Let $x \in \overline{\Omega}$. Then there is a triple $(\eta, v, l) \in \text{SP}(x)$ such that for any $t > 0$,*

$$\phi(x) - \phi(\eta(t)) = \int_0^t (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds. \tag{70}$$

Moreover, $(\eta, v, l) \in \text{Lip}(\overline{\mathbb{R}}_+) \times L^\infty(\mathbb{R}_+) \times L^\infty(\mathbb{R}_+)$.

Let ϕ and $(\eta, v, l) \in \text{SP}$. Following [8], we call a triple $(\eta, v, l) \in \text{SP}$ a *calibrated extremal* associated with ϕ if (70) holds for all $t > 0$.

Proof. Note that the function $u(x, t) = \phi(x)$ is a solution of (3), (4). Using Theorem 7.2, we define inductively the sequence $\{(\eta_k, v_k, l_k)\} \subset \text{SP}$ as follows. We choose first an $(\eta_1, v_1, l_1) \in \text{SP}(x)$ so that

$$\phi(\eta(0)) - \phi(\eta(1)) = \int_0^1 (L(\eta_1(s)) + g(\eta_1(s))l_1(s)) \, ds.$$

We next assume that $\{(\eta_k, v_k, l_k)\}_{k \leq j-1}$, with $j \geq 2$, is given and choose an $(\eta_j, v_j, l_j) \in \text{SP}(\eta_{j-1}(1))$ so that

$$\phi(\eta_j(1)) - \phi(\eta_j(0)) = \int_0^1 (L(\eta_j(s), -v_j(s)) + g(\eta_j(s))l_j(s)) \, ds.$$

Once the sequence $\{(\eta_k, v_k, l_k)\}_{k \in \mathbb{N}} \subset \text{SP}$ is given, we define the $(\eta, v, l) \in \text{SP}(x)$ by setting $(\eta(s+k-1), v(s+k-1), l(s+k-1)) = (\eta_k(s), v_k(s), l_k(s))$ for $k \in \mathbb{N}$ and $s \in [0, 1)$. It is clear that the triple (η, v, l) satisfies (70). Thanks to Theorem 7.2, we have $(\eta_k, v_k, l_k) \in \text{Lip}([0, 1]) \times L^\infty(0, 1) \times L^\infty(0, 1)$ for $k \in \mathbb{N}$. Reviewing the proof of Theorem 7.2, we see easily that $\sup_{k \in \mathbb{N}} \|v_k\|_{L^\infty(0,1)} < \infty$, from which we conclude that $(\eta, v, l) \in \text{Lip}(\overline{\mathbb{R}_+}) \times L^\infty(\mathbb{R}_+) \times L^\infty(\mathbb{R}_+)$. \square

Theorem 7.4. Let $\phi \in \text{Lip}(\overline{\Omega})$ be a solution of (1), (2), with $a := 0$ and $(\eta, v, l) \in \text{SP}$ a calibrated extremal associated with ϕ . Then

$$\lim_{t \rightarrow \infty} \text{dist}(\eta(t), \mathcal{A}) = 0.$$

Proof. According to Lemma 6.7, there are functions $\psi \in \text{Lip}(\overline{\Omega})$ and $f \in C(\overline{\Omega})$ such that $f(x) < 0$ for $x \in \overline{\Omega} \setminus \mathcal{A}$, $f(x) \leq 0$ for $x \in \overline{\Omega}$ and ψ is a subsolution of (7), with $U = \mathbb{R}^n$. Then $u(x, t) := \psi(x)$ is a subsolution of (3), (4), with H replaced by $H - f$. By comparison, if $w \in C(\overline{\Omega} \times \overline{\mathbb{R}_+})$ is a solution of (3)–(5), with H replaced by $H - f$ and $u_0 := \psi$, then we get $u \leq w$ on $\overline{\Omega} \times \overline{\mathbb{R}_+}$. Hence, using Theorem 5.1, with H replaced by $H - f$, we find that for any $t > 0$,

$$\begin{aligned} \psi(\eta(0)) &\leq \int_0^t (L(\eta(s), -v(s)) + f(\eta(s)) + g(\eta(s))l(s)) \, ds + \psi(\eta(t)) \\ &= \phi(\eta(0)) - \phi(\eta(t)) + \psi(\eta(t)) + \int_0^t f(\eta(s)) \, ds. \end{aligned} \tag{71}$$

From this we find that

$$\inf_{t > 0} \int_0^t f(\eta(s)) \, ds > -\infty \quad \text{or equivalently} \quad \int_0^\infty |f(\eta(s))| \, ds < \infty,$$

which yields,

$$\lim_{t \rightarrow \infty} \int_t^{t+1} |f(\eta(s))| \, ds = 0. \tag{72}$$

Reviewing the proof of Lemma 7.1 up to (63), since

$$\begin{aligned} \int_t^{t+1} (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds &= \phi(\eta(t)) - \phi(\eta(t+1)) \\ &\leq 2\|\phi\|_\infty \quad \text{for } t \geq 0, \end{aligned}$$

we deduce that for any $A > 0$ and $t \geq 0$,

$$(A - C_0 \|g\|_\infty) \int_t^{t+\varepsilon} |v(s)| \, ds \leq C_A \varepsilon + 2\|\phi\|_\infty + C_1,$$

where the constants C_0, C_1, C_A are selected as in the proof of Lemma 7.1. This estimate together with Proposition 4.4 guarantees that η is uniformly continuous on $\overline{\mathbb{R}}_+$. Now, (72) ensures that $\lim_{t \rightarrow \infty} f(\eta(t)) = 0$ and hence $\lim_{t \rightarrow \infty} \text{dist}(\eta(t), \mathcal{A}) = 0$. \square

Let $\text{SP}_{-\infty}$ denote the set of all triples $(\eta, v, l) : \mathbb{R} \rightarrow \overline{\Omega} \times \mathbb{R}^n \times \overline{\mathbb{R}}_+$ such that for every $T \geq 0$, the triple (η_T, v_T, l_T) defined on $\overline{\mathbb{R}}_+$ by $(\eta_T(t), v_T(t), l_T(t)) = (\eta(t - T), v(t - T), l(t - T))$ belongs to SP .

Theorem 7.5. *For any $y \in \mathcal{A}$ there exists a triple $(\eta, v, l) \in \text{SP}_{-\infty}$ such that $\eta(0) = y$, $\eta(t) \in \mathcal{A}$ for $t \in \mathbb{R}$ and for any $-\infty < \sigma < \tau < \infty$,*

$$\int_\sigma^\tau (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds = d(\eta(\sigma), \eta(\tau)),$$

where d is the function on $\overline{\Omega} \times \overline{\Omega}$ given by (51).

Proof. Fix $y \in \mathcal{A}$. By Theorem 6.4, for any $k \in \mathbb{N}$ there is a triple $(\bar{\eta}_k, \bar{v}_k, \bar{l}_k) \in \text{SP}$ such that $\bar{\eta}_k(0) = \bar{\eta}_k(\tau_k) = y$ for some $\tau_k > k$, and

$$\frac{1}{k} > \int_0^{\tau_k} (L(\bar{\eta}_k(s), -\bar{v}_k(s)) + g(\bar{\eta}_k(s))\bar{l}_k(s)) \, ds. \tag{73}$$

For $k \in \mathbb{N}$ we set

$$(\eta_k(t), v_k(t), l_k(t)) = \begin{cases} (\bar{\eta}_k(t), \bar{v}_k(t), \bar{l}_k(t)) & \text{for } t \in [0, \tau_k], \\ (\bar{\eta}_k(t + \tau_k), \bar{v}_k(t + \tau_k), \bar{l}_k(t + \tau_k)) & \text{for } t \in [-\tau_k, 0]. \end{cases}$$

In view of Proposition 6.3, using (73), we see that if $-\tau_k \leq \sigma \leq \tau \leq \tau_k$, then

$$\begin{aligned} d(y, \eta_k(\sigma)) &\leq \int_{-\tau_k}^\sigma (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s)) \, ds, \\ d(\eta_k(\sigma), \eta_k(\tau)) &\leq \int_\sigma^\tau (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s)) \, ds, \\ d(\eta_k(\tau), y) &\leq \int_\tau^{\tau_k} (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s)) \, ds, \\ \left(\int_{-\tau_k}^\sigma + \int_\sigma^\tau + \int_\tau^{\tau_k} \right) (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s)) \, ds \\ &< \frac{2}{k} = \frac{2}{k} + d(y, y) \\ &\leq \frac{2}{k} + d(y, \eta_k(\sigma)) + d(\eta_k(\sigma), \eta_k(\tau)) + d(\eta_k(\tau), y). \end{aligned}$$

Consequently we get for $-\tau_k < \sigma < \tau < \tau_k$,

$$\begin{aligned}
d(\eta_k(\sigma), \eta_k(\tau)) &\leq \int_{\sigma}^{\tau} (L(\eta_k(s), -v_k(s)) + g(\eta_k(s))l_k(s)) \, ds \\
&< d(\eta_k(\sigma), \eta_k(\tau)) + \frac{2}{k}, \\
0 &\leq d(y, \eta_k(\tau)) + d(\eta_k(\tau), y) < \frac{2}{k}.
\end{aligned}$$

Hence, applying Lemma 7.1, we find a triple $(\eta, v, l) \in \text{SP}_{-\infty}$ such that $\eta(0) = y$ and for any $-\infty < \sigma < \tau < \infty$,

$$\begin{aligned}
d(y, \eta(\tau)) + d(\eta(\tau), y) &= 0, \\
\int_{\sigma}^{\tau} (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds &\leq d(\eta(\sigma), \eta(\tau)). \tag{74}
\end{aligned}$$

The last inequality yields for any $-\infty < \sigma < \tau < \infty$,

$$d(\eta(\sigma), \eta(\tau)) = \int_{\sigma}^{\tau} (L(\eta(s), -v(s)) + g(\eta(s))l(s)) \, ds.$$

Theorem 6.4 and (74) together guarantee that $\eta(t) \in \mathcal{A}$ for all $t \in \mathbb{R}$. The proof is complete. \square

Acknowledgements

The author was supported in part by KAKENHI #18204009, #20340026 and #21224001, JSPS. He would like to thank Professor P. Cardaliaguet for his kind help in preparing the resume.

References

- [1] P.-L. Lions, Neumann type boundary conditions for Hamilton–Jacobi equations, *Duke Math. J.* 52 (4) (1985) 793–820.
- [2] P.-L. Lions, N.S. Trudinger, Linear oblique derivative problems for the uniformly elliptic Hamilton–Jacobi–Bellman equation, *Math. Z.* 191 (1) (1991) 1–15.
- [3] G. Barles, P.-L. Lions, Fully nonlinear Neumann type boundary conditions for first-order Hamilton–Jacobi equations, *Nonlinear Anal.* 16 (2) (1991) 143–153.
- [4] P. Dupuis, H. Ishii, On oblique derivative problems for fully nonlinear second-order elliptic partial differential equations on nonsmooth domains, *Nonlinear Anal.* 15 (12) (1990) 1123–1138.
- [5] M.G. Crandall, H. Ishii, P.-L. Lions, User’s guide to viscosity solutions of second order partial differential equations, *Bull. Amer. Math. Soc. (N.S.)* 27 (1) (1992) 1–67.
- [6] G. Barles, Fully nonlinear Neumann type boundary conditions for second-order elliptic and parabolic equations, *J. Differential Equations* 106 (1) (1993) 90–106.
- [7] P.-L. Lions, G. Papanicolaou, S.R.S. Varadhan, Homogenization of Hamilton–Jacobi equations, unpublished.
- [8] A. Fathi, Weak KAM theorem in Lagrangian dynamics, preliminary version number 10, 2008.
- [9] M. Bardi, I. Capuzzo-Dolcetta, *Optimal Control and Viscosity Solutions of Hamilton–Jacobi–Bellman Equations*. With Appendices by Maurizio Falcone and Pierpaolo Soravia, Birkhäuser Inc., Boston, MA, 1997.
- [10] A. Fathi, Théorème KAM faible et théorie de Mather sur les systèmes lagrangiens, *C. R. Acad. Sci. Paris Sér.* 324 (9) (1997) 1043–1046.
- [11] W. E, Aubry–Mather theory and periodic solutions of the forced Burgers equation, *Comm. Pure Appl. Math.* 52 (7) (1999) 811–828.
- [12] L.C. Evans, A survey of partial differential equations methods in weak KAM theory, *Comm. Pure Appl. Math.* 57 (4) (2004) 445–480.
- [13] H. Ishii, H. Mitake, Representation formulas for solutions of Hamilton–Jacobi equations with convex Hamiltonians, *Indiana Univ. Math. J.* 56 (5) (2007) 2159–2183.
- [14] A. Fathi, Sur la convergence du semi-groupe de Lax–Oleinik, *C. R. Acad. Sci. Paris Sér. I Math.* 327 (3) (1998) 267–270.
- [15] J.-M. Roquejoffre, Convergence to steady states or periodic solutions in a class of Hamilton–Jacobi equations, *J. Math. Pures Appl.* (9) 80 (1) (2001) 85–104.
- [16] A. Davini, A. Siconolfi, A generalized dynamical approach to the large time behavior of solutions of Hamilton–Jacobi equations, *SIAM J. Math. Anal.* 38 (2) (2006) 478–502.
- [17] H. Ishii, Asymptotic solutions for large time of Hamilton–Jacobi equations in Euclidean n space, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 25 (2) (2008) 231–266.
- [18] N. Ichihara, H. Ishii, Long-time behavior of solutions of Hamilton–Jacobi equations with convex and coercive Hamiltonians, *Arch. Ration. Mech. Anal.* 194 (2) (2009) 383–419.

- [19] H. Mitake, Asymptotic solutions of Hamilton–Jacobi equations with state constraints, *Appl. Math. Optim.* 58 (3) (2008) 393–410.
- [20] H. Mitake, The large-time behavior of solutions of the Cauchy–Dirichlet problem for Hamilton–Jacobi equations, *NoDEA Nonlinear Differential Equations Appl.* 15 (3) (2008) 347–362.
- [21] A. Fathi, A. Siconolfi, Existence of C^1 critical subsolutions of the Hamilton–Jacobi equation, *Invent. Math.* 155 (2) (2004) 363–388.
- [22] A. Fathi, A. Siconolfi, PDE aspects of Aubry–Mather theory for quasiconvex Hamiltonians, *Calc. Var. Partial Differential Equations* 22 (2) (2005) 185–228.
- [23] O.-S. Serea, Viscosity solutions for partial differential equations with Neumann type boundary conditions and some aspects of Aubry–Mather theory, *J. Math. Anal. Appl.* 336 (1) (2007) 664–682.
- [24] I. Capuzzo Dolcetta, P.-L. Lions, Hamilton–Jacobi equations with state constraints, *Trans. Amer. Math. Soc.* 318 (2) (1990) 643–683.
- [25] E.N. Barron, R. Jensen, Semicontinuous viscosity solutions for Hamilton–Jacobi equations with convex Hamiltonians, *Comm. Partial Differential Equations* 15 (12) (1990) 1713–1742.
- [26] P.-L. Lions, A.-S. Sznitman, Stochastic differential equations with reflecting boundary conditions, *Comm. Pure Appl. Math.* 37 (4) (1984) 511–537.
- [27] J.E. Jayne, C.A. Rogers, *Selectors*, Princeton University Press, Princeton, NJ, 2002.