

NEW SPECIES

Discovery of a new species within *Bacillus* genus in a stool sample from Gabon: “*Bacillus massiliogabonensis*” sp. nov.

G. Mourembou^{1,2}, A. Ndjoyi-Mbiguino⁴, J. B. Lekana-Douki^{3,5}, P.-E. Fournier¹, D. Raoult^{1,6} and F. Bittar¹

1) Aix-Marseille Université, URMITE, UM63, CNRS 7278, IRD 198, INSERM 1095, Marseille, France, 2) Ecole Doctorale Régionale d'Afrique Centrale, 3) Unité de Parasitologie Médicale (UPARAM) CIRMF, Franceville, 4) Département de Microbiologie, Laboratoire national de référence IST/sida, Faculté de Médecine, 5) Département de Parasitologie Mycologie et de Médecine Tropicale, Université des Sciences de la Santé, Libreville, Gabon and 6) Special Infectious Agents Unit, King Fahd Medical Research Center, King Abdulaziz University, Jeddah, Saudi Arabia

Abstract

The discovery of new bacteria from the human gut using a culturomics method is a novel field of increasing interest in microbiology. Here the main characteristics of “*Bacillus massiliogabonensis*” strain Marseille P2639, a new Gram-negative bacterium isolated from the stool sample of a healthy 16-year-old Gabonese boy, are reported.

© 2016 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases.

Keywords: 16S rRNA gene, “*Bacillus massiliogabonensis*”, culturomics, human gut microbiota, taxonomy

Original Submission: 15 September 2016; **Revised Submission:** 26 September 2016; **Accepted:** 28 September 2016

Article published online: 5 October 2016

Corresponding author: F. Bittar, Aix-Marseille Université, URMITE, UM63, CNRS 7278, IRD 198, INSERM 1095, Marseille, France

E-mail: fadi.bittar@univ-amu.fr

Recent advances in microbiology enhance the culture strategies of bacteria through a new method called culturomics, which is based on the multiplication of bacterial culture conditions [1,2]. After receiving the approval of the National Ethics Committee of Gabon (no. 0023/2013/SG/CNE) and IFR48 of Marseille, France (no. 09-022), a stool sample was collected from a healthy 16-year-old Gabonese boy (body mass index, 19.03 kg·m⁻²) (in Lebamba, Gabon) in January 2015 and sent to Unité des Maladies Infectieuses et Tropicales Emergentes (URMITE, Marseille, France) for microbiologic analysis. Coupled with taxonogenomics, culturomics detected and isolated strain Marseille P2639. Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) failed to identify this bacterial strain using a Microflex spectrometer (Bruker Daltonics, Bremen, Germany).

Succinctly, strain Marseille P2639 was isolated using a Marine medium culture bottle (bioMérieux, Marcy l'Etoile, France)

followed by subculture in 5% sheep's blood-enriched Columbia agar (bioMérieux) in an aerobic atmosphere at 37°C [3–5]. The Gram-negative bacillus, strain Marseille P2639, is motile and catalase positive. It does not exhibit oxidase activity. Endospore forming, individual colonies of this bacterium are whitish with a diameter of 3 mm on a 5% sheep's blood-enriched Columbia agar plate. Additionally, its individual cell is 3.7 µm (range, 3.3–4.2 µm) in length with a diameter of 0.8 µm (range, 0.7–1 µm). Strain Marseille P2639 is strictly aerobic and grows within 24 hours. Its growth temperature ranges from 23°C to 45°C with an optimum temperature of 37°C. Strain Marseille P2639 can grow under salinity ranging from 0 to 50 g/L and pH varying from 6 to 10.

To complete the identification of strain Marseille P2639, 16S rRNA gene sequencing was performed using a 3130-XL sequencer (Applied Biosciences, Saint Aubin, France). The set of primers used included fDI-rP2 [3–5]. Correction and assembly of the 16S rRNA sequences were done using CodonCode Aligner. Nucleotide Basic Local Alignment Search Tool (BLAST) analysis performed using the National Center for Biotechnology Information database showed that the 16S rRNA gene sequence of strain Marseille P2639 has a 97.2% similarity with the closest species, *Bacillus acidicola* strain 105-2 [6]. This similarity value is lower than the percentage of 16S rRNA gene sequence threshold recommended by Meier-Kolthoff et al. [7] for Firmicutes to




FIG. 1. Phylogenetic tree showing position of "Bacillus massiliogabonensis" strain Marseille P2639 relative to other phylogenetically closest species. Sequences were aligned using CLUSTALW and phylogenetic inferences were obtained using maximum-likelihood method within MEGA software. Numbers at nodes depict bootstrap percentages values obtained by repeating analysis 500 times to yield consensus tree. Only bootstrap values >95% were retained. Scale bar indicates 1% nucleotide sequence divergence.

delineate a new bacterial species without carrying out DNA-DNA hybridization, with a maximum error probability of 0.01%. Therefore, strain Marseille P2639 was putatively considered to be representative of a new bacterial species named "Bacillus massiliogabonensis" strain Marseille P2639 (Fig. 1).

The *Bacillus* genus was created in 1872 by Ferdinand Julius Cohn [8]. The closest neighbouring species to "Bacillus massiliogabonensis" sp. nov. by 16S rRNA gene sequence include *Bacillus acidicola* [6], *Bacillus eiseniae* [9], *Bacillus shackletonii* [10], *Bacillus homecckiae* [11] and *Bacillus subtilis* [12]. They are aerobic, catalase positive, motile and endospore forming. Of these species, Gram staining is negative for "Bacillus massiliogabonensis" sp. nov. and *B. eiseniae*. Oxidase activity is positive only for *B. subtilis* and *B. shackletonii*.

We propose the creation of "Bacillus massiliogabonensis" sp. nov. (named for Marseille and Gabon, the city and the country where the stool specimen was analysed and collected, respectively) as a new bacterial species. Strain Marseille P2639 is the type strain of this new species.

MALDI-TOF MS spectrum accession number

The MALDI-TOF MS spectrum of "Bacillus massiliogabonensis" is available at <http://www.mediterrane-infection.com/article.php?laref=256&titre=urms-database>.

Nucleotide sequence accession number

The 16S rRNA gene sequence of strain Marseille P2639 was deposited in GenBank database under accession number LT598571.

Deposit in culture collection

Strain Marseille P2639 was deposited in the Collection de Souches de l'Unité des Rickettsies (CSUR) and registered under number P2639.

Acknowledgements

We thank the Fondation Méditerranée Infection for financial support and K. Griffiths for English-language editorial work.

Conflict of Interest

None declared.

References

- [1] Lagier JC, Armougou F, Million M, Hugon P, Pagnier I, Robert C, et al. Microbial culturomics: paradigm shift in the human gut microbiome study. *Clin Microbiol Infect* 2012;18:1185–93.
- [2] Lagier JC, Hugon P, Khelaifia S, Fournier PE, La Scola B, Raoult D. The rebirth of culture in microbiology through the example of culturomics to study human gut microbiota. *Clin Microbiol Rev* 2015;28:237–64.
- [3] Mourembou G, Yasir M, Azhar El, Lagier JC, Bibi F, Jiman-Fatani AA, et al. Rise of microbial culturomics: non-contiguous finished genome sequence and description of *Beduini massiliensis* gen. nov., sp. nov. *Oncodes* 2015;19:766–76.
- [4] Mourembou G, Rathored J, Ndjoyi-Mbiguino A, Lekana-Douki JB, Fenollar F, Robert C, et al. Non-contiguous finished genome sequence

- and description of *Gabonia massiliensis* gen. nov., sp. nov. New Microbes New Infect 2015;9:35–44.
- [5] Mourembou G, Rathored J, Lekana-Douki JB, Ndjoyi-Mbigno A, Fenollar F, Robert C, et al. Non-contiguous finished genome sequence and description of *Kallipyga gabonensis* sp. nov. New Microbes New Infect 2015;9:15–23.
- [6] Albert RA, Archambault J, Rosselló-Mora R, Tindall BJ, Matheny M. *Bacillus acidicola* sp. nov., a novel mesophilic, acidophilic species isolated from acidic sphagnum peat bogs in Wisconsin. Int J Syst Evol Microbiol 2005;55(Pt 5):2125–30.
- [7] Meier-Kolthoff JP, Göker M, Spröer C, Klenk HP. When should a DDH experiment be mandatory in microbial taxonomy? Arch Microbiol 2013;195:413–8.
- [8] Skerman VBD, McGowan Vicki, Sneath PHA. Approved lists of bacterial names. Int J Syst Evol Microbiol 1980;30:225–420.
- [9] Hong SW, Park JM, Kim SJ, Chung KS. *Bacillus eiseniae* sp. nov., a swarming, moderately halotolerant bacterium isolated from the intestinal tract of an earthworm (*Eisenia fetida* L.). Int J Syst Evol Microbiol 2012;62(Pt 9):2077–83.
- [10] Logan NA, Lebbe L, Verhelst A, Goris J, Forsyth G, Rodríguez-Díaz M, et al. *Bacillus shackletonii* sp. nov., from volcanic soil on Candlemas Island, South Sandwich archipelago. Int J Syst Evol Microbiol 2004;54(Pt 2):373–6.
- [11] Vaishampayan P, Probst A, Krishnamurthi S, Ghosh S, Osman S, McDowall A, et al. *Bacillus horneckiae* sp. nov., isolated from a spacecraft-assembly clean room. Int J Syst Evol Microbiol 2010;60(Pt 5):1031–7.
- [12] Kampfer P. Whole-cell fatty acid analysis in the systematics of *Bacillus* and related genera. In: Berkeley R, Heyndrickx M, Logan N, De Vos P, editors. Applications and systematics of *Bacillus* and relatives. Oxford: Blackwell Science; 2002. p. 271–99.