




ELSEVIER

Contents lists available at [ScienceDirect](http://ScienceDirect)

## Virology

journal homepage: [www.elsevier.com/locate/yviro](http://www.elsevier.com/locate/yviro)

## Review

## viral silencing suppressors: Tools forged to fine-tune host-pathogen coexistence



Tibor Csorba\*, Levente Kontra, József Burgyán\*

National Agricultural Research and Innovation Center – Agricultural Biotechnology Center (NARIC – ABC), Gödöllő, Szent-Györgyi A.u.4., Pest 2100, Hungary

## ARTICLE INFO

## Article history:

Received 18 December 2014  
 Returned to author for revisions  
 31 January 2015  
 Accepted 16 February 2015  
 Available online 9 March 2015

## Keywords:

Plants  
 Viruses  
 Antiviral RNA interference  
 Viral suppressors of silencing  
 Defense  
 Pathogenesis

## ABSTRACT

RNA silencing is a homology-dependent gene inactivation mechanism that regulates a wide range of biological processes including antiviral defense. To deal with host antiviral responses viruses evolved mechanisms to avoid or counteract this, most notably through expression of viral suppressors of RNA silencing. Besides working as silencing suppressors, these proteins may also fulfill other functions during infection. In many cases the interplay between the suppressor function and other “unrelated” functions remains elusive. We will present host factors implicated in antiviral pathways and summarize the current status of knowledge about the diverse viral suppressors’ strategies acting at various steps of antiviral silencing in plants. Besides, we will consider the multi-functionality of these versatile proteins and related biochemical processes in which they may be involved in fine-tuning the plant-virus interaction. Finally, we will present the current applications and discuss perspectives of the use of these proteins in molecular biology and biotechnology.

© 2015 Elsevier Inc. All rights reserved.

## Contents

Plant viruses . . . . .	86
RNA silencing pathways in plants . . . . .	86
Antiviral silencing host factors . . . . .	86
Initiation of antiviral silencing . . . . .	86
Effector phase of antiviral silencing . . . . .	90
Amplification of antiviral silencing . . . . .	90
Actions of viral suppressors of RNA silencing . . . . .	91
Blocking initiation of antiviral response . . . . .	91
Arrest of functional RISC assembly through AGO interaction . . . . .	92
Inactivation of programmed antiviral RISC complex . . . . .	92
VSR activities downstream of RISC and RITS . . . . .	93
Modulation of AGO1 homeostasis . . . . .	93
Plant RDR-based activity suppression . . . . .	93
Targeting multiple steps of antiviral pathways . . . . .	93
VSR interactions with host factors . . . . .	93
Driving factors in VSRs’ evolution . . . . .	94
The control of pathogen impact on host . . . . .	94
Limitation of VSRs’ suppressor strength . . . . .	94
VSRs as links between RNA-based and protein-based immunity . . . . .	95
Connecting antiviral silencing to hormone signaling . . . . .	95
VSRs as tools . . . . .	96
Unraveling molecular basis of silencing itself . . . . .	96

\* Corresponding authors. Tel.: +36 28526100; fax: +36 28526101.

E-mail addresses: [csorba@abc.hu](mailto:csorba@abc.hu) (T. Csorba), [burgyan@abc.hu](mailto:burgyan@abc.hu) (J. Burgyán).

VSRs employed in molecular research . . . . .	96
VSRs as biotechnological and medical tools . . . . .	96
Conclusions and perspectives . . . . .	96
Acknowledgments . . . . .	97
References . . . . .	97

## Plant viruses

Plant viruses are amongst the most important pathogens causing huge economic losses worldwide by reducing crop quality and quantity. A better understanding of the viral infection processes and plant defense strategies is important for crop improvement.

Based on their genome organization viruses can be classified into positive-sense-, negative-sense-, double-stranded-RNA viruses and single-stranded or double-stranded DNA viruses (Hull, 2002). Difference in the genome organization implies difference in the replication strategy. Generally, the genetic information embedded into the viral RNA or DNA encode for a surprisingly restricted number of proteins that coordinate the infection process. Viral proteins interact with host factors to manipulate biochemical events and molecular interactions required for the virus replication and movement. Viruses can spread within the plants through plasmodesmata on short distance (cell-to-cell movement) or through phloem (systemic movement). During host-pathogen co-evolution a set of complex interactions involving virus attack and host defense has been developed. These include hypersensitive reaction (HR) (Mandadi and Scholthof, 2013), systemic acquired resistance (SAR) (Kachroo and Robin, 2013), activation of ubiquitin/26S proteasome system (UPS) (Dielen et al., 2010) or RNA silencing (RNA interference, RNAi) (Pumplin and Voinnet, 2013).

## RNA silencing pathways in plants

RNA silencing is a fundamental genetic regulatory mechanism conserved in eukaryotic organisms. RNAi can act at transcriptional (Transcriptional Gene Silencing, TGS) or at post-transcriptional levels (Post-Transcriptional Gene Silencing, PTGS), and has many diverse roles including developmental regulation, stress response or defense against invading nucleic acids like transposons or viruses. The antiviral function of RNA silencing was demonstrated in plants and invertebrates (Bronkhorst and van Rij, 2014; Pumplin and Voinnet, 2013), however recent reports have further provided evidence for a similar function in mammals (Cullen et al., 2013; Li et al., 2013; Maillard et al., 2013).

Mechanistically, the RNA silencing process consists of initiation phase, effector phase and amplification phase. During silencing initiation double-stranded RNAs (dsRNA) of different origins are processed by an RNase III type enzyme Dicer (DCR, in plants DICER-LIKE proteins, DCLs) into short, 21–24 nt long, small RNA (sRNA) duplexes (Bernstein et al., 2001; Hamilton and Baulcombe, 1999; Hutvagner et al., 2001). DICERs require DOUBLE-STRANDED RNA BINDING (DRB) proteins for accurate sRNA production (Eamens et al., 2012a,b; Hiraguri et al., 2005). The sRNAs are stabilized at their 3' end by the HUA Enhancer 1 (HEN1)-dependent methylation (a process found in plants and flies so far (Boutet et al., 2003; Yang et al., 2006) and exported from nucleus by HASTY (HST) (Park et al., 2005; Peragine et al., 2004) to be loaded onto Argonaute proteins (Fagard et al., 2000; Hammond et al., 2001; Liu et al., 2004), the effectors of the RNA-Induced Silencing Complex (RISC) (Lee et al., 2004; Pham et al., 2004; Tomari et al., 2004) or RNA Induced Transcriptional Silencing complex (RITS) (Ekwall, 2004). Guided by the sRNA sequence, RISC induces slicing or translational repression of its target RNAs (during PTGS) in a sequence-specific manner (Kim et al., 2014), whereas RITS complex causes histone and/


or DNA methylation, resulting in transcriptional gene silencing (TGS) of the homologous gene (Creamer and Partridge, 2011). In plants and worms the effector step can result in amplification of silencing response involving RNA-dependent RNA polymerases (RDRs) proteins (Mourrain et al., 2000; Dalmay et al., 2000; Sijen et al., 2001; Vaistij et al., 2002; Voinnet et al., 1998). Amplification of RNA silencing has been implicated in the spread of an RNA silencing signal, a non-cell-autonomous process (Kalantidis et al., 2008; Schwach et al., 2005).

The best studied plant model, *Arabidopsis thaliana* genome encodes 4 members of DCLs (DCL1–4) (Bologna and Voinnet, 2014), five DRBs (HYL1/DRB1, DRB2, 3, 4, 5) (Hiraguri et al., 2005), 10 AGOs (AGO1–10) (Mallory and Vaucheret, 2010) and 6 RDRs (RDR1, 2, 3a, 3b, 3c and 6) (Wassenegger and Krczal, 2006). These proteins have partially redundant roles and combine with each other to result in diverse classes of small RNAs and different effector outputs of the RNA silencing pathways. The small RNA classes identified in plants include microRNAs (miRNAs), transacting small interfering RNAs (ta-siRNAs), natural-antisense RNAs (nat-siRNAs), repeat-associated siRNAs (ra-siRNAs), viral siRNAs (vsiRNAs) and virus-activated siRNAs (vasiRNAs). These classes possess specialized roles during development, stress responses, heterochromatic silencing, viral infection and host-pathogen interplay, respectively (Bologna and Voinnet, 2014).

## Antiviral silencing host factors

### Initiation of antiviral silencing

The hallmark of the antiviral silencing response is the Dicer-dependent production of viral siRNAs (vsiRNAs) (Hamilton and Baulcombe, 1999). In uninfected cells long dsRNA is not detectable, however upon virus infection, viral dsRNA molecules of different sources becomes available. Highly structured regions of viral single-stranded RNAs (ssRNA), replicative intermediates (RI) or overlapping bidirectional read-through transcripts from DNA virus genome may all contribute to vsiRNA production (Aregger et al., 2012; Blevins et al., 2011; Donaire et al., 2008; Molnar et al., 2005). Although the viral dsRNA structures are likely accessible to all of the DCLs, a strong hierarchy exists between them regarding vsiRNA production: RNA virus infections are mainly affected by DCL4, while DCL2 becomes critical in a *dcl4* mutant background (Deleris et al., 2006; Donaire et al., 2008; Garcia-Ruiz et al., 2010; Qu et al., 2008) (Fig. 1). DCL3 has only a minor role against RNA viruses (Qu et al., 2008; Raja et al., 2008). Recent report suggests additional functional diversity between DCL4 and DCL2, such as that DCL2 stimulates transitivity and secondary siRNA production, while DCL4 is sufficient for silencing on its own (Parent et al., 2015). The fact that silencing suppressors of RNA viruses interfere with DCL3 pathway suggests it could have important antiviral gene regulatory functions (Azevedo et al., 2010; Hamera et al., 2012; Lacombe et al., 2010). DCL3 is essential against DNA viruses (Akbergenov et al., 2006) and works presumably by inducing DNA methylation (Blevins et al., 2006; Raja et al., 2014) (Fig. 1). Finally, DCL1 acts as a positive regulator in the production of vsiRNAs by making viral dsRNAs available to other DCLs both in RNA and DNA virus infections (Blevins et al., 2006; Moissiard and Voinnet, 2006) but also as a



**Fig. 1.** The model of antiviral RNA silencing and the diverse suppressor strategies that interfere with the pathway. Silencing response against RNA viruses is initiated by DCL4 and DCL2 through production of 21–22 nt viral siRNAs and to a lesser extent DCL3 (DCL1 that also has roles in antiviral defense was omitted for simplicity, please see main text). Viral siRNAs are methylated by HEN1 and subsequently loaded into AGOs. AGO1 and AGO2 are the two main effectors, but AGO5 and 7 may also have antiviral roles. Loaded-AGO containing RISC complexes complete the effector step of post-transcriptional gene silencing (PTGS) through viral RNA cleavage or translational inhibition (left side). AGO-sliced products, aberrant RNAs or aborted viral transcripts serve as templates for RDR complexes to amplify the antiviral response. SGS3 and SDE5 cofactors are required for RDR6 activity. (RDR1 and RDR2 also have antiviral functions, omitted for simplicity). RDR activities may be primed by DCL2-derived 22 nt vsiRNAs (middle). In DNA virus infections DCL4 and DCL3-derived viral siRNA initiate PTGS or transcriptional gene silencing (TGS) respectively. (DCL1 recognizes highly folded structures like the CaMV 35S leader, omitted for simplicity). DCL3-generated 24 nt vsiRNA following the HEN1-methylation are loaded into AGO4. AGO4-RITS causes the hypermethylation of viral genome to complete TGS (right side). DNA-virus transcripts are poor templates for RDR-based amplification process. The diverse viral suppressors inhibiting PTGS and/or TGS are shown in black boxes along the sides of the figure. The dashed lines point to the interference place where the VSRs interact with the antiviral pathway.

negative regulator limiting DCL4 and DCL3 through miRNA pathway (Azevedo et al., 2010; Qu et al., 2008). DCLs interact with DRBs to produce small RNAs. DCL4 partner DRB4 is important in antiviral

defense against RNA viruses *Turnip crinkle virus* (TCV) (Curtin et al., 2008) or *Turnip yellows mosaic virus* (TYMV) (Jakubiec et al., 2012). Evidence show that *Cauliflower mosaic virus* (CaMV) suppressor P6

**Table 1**  
The summary of identified VSRs, their mode of suppression and alternative functions. For suppressors encoded by different viruses within the genus and having the same name (e.g. cucumovirus 2b, potyvirus HC-Pro etc.) the alternative functions are displayed in a combined manner. For details on each individual function please see the references within.

Genus	Species	VSR	Function	Reference
Phytoreo virus	RDV	PNS10	siRNA binding, RDR6 downregulation	Ren et al. (2010) and Cao et al. (2005)
	RGDV	PNS11	Unknown, miRNA pathway interference	Shen et al. (2012) and Liu et al. (2008)
	RGDV	PNS12	Unknown (nucl. localization)	Guo et al. (2011) and Wu et al. (2011)
Oryza virus	RRSV	PNS6	Unknown	Wu et al. (2010)
Tospo virus	TSWV, GBNV	NSs	siRNA, miRNA and dsRNA binding, avirulence factor	Zhai et al. (2014), Ronde et al. (2014), Schnettler et al. (2010) and Goswami et al. (2012)
Cucumovirus	CMV, TAV	2b	sRNA binding, AGO1 binding, AGO4 binding, aphid interaction, RDR6 downregulation, AGO1 downregulation via miR168 upregulation, downregulation of AGOs and DCL1, impairing 5' secondary siRNA genesis, cell-to-cell movement, interaction with CAT3, induction of host drought resistance, SA/JA pathway disruption, HR elicitor	Nemes et al. (2014), Du et al. (2014), Gonzalez et al. (2012), Duan et al. (2012), Diaz-Pendon et al. (2007), Lewsey et al. (2007), Ahn et al. (2010), Chen et al. (2008), Zhang et al. (2006), Feng et al. (2013), Gonzalez et al. (2012), Hamera et al. (2012), Westwood et al. (2013a,b), Ziebell et al. (2011), Zhang et al. (2008), Inaba et al. (2011), Lewsey et al. (2010), Várallyay and Havelda (2013), Zhou et al. (2014), Ji and Ding (2001) and Li et al. (1999)
Illarivirus	AV-2	2b	Unknown (no local suppression)	Shimuran et al. (2013)
Como virus	CPMV	CP (S)	Unknown	Canizares et al. (2004)
Nepo virus	ToRSV	CP	Mediating AGO1 degradation	Karran and Sanfacon (2014)
Rymo virus	AgMV, HoMV	Hc-Pro	unknown	Young et al. (2012)
Poty virus	TEV, ZYMV, PVY, TuMV, SCMV, SMV, TVY, PVA, PRSV	Hc-Pro	ds-siRNA binding, blocking HEN1 methyltransferase, HEN1 binding, blocking primary siRNA biogenesis by RAV2 interaction, RDR6 downregulation, impairs 3' secondary siRNA genesis, JA pathway disruption, SAHH-interaction, Hip2 interaction, PaCRT interaction, ferredoxin-5 interaction, NtMinD interaction, interfering with miRNA pathway, interference with host gene expression levels, AGO1 downregulation via miR168 upregulation	Sahana et al. (2014), Torres-Barcelo et al. (2010), –Ruiz et al. (2010), Torres-Barcelo et al. (2008), Goto et al. (2007), Shibolet et al. (2007), Yu et al. (2006), Merai et al. (2006), Dunoyer et al. (2004), Llave et al. (2000), Mallory et al. (2001), Lozsa et al. (2008), Ebhardt et al. (2005), Jamous et al. (2011), Endres et al. (2010), Zhang et al. (2008), Seo et al. (2010), Westwood et al. (2014), Canizares et al. (2013), Haikonen et al. (2013a), Shen et al. (2010), Cheng et al. (2008), Jin et al. (2007), Chapman et al. (2004), Kasschau et al. (2003), Soitamo et al. (2011) and Várallyay and Havelda (2013)
	PPV PVA	HcPro-P1 VPg	Unknown SGS3 interaction	Valli et al. (2006) Rajamäki et al. (2014) and Rajamäki and Valkonen (2009)
Tritimo viruses	WSMV, ONMV	P1	Unknown	Young et al. (2012)
Poace virus	TriMV, SCSMV	P1	Unknown	Tatineni et al. (2012)
Ipomo virus	SPMMV	P1	AGO binding	Szabo et al. (2012) and Giner et al. (2010)
	CVYV	P1b	21 and 22 nt siRNA binding, protease activity, putativ Zn-finger	Valli et al. (2011) and Valli et al. (2008)
Tombus virus	CymRSV, CIRV, TBSV	P19	ds-sRNA binding, interfering with sRNA 3' methylation, mir168 upregulation mediated AGO1 downregulation, HR elicitor, Hin19 interaction, ALY interaction	Law et al. (2013), Rawlings et al. (2011), Cheng et al. (2009), Xia et al. (2009), Koukiekoloa et al. (2007), Merai et al. (2006), Lakatos et al. (2006), Omarov et al. (2006), Havelda et al. (2005), Dunoyer et al. (2004), Silhavy et al. (2002), Ye et al. (2003), Vargason et al. (2003), Lozsa et al. (2008), Yu et al. (2006), Chapman et al. (2004), Várallyay et al. (2010, 2014), Várallyay and Havelda (2013), Angel and Schoelz (2013), Hsieh et al. (2009), Park et al. (2004) and Uhrig et al. (2004)
	CNV	P20	Unknown	Hao et al. (2011)
Aureus virus	PoLV	P14	dsRNA binding	Merai et al. (2005, 2006)
Carmo virus	TCV	P38	AGO1 and 2 binding, DCL1 upregulation to antagonize DCL4 and DCL 3, dsRNA binding, blocking primary siRNA biogenesis by RAV2 interaction, TIP-interaction, DRB-HRT mediated HR elicitor, AGO1 downregulation via miR168 upregulation	Azevedo et al. (2010), Jin and Zhu (2010), Várallyay and Havelda (2013), Zhang J. et al. (2012), Merai et al. (2006), Endres et al. (2010), Donze et al. (2014), Choi et al. (2004), Ren et al. (2000), Zhu et al. (2013, 2014), Jeong et al. (2008) and Pérez-Cañamás and Hernández (2015)
	PFVB, HCRSV, PLPV	P37	siRNA binding	Martinez-Turino and Hernandez (2009), Meng et al. (2006) and Pérez-Cañamás and Hernández (2015)
	MNSV MNSV	P7B P42	Unknown, movement Protein Unknown	Genovés et al. (2011) and Genoves et al. (2006) Genoves et al. (2006)
Diantho virus	RCNMV RCNMV	replication MP	Unknown (DCL1 dependent), miRNA pathway interference Unknown	Takeda et al. (2005) Powers et al. (2008)
Clostero virus	BYV	P21	ds-sRNA binding, blocking HEN1 methyltransferase	Merai et al. (2006), Yu et al. (2006), Reed et al. (2003) and Chapman et al. (2004)
	CTV	P20	Unknown	Lu et al. (2004)

Table 1 (continued)

Genus	Species	VSR	Function	Reference
	CTV	P23	Unknown (nucleolar localization)	Ruiz-Ruiz et al. (2013) and Lu et al. (2004)
	CTV	CP	Unknown	Lu et al. (2004)
Crini virus	SPCSV, SPVD	RNase3	Endonuclease activity	Cuellar et al. (2009) and Kreuze et al. (2002, 2005)
	SPCSV	P22	Unknown	Kreuze et al. (2005)
	CYSDV	P25	Unknown (down stream of siRNA biogenesis)	Kataya et al. (2009)
	ToCV	P22	Unknown (local silencing)	Canizares et al. (2008)
	ToCV	CP	Unknown, SAHH-interaction	Canizares et al. (2008, 2013)
	ToCV	Cpm	Unknown	Canizares et al. (2008)
Ampelo virus	GLRaV-3	P19,7	siRNA and miRNA pathway interference	Gouveia and Nolasco (2012) and Gouveia et al. (2012)
Polero virus	BWYV, PLRV, CYDV, BMYV, TuYV, MABYV, ScYLV, CLRDV, CABYV	P0	Destabilizing AGOs, HR elicitor	Hendelman et al. (2013), Fusaro et al. (2012), Derrien et al. (2012), Kozłowska-Makulska et al. (2010), Han et al. (2010), Csorba et al. (2010), Mangwende et al. (2009), Bortolamiol et al. (2007), Pazhouhandeh et al. (2006), Delfosse et al. (2014) and Wang et al. (2014a,b)
Enamo virus	PEMV-1	P0	Destabilizing AGO1	Fusaro et al. (2012)
Tymo virus	TYMV	P69	Unknown (inhibition of DNA methylation, miRNA and DCL1 upregulation)	Chen et al. (2004)
	CVB	P12	Unknown (ZF domain dependent but NLS independent)	Lukhovitskaya et al. (2014)
Potex virus	PVX, PIAMV, AV3, WclMV, TVX, PepMV	P25 (TGBp1)	AGO1 degradation, coaggregation with SGS3/RDR6, CAT1 interaction, X-Body organization	Chiu et al. (2010), Yan et al. (2012), Senshu et al. (2009), Okano et al. (2014), Mathioudakis et al. (2013) and Tilsner et al. (2012)
	PepMV	CP	Unknown (blocks systemic signaling)	Mathioudakis et al. (2014)
Carla virus	PVM	TGBp1	Unknown (no local suppression)	Senshu et al. (2011)
	PVM, SPCFV, PIAMV	CRP (NaBp)	Unknown (local and systemic silencing)	Deng et al. (2014), Senshu et al. (2011) and Okano et al. (2014)
Tricho virus	ACLSV	P50	Unknown (no local suppression)	Yaegashi et al. (2007, 2008)
Vitivirus	GVA	P10	siRNA binding	Zhou et al. (2006)
Citri virus	CLBV	MP	Unknown	Renovell et al. (2012)
Tobamo virus	TMV, YoMV(ORMV)	P126	Unknown, interference with HEN1-mediated methylation, accumulation of novel miRNA-like sRNAs	Wang et al. (2012), Vogler et al. (2007), Ding et al. (2004) and Hu et al. (2011)
	ToMV	P130	siRNA binding	Kubota et al. (2003)
	TMV	P122	siRNA and miRNA binding, AGO1 downregulation via miR168 upregulation	Csorba et al. (2007) and Várallyay and Havelda (2013)
Tenui virus	RHBV, RSV	NS3	Binding to RNA/RNA or RNA/DNA duplex, larger than 9 nt and long ssRNA	Hemmes et al. (2007), Shen et al. (2010) and Xiong et al. (2009)
	RSV	P2	Interaction with SGS3	Du et al. (2011)
Tobra virus	TRV	16K	Downstream of dsRNA biogenesis	Andika et al. (2012), Ghazala et al. (2008), Martínez-Priego et al. (2008) and Reavy et al. (2004)
	TRV	29K	Unknown	Deng et al. (2013)
	PepRSV	12K	Unknown	Jaubert et al. (2011)
Furo virus	SBWMV, CWMV	19K	Unknown, interaction with N-ext/CP	Sun et al. (2013a,b) and Te et al. (2005)
Peclu virus	PCV	P15	siRNA binding, miRNA pathway interference	Merai et al. (2006), Dunoyer et al. (2002, 2004)
Beny virus	BNYVV, BSBMV	P14	Unknown	Andika et al. (2012), Guilley et al. (2009), Chiba et al. (2013) and Kozłowska-Makulska et al. (2010)
	BdMoV	P13	Unknown	Andika et al. (2012) and Guilley et al. (2009)
	BNYVV	p31	Unknown, interaction with PR-10	Rahim et al. (2007) and Wu et al. (2014)
Hordei virus	BSMV, PSLV	γB	siRNA binding	Merai et al. (2006) and Yelina et al. (2002)
Sobemo virus	CfMV, RYMV	P1	Unknown (siRNA binding ruled out), dcl4 dep.21 nt siRNA delocalization, reduction of 24 nt siRNAs, Zn-finger like binding	Lacombe et al. (2010), Gillet et al. (2013), Sarmientoa et al. (2007) and Weinheimer et al. (2010)
	CfMV	CP	Unknown, nuclear localization	Olsper et al. (2010, 2014)
Nucleo rhabdo virus	RYSV	P6	Blocking RDR6-mediated secondary siRNA biogenesis	Guo et al. (2013)
Curto virus	BCTV	L2	Methylation interference by ADK inhibition	Buchmann et al. (2009), Raja et al. (2008), Yang et al. (2007), Wang et al. (2003, 2005) and Hao et al. (2003)
Begomo virus	ACMV	AC4	ss-sRNA binding	Chellappan et al. (2005)
	CaLCuV, MYMV, ACMV	AL2/ AC2	Methylation interference by ADK inhibition, tran activation of host genes, interference with host gene expression levels	

Table 1 (continued)

Genus	Species	VSR	Function	Reference
	EACMCV	AV2	Unknown (downstream of siRNA biogenesis)	Buchmann et al. (2009), Raja et al. (2008), Yang et al. (2007), Wang et al. (2003, 2005), Hao et al. (2003), Trinks et al. (2005) and Soitamo et al. (2012)
	BSCTV, TYLCV-C, AYVV, ToLCJAV	C2	Methylation interference by SAMDC1 degradation	Chowda-Reddy et al. (2008)
	AYVV, CLCuMV, TYLCV, ToLCJAV-A	V2	Interaction with SGS3, HR elicitor, interaction with PLCPs	Zhang et al. (2011), Wezel et al. (2002), Sharma et al. (2010) and Kon et al. (2007)
	AYVV	C4	Unknown	Zhang J. et al. (2012), Glick et al. (2008), Sharma et al. (2010), Sharma and Ikegami (2010), Bar-Ziv et al. (2012), Zrachya et al. (2007) and Amin et al. (2011)
	ToLCJB, ToLCJAV, TYLCCNV, TYLCCNV-Y10, TbCSV-Y35, CLCuMV, GDARSLA, GmusSLA	$\beta$ C1	Unknown, rgs-CAM mediated RDR6 disruption, SAHH inhibition	Sharma et al. (2010), Sharma et al. (2011), Kon et al. (2007), Cui et al. (2005), Einia et al. (2012), Li et al. (2014) and Yang et al. (2011), Amin et al. (2011)
		Alpha-Rep	Unknown	Nawaz-ul-Rehman et al. (2010)
Mastre virus	WDV	Rep	siRNA binding	Wang et al. (2014a,b) and Liu et al. (2014)
	WDV	RepA	Unknown	Liu et al. (2014)
Caulimo virus	CaMV	P6 (TAV)	DRB4 inactivation, interference with NPR1 (SA/JA crosstalk regulator), cell-to-cell movement, translational trans activation	Laird et al. (2013), Haas et al. (2008), Shivaprasad et al. (2008), Love et al. (2007, 2012) and Rodriguez et al. (2014)
	CaMV	Decoy RNA	Overloading DCL capacity	Blevins et al. (2011)

protein binds to and inhibit DRB4 activity strongly suggesting for an antiviral role of DRB4 upon DNA virus infections (discussed latter). DCL3 cooperates with DRB3 (and AGO4) in antiviral defense through genome methylation against DNA viruses *Cabbage leaf curl virus* (CaLCuV) and *Beet curly top virus* (BCTV) (Raja et al., 2014). HEN1 is also required for the antiviral defense through methylation and stabilization of vsiRNAs (Vogler et al., 2007). Indeed, *Hen1* mutants are more susceptible to *Cucumber mosaic virus* (CMV) and TCV virus infections (Boutet et al., 2003; Zhang J. et al., 2012).

#### Effector phase of antiviral silencing

In theory the initiation phase of silencing could be enough to limit virus replication and spread through processing of the viral RNA into vsiRNAs. Dicing *per se*, however, is not sufficient to restrict or limit the viral infection (Wang et al., 2011), suggesting that the DCLs' substrates may be the byproducts or aborted transcripts of the viral replication process. It has been shown that the effector step of silencing involving AGO-dependent activity is required to restrict virus replication and spread (Wang et al., 2011). Indeed, AGO proteins are essential in antiviral defense against both RNA and DNA viruses (Azevedo et al., 2010; Pantaleo et al., 2007; Qu et al., 2008; Raja et al., 2008, 2014; Carbonell et al., 2012; Harvey et al., 2011; Wang et al., 2011). sRNA loading into AGOs is governed mostly by their 5' terminal nucleotides but length, thermodynamical properties of sRNA duplex ends and duplex structure are also important factors (Mi et al., 2008; Schuck et al., 2013; Schwarz et al., 2003; Zhang et al., 2014). *ago1* and *ago2* mutants are hypersusceptible to virus infections like CMV, *Turnip mosaic virus* (TuMV) or TCV (Carbonell et al., 2012; Harvey et al., 2011; Morel et al., 2002). AGO1 participate in removal of viral RNA through slicing activity (Carbonell et al., 2012), although translational repression activity was also found to play a role (Ghoshal and Sanfacon, 2014) (Fig. 1). It was shown that during RNA virus infections AGO1 homeostasis (Mallory et al., 2008) is disrupted and AGO1 protein levels are decreased probably through translational repression of AGO1 mRNA by miR168 activity (Varallyay et al., 2010). As AGO1 is the negative regulator of AGO2 through miR403 action in absence of AGO1 activity AGO2 levels are elevated (Azevedo et al., 2010; Harvey et al., 2011). AGO2 therefore emerges as a second layer in antiviral pathways. AGO2

was shown to be important in defense against CMV, TCV and *Potato virus X* (PVX) viruses in *A. thaliana* (Harvey et al., 2011; Jaubert et al., 2011) and against *Tomato bushy stunt virus* (TBSV) infection in *Nicotiana benthamiana* (Scholthof et al., 2011). In this scenario AGO1 is both a sensor of the infection that activates antiviral pathways such as AGO2 activity but also a direct effector of silencing. The phenotype of *ago1ago2* double mutant indicates that the two proteins act in a synergistic manner and have non-overlapping functions, as supported by their phylogenetic distance (Mallory and Vaucheret, 2010; Wang et al., 2011). Besides AGO1 and AGO2, AGO5 and AGO7 were also proposed to possess antiviral activities against RNA viruses (Qu et al., 2008; Takeda et al., 2008). AGO7 seems to work as a surrogate of AGO1 but with a preference for the less structured RNA targets (Qu et al., 2008; Takeda et al., 2008). The nuclear localized AGO4 has been shown to possess important antiviral functions against geminiviruses. *dcl3*, *drb3* and *ago4* mutants fail to hypermethylate the viral genome that is required for host recovery (Raja et al., 2014). Besides, AGO4 was proposed be important in transcriptional regulation of host transcriptional response during CMV virus infection (Hamera et al., 2012). The knowledge about RISC cofactors that cooperate with AGOs in plants is very limited. Heat shock protein 70 and 90 (HSP70, HSP90) have been found to be important players in AGO loading by using an *in vitro* cell-free system that recapitulates the loading process (Iki et al., 2010). Further understanding of RISC components, assembly and function may be helped by this and similar *in vitro* systems (Iki et al., 2010; Schuck et al., 2013).

#### Amplification of antiviral silencing

In many plant-virus combinations RDR activities contribute to the amplification of the antiviral response in order to achieve a robust defense response (Bologna and Voinnet, 2014; Wassenegger and Krczal, 2006) (Fig. 1). RDR polymerase activity is stimulated by the presence of aberrant RNAs lacking *bona fide* features like cap or polyA tail (Gazzani et al., 2004; Moreno et al., 2013). In some host-virus interactions (like during VSR-deficient CMV infection) sliced products of AGO1 or AGO2 were not required for secondary vsiRNA production (Wang et al., 2011) suggesting that the process may be different from that of ta-siRNA biogenesis. RDR-derived dsRNAs are

processed by DCL4 and DCL2 into 21–22 nt long vsRNAs, respectively. Both 21 and 22 nt long vsRNA are effective in antiviral response as has been shown in case of many virus infections (CMV, *Oilseed rape mosaic virus* (ORMV), TCV, *Tobacco rattle virus* (TRV), CaLCuV, CaMV) (Xie et al., 2004; Bouché et al., 2006; Deleris et al., 2006; Blevins et al., 2006; Donaire et al., 2008). DCL2 activity becomes more pronounced under higher temperature (Zhang X. et al., 2012b). However, Wang et al. (2011) have found that the 21- but not 22-nucleotide long vsRNAs guide efficient silencing through AGO1 and AGO2 effectors in CMV infection although AGO1 efficiently incorporates both of them. 22 nt long vsRNAs contribute to secondary siRNA production, as was shown for 22 nt long miRNAs and ta-siRNAs or mediate systemic silencing (García-Ruiz et al., 2010; Wang et al., 2011). It is assumed that the secondary vsRNA are able to move on short and long distances within the plant to immunize distant tissues ahead of the viral infection, however this awaits experimental validation. The requirement of RDR6-activity for systemic movement of a silencing signal suggesting that RNA silencing amplification has antiviral roles in uninfected distant tissues. RDR1, RDR2 and RDR6/SDE1/SGS2 were all found to be crucial factors in secondary vsRNA production during RNA virus (PVX, CMV, *Tobacco mosaic virus* (TMV), *Sugarcane mosaic virus* (SCMV), TuMV, TRV infections (Diaz-Pendon et al., 2007; Donaire et al., 2008; García-Ruiz et al., 2010; Qu et al., 2008; Schwach et al., 2005). Host RDR involvement in secondary vsRNA production could not be verified in certain virus-host interactions. Indeed, upon tombusvirus infections the overwhelming part of vsRNA derive from the positive RNA strand of the virus genome suggesting that they are primary DCL products (Aregger et al., 2012; Blevins et al., 2011; Donaire et al., 2008; Molnar et al., 2005; Szittyá et al., 2010). In case of DNA viruses, viral transcripts appear to be poor templates of RDRs. Majority of viral siRNAs accumulating during CaLCuV geminivirus infection were RDR1/2/6-independent primary siRNAs (Aregger et al., 2012).

RDR6 activity is facilitated by protein cofactors like Suppressor of GENE SILENCING 3 (SGS3) (Mourrain et al., 2000), SILENCING DEFECTIVE 5 (SDE5) (Hernandez-Pinzon et al., 2007) and SILENCING DEFECTIVE 3 (SDE3) (Dalmay et al., 2001). SGS3, a plant specific protein, was found associated with RISC complex and to be important in ta-siRNA biogenesis (Allen et al., 2005; Yoshikawa et al., 2005). It was proposed that SGS3 stabilize the RISC-cleavage product following slicing and enhance its conversion into dsRNA by RDR6 activity (Yoshikawa et al., 2013). Elimination of SGS3 not only abolishes ta-siRNA biogenesis but also leads to enhanced susceptibility to infection, at least in certain virus-host combinations. *sgs3* mutants have severe symptoms when challenged with CMV but do not show any difference in their viral symptoms during TuMV or *Turnip vein-clearing virus* (TVCV) infections (Adenot et al., 2006; Yoshikawa et al., 2013). SGS3 was shown to be required for CaLCuV virus induced VIGS of endogenous genes and further suggested that may be involved in the antiviral response against DNA viruses (Muangsan et al., 2004). This is supported by the fact that *Tomato yellow leaf curl virus* (TYLCV) encodes a VSR to compromise its activity (discussed below) (Glick et al., 2008; Fukunaga and Doudna, 2009). SDE5 is an RNA trafficking protein homologue of human mRNA export factor. It was proposed that SDE5 acts together with RDR6 in converting specific ssRNAs into dsRNA. *sde5* mutant plants are hypersusceptible to CMV but not TuMV infection (Hernandez-Pinzon et al., 2007). The amplification process is facilitated by the SDE3 an RNA-helicase like protein. SDE3 was shown to associate to AGOs through its GW motifs (García et al., 2012). *sde3* mutant plants are affected in defense response against CMV or PVX but not TRV infections (Dalmay et al., 2001). SDE3 activity occurs downstream to RDR6 and requires AGO1 and AGO2 activities (García et al., 2012). SDE3 was proposed therefore to facilitate the amplification process by unwinding a fraction of RDR6-synthetized dsRNA products using

its helicase activity. Furthermore, SDE3 could increase the efficiency of AGO-targeting of ssRNA targets (García et al., 2012).

### Actions of viral suppressors of RNA silencing

Some viruses avoid RNA silencing by replicating within well-defined subcellular compartments/structures like ER spherules (Schwartz et al., 2002) or by replicating and moving fast enough to outrun the mobile silencing signal. However, the most common way to protect viral genome against RNA silencing-mediated inactivation is to encode proteins that act as suppressors of RNA silencing (viral suppressors of RNA silencing, VSRs) (Lakatos et al., 2006). In fact, the strongest support of RNA silencing having antiviral roles was the discovery of VSRs. In an early work Ding and coworkers identified the cucumovirus 2b protein as responsible for induction of a non-conventional virus synergistic disease. Although at that time it was thought that 2b facilitate hypervirulence through its movement protein function (Ding et al., 1996), latter it was demonstrated to be a potent VSR (Brigneti et al., 1998). Another hint regarding viral suppression of plant RNA-based defense was the synergistic interaction of potyviruses with other viruses that relayed on potyviral Hc-Pro protein activity (Pruss et al., 1997). Whilst Hc-Pro was shown to block PTGS at tissue level, CMV 2b was shown to prevent systemic silencing (Brigneti et al., 1998). Since these first observations more than a decade ago, numerous VSRs were discovered and characterized (Table 1). Available evidences suggest that most viruses encode at least one VSR that in most cases is essential for successful viral life cycle. Although plant virus' VSRs are more studied, silencing suppression has been documented on insect and fungus-infecting viruses as well (Bronkhorst and van Rijn, 2014). The extraordinary diversity in sequence and structure of VSRs within and across kingdoms indicates that they have evolved independently. Diversity of VSRs implies diverse mechanistic activities, and indeed, VSRs were shown to block virtually all steps of RNA silencing (Fig. 1) such as dicing, effector assembly, targeting, amplification, transcriptional regulation of endogenous factors that control RNA silencing and its connections with protein-based immunity and hormone signaling.

### Blocking initiation of antiviral response

One strategy used by VSRs is to hinder mounting of antiviral silencing by blocking the silencing initiation step. This can be achieved through multiple ways like dicer protein or co-factor activity inhibition, dsRNA/siRNA-sequestration or AGO protein destabilization prior of RISC assembly. A widespread suppressor strategy is the ds-siRNA sequestration that is used by several VSRs encoded by diverse virus genera (P19, Hc-Pro, P21, p15, p130/p126/p122,  $\gamma$ B, NS3, Pns10, NSs etc.) (Csorba et al., 2007; Harries et al., 2008; Hemmes et al., 2007; Kubota et al., 2003; Lakatos et al., 2006; Merai et al., 2005, 2006; Silhavy et al., 2002) (Table 1 and Fig. 1). Probably the most characterized siRNA binder is the tombusviral p19 protein (Silhavy et al., 2002). Crystallographic studies have shown that p19 head-to-tail homodimer acts as a molecular caliper to size-select and sequester siRNA duplexes in a sequence-independent manner (Silhavy et al., 2002; Vargason et al., 2003). P19 siRNA sequestration prevents RISC assembly as shown by the heterologous *in vitro* Drosophila embryo extract system (Lakatos et al., 2006). Similarly p19 sequestration of miRNAs is efficient to prevent RISC-loading in p19-transgenic *A. thaliana* (Schott et al., 2012) and *N. benthamiana* plants (Kontra and Burgyan unpublished results). However, during authentic virus infections miRNA sequestration by p19 is not efficient (Lozsa et al., 2008), suggesting that, miRNA binding by P19 may depend on spatial and temporal co-expression of miRNA duplex and the virus encoded suppressor protein (Lozsa et al., 2008; Schott et al., 2012). A consequence of siRNA binding by VSRs is the block of HEN1-

dependent methylation of sRNAs (Csorba et al., 2007; Lozsa et al., 2008; Vogler et al., 2007), however this also depends on the co-expression of sRNA and the suppressor (Lozsa et al., 2008). In addition to blocking silencing activation at cellular level, p19 interferes with the systemic spread of mobile silencing signal as well (Dunoyer et al., 2010; Molnar et al., 2010). This latter characteristic is an excellent indicator whether a VSR acts indeed as a *bona fide* siRNA sequester during authentic viral infection. Some VSRs bind dsRNAs in a size-independent manner: *Pothos latent aureusvirus* (PoIV) P14, TCV p38 and CMV 2b (Deleris et al., 2006; Goto et al., 2007; Merai et al., 2005) have been all described to bind dsRNA and suggested therefore to block vsiRNA maturation. In turn nuclear localized P6 suppressor of CaMV diminish dicing efficiency through protein-protein interaction. The two importin- $\alpha$  dependent nuclear localization signals of P6 are mandatory for CaMV infectivity. P6 genetically and physically interacts with the nuclear DRB4, a cofactor required for DCL4-dependent vsiRNA processing (Haas et al., 2008). (TCV p38, CMV 2b and CaMV P6 are discussed latter). Similar strategy was described in insect infecting viruses: *Flock house virus*, *Nodamura virus* and *Wuhan nodavirus* encoded B2, *Drosophila C virus* (DCV) encoded canonical dsRBD, the VP3 proteins of *Drosophila X virus* or the dsDNA virus *Invertebrate iridescent virus 6* (IIV-6)-encoded 340R suppressors can block Dicer-2 activity and prevent RISC loading via size-independent dsRNA/siRNA-binding (Bronkhorst and van Rij, 2014).

A completely different strategy (to siRNA-binding) but with very similar outcome is used by *Sweet potato chlorotic stunt crinivirus* (SPCSV) suppressor RNase3. In this case the build-up of an efficient antiviral silencing complex is prevented by endonuclease activity of RNase3 that cleaves the 21–24 nt vsiRNAs into 14 bp products rendering them inactive (Cuellar et al., 2009; Kreuze et al., 2005). Insect infecting *Heliothis virescens ascovirus-3e* orf27 (RNase3 enzyme) suppressor works in analogous manner: it competes with Dcr-2 for dsRNAs and degrades dsRNAs and siRNAs (Hussain et al., 2010).

#### Arrest of functional RISC assembly through AGO interaction

The arrest in the assembly of a functional RISC can be carried out also through direct binding the protein component of minimal RISC, AGO protein. P0 protein (Mayo and Ziegler-Graff, 1996; Sadowy et al., 2001) the suppressor of *Ploverviruses* does not possess RNA binding activity (Csorba et al., 2010; Zhang et al., 2006), instead interacts with E3-ligase S-phase kinase regulated protein 1 (SKP1) through its F-box motif. P0 was shown to enhance the degradation of multiple AGOs (AGO1, 2, 4–6, 9) before holo-RISC assembly. Mutations in the F-box motif abolished P0 suppressor activity (Baumberger et al., 2007; Bortolamiol et al., 2007; Csorba et al., 2010; Derrien et al., 2012; Pazhouhandeh et al., 2006). These suggested a model where P0 destabilizes AGOs through the proteasome pathway. Instead, it was shown that P0-mediated AGO destabilization is not sensitive to proteasome inhibitor and that AGO degradation occurs through autophagy pathway (Baumberger et al., 2007; Bortolamiol et al., 2007; Csorba et al., 2010; Derrien et al., 2012; Pazhouhandeh et al., 2006). Massive accumulation of K48-linked ubiquitinated host proteins accumulated in the presence of P0, including the N-terminal cleavage product of AGO1. P0 therefore may have a role in promoting degradation of AGOs and other host proteins through ubiquitination-proteasome pathways (Csorba et al., 2010). *Tomato ringspot virus* (ToRSV) suppressor CP binds to AGO1 to suppress its translational inhibitory activity and to enhance AGO1 degradation through autophagy (Karran and Sanfacon, 2014). It was shown that PVX p25 physically interacts with AGOs (AGO1, 2, 3 4 but not AGO 5 and AGO9) to promote their destabilization in a proteasome-dependent manner. Consistently with these, plants treated with proteasome inhibitor were less susceptible to PVX (Chiu et al., 2010).

#### Inactivation of programmed antiviral RISC complex

VSRs may mimic cellular protein cofactors to inactivate programmed RISC. In *Sweet potato mild mottle ipomovirus* (SPMMV) the role of suppressor is played by P1 protein, a serine protease, in spite of the presence of HC-Pro (the suppressor of *Potyvirus*). P1 interacts directly with siRNA and/or miRNA-loaded AGO1 present in the high molecular weight holo-RISC but not minimal-RISC through GW/WG-motifs (AGO-hook). The conserved GW/WG-motif containing protein family (GW182 family) has been shown to bind to AGOs and to be required for diverse RISC function (Eulalio et al., 2009). Site-directed mutagenesis in the P1 protein proved that the conserved AGO-hook motifs (three GW/WG domains) located at the N-terminal part of P1 are absolutely necessary for both binding and suppression of AGO1 function (Giner et al., 2010). SPMMV P1 not just inhibits de novo RISC assembly but also block si/miRNA-loaded RISC activity. The role GW/WG-motif in silencing suppression was further evidenced by the restoration of naturally inactive P1 protein suppressor activity through the introduction of two additional GW/WG motifs (Szabo et al., 2012).

Coat protein of TCV also known as p38 may suppress silencing at multiple levels including vsiRNA generation and assembled RISC activity block. Since p38 possesses dsRNA-binding activity, and in its presence siRNAs are undetectable it was proposed that p38 acts to suppress Dicer's activity (Qu et al., 2003). Genetic evidences supported the role of p38 in inhibiting DCL4 but not DCL2 (Deleris et al., 2006). Although 22 nt vsiRNA were produced, they were inactive in the presence of p38 because  $\Delta$ p38 mutant TCV was not able to infect systemically the single *dcl*, *dcl2dcl3* or *dcl3dcl4* double mutant plants except *dcl2dcl4* mutant (Deleris et al., 2006). In a later study however, p38 suppressor impact on DCL4 was attributed to an indirect effect of AGO1-mediated DCL-homeostasis and has been shown that p38 blocks AGO1 but not AGO4 activity through its GW-motif binding. DCL2-derived 22 nt vsiRNA production, HEN1 and AGO2-activity pathway is crucial against TCV infection at elevated temperature (Zhang X. et al., 2012). Of note, p38 is also capable to bind and inactivate AGO2 (Zhang X. et al., 2012). P38 interacts with unloaded AGOs. Site-directed mutagenesis (GW-to-GA) in the p38 proved that GW motif is absolutely required for both binding and suppression of AGO1 function. GA mutation was enough to abolish TCV virulence that is restored in *ago1* hypomorph plants showing GW motif functionality during authentic virus infection (Azevedo et al., 2010). The coat protein (p37) of *Pelargonium line pattern virus* (PLPV) is a GW-containing protein that also functions as a VSR. It was shown that P37 mainly suppress silencing through siRNA sequestration. Mutations within its GW-motif concurrently affected p37 localization, its interaction with AGO1 and its sRNA-binding ability. Furthermore, binding assays have shown that also in case of TCV p38, the GW-mutation abolished p38 sRNA and long dsRNA-binding capacity (Pérez-Cañamás and Hernández, 2015). The overlapping multiple functions of p38 may lead to misinterpretation of experimental data. However, the data may suggest that these functions could cooperate: VSR interaction to AGO could enhance sRNA duplex sequestration in order to prevent RISC programming.

Silencing effector complex activity block can be achieved also through targeting holo-RISC's RNA component, the guide RNA. *African cassava mosaic virus* (ACMV) encoded AC4 was shown to bind to the ss-sRNAs but not dsRNA forms *in vitro*. Transgenic expression of AC4 correlated with decreased accumulation of miRNAs and upregulation of target mRNAs. AC4 acts downstream of the unwinding process: to bind mature miRNAs presumably loaded into AGO protein (Chellappan et al., 2005; Xiong et al., 2009; Zhou et al., 2006). *Rice stripe virus* suppressor NS3 is able to suppress and reverse GFP silencing and also prevent long distance spread of silencing signal. NS3 was found to bind to various RNA



forms like ss-siRNA, ds-siRNA or long ssRNA but not long dsRNA (Chellappan et al., 2005; Xiong et al., 2009; Zhou et al., 2006). *Grapevine virus A* p10 suppressor was also suggested to act through RNA sequestration: recombinant p10 was able to bind to both ss- and ds-si/miRNA species (Chellappan et al., 2005; Xiong et al., 2009; Zhou et al., 2006).

#### VSR activities downstream of RISC and RITS

VSRs may inactivate host defense downstream to RISC or RITS. Several DNA viruses encode VSRs that have been described to alter DNA/histone methylation, the effector step of TGS. *Tomato golden mosaic virus* (TGMV) suppressor AL2 and *Beet curly top virus* (BCTV) suppressor L2 inhibit adenosine kinase (ADK) activity that plays crucial role in adenosine and methyl-cycle maintenance or cytokinin regulation. In the presence of AL2 and L2 global reduction in cytosine methylation was observed that leads to inactivation and reversal of antiviral silencing (Buchmann et al., 2009; Wang et al., 2003, 2005). *In vitro* methylated TGMV cannot replicate in protoplasts suggesting that viral genome methylation is a bona fide defense against geminiviruses that is worth to be suppressed (Bisaro, 2006). Similarly, *Tomato yellow leaf curl China virus* (TYLCCNV) another *Begomovirus* encodes  $\beta$ C1 that interacts and inhibit activity of S-adenosyl-homocystein-hydrolase (SAHH) that is involved in methyl-cycle and therefore indirectly affects TGS (Yang et al., 2011).

#### Modulation of AGO1 homeostasis

Similarly to the aforementioned examples of DNA virus encoded VSRs, RNA virus-encoded VSRs were also described to modulate host gene expression on transcriptional level to their benefit. During tombusviral infection AGO1 transcription is induced as part of the host antiviral arsenal. AGO1 homeostasis in plants depends on the miR168-guided AGO1 mRNA cleavage and translational inhibition (Rhoades et al., 2002). To counteract AGO1-based defense, the virus promotes miR168 transcriptional induction that results in miR168-guided AGO1 down-regulation. The miR168 accumulation spatially correlates with the virus localization and depends on its p19 VSR (Varallyay et al., 2010) (Fig. 1). Similarly to p19 all VSRs, which are very heterogeneous in protein sequence but bind vsRNA, promote miR168 transcriptional induction and AGO1 down-regulation suggesting that VSR-siRNA complexes are effectors and recognized by the plant surveillance system (Várallyay and Havelda, 2013).

#### Plant RDR-based activity suppression

Host RDRs (RDR1, 2 and 6) contribute to amplification of RNA silencing and spread of a systemic signal by synthesis of vsRNAs (Schwach et al., 2005). Interestingly, plant RDR1 itself was suggested to have adverse functions. RDR1 is an antagonist of RDR6-mediated sense-PTGS silencing therefore behaves as an endogenous silencing suppressor (Ying et al., 2010). Suppression of RDR activities may constitute a target point for VSRs since it dampens cell-autonomous silencing amplification and systemic signal movement in distant tissues to facilitate the virus replication and spread. It was shown that V2 protein of *Tomato yellow leaf curl virus* (TYLCV) directly interacts with SGS3, the cofactor of RDR6, to block silencing amplification (Glick et al., 2008). Another *in vitro* study has shown that V2 competes with SGS3 for dsRNA having 5' overhang ends that may be an RDR6/SGS3 intermediate in vsRNA amplification (Fukunaga and Doudna, 2009; Kumakura et al., 2009). This may suggest that a 5'-overhanged intermediate may be the RDR6-complex template. Similar structures may be present on viral RNAs. Similarly, potyviral TRIPLE GENE BOX PROTEIN1 (TGBp1) was also shown to inhibit RDR6/SGS3-dependent dsRNA synthesis (Okano et al., 2014).  $\beta$ C1

suppressor of *Tomato yellows leaf curl China virus* (TYLCCNV) DNA satellite interacts with the endogenous suppressor of silencing calmodulin-like protein (rgsCAM) in *N. benthamiana* to repress RDR6 expression and secondary siRNA production (Li et al., 2014). *Sugarcane mosaic virus* (SCMV) encoded HC-Pro and TAV 2b were shown to downregulate RDR6 mRNA in *N. benthamiana* transient assay (Zhang et al., 2008). Pns10 suppressor of *Rice dwarf phyto-reovirus* (RDV) downregulate RDR6 to enhance viral invasion of shoot apices (Ren et al., 2010). RDR6-based activity suppression therefore is a widely used strategy that effectively blocks antiviral silencing but has only a limited impact on endogenous silencing pathways (discussed latter).

#### Targeting multiple steps of antiviral pathways

VSRs may act at multiple points to suppress silencing (Fig. 1). There is substantial evidence that the cucumoviral 2b protein can interfere with silencing at many steps. 2b prevents the spread of long-range silencing signal (Guo and Ding, 2002). N-terminal domain of 2b contains a dsRNA-binding domain that exhibits high affinity for short and long dsRNAs (Duan et al., 2012). CM95R strain of CMV and the related *Tomato aspermy virus* (TAV) 2b were also shown to bind siRNAs and ds-miRNAs *in vivo* and *in vitro* (Chen et al., 2008; Gonzalez et al., 2012; Goto et al., 2007). 2b (Fny and SD strains) was found to interact with AGO through the PAZ- and partly PIWI domains and blocks AGO1/RISC slicer activity (Duan et al., 2012; Zhang et al., 2006). FnyCMV 2b transgenic expression phenocopies *ago1* mutant plants (Duan et al., 2012; Zhang et al., 2006). Additionally, CMV 2b encoded by the SD strain alters RdDM pathway as well. 2b facilitates cytosine methylation through the transport of siRNAs into the nucleus (Kanazawa et al., 2011). 2b interacts both with AGO4-related siRNAs and with AGO4 protein through PAZ and PIWI domains. Interaction of 2b with AGO4 reduces AGO4 access to endogenous target loci and consequently modulates endogenous transcription to create a favorable cellular niche for CMV proliferation (Duan et al., 2012; Gonzalez et al., 2010, 2012; Hamera et al., 2012).

#### VSR interactions with host factors

There are emerging evidences that besides the “canonical” block of RNA silencing (through ds-, si-, mi-ssRNA-binding e.g. p19, RNase3 etc. or manipulating silencing-related protein activities via direct/ indirect interactions e.g. P0, V2, P1 etc.) some suppressors may target endogenous regulators of the silencing to modulate host defense. Potyviral helper-component protease (HC-Pro) is a multifunctional protein involved in many aspects of virus infection (Anandalakshmi et al., 1998; Carrington et al., 1989; Guo et al., 2011; Kasschau et al., 1997; Lakatos et al., 2006; Mallory et al., 2001). *Tobacco etch virus* (TEV) HC-Pro suppresses silencing through vsRNA-sequestration (Lakatos et al., 2006) and interferes with vsRNA methylation (Lozsa et al., 2008). *Zucchini yellows mosaic virus* (ZYMV) Hc-Pro interacts with HEN1 directly in *in vitro* assays (Jamous et al., 2011). TEV HC-Pro was found to interact with rgsCAM in a yeast two-hybrid system. RgsCAM itself is a host suppressor of RNA silencing (Li et al., 2014; Anandalakshmi et al., 2000; Endres et al., 2010). Interestingly, geminivirus AL2 protein induces expression and interacts with rgsCAM. Overexpression of rgsCAM leads to increased virus susceptibility (Yong Chung et al., 2014). Contradictory, in another study it was shown that rgsCAM counteracts HC-Pro through binding to its positively charged dsRNA-binding surface. RgsCAM binding prevents HC-Pro ds-siRNA binding activity and promotes its degradation through autophagy pathway (Nakahara et al., 2012). Multiple host interactors may further modulate activities of HC-Pro. It was proposed that RAV2, a transcription factor is required for suppression of

silencing mediated by TuMV HC-Pro. RAV2 downstream targets include *FIERY1* an endogenous silencing suppressor in *Arabidopsis* (Gy et al., 2007) and *CML38*, the likely homologue of *rgsCAM* in *Arabidopsis* (Anandalakshmi et al., 2000; Endres et al., 2010). Genome expression analysis reveals that RNA silencing related genes were unaltered in HC-Pro transgenic lines, instead RAV2 was required for HC-Pro-mediated induction of stress and defense-related genes (Endres et al., 2010). The unrelated *Carmoviral* p38 had a very similar effect on RAV2-mediated changes. These findings may suggest that RAV2 is a cross-talk point between pathogen stress-defense and silencing pathways that is used by VSRs to manipulate host reactions. *Papaya ringspot virus* (PRSV) HC-Pro interacts with papaya calreticulin to modulate host defense to virus infection through calcium signaling (Shen et al., 2010a; Shen et al., 2010b). HC-Pro of *Potato virus A* (PVA), *Potato virus Y* (PVY) and TEV interacts also with microtubule-associated protein (HIP2) through its highly variable region (HVR). HIP2 depletion reduces virus accumulation, whereas mutations affecting HC-Pro HVR domain induces necrosis and ethylene- and jasmonic acid-mediated systemic induction of host pathogen-related defense genes (Haikonen et al., 2013a,b).

Another example of how the endogenous factors may be used to modulate silencing efficiency is provided by *Red clover necrotic mosaic virus* (RCNMV). RCNMV recruits DCL enzymes into its replication complex and therefore deprives them from the silencing machinery. *dcl1* mutant plants show reduced susceptibility to RCNMV (Takeda et al., 2005). A similar strategy is used by CaMV pararetrovirus: massive amounts of vsRNAs derive from the 35S leader sequence produced by all four DCLs. These do not restrict viral replication but instead may serve as decoy RNAs to divert the effectors of the silencing machinery from important viral features like promoter and coding sequences (Blevins et al., 2011) (Fig. 1).

#### Driving factors in VSRs' evolution

The high diversity in structure and function, the various position of their gene-code within the viral genome, the alternative expression strategies like transcriptional read-through, leaky ribosomal scanning, proteolytic maturation and being often encoded by out-of-frame ORFs within conserved viral genes suggests that VSRs are of recent evolutionary origin (Ding and Voinnet, 2007). Therefore, in most cases, the suppressor function of VSRs may have evolved after the ancient role as replicase, coat protein, movement protein, protease, transcriptional regulator etc. or co-evolved with these to combine within the suppressor role and other essential roles important for viral life cycle. The different VSRs can inhibit all steps of the antiviral RNA silencing pathway, including cell-autonomous and non-cell autonomous aspects of it. Using mathematical modeling of dynamics of suppression has been shown that the different strategies employed result in slightly different outcomes regarding suppression of antiviral silencing. Suppressors targeting effector step are more potent at single cell level whereas siRNA binding is more effective at tissue level (Groenenboom and Hogeweg, 2012). Besides this however, an important driving factor in the suppressors' evolution was probably the availability of ancient/original viral protein activities that could be selected from with a minimum number of changes to acquire an additional suppressor features. Many of viral proteins have RNA binding capacity (like replicase, coat protein, movement protein). This may explain why an overwhelming number of suppressors act through RNA binding. Silencing functions could have become established in the cases when the tradeoff between its positive effect on viral life cycle and negative effects on host were worth it. The suppressors being too weak or too strong were out-selected through evolution.

#### The control of pathogen impact on host

Antiviral and endogenous silencing pathways share common elements. The ability of viruses to block antiviral silencing may have an impact on endogenous silencing pathway that results in alteration in short RNAs expression profile/activity and changes in gene expression both in a direct and in an indirect manner. vsRNA-binding VSRs can bind endogenous si- and miRNAs that could result in alteration of their downstream targets as was shown previously (Chapman et al., 2004; Kasschau et al., 2003; Lozsa et al., 2008). In case of miRNAs that target RNA silencing target components an unpredicted number of genes will be altered indirectly (e.g. miR162-mediated DCL1 negative feed-back loop, DCL1-dependent suppression of DCL3 and DCL4, miR168 and AGO1 mRNA-derived siRNA control of AGO1, miR403 control of AGO2) (Allen et al., 2005; Mallory and Vaucheret, 2009; Qu et al., 2008; Rajagopalan et al., 2006; Vaucheret et al., 2006; Xie et al., 2005). The situation is similar in case of AGO-targeting VSRs (P0, P1, P38) (Azevedo et al., 2010; Baumberger et al., 2007; Derrien et al., 2012; Giner et al., 2010). An elegant demonstration of this was described for TCV p38. AGO1 quenching by p38 had a profound impact on DCLs' homeostasis uncovering the strong interconnection of the silencing components into a functional network (Azevedo et al., 2010). VSRs' presence therefore may have a big impact and lead to an altered developmental program of host organism and symptom development. In support of VSRs as contributors to the viral symptoms, VSR-transgenic lines were created and analyzed. In many cases the VSR-expressing transgenic plants display phenotypes similar to viral infections (Dunoyer et al., 2004; Jay et al., 2011; Kasschau et al., 2003; Lewsey et al., 2007; Zhang et al., 2006). However, transgenic expression of VSR does not recapitulate the expression pattern in time and space of an authentic viral infection, therefore conclusions need to be drawn very carefully.

#### Limitation of VSRs' suppressor strength

Some of the VSRs differentially impact the antiviral and the endogenous pathways. For example vsRNA binding *Tombusviral* p19 protein binds more efficiently to free ds-siRNA forms. It was shown previously that p19 blocks very efficiently HEN1-dependent methylation of vsRNA but not miR159 (Lozsa et al., 2008). Furthermore, vsRNA but not miR159 are bound into p19-dimer:siRNA nucleoprotein complex whereas miR159 incorporates efficiently to AGO/RISC complex (Varallyay et al., 2010). The miR168, a particular miRNA that has a weak AGO-loading rate (Mallory and Vaucheret, 2009), was also available for *in vivo* p19-binding (Varallyay et al., 2010). All these suggests that p19 affects more potently vsRNA- but at less extent miRNA-pathway (Lozsa, Kontra and Burgyán, unpublished.). These differences may rise probably from the differences in the DCL1-dependent and DCL4/2-dependent si/miRNA-maturation pathways and distinct sub-cellular localization of silencing components. Conversely to p19, the TEV HC-Pro significantly affected 3'-methylation of both si- and miRNAs. miRNA 3'-methylation may take place in the nucleus where HC-Pro could access them (Lozsa et al., 2008). Similarly, TMV suppressor localizes to the nucleus as well (dos Reis Figueira et al., 2002). This may explain the more efficient sequestration of ds-miRNAs as assessed by the miRNA-star strand accumulation (Csorba et al., 2007). Therefore modulating sub-cellular localization of VSR will result in differential impact on antiviral and endogenous pathways respectively and therefore may constitute a regulatory point (Papp et al., 2003).

P6 is mostly present in the cytoplasm consistent with its transactivator function, but a small fraction of it is imported into the nucleus (where it blocks DRB4). It was shown that cytoplasmic-nuclear shuttling is prerequisite for VSR function and successful virus infection

(Haas et al., 2008). Although P6 contains a dsRNA-binding domain, this is not required for its suppressor function. In fact, perhaps surprisingly, the dsRNA-binding domain has an indirect negative effect on P6 suppressor function, presumably because nucleoplasmic localization of P6 is decreased by the dsRNA-binding domain interaction with ribosomal proteins L13 and L18 in the nucleolus and consequently P6 localizes more effectively to the nucleoplasm. Sub-cellular localization of P6 therefore alters its impact on host silencing. Similarly, enhanced nuclear targeting of CMV Fny strain 2b by addition of a nuclear localization signal compromises its suppressor activity but in the same time enhances 2b-mediated pathogenicity and CMV virulence, making the virus necrotic and accentuating the disease symptoms in *Arabidopsis*. Enhanced virulence was unrelated to effects of nuclear localized 2b on vsi- or miRNA-regulated target functions (Du et al., 2014). Consequently, CMV Q strain 2b mild suppressor activity could be explained by its preferential accumulation in the nucleus in contrary to cytoplasmic Fny 2b.

The suppression of antiviral pathways could additionally also protect the host endogenous targets from the vsiRNA-“off-target” effects. Recently it was shown that in the absence of 2b suppressor of CMV virus-activated siRNAs (vasiRNAs) are produced from endogenous transcripts in a DCL4- and RDR1-dependent manner. AGO2 is essential for the silencing activity of vasiRNAs (Cao et al., 2014).

Another VSR strength-limitation strategy is exemplified by the *Poleroviruses*. *Poleroviral* P0 protein expression is restricted by its suboptimal translation initiation codon, even though the use of such a weak suppressor-strategy results in phloem restriction of the virus. Attempts to restore optimal P0 translation lead to secondary mutations indirectly affecting P0 (Pfeffer et al., 2002). This shows that P0's impact on endogenous pathways is so devastating that is unfavorable for the virus.

Suppressor strength of HC-Pro was tested on a collection of point mutants in a GFP co-agroinfiltration assay and parallel the virulence of mutant alleles encoding TEV viruses *in vivo* (Torres-Barcelo et al., 2008). Both hyposuppressor and hypersuppressor HC-Pro alleles were found suggesting that the wild type protein has intermediate suppressor activity strength. Interestingly, while viruses carrying hyposuppressor alleles induced milder symptoms and accumulated to a lower level, hypersuppressor mutant viruses produced symptoms and accumulated to levels characteristic to wild type virus (Torres-Barcelo et al., 2008). This suggests that the suppressor strength could have been optimized by natural selection.

Since antiviral pathways strongly relay on amplification and systemic spread of silencing, those suppressors that target amplification of silencing (e.g. V2 or TGBp1 inhibition of SGS3) will have a definite impact to fight off antiviral pathway and promote virus spread (Fukunaga and Doudna, 2009; Glick et al., 2008; Kumakura et al., 2009; Okano et al., 2014). In certain cases however, in order to keep a tight control and spare the host, virus' systemic spread is limited by enhancing systemic silencing signal by the pathogen itself. *Rice yellow mottle virus* (RYMV) P1 was described originally as the movement protein of the virus. Recently it was shown that P1 suppresses local transgenic silencing and inhibits DCL4-dependent endogenous siRNA pathways in *Nicotiana* plants and its expression caused similar developmental defects in rice as *dcl4* mutation. In the same time P1 facilitated the short- and long distance spread of silencing signal in *Nicotiana* plants. P1 therefore may have a dual role, both to suppress and to elicit silencing of host (Lacombe et al., 2010). Although the precise mechanism of RYMV P1 activities are not known and need to be confirmed during authentic virus infections, it seems that P1 plays a fundamental role for an efficient infection process where the virus preserves the integrity of the host. Similarly to RYMV movement protein P1, the TMV MP (30kDa) has been shown to support the spread of antiviral silencing signal (Vogler et al., 2008) and therefore counter-balance the activity of p130 the suppressor of TMV to attenuate pathogen impact (Kubota et al., 2003).

Decreasing rather than eliminating specific antiviral silencing activities may allow setting a balance between blocking antiviral pathway and host transcriptome control through silencing. CaMV P6 protein was described as a symptom and host range determinant, a translational transactivator and a silencing suppressor (Bonneville et al., 1989; Haas et al., 2008; Love et al., 2007). P6 suppressor blocks DRB4 function that is required for DCL4 function therefore only diminishes (but not blocks) DCL4 activity. This maybe sufficient enough to dampen antiviral silencing but in the same time allow host gene control by the 21 nt 35S-derived vsiRNA. It was shown previously that several CaMV vsiRNAs exhibit near-complete complementarity to many endogenous targets (Moissiard and Voinnet, 2006). Besides, DRB4 targeting may allow accumulation of 24 nt vsiRNA that are required for heterochromatin silencing of CaMV minichromosomes (Al-Kaff et al., 1998).

#### VSRs as links between RNA-based and protein-based immunity

Alteration of silencing pathways (an RNA-based immunity) in the presence of the VSR and/or viral infection triggers the protein-based immunity in host as part of the counter-counter defense response. Resistance (R) genes present in the plant genome convey disease resistance against pathogens by producing R proteins and their action are main part of the protein-based immunity arsenal. Conserved miRNA family controls a plethora of R genes (Li et al., 2012; Shivaprasad et al., 2012; Zhai et al., 2011). It is assumed that R genes are silenced in the absence of the pathogen in order to minimize the cost for the plants and prevent autoimmunity reactions (Tian et al., 2003). It was found that the NBS-LRR genes (the main class of R proteins with nucleotide binding site (NBS) and leucine-rich repeat (LRR) motifs are silenced in a siRNA-regulated cascade similarly to ta-siRNA biogenesis scheme: RDR6-dependent secondary siRNAs are produced following the original 22 nt miRNA-mediated cleavage on a R-gene transcript. The secondary siRNA may target other R-genes. When tomato plants were infected with viruses (TCV, CMV, TRV) (or bacteria) accumulation of miR482 was reduced. In the absence of miR482 activity the resistance R gene targets are released and consequently R gene products accumulate to enhance immunity of the plants (Shivaprasad et al., 2012). In tobacco the R gene (TIR-NB-LRR immune receptor *N*) that confers resistance against TMV stays under nta-miR6019 and nta-miR6020 regulation. It was also shown that miRNA repression of *N* attenuates resistance to TMV (Li et al., 2012). Eight miRNA families were identified from *Solanaceae* (tobacco, tomato and potato) that may target R genes (Li et al., 2012). In summary miRNA-regulated R genes participate in a non-race-specific immunity mechanism where the miRNAs are the sensors of the infection. It is supposed that release of R-gene based defense may be the cause of the inhibitory action of pathogen-encoded suppressors of silencing (VSRs and bacterial suppressor proteins, BSR) on miRNA activity during infection, however this assumption needs to be experimentally tested in the future. It was shown, however, that in specific *Nicotiana* species the presence of p19 activates extreme resistance (ER) to protect tissues against TBSV and siRNA binding of p19 was necessary for ER (Sansregret et al., 2013). TAV 2b suppressor was found to elicit hypersensitive reaction (HR) when expressed in tobacco from the unrelated TMV genome (Li et al., 1999). Whether the R proteins are the monitors of the VSR presence or activity remains to be established.

#### Connecting antiviral silencing to hormone signaling

Several studies have shown that antiviral silencing might be connected to signal transduction pathways responsible for induction of salicylic acid (SA)-mediated resistance (Alamillo et al., 2006; Ji and Ding, 2001). SA is a plant hormone that is involved in local and systemic antiviral defense responses including systemic acquired resistance (SAR). SA induces expression of key antiviral silencing

factor RDR1 (Liao et al., 2013; Xie et al., 2001). In turn, RDR1 affects many jasmonic acid (JA)-regulated genes (Pandey et al., 2008). JA was implicated as a defense-related hormone (Lewsey et al., 2010). VSRs seem to interfere with hormone signaling-based responses, although the precise mechanisms are elusive. It was shown that CaMV P6 suppressor inhibits SA-induced gene expression through NPR1 (Laird et al., 2013; Love et al., 2012). Upon CMV but not CMV $\Delta$ 2b suppressor mutant strain infection the SA levels accumulated (Lewsey et al., 2010; Zhou et al., 2014). 2b was shown to have a dual action: it inhibited expression of few SA-regulated genes but in the same time enhanced the effect of SA on others. Remarkably, 2b changes the expression of 90% of jasmonic acid (JA)-regulated genes (Lewsey et al., 2010). 2b protein is known to affect RDR1 activity (Diaz-Pendon et al., 2007) that may explain partly the 2b effect on JA-signaling. HC-Pro was also shown to interfere with JA-regulated transcript expression (Endres et al., 2010). TCV p38 has been shown to interact with a NAC transcription factor called TIP in *Arabidopsis*. It was suggested that p38 interaction with TIP alters defense signaling to favor enhanced TCV invasion (Donze et al., 2014). VSRs therefore emerge as regulators of hormone-based signaling to create favorable conditions for the virus. Although at the moment the complex interplay between the RNA silencing and SA-mediated defense is elusive VSRs might be important coordinators of this crosstalk during infection.

## VSRs as tools

### *Unraveling molecular basis of silencing itself*

Due to their versatility and availability with a wide range of actions covering theoretically every aspect of RNA silencing, VSRs can be used to unravel the molecular mechanisms of both antiviral and endogenous RNA silencing pathways. Transgenic expression of cytoplasmic and nuclear p19 variants has brought evidence for nuclear DCL activity (Papp et al., 2003). Employing B2 and the cytoplasmic and nuclear variants of p19 it was suggested that a portion of cytoplasmically located endo-siRNAs are translocated back into nucleus to induce chromatin modifications. B2 or the nuclear p19 but not cytoplasmic p19 suppressed silencing of heterochromatic gene markers and altered H3K9 methylation and distribution of histone modifying complexes in flies (Fagegaltier et al., 2009). p19 size-specific dsRNA binding capacity was crucial to define the 21 nt ds-siRNAs but not long dsRNA precursors as mobile silencing signals (Dunoyer et al., 2010). The differential effects of transgenically expressed VSRs in *Arabidopsis* on miRNA and siRNA pathways have shown that these pathways are only partially overlapping (Dunoyer et al., 2004). Employing TCV p38 it was shown that DCLs work in a hierarchical manner and that there is a complex autoregulation of silencing pathways based on DCLs- and AGOs-activity feedback loops (Deleris et al., 2006; Mallory and Vaucheret, 2009). As TCV p38 could prevent siRNA but not miRNA loading, it was suggested that AGO1 is located in two distinct cellular pools (Dunoyer et al., 2010). The use of HC-Pro has led to discovery of endogenous regulators of silencing like rgsCAM following a yeast-two hybrid assay (Anandalakshmi et al., 2000). The study of poleroviral P0 and potexviral p25 has unraveled the degradation pathways responsible for AGO protein turnover both in a nonviral and viral context (Chiu et al., 2010; Derrien et al., 2012). By using P0, a VSR that destabilizes exclusively the unloaded AGO protein, the half-life of AGO/RISC complex was estimated to be in the range of 2–3 days (Csorba et al., 2010). Besides the study of VSRs helped to start understanding the complex multilateral connections of RNA- and protein based immunity and hormone regulation during host-pathogen interactions.

### *VSRs employed in molecular research*

Apart of these, VSRs can be also employed as molecular biology tools in research. P19 was used to probe the si- and miRNA content of cellular milieu following immunoprecipitations. The use of P19 provided sensitive detection without the need for amplification (Qavi et al., 2010). A surface plasmon resonance (SPR)-based miRNA sensing method was developed where the RNA probes are immobilized on gold and p19 recognize and binds the miRNA: probe duplexes. This allowed detection of miRNA at nanomolar range. Coupled to immunoassay p19 aided miRNA detection at femtomole sensitivity (Nasheri et al., 2011). Similarly P19 was used in protein-facilitated affinity capillary electrophoresis for miRNA detection in blood serum (Berezovski and Khan, 2013). These high-sensitivity methods are very powerful especially because there is no amplification bias in contrary to the PCR-based techniques.

### *VSRs as biotechnological and medical tools*

VSRs also emerge as important and useful tools for biotechnology and medicine. RNA silencing impairs or limits the use of transgenic plants in numerous biotechnological applications. VSRs can be employed efficiently to limit transgene silencing and attain consistently high-level expression of diverse products like vaccines and pharmaceuticals, high-nutritive foods, high-value products etc. (Kanagarajan et al., 2012; Naim et al., 2012).

## Conclusions and perspectives

VSRs regulate the multiple layers of the complex defense, counter-defense and counter-counter-defense arm race between host and pathogen. Although the study of VSRs was at the frontline of investigations for more than ten years, many aspects of VSR' molecular behaviors are still elusive. It is becoming more obvious now that VSRs are not just simply blockers of RNA silencing but serve as central hub regulators to dynamically integrate connections between antiviral silencing, protein-based immunity, hormone signaling, RNA metabolism and subcellular organization (Pumplin and Voinnet, 2013). As most suppressor proteins have parallel functions, the silencing function and the non-silencing activities (e.g. coat protein, movement protein, replicase, protease etc.) need to be synchronized in order to fulfill these multiple tasks and achieve "optimal" infection. In most cases, how the silencing function and the other functions of VSRs are selected and how the interconnections between the alternative functions are maintained is not known. We are just at the beginning to appreciate the complex regulatory network involving VSR multiple activities and there are still several unanswered questions. More detailed information is needed about viral replication, subcellular localization of viral proteins including VSRs and their interaction with host factors. Very little is known about not just "antiviral" – but the "endogenous" – RISC complex as well. The relatively recently developed *in vitro* system may help to better understand the RISC components and activities. In exploring RISC complex VSRs such as P0, p38 or P1 may be powerful tools in the future research. It was suggested that some VSRs facilitate viral long-distance movement by suppressing the spread of antiviral silencing (e.g. p19 sequesters the signal, 2b or p38 blocks the production of the signal). Understanding antiviral systemic signaling is a very important but still unknown aspect of host defense. The assumption that vsiRNA move throughout the plant to convey immunity still awaits direct experimental validation. Several studies have shown that antiviral acquired resistance can be passed transgenerationally to offspring through siRNAs and RdDM by the use of virus-induced TGS (VITGS) (Kanazawa et al., 2011; Kon and Yoshikawa, 2014; Martín-Hernández and Baulcombe, 2008; Otagaki et al., 2011). How VSR activities could modulate/interfere with

resistance “priming” in subsequent generations is not known. VITGS could provide a useful technology for plant breeding via epigenetic modifications that needs to be explored in the future.

In spite of its paramount economical importance viral symptom development is a poorly understood aspect of virus infection process. How does VSR presence/activities relate to viral symptom development? Due to their effect on accumulation and function of endogenous small RNAs it is supposed that VSRs could be major contributors to the development of viral symptoms. It was shown however that VSRs might be dispensable for disease induction (Diaz-Pendon et al., 2007), therefore, answering this question need more investigation. Most of the knowledge on VSRs and antiviral silencing comes from studies in model organisms like *Arabidopsis* that may miss important aspects of the authentic virus infection and host defense responses that occur in nature. The recent development of genetic tools such CRISPR/Cas9 technology will allow us to generate specific mutation in several other plant species including important virus hosts (e.g. *N. benthamiana*) and economically significant crops to extend our knowledge on RNA silencing.

The use of natural host-virus pairs would be an important track to follow in the future in order to be able to understand and ultimately control viruses that cause huge economical losses on crop plant worldwide.

## Acknowledgments

Országos Tudományos Kutatási Alapprogramok-NK105850 and the János Bolyai Research Scholarship of the Hungarian Academy of Sciences supported this work. We apologize to those colleagues whose work we were unable to cite because of space restrictions. We thank to Gy. Szittyá for helpful comments on the manuscript.

## References

- Adenot, X., Elmayan, T., Laressergues, D., Boutet, S., Bouche, N., Gascioli, V., Vaucheret, H., 2006. DRB4-dependent TAS3 trans-acting siRNAs control leaf morphology through AGO7. *Curr. Biol.* 16, 927–932.
- Ahn, J.W., Yin, C.J., Liu, J.R., Jeong, W.J., 2010. Cucumber mosaic virus 2b protein inhibits RNA silencing pathways in green alga *Chlamydomonas reinhardtii*. *Plant Cell Rep.* 29, 967–975.
- Akbergenov, R., Si-Ammour, A., Blevins, T., Amin, I., Kutter, C., Vanderschuren, H., Zhang, P., Gruissem, W., Meins Jr., F., Hohn, T., Pooggin, M.M., 2006. Molecular characterization of geminivirus-derived small RNAs in different plant species. *Nucleic Acids Res.* 34, 462–471.
- Al-Kaff, N.S., Covey, S.N., Kreike, M.M., Page, A.M., Pinder, R., Dale, P.J., 1998. Transcriptional and posttranscriptional plant gene silencing in response to a pathogen. *Science* 279, 2113–2115.
- Alamillo, J.M., Saenz, P., Garcia, J.A., 2006. Salicylic acid-mediated and RNA-silencing defense mechanisms cooperate in the restriction of systemic spread of plum pox virus in tobacco. *Plant J. Cell Mol. Biol.* 48, 217–227.
- Allen, E., Xie, Z., Gustafson, A.M., Carrington, J.C., 2005. microRNA-directed phasing during trans-acting siRNA biogenesis in plants. *Cell* 121, 207–221.
- Anandalakshmi, R., Marathe, R., Ge, X., Herr Jr., J.M., Mau, C., Mallory, A., Pruss, G., Bowman, L., Vance, V.B., 2000. A calmodulin-related protein that suppresses posttranscriptional gene silencing in plants. *Science* 290, 142–144.
- Anandalakshmi, R., Pruss, G.J., Ge, X., Marathe, R., Mallory, A.C., Smith, T.H., Vance, V.B., 1998. A viral suppressor of gene silencing in plants. *Proc. Natl. Acad. Sci. USA* 95, 13079–13084.
- Andika, I.B., Kondo, H., Nishiguchi, M., Tamada, T., 2012. The cysteine-rich proteins of beet necrotic yellow vein virus and tobacco rattle virus contribute to efficient suppression of silencing in roots. *J. Gen. Virol.* 93, 1841–1850.
- Angel, C.A., Schoelz, J.E., 2013. A survey of resistance to tomato bushy stunt virus in the genus *Nicotiana* reveals that the hypersensitive response is triggered by one of three different viral proteins. *Mol. Plant–Microbe Interact.* 26, 240–248.
- Aregger, M., Borah, B.K., Seguin, J., Rajeswaran, R., Gubaeva, E.G., Zvereva, A.S., Windels, D., Vazquez, F., Blevins, T., Farinelli, L., Pooggin, M.M., 2012. Primary and secondary siRNAs in geminivirus-induced gene silencing. *PLoS Pathog.* 8, e1002941.
- Azevedo, J., Garcia, D., Pontier, D., Ohnesorge, S., Yu, A., Garcia, S., Braun, L., Bergdoll, M., Hakimi, M.A., Lagrange, T., Voinnet, O., 2010. Argonaute quenching and global changes in dicer homeostasis caused by a pathogen-encoded GW repeat protein. *Genes Dev.* 24, 904–915.
- Bar-Ziv, A., Levy, Y., Hak, H., Mett, A., Belausov, E., Citovsky, V., Gafni, Y., 2012. The Tomato yellow leaf curl virus (TYLCV) V2 protein interacts with the host papain-like cysteine protease CYP1. *Plant Signal. Behav.* 7, 983–989.
- Baumberger, N., Tsai, C.H., Lie, M., Havecker, E., Baulcombe, D.C., 2007. The Ploverovirus silencing suppressor P0 targets ARGONAUTE proteins for degradation. *Curr. Biol.* 17, 1609–1614.
- Berezovski, M.V., Khan, N., 2013. Quantitative analysis of microRNA in blood serum with protein-facilitated affinity capillary electrophoresis. *Methods Mol. Biol.* 1039, 245–259.
- Bernstein, E., Caudy, A.A., Hammond, S.M., Hannon, G.J., 2001. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* 409, 363–366.
- Bisaro, D.M., 2006. Silencing suppression by geminivirus proteins. *Virology* 344, 158–168.
- Blevins, T., Rajeswaran, R., Aregger, M., Borah, B.K., Schepetilnikov, M., Baerlocher, L., Farinelli, L., Meins Jr., F., Hohn, T., Pooggin, M.M., 2011. Massive production of small RNAs from a non-coding region of cauliflower mosaic virus in plant defense and viral counter-defense. *Nucleic Acids Res.* 39, 5003–5014.
- Blevins, T., Rajeswaran, R., Shivaprasad, P.V., Beknazariants, D., Si-Ammour, A., Park, H.S., Vazquez, F., Robertson, D., Meins Jr., F., Hohn, T., Pooggin, M.M., 2006. Four plant dicers mediate viral small RNA biogenesis and DNA virus induced silencing. *Nucleic Acids Res.* 34, 6233–6246.
- Bologna, N.G., Voinnet, O., 2014. The diversity, biogenesis, and activities of endogenous silencing small RNAs in *Arabidopsis*. *Annu. Rev. Plant Biol.* 65, 473–503.
- Bonneville, J.M., Sanfacon, H., Fütterer, J., Hohn, T., 1989. Posttranscriptional trans-acting in cauliflower mosaic virus. *Cell* 59, 1135–1143.
- Bortolamiol, D., Pazhouhandeh, M., Marrocco, K., Genschik, P., Ziegler-Graff, V., 2007. The Ploverovirus F box protein P0 targets ARGONAUTE1 to suppress RNA silencing. *Curr. Biol.* 17, 1615–1621.
- Bouché, N., Laressergues, D., Gascioli, V., Vaucheret, H., 2006. An antagonistic function for *Arabidopsis* DCL2 in development and a new function for DCL4 in generating viral siRNAs. *EMBO J.* 25, 3347–3356.
- Boutet, S., Vazquez, F., Liu, J., Beclin, C., Fagard, M., Grati, A., Morel, J.B., Crete, P., Chen, X., Vaucheret, H., 2003. *Arabidopsis* HEN1: a genetic link between endogenous miRNA controlling development and siRNA controlling transgene silencing and virus resistance. *Curr. Biol.* 13, 843–848.
- Brigneti, G., Voinnet, O., Li, W.X., Ji, L.H., Ding, S.W., Baulcombe, D.C., 1998. Viral pathogenicity determinants are suppressors of transgene silencing in *Nicotiana benthamiana*. *EMBO J.* 17, 6739–6746.
- Bronkhorst, A.W., van Rij, R.P., 2014. The long and short of antiviral defense: small RNA-based immunity in insects. *Curr. Opin. Virol.* 7, 19–28.
- Buchmann, R.C., Asad, S., Wolf, J.N., Mohannath, G., Bisaro, D.M., 2009. Geminivirus AL2 and L2 proteins suppress transcriptional gene silencing and cause genome-wide reductions in cytosine methylation. *J. Virol.* 83, 5005–5013.
- Canizares, M.C., Navas-Castillo, J., Moriones, E., 2008. Multiple suppressors of RNA silencing encoded by both genomic RNAs of the crinivirus, Tomato chlorosis virus. *Virology* 379, 168–174.
- Canizares, M.C., Taylor, K.M., Lomonosoff, G.P., 2004. Surface-exposed C-terminal amino acids of the small coat protein of Cowpea mosaic virus are required for suppression of silencing. *J. Gen. Virol.* 85, 3431–3435.
- Canizares, M.C., Lozano-Durán, R., Canto, T., Bejarano, E.R., Bisaro, D.M., Navas-Castillo, J., Moriones, E., 2013. Effects of the crinivirus coat protein-interacting plant protein SAHH on post-transcriptional RNA silencing and its suppression. *Mol. Plant–Microbe Interact.* 26, 1004–1015.
- Cao, M., Du, P., Wang, X., Yu, Y.Q., Qiu, Y.H., Li, W., Gal-On, A., Zhou, C., Li, Y., Ding, S.W., 2014. Virus infection triggers widespread silencing of host genes by a distinct class of endogenous siRNAs in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 111, 14613–14618.
- Cao, X., Zhou, P., Zhang, X., Zhu, S., Zhong, X., Xiao, Q., Ding, B., Li, Y., 2005. Identification of an RNA silencing suppressor from a plant double-stranded RNA virus. *J. Virol.* 79, 13018–13027.
- Carbonell, A., Fahlgren, N., Garcia-Ruiz, H., Gilbert, K.B., Montgomery, T.A., Nguyen, T., Cuperus, J.T., Carrington, J.C., 2012. Functional analysis of three *Arabidopsis* ARGONAUTES using slicer-defective mutants. *Plant Cell* 24, 3613–3629.
- Carrington, J.C., Freed, D.D., Sanders, T.C., 1989. Autocatalytic processing of the potyvirus helper component proteinase in *Escherichia coli* and in vitro. *J. Virol.* 63, 4459–4463.
- Chapman, E.J., Prokhnovsky, A.I., Gopinath, Dolja, V.V., Carrington, J.C., 2004. Viral RNA silencing suppressors inhibit the microRNA pathway at an intermediate step. *Genes Dev.* 18, 1179–1186.
- Chellappan, P., Vanitharani, R., Fauquet, C.M., 2005. MicroRNA-binding viral protein interferes with *Arabidopsis* development. *Proc. Natl. Acad. Sci. USA* 102, 10381–10386.
- Chen, H.Y., Yang, J., Lin, C., Yuan, Y.A., 2008. Structural basis for RNA-silencing suppression by Tomato aspermy virus protein 2b. *EMBO Rep.* 9, 754–760.
- Chen, J., Li, W.X., Xie, D., Peng, J.R., Ding, S.W., 2004. Viral virulence protein suppresses RNA silencing-mediated defense but upregulates the role of microRNA in host gene expression. *Plant Cell* 16, 1302–1313.
- Cheng, J., Koukiokolo, R., Kieliszewicz, K., Sagan, S.M., Pezacki, J.P., 2009. Cysteine residues of Carnation Italian Ringspot virus p19 suppressor of RNA silencing maintain global structural integrity and stability for siRNA binding. *Biochim. Biophys. Acta* 1794, 1197–1203.
- Cheng, Y.Q., Liu, Z.M., Xu, J., Zhou, T., Wang, M., Chen, Y.T., Li, H.F., Fan, Z.F., 2008. HC-Pro protein of sugar cane mosaic virus interacts specifically with maize ferredoxin-5 in vitro and in planta. *J. Gen. Virol.* 89, 2046–2054.
- Chiba, S., Hleibieh, K., Delbianco, A., Klein, E., Ratti, C., Ziegler-Graff, V., Bouzoubaa, S., Gilmer, D., 2013. The benyvirus RNA silencing suppressor is essential for

- long-distance movement, requires both zinc-finger and NoLS basic residues but not a nucleolar localization for its silencing-suppression activity. *Mol. Plant Microbe Interact.* 26, 168–181.
- Chiu, M.H., Chen, I.H., Baulcombe, D.C., Tsai, C.H., 2010. The silencing suppressor P25 of Potato virus X interacts with Argonaute1 and mediates its degradation through the proteasome pathway. *Mol. Plant Pathol.* 11, 641–649.
- Choi, C.W., Qu, F., Ren, T., Ye, X., Morris, T.J., 2004. RNA silencing-suppressor function of Turnip crinkle virus coat protein cannot be attributed to its interaction with the Arabidopsis protein TIP. *J. Gen. Virol.* 85, 3415–3420.
- Chowda-Reddy, R.V., Achenjang, F., Felton, C., Etarock, M.T., Anangfac, M.T., Nugent, P., Fondong, V.N., 2008. Role of a geminivirus AV2 protein putative protein kinase C motif on subcellular localization and pathogenicity. *Virus Res.* 135, 115–124.
- Creamer, K.M., Partridge, J.F., 2011. RITS-connecting transcription, RNA interference, and heterochromatin assembly in fission yeast. *Wiley interdisciplinary reviews. RNA* 2, 632–646.
- Csorba, T., Bovi, A., Dalmay, T., Burgyan, J., 2007. The p122 subunit of Tobacco Mosaic Virus replicase is a potent silencing suppressor and compromises both small interfering RNA- and microRNA-mediated pathways. *J. Virol.* 81, 11768–11780.
- Csorba, T., Lozsa, R., Hutvagner, G., Burgyan, J., 2010. Polerovirus protein P0 prevents the assembly of small RNA-containing RISC complexes and leads to degradation of ARGONAUTE1. *Plant J. Cell Mol. Biol.* 62, 463–472.
- Cuellar, W.J., Kreuze, J.F., Rajamaki, M.L., Cruzado, K.R., Untiveros, M., Valkonen, J.P., 2009. Elimination of antiviral defense by viral RNase III. *Proc. Natl. Acad. Sci. USA* 106, 10354–10358.
- Cui, X., Li, G., Wang, D., Hu, D., Zhou, X., 2005. A Begomovirus DNA $\beta$ -encoded protein binds DNA, functions as a suppressor of RNA silencing, and targets the cell nucleus. *J. Virol.* 79, 10764–10775.
- Cullen, B.R., Cherry, S., tenOever, B.R., 2013. Is RNA interference a physiologically relevant innate antiviral immune response in mammals? *Cell Host Microbe* 14, 374–378.
- Curtin, S.J., Watson, J.M., Smith, N.A., Eamens, A.L., Blanchard, C.L., Waterhouse, P.M., 2008. The roles of plant dsRNA-binding proteins in RNAi-like pathways. *FEBS Lett.* 582, 2753–2760.
- Dalmay, T., Hamilton, A., Rudd, S., Angell, S., Baulcombe, D.C., 2000. An RNA-dependent RNA polymerase gene in Arabidopsis is required for posttranscriptional gene silencing mediated by a transgene but not by a virus. *Cell* 101, 543–553.
- Dalmay, T., Horsefield, R., Braumstein, T.H., Baulcombe, D.C., 2001. SDE3 encodes an RNA helicase required for post-transcriptional gene silencing in Arabidopsis. *EMBO J.* 20, 2069–2078.
- Deleris, A., Gallego-Bartolome, J., Bao, J., Kasschau, K.D., Carrington, J.C., Voinnet, O., 2006. Hierarchical action and inhibition of plant dicer-like proteins in antiviral defense. *Science* 313, 68–71.
- Delfosse, V.C., Agrofoglio, Y.C., Casse, M.F., Kresic, I.B., Hopp, H.E., Ziegler-Graff, V., Distéfano, A.J., 2014. The P0 protein encoded by cotton leafroll dwarf virus (CLRVD) inhibits local but not systemic RNA silencing. *Virus Res.* 180, 70–75.
- Deng, X., Kelloniemi, J., Haikonen, T., Vuorinen, A.L., Elomaa, P., Teeri, T.H., Valkonen, J.P.T., 2013. Modification of tobacco rattle virus RNA1 to serve as a VIGS vector reveals that the 29K movement protein is an RNA silencing suppressor of the virus. *Mol. Plant – Microbe Interact.* 26, 503–514.
- Deng, X.G., Peng, X.J., Zhu, F., Chen, Y.J., Zhu, T., Qin, S.B., Xi, D.H., Lin, H.H., 2014. A critical domain of sweet potato chlorotic fleck virus nucleotide-binding protein (NaBp) for RNA silencing suppression, nuclear localization and viral pathogenesis. *Mol. Plant Pathol.* , <http://dx.doi.org/10.1111/mpp.12186>.
- Derrien, B., Baumberger, N., Schepetilnikov, M., Viotti, C., De Cillia, J., Ziegler-Graff, V., Isono, E., Schumacher, K., Genschik, P., 2012. Degradation of the antiviral component ARGONAUTE1 by the autophagy pathway. *Proc. Natl. Acad. Sci. USA* 109, 15942–15946.
- Diaz-Pendon, J.A., Li, F., Li, W.X., Ding, S.W., 2007. Suppression of antiviral silencing by cucumber mosaic virus 2b protein in Arabidopsis is associated with drastically reduced accumulation of three classes of viral small interfering RNAs. *Plant Cell* 19, 2053–2063.
- Dielen, A.S., Badaoui, S., Candresse, T., German-Retana, S., 2010. The ubiquitin/26S proteasome system in plant-pathogen interactions: a never-ending hide-and-seek game. *Mol. Plant Pathol.* 11, 293–308.
- Ding, S.W., Shi, B.J., Li, W.X., Symons, R.H., 1996. An interspecies hybrid RNA virus is significantly more virulent than either parental virus. *Proc. Natl. Acad. Sci. USA* 93, 7470–7474.
- Ding, S.W., Voinnet, O., 2007. Antiviral immunity directed by small RNAs. *Cell* 130, 413–426.
- Ding, X.S., Liu, J., Cheng, N.H., Folimonov, A., Hou, Y.M., Bao, Y., Katagi, C., Carter, S.A., Nelson, R.S., 2004. The Tobacco mosaic virus 126-kDa protein associated with virus replication and movement suppresses RNA silencing. *Mol. Plant – Microbe Interact.* 17, 583–592.
- Donaire, L., Barajas, D., Martinez-Garcia, B., Martinez-Priego, L., Pagan, I., Llave, C., 2008. Structural and genetic requirements for the biogenesis of tobacco rattle virus-derived small interfering RNAs. *J. Virol.* 82, 5167–5177.
- Donze, T., Qu, F., Twigg, P., Morris, T.J., 2014. Turnip crinkle virus coat protein inhibits the basal immune response to virus invasion in Arabidopsis by binding to the NAC transcription factor TIP. *Virology* 449, 207–214.
- dos Reis Figueira, A., Golem, S., Goregaoker, S.P., Culver, J.N., 2002. A nuclear localization signal and a membrane association domain contribute to the cellular localization of the tobacco mosaic virus 126-kDa replicase protein. *Virology* 301, 81–89.
- Du, Z., Chen, A., Chen, W., Liao, Q., Zhang, H., Bao, Y., Roossinck, M.J., Carr, J.P., 2014. Nuclear-cytoplasmic partitioning of cucumber mosaic virus protein 2b determines the balance between its roles as a virulence determinant and an RNA-silencing suppressor. *J. Virol.* 88, 5228–5241.
- Du, Z., Xiao, D., Wu, J., Jia, D., Yuan, Z., Liu, Y., Hu, L., Han, Z., Wei, T., Lin, Q., Wu, Z., Xie, L., 2011. p2 of rice stripe virus (RSV) interacts with OsSGS3 and is a silencing suppressor. *Mol. Plant Pathol.* 12, 808–814.
- Duan, C.G., Fang, Y.Y., Zhou, B.J., Zhao, J.H., Hou, W.N., Zhu, H., Ding, S.W., Guo, H.S., 2012. Suppression of Arabidopsis ARGONAUTE1-mediated slicing, transgene-induced RNA silencing, and DNA methylation by distinct domains of the cucumber mosaic virus 2b protein. *Plant Cell* 24, 259–274.
- Dunoyer, P., Lecellier, C.H., Parizotto, E.A., Himber, C., Voinnet, O., 2004. Probing the microRNA and small interfering RNA pathways with virus-encoded suppressors of RNA silencing. *Plant Cell* 16, 1235–1250.
- Dunoyer, P., Pfeffer, S., Fritsch, C., Hemmer, O., Voinnet, O., Richards, K.E., 2002. Identification, subcellular localization and some properties of a cysteine-rich suppressor of gene silencing encoded by peanut clump virus. *Plant J. Cell Mol. Biol.* 29, 555–567.
- Dunoyer, P., Schott, G., Himber, C., Meyer, D., Takeda, A., Carrington, J.C., Voinnet, O., 2010. Small RNA duplexes function as mobile silencing signals between plant cells. *Science* 328, 912–916.
- Eamens, A.L., Kim, K.W., Curtin, S.J., Waterhouse, P.M., 2012a. DRB2 is required for microRNA biogenesis in *Arabidopsis thaliana*. *PLoS One* 7, e35933.
- Eamens, A.L., Wook Kim, K., Waterhouse, P.M., 2012b. DRB2, DRB3 and DRB5 function in a non-canonical microRNA pathway in *Arabidopsis thaliana*. *Plant Signal. Behav.* 7, 1224–1229.
- Ekwall, K., 2004. The RITS complex-A direct link between small RNA and heterochromatin. *Mol. Cell* 13, 304–305.
- Ehardt, H.A., Thi, E.P., Wang, M.B., Unrau, P.J., 2005. Extensive 3' modification of plant small RNAs is modulated by helper component-proteinase expression. *Proc. Natl. Acad. Sci. USA* 102, 13398–13403.
- Einiä, O., Dograd, S.C., Drye, I.B., Randles, J.W., 2012. Silencing suppressor activity of a begomovirus DNA  $\beta$  encoded protein and its effect on heterologous helper virus replication. *Virus Res.* 167, 97–101.
- Endres, M.W., Gregory, B.D., Gao, Z., Foreman, A.W., Mlotshwa, S., Ge, X., Pruss, G.J., Ecker, J.R., Bowman, L.H., Vance, V., 2010. Two plant viral suppressors of silencing require the ethylene-inducible host transcription factor RAV2 to block RNA silencing. *PLoS Pathog.* 6, e1000729.
- Eulalio, A., Tritschler, F., Izaurralde, E., 2009. The GW182 protein family in animal cells: new insights into domains required for miRNA-mediated gene silencing. *RNA* 15, 1433–1442.
- Fagard, M., Boutet, S., Morel, J.B., Bellini, C., Vaucheret, H., 2000. AGO1, QDE-2, and RDE-1 are related proteins required for post-transcriptional gene silencing in plants, quelling in fungi, and RNA interference in animals. *Proc. Natl. Acad. Sci. USA* 97, 11650–11654.
- Fagegaltier, D., Bouge, A.L., Berry, B., Poisot, E., Sismeiro, O., Coppee, J.Y., Theodore, L., Voinnet, O., Antoniewski, C., 2009. The endogenous siRNA pathway is involved in heterochromatin formation in drosophila. *Proc. Natl. Acad. Sci. USA* 106, 21258–21263.
- Feng, L., Duan, C.G., Guo, H.S., 2013. Inhibition of in vivo Slicer activity of Argonaute protein 1 by the viral 2b protein independent of its dsRNA-binding function. *Mol. Plant Pathol.* 14, 617–622.
- Fukunaga, R., Doudna, J.A., 2009. dsRNA with 5' overhangs contributes to endogenous and antiviral RNA silencing pathways in plants. *EMBO J.* 28, 545–555.
- Fusaro, A.F., Correa, R.L., Nakasugi, K., Jackson, C., Kawchuk, L., Vaslin, M.F., Waterhouse, P.M., 2012. The Enamovirus P0 protein is a silencing suppressor which inhibits local and systemic RNA silencing through AGO1 degradation. *Virology* 426, 178–187.
- Garcia, D., Garcia, S., Pontier, D., Marchais, A., Renou, J.P., Lagrange, T., Voinnet, O., 2012. Ago hook and RNA helicase motifs underpin dual roles for SDE3 in antiviral defense and silencing of nonconserved intergenic regions. *Mol. Cell* 48, 109–120.
- Garcia-Ruiz, H., Takeda, A., Chapman, E.J., Sullivan, C.M., Fahlgren, N., Brempelis, K.J., Carrington, J.C., 2010. Arabidopsis RNA-dependent RNA polymerases and dicer-like proteins in antiviral defense and small interfering RNA biogenesis during turnip mosaic virus infection. *Plant Cell* 22, 481–496.
- Gazzani, S., Lawrenson, T., Woodward, C., Headon, D., Sablowski, R., 2004. A link between mRNA turnover and RNA interference in Arabidopsis. *Science* 306, 1046–1048.
- Genoves, A., Navarro, J.A., Pallas, V., 2006. Functional analysis of the five melon necrotic spot virus genome-encoded proteins. *J. Gen. Virol.* 37, 2371–2380.
- Genoves, A., Pallas, V., Navarro, J.A., 2011. Contribution of topology determinants of a viral movement protein to its membrane association, intracellular traffic, and viral cell-to-cell movement. *J. Virol.* 85, 7797–7809.
- Ghazala, W., Waltermann, A., Pilot, R., Winter, S., Varrelmann, M., 2008. Functional characterization and subcellular localization of the 16K cysteine-rich suppressor of gene silencing protein of tobacco rattle virus. *J. Gen. Virol.* 89, 1748–1758.
- Ghoshal, B., Sanfacon, H., 2014. Temperature-dependent symptom recovery in *Nicotiana benthamiana* plants infected with tomato ringspot virus is associated with reduced translation of viral RNA2 and requires ARGONAUTE 1. *Virology* 456–457, 188–197.
- Gillet, F.X., Cattoni, D.I., Petiot-Bécard, S., Delalande, F., Poignavet, V., Brizard, J.P., Bessin, Y., Dorselaer, A., Declerck, N., Sanglier-Cianféran, S., Brugidou, C., Vignols, F., 2013. The RYMV-encoded viral suppressor of RNA silencing P1 is a zinc-binding protein with redox-dependent flexibility. *J. Mol. Biol.* 425, 2423–2435.
- Giner, A., Lakatos, L., Garcia-Chapa, M., Lopez-Moya, J.J., Burgyan, J., 2010. Viral protein inhibits RISC activity by argonaute binding through conserved WG/GW motifs. *PLoS Pathog.* 6, e1000996.

- Glick, E., Zrachya, A., Levy, Y., Mett, A., Gidoni, D., Belausov, E., Citovsky, V., Gafni, Y., 2008. Interaction with host SGS3 is required for suppression of RNA silencing by tomato yellow leaf curl virus V2 protein. *Proc. Natl. Acad. Sci. USA* 105, 157–161.
- Gonzalez, I., Martinez, L., Rakitina, D.V., Lewsey, M.G., Atencio, F.A., Llave, C., Kalinina, N.O., Carr, J.P., Palukaitis, P., Canto, T., 2010. Cucumber mosaic virus 2b protein subcellular targets and interactions: their significance to RNA silencing suppressor activity. *Mol. Plant–Microbe Interact.* 23, 294–303.
- Gonzalez, I., Rakitina, D., Semashko, M., Taliansky, M., Praveen, S., Palukaitis, P., Carr, J.P., Kalinina, N., Canto, T., 2012. RNA binding is more critical to the suppression of silencing function of cucumber mosaic virus 2b protein than nuclear localization. *RNA* 18, 771–782.
- Goto, K., Kobori, T., Kosaka, Y., Natsuaki, T., Masuta, C., 2007. Characterization of silencing suppressor 2b of cucumber mosaic virus based on examination of its small RNA-binding abilities. *Plant Cell Physiol.* 48, 1050–1060.
- Goswami, S., Sahana, N., Pandey, V., Doblas, P., Jain, R.K., Palukaitis, P., Canto, T., Praveen, S., 2012. Interference in plant defense and development by non-structural protein NSs of Groundnut bud necrosis virus. *Virus Res.* 163, 368–373.
- Gouveia, P., Nolasco, G., 2012. The p19.7 RNA silencing suppressor from grapevine leafroll-associated virus 3 shows different levels of activity across phylogenetic groups. *Virus Genes* 45, 333–339.
- Gouveia, P., Dandlen, S., Costa, A., Marques, N., Nolasco, G., 2012. Identification of an RNA silencing suppressor encoded by grapevine leafroll-associated virus 3. *Eur. J. Plant Pathol.* 133, 237–245.
- Groenenboom, M.A., Hogeweg, P., 2012. Modelling the dynamics of viral suppressors of RNA silencing. *J. R. Soc., Interface/R. Soc.* 9, 436–447.
- Guilley, H., Bortolamiol, D., Jonard, G., Bouzoubaa, S., Ziegler-Graff, V., 2009. Rapid screening of RNA silencing suppressors by using a recombinant virus derived from beet necrotic yellow vein virus. *J. Gen. Virol.* 90, 2536–2541.
- Guo, B., Lin, J., Ye, K., 2011. Structure of the autocatalytic cysteine protease domain of potyvirus helper-component proteinase. *J. Biol. Chem.* 286, 21937–21943.
- Guo, H., Song, X., Xie, C., Huo, Y., Zhang, F., Chen, X., Geng, Y., Fang, R., 2013. Rice yellow stunt rhabdovirus protein 6 suppresses systemic RNA silencing by blocking RDR6-mediated secondary siRNA synthesis. *Mol. Plant–Microbe Interact.* 26, 927–936.
- Guo, H.S., Ding, S.W., 2002. A viral protein inhibits the long range signaling activity of the gene silencing signal. *EMBO J.* 21, 398–407.
- Gy, I., Gascioli, V., Laussergues, D., Morel, J.B., Gombert, J., Proux, F., Proux, C., Vaucheret, H., Mallory, A.C., 2007. Arabidopsis FIERY1, XRN2, and XRN3 are endogenous RNA silencing suppressors. *Plant Cell* 19, 3451–3461.
- Haas, G., Azevedo, J., Moissiard, G., Geldreich, A., Himber, C., Bureau, M., Fukuhara, T., Keller, M., Voinnet, O., 2008. Nuclear import of CaMV P6 is required for infection and suppression of the RNA silencing factor DRB4. *EMBO J.* 27, 2102–2112.
- Haikonen, T., Rajamaki, M.L., Tian, Y.P., Valkonen, J.P., 2013a. Mutation of a short variable region in HCpro protein of potato virus A affects interactions with a microtubule-associated protein and induces necrotic responses in tobacco. *Mol. Plant–Microbe Interact.* 26, 721–733.
- Haikonen, T., Rajamaki, M.L., Valkonen, J.P., 2013b. Interaction of the microtubule-associated host protein HIP2 with viral helper component proteinase is important in infection with potato virus A. *Mol. Plant–Microbe Interact.* 26, 734–744.
- Hamera, S., Song, X., Su, L., Chen, X., Fang, R., 2012. Cucumber mosaic virus suppressor 2b binds to AGO4-related small RNAs and impairs AGO4 activities. *Plant J. Cell Mol. Biol.* 69, 104–115.
- Hamilton, A.J., Baulcombe, D.C., 1999. A species of small antisense RNA in posttranscriptional gene silencing in plants. *Science* 286, 950–952.
- Hammond, S.M., Boettcher, S., Caudy, A.A., Kobayashi, R., Hannon, G.J., 2001. Argonaute2, a link between genetic and biochemical analyses of RNAi. *Science* 293, 1146–1150.
- Han, Y.H., Xiang, H.Y., Wang, Q., Li, Y.Y., Wu, W.Q., Han, C.G., Li, D.W., Yu, J.L., 2010. Ring structure amino acids affect the suppressor activity of melon aphid-borne yellows virus P0 protein. *Virology* 406, 21–27.
- Hao, L., Wang, H., Sunter, G., Bisaro, D.M., 2003. Geminivirus AL2 and L2 proteins interact with and inactivate SNF1 kinase. *Plant Cell* 15, 71034–71048.
- Hao, X., Lu, A., Sokal, N., Bhagwat, B., Leung, E., Mao, R., Reade, R., Wu, Y., Rochon, D., Xiang, Y., 2011. Cucumber necrosis virus p20 is a viral suppressor of RNA silencing. *Virus Res.* 155, 423–432.
- Harries, P.A., Palanichelvam, K., Bhat, S., Nelson, R.S., 2008. Tobacco mosaic virus 126-kDa protein increases the susceptibility of *Nicotiana tabacum* to other viruses and its dosage affects virus-induced gene silencing. *Mol. Plant–Microbe Interact.* 21, 1539–1548.
- Harvey, J.J., Lewsey, M.G., Patel, K., Westwood, J., Heimstadt, S., Carr, J.P., Baulcombe, D.C., 2011. An antiviral defense role of AGO2 in plants. *PLoS One* 6, e14639.
- Havelda, Z., Hornyik, C., Valoczi, A., Burgyan, J., 2005. Defective interfering RNA hinders the activity of a tombusvirus-encoded posttranscriptional gene silencing suppressor. *J. Virol.* 79, 450–457.
- Hemmes, H., Lakatos, L., Goldbach, R., Burgyan, J., Prins, M., 2007. The NS3 protein of rice hoja blanca tenuivirus suppresses RNA silencing in plant and insect hosts by efficiently binding both siRNAs and miRNAs. *RNA* 13, 1079–1089.
- Hendelman, A., Kravchik, M., Stav, R., Zik, M., Lugassi, N., Arazi, T., 2013. The developmental outcomes of P0-mediated ARGONAUTE destabilization in tomato. *PLoS One* 8, e63377.
- Hernandez-Pinzon, I., Yelina, N.E., Schwach, F., Studholme, D.J., Baulcombe, D., Dalmay, T., 2007. SDE5, the putative homologue of a human mRNA export factor, is required for transgene silencing and accumulation of trans-acting endogenous siRNA. *Plant J. Cell Mol. Biol.* 50, 140–148.
- Hiraguri, A., Itoh, R., Kondo, N., Nomura, Y., Aizawa, D., Murai, Y., Koiki, H., Seki, M., Shinozaki, K., Fukuhara, T., 2005. Specific interactions between dicer-like proteins and HYL1/DRB-family dsRNA-binding proteins in *Arabidopsis thaliana*. *Plant Mol. Biol.* 57, 173–188.
- Hull, R., 2002. *Matthews' Plant Virology*, Fourth ed. Academic Press, San Diego, California, USA.
- Hussain, M., Abraham, A.M., Asgari, S., 2010. An Ascovirus-encoded RNase III autoregulates its expression and suppresses RNA interference-mediated gene silencing. *J. Virol.* 84, 3624–3630.
- Hsieh, Y.C., Omarov, R.T., Scholthof, H.B., 2009. Diverse and newly recognized effects associated with short interfering RNA binding site modifications on the tomato bushy stunt virus P19 silencing suppressor. *J. Virol.* 83, 2188–2200.
- Hu, Q., Hollunder, J., Niehl, A., Korner, C.J., Gereige, D., Windels, D., Arnold, A., Kuiper, M., Vazquez, F., Pooggin, M., Heinlein, M., 2011. Specific impact of tobamovirus infection on the Arabidopsis small RNA profile. *PLoS One* 6, e19549.
- Hutvagner, G., McLachlan, J., Pasquinelli, A.E., Balint, E., Tuschl, T., Zamore, P.D., 2001. A cellular function for the RNA-interference enzyme dicer in the maturation of the let-7 small temporal RNA. *Science* 293, 834–838.
- Iki, T., Yoshikawa, M., Nishikiori, M., Jaudal, M.C., Matsumoto-Yokoyama, E., Mitsuhashi, I., Meshi, T., Ishikawa, M., 2010. In vitro assembly of plant RNA-induced silencing complexes facilitated by molecular chaperone HSP90. *Mol. Cell* 39, 282–291.
- Inaba, J.I., Kim, B.M., Shimura, H., Masuta, C., 2011. Virus-induced necrosis is a consequence of direct protein–protein interaction between a viral RNA-silencing suppressor and a host catalase. *Plant Physiol.* 156, 2026–2036.
- Jakubiec, A., Yang, S.W., Chua, N.H., 2012. Arabidopsis DRB4 protein in antiviral defense against turnip yellow mosaic virus infection. *Plant J. Cell Mol. Biol.* 69, 14–25.
- Jamous, R.M., Boonrod, K., Fuellgrabe, M.W., Ali-Shtayeh, M.S., Krczal, G., Wassenegger, M., 2011. The helper component-proteinase of the Zucchini yellow mosaic virus inhibits the Hua Enhancer 1 methyltransferase activity in vitro. *J. Gen. Virol.* 92, 2222–2226.
- Jaubert, M., Bhattacharjee, S., Mello, A.F., Perry, K.L., Moffett, P., 2011. ARGONAUTE2 mediates RNA-silencing antiviral defenses against potato virus X in Arabidopsis. *Plant Physiol.* 156, 1556–1564.
- Jay, F., Wang, Y., Yu, A., Taconnat, L., Pelletier, S., Colot, V., Renou, J.P., Voinnet, O., 2011. Misregulation of AUXIN RESPONSE FACTOR 8 underlies the developmental abnormalities caused by three distinct viral silencing suppressors in Arabidopsis. *PLoS Pathog.* 7, e1002035.
- Jeong, R.D., Chandra-Shekar, A.C., Kachroo, A., Klessig, D.F., Kachroo, P., 2008. HRT-mediated hypersensitive response and resistance to turnip crinkle virus in Arabidopsis does not require the function of TIP, the presumed guard cell protein. *Mol. Plant–Microbe Interact.* 21, 1316–1324.
- Ji, L.H., Ding, S.W., 2001. The suppressor of transgene RNA silencing encoded by cucumber mosaic virus interferes with salicylic acid-mediated virus resistance. *Mol. Plant–Microbe Interact.* 14, 715–724.
- Jin, H., Zhu, J.K., 2010. A viral suppressor protein inhibits host RNA silencing by hooking up with Argonautes. *Genes Dev.* 24, 853–856.
- Jin, Y., Ma, D., Dong, J., Li, D., Deng, C., Jin, J., Wang, T., 2007. The HC-Pro protein of potato virus Y interacts with NtMinD of tobacco. *Mol. Plant–Microbe Interact.* 20, 1505–1511.
- Kachroo, A., Robin, G.P., 2013. Systemic signaling during plant defense. *Curr. Opin. Plant Biol.* 4, 527–533.
- Kalantidis, K., Schumacher, H.T., Alexiadis, T., Helm, J.M., 2008. RNA silencing movement in plants. *Biol. Cell* 100, 13–26.
- Kanazawa, A., Inaba, J., Shimura, H., Otagaki, S., Tsukahara, S., Matsuzawa, A., Kim, B. M., Goto, K., Masuta, C., 2011. Virus-mediated efficient induction of epigenetic modifications of endogenous genes with phenotypic changes in plants. *Plant J. Cell Mol. Biol.* 65, 156–168.
- Kanagarajan, S., Muthusamy, S., Gliszczynska, A., Lundgren, A., Brodelius, P.E., 2012. Functional expression and characterization of sesquiterpene synthases from *Artemisia annua* L. using transient expression system in *Nicotiana benthamiana*. *Plant Cell Rep.* 7, 1309–1319.
- Karran, R.A., Sanfacon, H., 2014. Tomato ringspot virus coat protein binds to ARGONAUTE 1 and suppresses the translation repression of a reporter gene. *Mol. Plant–Microbe Interact.* 27, 933–943.
- Kasschau, K.D., Cronin, S., Carrington, J.C., 1997. Genome amplification and long-distance movement functions associated with the central domain of tobacco etch potyvirus helper component-proteinase. *Virology* 228, 251–262.
- Kasschau, K.D., Xie, Z., Allen, E., Llave, C., Chapman, E.J., Krizan, K.A., Carrington, J.C., 2003. P1/HC-Pro, a viral suppressor of RNA silencing, interferes with Arabidopsis development and miRNA function. *Dev. Cell* 4, 205–217.
- Kataya, A.R., Suliman, M.N., Kalantidis, K., Livieratos, I.C., 2009. Cucurbit yellow stunting disorder virus p25 is a suppressor of post-transcriptional gene silencing. *Virus Res.* 145, 48–53.
- Kim, Y.J., Maizel, A., Chen, X., 2014. Traffic into silence: endomembranes and post-transcriptional RNA silencing. *EMBO J.* 33, 968–980.
- Kon, T., Sharma, P., Ikegami, M., 2007. Suppressor of RNA silencing encoded by the monopartite tomato leaf curl Java begomovirus. *Arch. Virol.* 152, 1273–1282.
- Kon, T., Yoshikawa, N., 2014. Induction and maintenance of DNA methylation in plant promoter sequences by apple latent spherical virus-induced transcriptional gene silencing. *Front. Microbiol.* 5, 595.
- Koukiokola, R., Sagana, S.M., Pezacki, J.P., 2007. Effects of pH and salt concentration on the siRNA binding activity of the RNA silencing suppressor protein p19. *FEBS Lett.* 581, 3051–3056.

- Kozłowska-Makulska, A., Guillel, H., Szyndel, M.S., Beuve, M., Lemaire, O., Herrbach, E., Bouzoubaa, S., 2010. P0 proteins of European beet-infecting poleroviruses display variable RNA silencing suppression activity. *J. Gen. Virol.* 91, 1082–1091.
- Kreuze, J.F., Savenkov, E.I., Cuellar, W., Li, X., Valkonen, J.P., 2005. Viral class 1 RNase III involved in suppression of RNA silencing. *J. Virol.* 79, 7227–7238.
- Kreuze, J.F., Savenkov, E.I., Valkonen, J.P.T., 2002. Complete genome sequence and analyses of the subgenomic RNAs of sweet potato chlorotic stunt virus reveal several new features for the genus crinivirus. *J. Virol.* 76, 9260–9270.
- Kubota, K., Tsuda, S., Tamai, A., Meshi, T., 2003. Tomato mosaic virus replication protein suppresses virus-targeted posttranscriptional gene silencing. *J. Virol.* 77, 11016–11026.
- Kumakura, N., Takeda, A., Fujioka, Y., Motose, H., Takano, R., Watanabe, Y., 2009. SGS3 and RDR6 interact and colocalize in cytoplasmic SGS3/RDR6-bodies. *FEBS Lett.* 583, 1261–1266.
- Lacombe, S., Bangratz, M., Vignols, F., Brugidou, C., 2010. The rice yellow mottle virus P1 protein exhibits dual functions to suppress and activate gene silencing. *Plant J. Cell Mol. Biol.* 61, 371–382.
- Laird, J., McNally, C., Carr, C., Doddiah, S., Yates, G., Chrysanthou, E., Khattab, A., Love, A.J., Geri, C., Sadanandom, A., Smith, B.O., Kobayashi, K., Milner, J.J., 2013. Identification of the domains of cauliflower mosaic virus protein P6 responsible for suppression of RNA silencing and salicylic acid signalling. *J. Gen. Virol.* 94, 2777–2789.
- Lakatos, L., Csorba, T., Pantaleo, V., Chapman, E.J., Carrington, J.C., Liu, Y.P., Dolja, V. V., Calvino, L.F., Lopez-Moya, J.J., Burgyan, J., 2006. Small RNA binding is a common strategy to suppress RNA silencing by several viral suppressors. *EMBO J.* 25, 2768–2780.
- Law, S.M., Zhang, B.W., Brooks, C.L., 2013. pH-sensitive residues in the P19 RNA silencing suppressor protein from Carnation Italian ringspot virus affect siRNA binding stability. *Prot. Sci. Publ. Prot. Soc.* 22, 595–604.
- Lee, Y.S., Nakahara, K., Pham, J.W., Kim, K., He, Z., Sontheimer, E.J., Carthew, R.W., 2004. Distinct roles for drosophila *dicer-1* and *dicer-2* in the siRNA/miRNA silencing pathways. *Cell* 117, 69–81.
- Lewsey, M., Robertson, F.C., Canto, T., Palukaitis, P., Carr, J.P., 2007. Selective targeting of miRNA-regulated plant development by a viral counter-silencing protein. *Plant J. Cell Mol. Biol.* 50, 240–252.
- Lewsey, M.G., Murphy, A.M., Maclean, D., Dalchau, N., Westwood, J.H., Macaulay, K., Bennett, M.H., Moulin, M., Hanke, D.E., Powell, G., Smith, A.G., Carr, J.P., 2010. Disruption of two defensive signaling pathways by a viral RNA silencing suppressor. *Mol. Plant–Microbe Interact.* 23, 835–845.
- Li, F., Pignatta, D., Bendix, C., Brunkard, J.O., Cohn, M.M., Tung, J., Sun, H., Kumar, P., Baker, B., 2012. MicroRNA regulation of plant innate immune receptors. *Proc. Natl. Acad. Sci. USA* 109, 1790–1795.
- Li, Y., Lu, J., Han, Y., Fan, X., Ding, S.W., 2013. RNA interference functions as an antiviral immunity mechanism in mammals. *Science* 342, 231–234.
- Li, F., Huang, C., Li, Z., Zhou, X., 2014. Suppression of RNA silencing by a plant DNA virus satellite requires a host calmodulin-like protein to repress RDR6 expression. *PLoS Pathog.* 10, e1003921.
- Li, H.W., Lucy, A.P., Guo, H.S., Li, W.X., Ji, L.H., Wong, S.M., Ding, S.W., 1999. Strong host resistance targeted against a viral suppressor of the plant gene silencing defence mechanism. *EMBO J.* 18, 2683–2691.
- Liao, Y.W., Sun, Z.H., Zhou, Y.H., Shi, K., Li, X., Zhang, G.Q., Xia, X.J., Chen, Z.X., Yu, J.Q., 2013. The role of hydrogen peroxide and nitric oxide in the induction of plant-encoded RNA-dependent RNA polymerase 1 in the basal defense against tobacco mosaic virus. *PLoS One* 8, e76090.
- Liu, F.X., Zhao, Q., Ruan, X.L., He, Y.W., Li, H.P., 2008. Suppressor of RNA silencing encoded by rice gall dwarf virus genome segment 11. *Chin. Sci. Bull.* 53, 362–369.
- Liu, J., Carmell, M.A., Rivas, F.V., Marsden, C.G., Thomson, J.M., Song, J.J., Hammond, S.M., Joshua-Tor, L., Hannon, G.J., 2004. Argonaute2 is the catalytic engine of mammalian RNAi. *Science* 305, 1437–1441.
- Liu, Y., Jin, W., Wang, L., Wang, X., 2014. Replication-associated proteins encoded by wheat dwarf virus act as RNA silencing suppressors. *Virus Res.* 190, 34–39.
- Llave, C., Kasschau, K.D., Carrington, J.C., 2000. Virus-encoded suppressor of posttranscriptional gene silencing targets a maintenance step in the silencing pathway. *Proc. Natl. Acad. Sci. USA* 97, 13401–13406.
- Love, A.J., Geri, C., Laird, J., Carr, C., Yun, B.W., Loake, G.J., Tada, Y., Sadanandom, A., Milner, J.J., 2012. Cauliflower mosaic virus protein P6 inhibits signaling responses to salicylic acid and regulates innate immunity. *PLoS One* 7, e47535.
- Love, A.J., Laird, J., Holt, J., Hamilton, A.J., Sadanandom, A., Milner, J.J., 2007. Cauliflower mosaic virus protein P6 is a suppressor of RNA silencing. *J. Gen. Virol.* 88, 3439–3444.
- Lozsa, R., Csorba, T., Lakatos, L., Burgyan, J., 2008. Inhibition of 3' modification of small RNAs in virus-infected plants require spatial and temporal co-expression of small RNAs and viral silencing-suppressor proteins. *Nucleic Acids Res.* 36, 4099–4107.
- Lu, R., Folimonov, A., Shintaku, M., Li, W.X., Falk, B.W., Dawson, W.O., Ding, S.W., 2004. Three distinct suppressors of RNA silencing encoded by a 20-kb viral RNA genome. *Proc. Natl. Acad. Sci. USA* 101, 15742–15747.
- Lukhovitskaya, N.I., Vetukuri, R.R., Sama, I., Thaduri, S., Solovyev, A.G., Savenkov, E. I., 2014. A viral transcription factor exhibits antiviral RNA silencing suppression activity independent of its nuclear localization. *J. Gen. Virol.* , <http://dx.doi.org/10.1099/vir.0.067884-0>.
- Mallory, A., Vaucheret, H., 2010. Form, function, and regulation of ARGONAUTE proteins. *Plant Cell* 22, 3879–3889.
- Maillard, P.V., Ciaudo, C., Marchais, A., Li, Y., Jay, F., Ding, S.W., Voinnet, O., 2013. Antiviral RNA interference in mammalian cells. *Science* 342, 235–238.
- Mallory, A.C., Elmayan, T., Vaucheret, H., 2008. MicroRNA maturation and action—the expanding roles of ARGONAUTES. *Curr. Opin. Plant Biol.* 11, 560–566.
- Mallory, A.C., Ely, L., Smith, T.H., Marathe, R., Anandalakshmi, R., Fagard, M., Vaucheret, H., Pruss, G., Bowman, L., Vance, V.B., 2001. HC-Pro suppression of transgene silencing eliminates the small RNAs but not transgene methylation or the mobile signal. *Plant Cell* 13, 571–583.
- Mallory, A.C., Vaucheret, H., 2009. ARGONAUTE 1 homeostasis invokes the coordinate action of the microRNA and siRNA pathways. *EMBO Rep.* 10, 521–526.
- Mangwende, T., Wang, M.L., Borth, W., Hu, J., Moore, P.H., Mirkov, T.E., Albert, H.H., 2009. The P0 gene of Sugarcane yellow leaf virus encodes an RNA silencing suppressor with unique activities. *Virology* 384, 38–50.
- Mandadi, K.K., Scholthof, K.B., 2013. Plant immune responses against viruses: how does a virus cause disease? *Plant Cell* 25, 1489–1505.
- Martín-Hernández, A.M., Baulcombe, D.C., 2008. Tobacco rattle virus 16-kilodalton protein encodes a suppressor of RNA silencing that allows transient viral entry in meristems. *J. Virol.* 82, 4064–4071.
- Martínez-Priego, L., Donaire, L., Barajas, D., Llave, C., 2008. Silencing suppressor activity of the Tobacco rattle virus-encoded 16-kDa protein and interference with endogenous small RNA-guided regulatory pathways. *Virology* 376, 346–356.
- Martinez-Turino, S., Hernandez, C., 2009. Inhibition of RNA silencing by the coat protein of Pelargonium flower break virus: distinctions from closely related suppressors. *J. Gen. Virol.* 90, 519–525.
- Mathioudakis, M.M., Rodríguez-Moreno, L., Sempere, R.N., Aranda, M.A., Livieratos, I., 2014. Multifaceted capsid proteins: multiple interactions suggest multiple roles for pepino mosaic virus capsid protein. *Mol. Plant–Microbe Interact.* 27, 1356–1369.
- Mathioudakis, M.M., Veiga, R.S., Cant, T., Medina, V., Mossialos, D., Makris, A.M., Livieratos, I., 2013. Pepino mosaic virus triple gene block protein 1 (TGBp1) interacts with and increases tomato catalase 1 activity to enhance virus accumulation. *Mol. Plant Pathol.* 14, 589–601.
- Mayo, M.A., Ziegler-Graff, V., 1996. Molecular biology of luteoviruses. *Adv. Virus Res.* 46, 413–460.
- Meng, C., Chen, J., Peng, J., Wong, S.M., 2006. Host-induced avirulence of hibiscus chlorotic ringspot virus mutants correlates with reduced gene-silencing suppression activity. *J. Gen. Virol.* 87, 451–459.
- Meraí, Z., Kerenyi, Z., Kertesz, S., Magna, M., Lakatos, L., Silhavy, D., 2006. Double-stranded RNA binding may be a general plant RNA viral strategy to suppress RNA silencing. *J. Virol.* 80, 5747–5756.
- Meraí, Z., Kerenyi, Z., Molnar, A., Barta, E., Valoczi, A., Bisztray, G., Havelda, Z., Burgyan, J., Silhavy, D., 2005. Aureusvirus P14 is an efficient RNA silencing suppressor that binds double-stranded RNAs without size specificity. *J. Virol.* 79, 7217–7226.
- Mi, S., Cai, T., Hu, Y., Chen, Y., Hodges, E., Ni, F., Wu, L., Li, S., Zhou, H., Long, C., Chen, S., Hannon, G.J., Qi, Y., 2008. Sorting of small RNAs into *Arabidopsis argonaute* complexes is directed by the 5' terminal nucleotide. *Cell* 133, 116–127.
- Moissiard, G., Voinnet, O., 2006. RNA silencing of host transcripts by cauliflower mosaic virus requires coordinated action of the four *Arabidopsis* *dicer*-like proteins. *Proc. Natl. Acad. Sci. USA* 103, 19593–19598.
- Molnar, A., Csorba, T., Lakatos, L., Varallyay, E., Lacomme, C., Burgyan, J., 2005. Plant virus-derived small interfering RNAs originate predominantly from highly structured single-stranded viral RNAs. *J. Virol.* 79, 7812–7818.
- Molnar, A., Melnyk, C.W., Bassett, A., Hardcastle, T.J., Dunn, R., Baulcombe, D.C., 2010. Small silencing RNAs in plants are mobile and direct epigenetic modification in recipient cells. *Science* 328, 872–875.
- Morel, J.B., Godon, C., Mourrain, P., Beclin, C., Boutet, S., Feuerbach, F., Proux, F., Vaucheret, H., 2002. Fertile hypomorphic ARGONAUTE (*ago1*) mutants impaired in post-transcriptional gene silencing and virus resistance. *Plant Cell* 14, 629–639.
- Moreno, A.B., Martínez de Alba, A.E., Bardou, F., Crespi, M.D., Vaucheret, H., Maizel, A., Mallory, A.C., 2013. Cytoplasmic and nuclear quality control and turnover of single-stranded RNA modulate post-transcriptional gene silencing in plants. *Nucleic Acids Res.* 41, 4699–4708.
- Mourrain, P., Beclin, C., Elmayan, T., Feuerbach, F., Godon, C., Morel, J.B., Jouette, D., Lacombe, A.M., Nikic, S., Picault, N., Remoue, K., Sanial, M., Vo, T.A., Vaucheret, H., 2000. *Arabidopsis* *SGS2* and *SGS3* genes are required for posttranscriptional gene silencing and natural virus resistance. *Cell* 101, 533–542.
- Muangsan, N., Beclin, C., Vaucheret, H., Robertson, D., 2004. Geminivirus VIGS of endogenous genes requires *SGS2/SDE1* and *SGS3* and defines a new branch in the genetic pathway for silencing in plants. *Plant J. Cell Mol. Biol.* 38, 1004–1014.
- Naim, F., Nakasugi, K., Crowhurst, R.N., Hilario, E., Zwart, A.B., Hellens, R.P., Taylor, J. M., Waterhouse, P.M., Wood, C.C., 2012. Advanced engineering of lipid metabolism in *Nicotiana benthamiana* using a draft genome and the V2 viral silencing-suppressor protein. *PLoS One* 7, e52717.
- Nakahara, K.S., Masuta, C., Yamada, S., Shimura, H., Kashihara, Y., Wada, T.S., Meguro, A., Goto, K., Tadamura, K., Sueda, K., Sekiguchi, T., Shao, J., Itchoda, N., Matsumura, T., Igarashi, M., Ito, K., Carthew, R.W., Uyeda, I., 2012. Tobacco calmodulin-like protein provides secondary defense by binding to and directing degradation of virus RNA silencing suppressors. *Proc. Natl. Acad. Sci. USA* 109, 10113–10118.
- Nasher, N., Cheng, J., Singaravelu, R., Wu, P., McDermott, M.T., Pezacki, J.P., 2011. An enzyme-linked assay for the rapid quantification of microRNAs based on the viral suppressor of RNA silencing protein p19. *Anal. Biochem.* 412, 165–172.



- Nawaz-ul-Rehman, M.S., Nahid, N., Mansoor, S., Briddon, R.W., Fauquet, C.M., 2010. Post-transcriptional gene silencing suppressor activity of two non-pathogenic alphasatellites associated with a begomovirus. *Virology* 405, 300–308.
- Nemes, K., Gellert, A., Balazs, E., Salanki, K., 2014. Alanine scanning of cucumber mosaic virus (CMV) 2B protein identifies different positions for cell-to-cell movement and gene silencing suppressor activity. *PLoS One* 9, e11209.
- Okano, Y., Senshu, H., Hashimoto, M., Neriya, Y., Netsu, O., Minato, N., Yoshida, T., Maejima, K., Oshima, K., Komatsu, K., Yamaji, Y., Namba, S., 2014. In planta recognition of a double-stranded RNA synthesis protein complex by a potexviral RNA silencing suppressor. *Plant Cell* 26, 2168–2183.
- Olsper, A., Kamsol, K., Sarmiento, C., Gerassimenko, J., Truve, E., 2014. Cocksfoot mottle virus coat protein is dispensable for the systemic infection. *Virology* 471, 19.
- Olsper, A., Paves, H., Toomela, R., Tamm, T., Truve, E., 2010. Cocksfoot mottle virus coat protein contains two nuclear localization signals. *Virus Genes* 40, 423–431.
- Omarov, R., Sparks, K., Smith, L., Zindovic, J., Scholthof, H.B., 2006. Biological relevance of a stable biochemical interaction between the Tombusvirus-encoded P19 and short interfering RNAs. *J. Virol.* 80, 3000–3008.
- Otagaki, S., Kawai, M., Masuta, C., Kanazawa, A., 2011. Size and positional effects of promoter RNA segments on virus-induced RNA-directed DNA methylation and transcriptional gene silencing. *Epigenetics: Off. J. DNA Methylation Soc.* 6, 681–691.
- Pandey, S.P., Shahi, P., Gase, K., Baldwin, I.T., 2008. Herbivory-induced changes in the small-RNA transcriptome and phytohormone signaling in *Nicotiana attenuata*. *Proc. Natl. Acad. Sci. USA* 105, 4559–4564.
- Pantaleo, V., Szittyá, G., Burgyan, J., 2007. Molecular bases of viral RNA targeting by viral small interfering RNA-programmed RISC. *J. Virol.* 81, 3797–3806.
- Papp, I., Mette, M.F., Aufsatz, W., Daxinger, L., Schauer, S.E., Ray, A., van der Winden, J., Matzke, M., Matzke, A.J., 2003. Evidence for nuclear processing of plant micro RNA and short interfering RNA precursors. *Plant Physiol.* 132, 1382–1390.
- Parent, J.S., Bouteiller, N., Elmayer, T., Vaucheret, H., 2015. Respective contributions of *Arabidopsis* DCL2 and DCL4 to RNA silencing. *Plant J.* 81, 223–232.
- Park, J.W., Faure-Rabasse, S., Robinson, M.A., Desvoyes, B., Scholthof, H.B., 2004. The multifunctional plant viral suppressor of gene silencing P19 interacts with itself and an RNA binding host protein. *Virology* 323, 49–58.
- Park, M.Y., Wu, G., Gonzalez-Sulser, A., Vaucheret, H., Poethig, R.S., 2005. Nuclear processing and export of microRNAs in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 102, 3691–3696.
- Pazhouhandeh, M., Dieterle, M., Marrocco, K., Lechner, E., Berry, B., Brault, V., Hemmer, O., Kretsch, T., Richards, K.E., Genschik, P., Ziegler-Graff, V., 2006. F-box-like domain in the poliovirus protein P0 is required for silencing suppressor function. *Proc. Natl. Acad. Sci. USA* 103, 1994–1999.
- Peragine, A., Yoshikawa, M., Wu, G., Albrecht, H.L., Poethig, R.S., 2004. SGS3 and SGS2/SDE1/RDR6 are required for juvenile development and the production of trans-acting siRNAs in *Arabidopsis*. *Genes Dev.* 18, 2368–2379.
- Pérez-Cañamás, M., Hernández, C., 2015. Key importance of small RNA binding for the activity of a glycine–tryptophan (GW) motif-containing viral suppressor of RNA silencing. *J. Biol. Chem.* 290, 3106–3120.
- Pfeffer, S., Dunoyer, P., Heim, F., Richards, K.E., Jonard, G., Ziegler-Graff, V., 2002. PO of beet Western yellows virus is a suppressor of posttranscriptional gene silencing. *J. Virol.* 76, 6815–6824.
- Pham, J.W., Pellino, J.L., Lee, Y.S., Carthew, R.W., Sontheimer, E.J., 2004. A dicer-2-dependent 80s complex cleaves targeted mRNAs during RNAi in *Drosophila*. *Cell* 117, 83–94.
- Powers, J.G., Sit, T.L., Heinsohn, C., George, C.G., Kim, K.H., Lommel, S.A., 2008. The red clover necrotic mosaic virus RNA-2 encoded movement protein is a second suppressor of RNA silencing. *Virology* 381, 277–286.
- Pruss, G., Ge, X., Shi, X.M., Carrington, J.C., Bowman Vance, V., 1997. Plant viral synergism: the potyviral genome encodes a broad-range pathogenicity enhancer that transactivates replication of heterologous viruses. *Plant Cell* 9, 859–868.
- Pumplin, N., Voinnet, O., 2013. RNA silencing suppression by plant pathogens: defence, counter-defence and counter–counter-defence. *Nat. Rev. Microbiol.* 11, 745–760.
- Qavi, A.J., Kindt, J.T., Bailey, R.C., 2010. Sizing up the future of microRNA analysis. *Anal. Bioanal. Chem.* 398, 2535–2549.
- Qu, F., Ren, T., Morris, T.J., 2003. The coat protein of turnip crinkle virus suppresses posttranscriptional gene silencing at an early initiation step. *J. Virol.* 77, 511–522.
- Qu, F., Ye, X., Morris, T.J., 2008. *Arabidopsis* DRB4, AGO1, AGO7, and RDR6 participate in a DCL4-initiated antiviral RNA silencing pathway negatively regulated by DCL1. *Proc. Natl. Acad. Sci. USA* 105, 14732–14737.
- Rahim, M.D., Andika, I.B., Han, C., Kondo, H., Tamada, T., 2007. RNA4-encoded p31 of beet necrotic yellow vein virus is involved in efficient vector transmission, symptom severity and silencing suppression in roots. *J. Gen. Virol.* 88, 1611–1619.
- Raja, P., Jackel, J.N., Li, S., Heard, I.M., Bisaro, D.M., 2014. *Arabidopsis* double-stranded RNA binding protein DRB3 participates in methylation-mediated defense against geminiviruses. *J. Virol.* 88, 2611–2622.
- Raja, P., Sanville, B.C., Buchmann, R.C., Bisaro, D.M., 2008. Viral genome methylation as an epigenetic defense against geminiviruses. *J. Virol.* 82, 8997–9007.
- Rajagopalan, R., Vaucheret, H., Trejo, J., Bartel, D.P., 2006. A diverse and evolutionarily fluid set of microRNAs in *Arabidopsis thaliana*. *Genes Dev.* 20, 3407–3425.
- Rajamäki, M.L., Valkonen, J.P.T., 2009. Control of nuclear and nucleolar localization of nuclear inclusion protein A of picorna-like potato virus A in *Nicotiana* species. *Plant Cell* 21, 2485–2502.
- Rajamäki, M.L., Streng, J., Valkonen, J.P.T., 2014. Silencing suppressor protein VPg of a potyvirus interacts with the plant silencing-related protein SGS3. *Mol. Plant–Microbe Interact.* 27, 1199–1210.
- Rawlings, R.A., Krishnan, V., Walter, N.G., 2011. Viral RNAi suppressor reversibly binds siRNA to outcompete dicer and RISC via multiple-turnover. *J. Mol. Biol.* 408, 262–276.
- Reavy, B., Dawson, S., Canto, T., MacFarlane, S.A., 2004. Heterologous expression of plant virus genes that suppress post-transcriptional gene silencing results in suppression of RNA interference in *Drosophila* cells. *BMC Biotechnol.* 4, 18.
- Reed, J.C., Kasschau, K.D., Prokhnovsky, A.I., Gopinath, K., Pogue, G.P., Carrington, J.C., Dolja, V.V., 2003. Suppressor of RNA silencing encoded by beet yellows virus. *Virology* 15, 203–209.
- Ren, B., Guo, Y., Gao, F., Zhou, P., Wu, F., Meng, Z., Wei, C., Li, Y., 2010. Multiple functions of rice dwarf phytoevirus Pns10 in suppressing systemic RNA silencing. *J. Virol.* 84, 12914–12923.
- Ren, T., Qu, F., Morris, T.J., 2000. HRT gene function requires interaction between a NAC protein and viral capsid protein to confer resistance to turnip crinkle virus. *Plant Cell* 12, 1917–1925.
- Renovell, A., Vives, M.C., Ruiz-Ruiz, S., Navarro, L., Moreno, P., Guerri, J., 2012. The citrus leaf blotch virus movement protein acts as silencing suppressor. *Virus Genes* 44, 131–140.
- Rhoades, M.W., Reinhart, B.J., Lim, L.P., Burge, C.B., Bartel, B., Bartel, D.P., 2002. Prediction of plant microRNA targets. *Cell* 110, 513–520.
- Rodríguez, A., Angel, C.A., Lutz, L., Leisner, S.M., Nelson, R.S., Schoelz, J.E., 2014. Association of the P6 protein of cauliflower mosaic virus with plasmodesmata and plasmodesmal proteins. *Plant Physiol.* 166, 1345–1358.
- Ronde, D., Pasquier, A., Ying, S., Butterbach, P., Lohuis, D., Kormelink, R., 2014. Analysis of tomato spotted wilt virus NSs protein indicates the importance of the N-terminal domain for avirulence and RNA silencing suppression. *Mol. Plant Pathol.* 15, 185–195.
- Ruiz-Ruiz, S., Soler, N., Sanchez-Navarro, J., Fagoaga, C., Lopez, C., Navarro, L., Moreno, P., Pena, L., Flores, R., 2013. Citrus tristeza virus p23: determinants for nucleolar localization and their influence on suppression of RNA silencing and pathogenesis. *Mol. Plant–Microbe Interact.* 26, 306–318.
- Sadowy, E., Maasen, A., Juszczak, M., David, C., Zagórski-Ostojka, W., Gronenborn, B., Hulanicka, M.D., 2001. The ORFO product of potato leafroll virus is indispensable for virus accumulation. *J. Gen. Virol.* 82, 1529–1532.
- Sahana, N., Kaur, H., Jain, R.K., Palukaitis, P., Canto, T., Praveen, S., 2014. The asparagine residue in the FRNK box of potyviral helper-component protease is critical for its small RNA binding and subcellular localization. *J. Gen. Virol.* 95, 1167–1177.
- Sansregret, R., Dufour, V., Langlois, M., Daayf, F., Dunoyer, P., Voinnet, O., Bouarab, K., 2013. Extreme resistance as a host counter-counter defense against viral suppression of RNA silencing. *PLoS Pathog.* 9, e1003435.
- Sarmiento, C., Gomez, E., Meier, M., Kavanagh, T.A., Truve, E., 2007. Cocksfoot mottle virus P1 suppresses RNA silencing in *Nicotiana benthamiana* and *Nicotiana tabacum*. *Virus Res.* 123, 95–99.
- Schnettler, E., Hemmes, H., Huisman, R., Goldbach, R., Prins, M., Kormelink, R., 2010. Diverging affinity of tospovirus RNA silencing suppressor proteins, NSs, for various RNA duplex molecules. *J. Virol.* 84, 11542–11554.
- Scholthof, H.B., Alvarado, V.Y., Vega-Arrequin, J.C., Ciomperlik, J., Odokonyero, D., Brosseau, C., Jaubert, M., Zamora, A., Moffett, P., 2011. Identification of an ARGONAUTE for antiviral RNA silencing in *Nicotiana benthamiana*. *Plant Physiol.* 156, 1548–1555.
- Schott, G., Mari-Ordóñez, A., Humber, C., Alioua, A., Voinnet, O., Dunoyer, P., 2012. Differential effects of viral silencing suppressors on siRNA and miRNA loading support the existence of two distinct cellular pools of ARGONAUTE1. *EMBO J.* 31, 2553–2565.
- Schuck, J., Gursinsky, T., Pantaleo, V., Burgyan, J., Behrens, S.E., 2013. AGO/RISC-mediated antiviral RNA silencing in a plant in vitro system. *Nucleic Acids Res.* 41, 5090–5103.
- Schwach, F., Vaistij, F.E., Jones, L., Baulcombe, D.C., 2005. An RNA-dependent RNA polymerase prevents meristem invasion by potato virus X and is required for the activity but not the production of a systemic silencing signal. *Plant Physiol.* 138, 1842–1852.
- Schwartz, M., Chen, J., Janda, M., Sullivan, M., den Boon, J., Ahlquist, P., 2002. A positive-strand RNA virus replication complex parallels form and function of retrovirus capsids. *Mol. Cell* 9, 505–514.
- Schwarz, D.S., Hutvagner, G., Du, T., Xu, Z., Aronin, N., Zamore, P.D., 2003. Asymmetry in the assembly of the RNAi enzyme complex. *Cell* 115, 199–208.
- Senshu, H., Ozeki, J., Komatsu, K., Hashimoto, M., Hatada, K., Aoyama, M., Kagiwada, S., Yamaji, Y., Namba, S., 2009. Variability in the level of RNA silencing suppression caused by triple gene block protein 1 (TGBp1) from various potexviruses during infection. *J. Gen. Virol.* 90, 1014–1024.
- Senshu, H., Yamaji, Y., Minato, N., Shiraiishi, T., Maejima, K., Hashimoto, M., Miura, C., Neriya, Y., Namba, S., 2011. A dual strategy for the suppression of host antiviral silencing: two distinct suppressors for viral replication and viral movement encoded by potato virus M. *J. Virol.* 85, 10269–10278.
- Seo, J.K., Kang, S.H., Seo, B.Y., Jung, J.K., Kim, K.H., 2010. Mutational analysis of interaction between coat protein and helper component-proteinase of Soybean mosaic virus involved in aphid transmission. *Mol. Plant Pathol.* 11, 265–276.
- Sharma, P., Ikegami, M., Kon, T., 2010. Identification of the virulence factors and suppressors of posttranscriptional gene silencing encoded by *Ageratum* yellow vein virus, a monopartite begomovirus. *Virus Res.* 149, 19–27.
- Sharma, P., Ikegami, M., 2010. Tomato leaf curl Java virus V2 protein is a determinant of virulence, hypersensitive response and suppression of post-transcriptional gene silencing. *Virology* 396, 85–93.

- Sharma, P., Matsuda, N., Bajet, N.B., Ikegami, M., 2011. Molecular analysis of new isolates of Tomato leaf curl Philippines virus and an associated betasatellite occurring in the Philippines. *Arch. Virol.* 156, 305–312.
- Shen, M., Xu, Y., Jia, R., Zhou, X., Ye, K., 2010. Size-independent and noncooperative recognition of dsRNA by the rice stripe virus RNA silencing suppressor NS3. *J. Mol. Biol.* 404, 665–679.
- Shen, W.J., Ruan, X.L., Li, X.S., Zhao, Q., Li, H.P., 2012. RNA silencing suppressor Pns11 of rice gall dwarf virus induces virus-like symptoms in transgenic rice. *Arch. Virol.* 157, 1531–1539.
- Shen, W., Yan, P., Gao, L., Pan, X., Wu, J., Zhou, P., 2010a. Helper component-proteinase (HC-Pro) protein of papaya ringspot virus interacts with papaya calreticulin. *Mol. Plant Pathol.* 11, 335–346.
- Shen, W.T., Wang, M.Q., Yan, P., Gao, L., Zhou, P., 2010b. Protein interaction matrix of papaya ringspot virus type P based on a yeast two-hybrid system. *Acta Virol.* 54, 49–54.
- Shiboleth, Y.M., Haronsky, E., Leibman, D., Arazi, T., Wassenegger, M., Whitham, S. A., Gaba, V., Gal-On, A., 2007. The conserved FRNK box in HC-Pro, a plant viral suppressor of gene silencing, is required for small RNA binding and mediates symptom development. *J. Virol.* 81, 13135–13148.
- Shimuram, H., Masuta, C., Yoshida, N., Sueda, K., Suzuki, M., 2013. The 2b protein of Asparagus virus 2 functions as an RNA silencing suppressor against systemic silencing to prove functional synteny with related cucumoviruses. *Virology* 442, 180–188.
- Shivaprasad, P.V., Chen, H.M., Patel, K., Bond, D.M., Santos, B.A., Baulcombe, D.C., 2012. A microRNA superfamily regulates nucleotide binding site-leucine-rich repeats and other mRNAs. *Plant Cell* 24, 859–874.
- Shivaprasad, P.V., Rajeswaran, R., Blevins, T., Schoelz, J., Meins, F., Hohn, T., Pooggin, M.M., 2008. The CaMV transactivator/viroplasm interferes with RDR6-dependent trans-acting and secondary siRNA pathways in Arabidopsis. *Nucleic acid Res.* 36, 5896–5909.
- Sijen, T., Fleenor, J., Simmer, F., Thijsen, K.L., Parrish, S., Timmons, L., Plasterk, R.H., Fire, A., 2001. On the role of RNA amplification in dsRNA-triggered gene silencing. *Cell* 107, 465–476.
- Silhavy, D., Molnar, A., Lucieli, A., Szitty, G., Hornyik, C., Tavazza, M., Burgyan, J., 2002. A viral protein suppresses RNA silencing and binds silencing-generated, 21- to 25-nucleotide double-stranded RNAs. *EMBO* 21, 3070–3080.
- Soitamo, A.J., Jada, B., Lehto, K., 2011. HC-Pro silencing suppressor significantly alters the gene expression profile in tobacco leaves and flowers. *BMC Plant Biol.* 11, 68.
- Soitamo, A.J., Jada, B., Lehto, K., 2012. Expression of geminiviral AC2 RNA silencing suppressor changes sugar and jasmonate responsive gene expression in transgenic tobacco plants. *BMC Plant Biol.* 12, 204.
- Sun, L., Andika, I.B., Kondo, H., Chen, J., 2013a. Identification of the amino acid residues and domains in the cysteine-rich protein of Chinese wheat mosaic virus that are important for RNA silencing suppression and subcellular localization. *Mol. Plant Pathol.* 14, 265–278.
- Sun, L., Andika, I.B., Shen, J., Yang, D., Ratti, C., Chen, J., 2013b. The CUG-initiated larger form coat protein of Chinese wheat mosaic virus binds to the cysteine-rich RNA silencing suppressor. *Virus Res.* 177, 66–74.
- Szabo, E.Z., Manczinger, M., Goblos, A., Kemeny, L., Lakatos, L., 2012. Switching on RNA silencing suppressor activity by restoring argonaute binding to a viral protein. *J. Virol.* 86, 8324–8327.
- Szitty, G., Moxon, S., Pantaleo, V., Toth, G., Rusholme Pilcher, R.L., Moulton, V., Burgyan, J., Dalmay, T., 2010. Structural and functional analysis of viral siRNAs. *PLoS Pathog.* 6, e1000838.
- Takeda, A., Iwasaki, S., Watanabe, T., Utsumi, M., Watanabe, Y., 2008. The mechanism selecting the guide strand from small RNA duplexes is different among argonaute proteins. *Plant Cell Physiol.* 49, 493–500.
- Takeda, A., Tsukuda, M., Mizumoto, H., Okamoto, K., Kaido, M., Mise, K., Okuno, T., 2005. A plant RNA virus suppresses RNA silencing through viral RNA replication. *EMBO* 24, 3147–3157.
- Tatineni, S., Qu, F., Li, R., Morris, T.J., French, R., 2012. Triticum mosaic poacevirus enlists P1 rather than HC-Pro to suppress RNA silencing-mediated host defense. *Virology* 433, 104–115.
- Te, J., Melcher, U., Howard, A., Verchot-Lubicz, J., 2005. Soilborne wheat mosaic virus (SBWMV) 19K protein belongs to a class of cysteine rich proteins that suppress RNA silencing. *Virol. J.* 2, 18.
- Tian, D., Traw, M.B., Chen, J.Q., Kreitman, M., Bergelson, J., 2003. Fitness costs of R-gene-mediated resistance in *Arabidopsis thaliana*. *Nature* 423, 74–77.
- Tilsner, J., Linnik, O., Wright, K.M., Bell, K., Roberts, A.G., Lacomme, C., Cruz, S.S., Oparka, K.J., 2012. The TGB1 movement protein of potato virus X reorganizes actin and endomembranes into the X-body, a viral replication factory. *Plant Physiol.* 158, 1359–1370.
- Tomari, Y., Du, T., Haley, B., Schwarz, D.S., Bennett, R., Cook, H.A., Koppetsch, B.S., Theurkauf, W.E., Zamore, P.D., 2004. RISC assembly defects in the drosophila RNAi mutant armitage. *Cell* 116, 831–841.
- Torres-Barcelo, C., Daros, J.A., Elena, S.F., 2010. HC-Pro hypo- and hypersuppressor mutants: differences in viral siRNA accumulation in vivo and siRNA binding activity in vitro. *Arch. Virol.* 155, 251–254.
- Torres-Barcelo, C., Martin, S., Daros, J.A., Elena, S.F., 2008. From hypo- to hypersuppression: effect of amino acid substitutions on the RNA-silencing suppressor activity of the tobacco etch potyvirus HC-Pro. *Genetics* 180, 1039–1049.
- Trinks, D., Rajeswaran, R., Shivaprasad, P.V., Akbergenov, R., Oakeley, E.J., Veluthambi, K., Hohn, T., Pooggin, M.M., 2005. Suppression of RNA silencing by a geminivirus nuclear protein, AC2, correlates with transactivation of host genes. *J. Virol.* 79, 2517–2527.
- Uhrig, J.F., Canto, T., Marshall, D., MacFarlane, S.A., 2004. Relocalization of nuclear ALY proteins to the cytoplasm by the tomato bushy stunt virus P19 pathogenicity protein. *Plant Physiol.* 135, 2411–2423.
- Vaistij, F.E., Jones, L., Baulcombe, D.C., 2002. Spreading of RNA targeting and DNA methylation in RNA silencing requires transcription of the target gene and a putative RNA-dependent RNA polymerase. *Plant Cell* 14, 857–867.
- Valli, A., Dujovny, G., Garcia, J.A., 2008. Protease activity, self interaction, and small interfering RNA binding of the silencing suppressor P1b0 from cucumber vein yellowing Ipomovirus. *J. Virol.* 82, 974–986.
- Valli, A., Martin-Hernandez, A.M., Lopez-Moya, J.J., Garcia, J.A., 2006. RNA silencing suppression by a second copy of the P1 serine protease of cucumber vein yellowing ipomovirus, a member of the family Potyviridae that lacks the cysteine protease HCPro. *J. Virol.* 80, 10055–10063.
- Valli, A., Oliveros, J.C., Molnar, A., Baulcombe, D.C., Garcia, J.A., 2011. The specific binding to 21-nt double-stranded RNAs is crucial for the anti-silencing activity of cucumber vein yellowing virus P1b and perturbs endogenous small RNA populations. *RNA* 17, 1148–1158.
- Varallyay, E., Olah, E., Havelda, Z., 2014. Independent parallel functions of p19 plant viral suppressor of RNA silencing required for effective suppressor activity. *Nucleic Acid Res.* 42, 599–608.
- Várallyay, E., Havelda, Z., 2013. Unrelated viral suppressors of RNA silencing mediate the control of ARGONAUTE1 level. *Mol. Plant Pathol.* 14, 567–575.
- Varallyay, E., Valoczi, A., Agyi, A., Burgyan, J., Havelda, Z., 2010. Plant virus-mediated induction of miR168 is associated with repression of ARGONAUTE1 accumulation. *EMBO J.* 29, 3507–3519.
- Vargason, J.M., Szitty, G., Burgyan, J., Hall, T.M., 2003. Size selective recognition of siRNA by an RNA silencing suppressor. *Cell* 115, 799–811.
- Vaucheret, H., Mallory, A.C., Bartel, D.P., 2006. AGO1 homeostasis entails coexpression of MIR168 and AGO1 and preferential stabilization of miR168 by AGO1. *Mol. Cell* 22, 129–136.
- Vogler, H., Akbergenov, R., Shivaprasad, P.V., Dang, V., Fasler, M., Kwon, M.O., Zhanybekova, S., Hohn, T., Heinlein, M., 2007. Modification of small RNAs associated with suppression of RNA silencing by tobamovirus replicase protein. *J. Virol.* 81, 10379–10388.
- Vogler, H., Kwon, M.O., Dang, V., Sambade, A., Fasler, M., Ashby, J., Heinlein, M., 2008. Tobacco mosaic virus movement protein enhances the spread of RNA silencing. *PLoS Pathog.* 4, e1000038.
- Voinnet, O., Vain, P., Angell, S., Baulcombe, D.C., 1998. Systemic spread of sequence-specific transgene RNA degradation in plants is initiated by localized introduction of ectopic promoterless DNA. *Cell* 95, 177–187.
- Wang, H., Buckley, K.J., Yang, X., Buchmann, R.C., Bisaro, D.M., 2005. Adenosine kinase inhibition and suppression of RNA silencing by geminivirus AL2 and L2 proteins. *J. Virol.* 79, 7410–7418.
- Wang, H., Hao, L., Shung, C.Y., Sunter, G., Bisaro, D.M., 2003. Adenosine kinase is inactivated by geminivirus AL2 and L2 proteins. *Plant Cell* 15, 3020–3032.
- Wang, K.D., Empleo, R., Nguyen, T.T., Moffett, P., Sacco, M.A., 2014a. Elicitation of hypersensitive responses in *Nicotiana glauca* by the suppressor of RNA silencing protein P0 from poleroviruses. *Mol. Plant Pathol.* , <http://dx.doi.org/10.1111/mpp.12201>.
- Wang, L.Y., Lin, S.S., Hung, T.H., Li, T.K., Lin, N.C., Shen, T.L., 2012. Multiple domains of the tobacco mosaic virus p126 protein can independently suppress local and systemic RNA silencing. *Mol. Plant – Microbe Interact.* 25, 648–657.
- Wang, X.B., Jovel, J., Udornporn, P., Wang, Y., Wu, Q., Li, W.X., Gascioli, V., Vaucheret, H., Ding, S.W., 2011. The 21-nucleotide, but not 22-nucleotide, viral secondary small interfering RNAs direct potent antiviral defense by two cooperative argonautes in *Arabidopsis thaliana*. *Plant Cell* 23, 1625–1638.
- Wang, Y., Dang, M., Hou, H., Mei, Y., Qian, Y., Zhou, X., 2014b. Identification of an RNA silencing suppressor encoded by a mastrevirus. *J. Gen. Virol.* 95, 2082–2088.
- Wassenegger, M., Krczal, G., 2006. Nomenclature and functions of RNA-directed RNA polymerases. *Trends Plant Sci.* 11, 142–151.
- Weinheimer, I., Boonrod, K., Moser, M., Zwiebel, M., Füllgrabe, M., Krczal, G., Wassenegger, M., 2010. Analysis of an autolytic activity of rice yellow mottle virus silencing suppressor P1. *Biol. Chem.* 391, 271–281.
- Westwood, J.H., McCann, Lucy, Naish, Matthew, Dixon, Heather, Murphy, Alex M., Stancombe, Matthew A., Bennett, Mark H., Powell, Glen, Webb, Alex A.R., John, P., 2013a. Carr: a viral RNA silencing suppressor interferes with abscisic acid-mediated signalling and induces drought tolerance in *Arabidopsis thaliana*. *Mol. Plant Pathol.* 14, 158–170.
- Westwood, J.H., Lewsey, M.G., Murphy, A.M., Tungadi, T., Bates, A., Gilligan, C.A., Carr, J.P., 2014. Interference with jasmonic acid-regulated gene expression is a general property of viral suppressors of RNA silencing but only partly explains virus-induced changes in plant-aphid interactions. *J. Gen. Virol.* 95, 733–739.
- Westwood, J.H., Groen, Simon C., Du, Zhiyou, Murphy, Alex M., Tri Anggoro, Damar, Tungadi, Trisna, Luang-In, Vijtra, Lewsey, Mathew G., Rossiter, John T., Powell, Glen, Mith, Alison G., John, P., 2013b. Carr: a trio of viral proteins tunes aphid-plant interactions in *Arabidopsis thaliana*. *PLOS One* 8 (12), e83066.
- Wezel, R., Dong, X., Liu, D., Tien, P., Stanley, J., Hong, Y., 2002. Mutation of three cysteine residues in tomato yellow leaf curl virus-China C2 protein causes dysfunction in pathogenesis and posttranscriptional gene-silencing suppression. *Mol. Plant – Microbe Interact.* 15, 203–208.
- Wu, J., Du, Z., Wang, C., Cai, L., Hu, M., Lin, Q., Wu, Z., Li, Y., Xie, L., 2010. Identification of Pns6, a putative movement protein of RRSV, as a silencing suppressor. *Virol. J.* 7, 335.

- Wu, J.G., Wang, C., Du, Z., Cai, L., Hu, M., Wu, Z., Li, Y., Xie, L., 2011. Identification of Pns12 as the second silencing suppressor of rice gall dwarf virus. *Sci. China. Life Sci.* 54, 201–208.
- Wu, W.Q., Fan, H.Y., Jiang, N., Wang, Y., Zhang, Z.Y., Zhang, Y.L., Wang, X.B., Li, D.W., Yu, J.L., Han, C.G., 2014. Infection of Beet necrotic yellow vein virus with RNA4-encoded P31 specifically up-regulates pathogenesis-related protein 10 in *Nicotiana benthamiana*. *Virology* 479, 111–118.
- Xia, Z., Zhu, Z., Zhou, J., Zhou, R., 2009. Recognition mechanism of siRNA by viral p19 suppressor of RNA silencing: a molecular dynamics study. *Biophys. J.* 96, 1761–1769.
- Xie, Z., Allen, E., Fahlgren, N., Calamar, A., Givan, S.A., Carrington, J.C., 2005. Expression of Arabidopsis MIRNA genes. *Plant Physiol.* 138, 2145–2154.
- Xie, Z., Johansen, L.K., Gustafson, A.M., Kasschau, K.D., Lellis, A.D., Zilberman, D., Jacobsen, S.E., Carrington, J.C., 2004. Genetic and functional diversification of small RNA pathways in plants. *PLoS Biol.* 2, E104.
- Xie, Z., Fan, B., Chen, C., Chen, Z., 2001. An important role of an inducible RNA-dependent RNA polymerase in plant antiviral defense. *Proc. Natl. Acad. Sci. USA* 98, 6516–6521.
- Xiong, R., Wu, J., Zhou, Y., Zhou, X., 2009. Characterization and subcellular localization of an RNA silencing suppressor encoded by rice stripe tenuivirus. *Virology* 387, 29–40.
- Yaegashi, H., Tamura, A., Isogai, M., Yoshikawa, N., 2008. Inhibition of long-distance movement of RNA silencing signals in *Nicotiana benthamiana* by apple chlorotic leaf spot virus 50 kDa movement protein. *Virology* 382, 199–206.
- Yaegashi, H., Takahashi, T., Isogai, M., Kobori, T., Ohki, S., Yoshikawa, N., 2007. Apple chlorotic leaf spot virus 50 kDa movement protein acts as a suppressor of systemic silencing without interfering with local silencing in *Nicotiana benthamiana*. *J. Gen. Virol.* 88, 316–324.
- Yan, F., Lu, Y., Lin, L., Zheng, H., Chen, J., 2012. The ability of PVX p25 to form RL structures in plant cells is necessary for its function in movement, but not for its suppression of RNA silencing. *PLoS One* 2012 (7), e43242.
- Yang, X., Baliji, S., Buchmann, R.C., Wang, H., Lindbo, J.A., Sunter, G., Bisaro, D.M., 2007. Functional modulation of the geminivirus AL2 transcription factor and silencing suppressor by self-interaction. *J. Virol.* 81, 11972–11981.
- Yang, X., Xie, Y., Raja, P., Li, S., Wolf, J.N., Shen, Q., Bisaro, D.M., Zhou, X., 2011. Suppression of methylation-mediated transcriptional gene silencing by betaC1-SAHH protein interaction during geminivirus-beta satellite infection. *PLoS Pathog.* 7, e1002329.
- Yang, Z., Ebright, Y.W., Yu, B., Chen, X., 2006. HEN1 recognizes 21–24 nt small RNA duplexes and deposits a methyl group onto the 2' OH of the 3' terminal nucleotide. *Nucleic Acids Res.* 34, 667–675.
- Ye, K., Malinin, L., Patel, D.J., 2003. Recognition of small interfering RNA by a viral suppressor of RNA silencing. *Nature* 426, 874–878.
- Yelina, N.E., Savenkov, E.I., Solovyev, A.G., Morozov, S.Y., Valkonen, J.P.T., 2002. Long-distance movement, virulence, and RNA silencing suppression controlled by a single protein in hordeid- and potyvirus: complementary functions between virus families. *J. Virol.* 76, 12981–12991.
- Ying, X.B., Dong, L., Zhu, H., Duan, C.G., Du, Q.S., Lv, D.Q., Fang, Y.Y., Garcia, J.A., Fang, R.X., Guo, H.S., 2010. RNA-dependent RNA polymerase 1 from *Nicotiana tabacum* suppresses RNA silencing and enhances viral infection in *Nicotiana benthamiana*. *Plant Cell* 22, 1358–1372.
- Yong Chung, H., Lacatus, G., Sunter, G., 2014. Geminivirus AL2 protein induces expression of, and interacts with, a calmodulin-like gene, an endogenous regulator of gene silencing. *Virology* 460–461, 108–118.
- Yoshikawa, M., Iki, T., Tsutsui, Y., Miyashita, K., Poethig, R.S., Habu, Y., Ishikawa, M., 2013. 3' fragment of miR173-programmed RISC-cleaved RNA is protected from degradation in a complex with RISC and SGS3. *Proc. Natl. Acad. Sci. USA* 110, 4117–4122.
- Yoshikawa, M., Peragine, A., Park, M.Y., Poethig, R.S., 2005. A pathway for the biogenesis of trans-acting siRNAs in Arabidopsis. *Genes Dev.* 19, 2164–2175.
- Young, B.A., Stenger, D.C., Qu, F., Morris, T.J., Tatini, S., French, R., 2012. Triticum virus P1 functions as a suppressor of RNA silencing and an enhancer of disease symptoms. *Virus Res.* 163, 672–677.
- Yu, B., Chapman, E.J., Yang, Z., Carrington, J.C., Chen, X., 2006. Transgenically expressed viral RNA silencing suppressors interfere with microRNA methylation in Arabidopsis. *FEBS Lett.* 580, 3117–3120.
- Zhai, Y., Bag, S., Mitter, N., Turina, M., Pappu, H.R., 2014. Mutational analysis of two highly conserved motifs in the silencing suppressor encoded by tomato spotted wilt virus (genus *Tospovirus*, family *Bunyaviridae*). *Arch. Virol.* 159, 1499–1504.
- Zhai, J., Jeong, D.H., De Paoli, E., Park, S., Rosen, B.D., Li, Y., Gonzalez, A.J., Yan, Z., Kitto, S.L., Grusak, M.A., Jackson, S.A., Stacey, G., Cook, D.R., Green, P.J., Sherrier, D.J., Meyers, B.C., 2011. MicroRNAs as master regulators of the plant NB-LRR defense gene family via the production of phased, trans-acting siRNAs. *Genes Dev.* 25, 2540–2553.
- Zhang, J., Dong, J., Xu, Y., Wu, J., 2012. V2 protein encoded by Tomato yellow leaf curl China virus is an RNA silencing suppressor. *Virus Res.* 163, 51–58.
- Zhang, X., Du, P., Lu, L., Xiao, Q., Wang, W., Cao, X., Ren, Bo., Wei, C., Li, Y., 2008. Contrasting effects of HC-Pro and 2b viral suppressors from Sugarcane mosaic virus and tomato aspermy cucumovirus on the accumulation of siRNAs. *Virology* 374, 351–360.
- Zhang, X., Yuan, Y.R., Pei, Y., Lin, S.S., Tuschl, T., Patel, D.J., Chua, N.H., 2006. Cucumber mosaic virus-encoded 2b suppressor inhibits Arabidopsis Argonaute1 cleavage activity to counter plant defense. *Genes Dev.* 20, 3255–3268.
- Zhang, X., Zhang, X., Singh, J., Li, D., Qu, F., 2012. Temperature-dependent survival of turnip crinkle virus-infected Arabidopsis plants relies on an RNA silencing-based defense that requires dcl2, AGO2, and HEN1. *J. Virol.* 86, 6847–6854.
- Zhang, Z., Chen, H., Huang, X., Xia, R., Zhao, Q., Lai, J., Teng, K., Li, Y., Liang, L., Du, Q., Zhou, X., Guo, H., Xie, Q., 2011. BSCTV C2 attenuates the degradation of SAMDC1 to suppress DNA methylation-mediated gene silencing in Arabidopsis. *Plant Cell* 23, 273–288.
- Zhang, X., Niu, D., Carbonell, A., Wang, A., Lee, A., Tun, V., Wang, Z., Carrington, J.C., Chang, C.E., Jin, H., 2014. ARGONAUTE PIWI domain and microRNA duplex structure regulate small RNA sorting in Arabidopsis. *Nat. Commun.* 5, 5468.
- Zhou, T., Murphy, A.M., Lewsey, M.G., Westwood, J.H., Zhang, H.M., Gonzalez, I., Canto, T., Carr, J.P., 2014. Domains of the cucumber mosaic virus 2b silencing suppressor protein affecting inhibition of salicylic acid-induced resistance and priming of salicylic acid accumulation during infection. *J. Gen. Virol.* 95, 1408–1413.
- Zhou, Z., Dell'Orco, M., Saldarelli, P., Turturo, C., Minafra, A., Martelli, G.P., 2006. Identification of an RNA-silencing suppressor in the genome of Grapevine virus A. *J. Gen. Virol.* 87, 2387–2395.
- Zhu, S., Lim, G.H., Yu, K., Jeong, R.D., Kachroo, A., Kachroo, P., 2014. RNA silencing components mediate resistance signaling against turnip crinkle virus. *Plant Signal. Behav.* 9, e28435.
- Zhu, S., Jeong, R.D., Lim, G.H., Yu, K., Wang, C., Chandra-Shekhara, A.C., Navarre, D., Klessig, D.F., Kachroo, A., Kachroo, P., 2013. Double-stranded RNA-binding protein 4 is required for resistance signaling against viral and bacterial pathogens. *Cell Rep.* 4, 1168–1184.
- Ziebell, H., Murphy, A.M., Groen, S.C., Tungadi, T., Westwood, J.H., Lewsey, M.G., Moulin, M., Kleczkowski, A., Smith, A.G., Stevens, M., Powell, G., Carr, J.P., 2011. Cucumber mosaic virus and its 2b RNA silencing suppressor modify plant-aphid interactions in tobacco. *Sci. Rep.* 1, 187.
- Zrachya, A., Glick, E., Levy, Y., Arazi, T., Citovsky, V., Gafni, Y., 2007. Suppressor of RNA silencing encoded by tomato yellow leaf curl virus-Israel. *Virology* 358, 159–165.