

Evidence for Altered Wnt Signaling in Psoriatic Skin

Johann E. Gudjonsson¹, Andrew Johnston¹, Stefan W. Stoll¹, Mary B. Riblett¹, Xianying Xing¹, James J. Kochkodan¹, Jun Ding², Rajan P. Nair¹, Abhishek Aphale¹, John J. Voorhees¹ and James T. Elder^{1,3}

The Wnt gene family encodes a set of highly conserved secreted signaling proteins that have major roles in embryogenesis and tissue homeostasis. Yet the expression of this family of important mediators in psoriasis, a disease characterized by marked changes in keratinocyte growth and differentiation, is incompletely understood. We subjected 58 paired biopsies from lesional and uninvolved psoriatic skin and 64 biopsies from normal skin to global gene expression profiling. *WNT5A* transcripts were upregulated fivefold in lesional skin, accompanied by increased Wnt-5a protein levels. Notably, *WNT5A* mRNA was markedly induced by IL-1 α , tumor necrosis factor- α , IFN- γ , and transforming growth factor- α in cultured keratinocytes. Frizzled 2 (*FZD2*) and *FZD5*, which encode receptors for Wnt5A, were also increased in lesional psoriatic skin. In contrast, expression of *WIF1* mRNA, encoding a secreted antagonist of the Wnt proteins, was downregulated >10-fold in lesional skin, along with decreased WNT inhibitory factor (WIF)-1 immunostaining. Interestingly, pathway analysis along with reduced *AXIN2* expression and lack of nuclear translocation of β -catenin indicated a suppression of canonical Wnt signaling in lesional skin. The results of our study suggest a shift away from canonical Wnt signaling toward noncanonical pathways driven by interactions between Wnt-5a and its cognate receptors in psoriasis, accompanied by impaired homeostatic inhibition of Wnt signaling by WIF-1 and dickkopf.

Journal of Investigative Dermatology (2010) **130**, 1849–1859; doi:10.1038/jid.2010.67; published online 8 April 2010

INTRODUCTION

The Wnt family of signaling proteins are highly conserved, lipid-modified, secreted molecules that participate in multiple developmental events during embryogenesis (Logan and Nusse, 2004). The Wnt proteins have also been shown to have fundamental roles in controlling cell proliferation, cell-fate determination, and differentiation during adult homeostasis (van Amerongen *et al.*, 2008). Classically, Wnt signaling has been divided into two major pathways. First, the canonical signaling pathway involves the use of Frizzled (Fz) receptors paired with low-density lipoprotein receptor-related proteins (LRPs) 5 and 6 as co-receptors (Wehrli *et al.*, 2000). This leads to the activation and nuclear translocation of β -catenin and is typically linked to cell-fate determination and stem cell maintenance. Alternatively, Wnt signaling occurs through the noncanonical pathway involving Fz receptors, independent of the β -catenin activation

cascade (Kuhl *et al.*, 2000). In mammals 19 Wnt proteins and 10 Fz transmembrane receptors are known (van Amerongen *et al.*, 2008). To date, several noncanonical pathways have been described, involving the receptor tyrosine kinase Ror2 (Oishi *et al.*, 2003), the atypical tyrosine kinase Ryk (Lu *et al.*, 2004), and Wnt-Ca²⁺ signaling pathways (Kohn and Moon, 2005), all signaling modes associated with controlling cell adhesion and movement. On the basis of this concept, Wnt proteins have been divided into two main categories depending on which pathway they activate (Sen and Ghosh, 2008). Wnt-1, Wnt-3A, and Wnt-8 have been classified as canonical Wnts whereas others such as Wnt-5a and Wnt-11 have been classified as noncanonical Wnts (van Amerongen *et al.*, 2008). However, it has recently become evident that this is probably an oversimplification, as typical noncanonical Wnt ligands such as Wnt-5a (Liu *et al.*, 2005) and Wnt-11 (Tao *et al.*, 2005) have been shown to be able to activate β -catenin signaling. Thus, it is likely that individual Wnt proteins may activate multiple pathways, depending on which Fz receptors are expressed on the cell surface (van Amerongen *et al.*, 2008).

Three families of secreted proteins are known to inhibit Wnt signaling activity. These are the secreted Frizzled-related protein family that bind Wnt proteins and prevent them from binding to the Fz receptors (Kawano and Kypta, 2003); the Dickkopf (Dkk) protein family that promote the internalization of LRP making it unavailable for Wnt binding (Logan and Nusse, 2004); and finally WNT inhibitory factor (WIF)-1, a secreted protein that binds to Wnt proteins and inhibits their activity (Hsieh *et al.*, 1999). As is evident from the description above, this combination of multiple ligands along with multiple receptors and soluble inhibitors creates an extremely complex system.

¹Department of Dermatology, University of Michigan Medical Center, Ann Arbor, Michigan, USA; ²Department of Biostatistics, Center for Statistical Genetics, School of Public Health, University of Michigan, Ann Arbor, Michigan, USA and ³Ann Arbor Veterans Affairs Health System, Ann Arbor, Michigan, USA

Correspondence: Johann E. Gudjonsson, Department of Dermatology, University of Michigan, Ann Arbor, Michigan 48109, USA.
E-mail: johanng@med.umich.edu

Abbreviations: DKK, dickkopf; Fzd, Frizzled; LRP, lipoprotein receptor-related protein; KRT, keratin; NHK, normal human keratinocyte; QRT-PCR, quantitative real-time PCR; SFRP, secreted frizzled-related protein; TGF, transforming growth factor; TNF, tumor necrosis factor; WIF, WNT inhibitory factor

Received 6 January 2009; revised 31 January 2010; accepted 4 February 2010; published online 8 April 2010

Psoriasis is a disease characterized by chronic inflammation and altered differentiation and hyperproliferation of keratinocytes. In normal skin the fraction of proliferating keratinocytes is probably around 20% (Wright and Camplejohn, 1983), whereas in psoriasis it is almost 100%, and the mean cell cycle time is reduced from 13 days to 36 hours (Weinstein *et al.*, 1985). Moreover, it has been suggested that this hyperproliferation is not restricted to the basal epidermal layer containing keratinocyte stem cells, but may also involve suprabasal cells (Leigh *et al.*, 1985). Despite the fundamental functions of Wnt proteins in controlling cell proliferation and differentiation, surprisingly little is known about the state of Wnt signaling in psoriasis. Of the Wnt proteins, only Wnt-5a has been described to be upregulated in lesional psoriatic skin as determined by gene expression (Reischl *et al.*, 2007) and was recently shown to synergize with type 1 IFNs (Romanowska *et al.*, 2009). However, the pathogenic role of this molecule in psoriasis is presently unknown. The aims of this study were to determine the cellular source of the increased expression of Wnt-5a in psoriasis and to characterize the expression of other mediators of canonical versus noncanonical Wnt signaling.

RESULTS

Differential Expression of Wnt Pathway Genes in Psoriasis Lesions

We have previously used a microarray data set to explore the differences between normal skin and uninvolved psoriatic (Gudjonsson *et al.*, 2009b) skin, to assess the expression of candidate risk genes in psoriasis (Nair *et al.*, 2009) and the activity of the sonic hedgehog pathway in psoriasis (Gudjonsson *et al.*, 2009a). Here we show that this data set contains a differential expression of Wnt pathway genes in psoriatic lesions. Global gene expression analysis revealed significant changes in several members of the Wnt ligand family and several of the Fz receptors in lesional psoriatic skin compared with both normal and uninvolved psoriatic skin (Figure 1). Of the Wnt ligands, *WNT5A* showed a 5.0-fold upregulation ($P < 0.0001$) and *WNT10A* had a 1.3-fold upregulation ($P < 0.0001$). In contrast, *WNT2*, *WNT2B*, *WNT5B*, and *WNT7B* were all downregulated (1.3-, 1.3-, 1.2-, and 1.3-fold, respectively, all P 's < 0.0001 ; Table 1). Of the Fz receptor genes, Frizzled 2 (*FZD2*) and *FZD5*, which encode receptors for Wnt-5a, had a 1.2- and 1.3-fold upregulation, respectively ($P < 0.0001$), whereas *FZD1*, *FZD4*, *FZD7*, *FZD8*, and *FZD10* were decreased (1.3-, 1.5-, 1.7-, 1.4-, and 1.5-fold, respectively, all P 's < 0.0001 ; Table 1). The Fz homologues act in concert with the low-density lipoprotein receptor-related proteins LRP5 or LRP6. *LRP6* demonstrated a 1.6-fold downregulation and data for *LRP5* was inconclusive due to a limited probe set.

Among the soluble inhibitors and modulators of Wnt signaling, *WIF1* was most strongly decreased in psoriatic skin, being expressed at 14-fold less than in uninvolved skin ($P < 0.0001$). The secreted frizzled-related protein transcripts *SFRP1*, *SFRP2*, *SFRP4* and *SFRP5* were all downregulated (1.3-, 1.3-, 1.3- and 1.1-fold, respectively, $P < 0.001$). In addition, the Dkk homologue genes *DKK1*, *DKK2*, and *DKK3*

were downregulated by 1.6-, 2.2-, and 1.3-fold, respectively ($P < 0.0001$), whereas *DKK4* showed a modest 1.2-fold upregulation ($P < 0.0001$).

Downstream members of the Wnt-canonical pathway, such as β -catenin 1 (*CTNNB1*) and β -catenin-interacting protein 1 (*CTNBP1*) were downregulated by 1.5- and 1.9-fold, respectively ($P < 0.0001$). Consistent with previously published studies (Belso *et al.*, 2008), cyclin D1 (*CCND1*), which is downstream of β -catenin (Prasad *et al.*, 2007), was downregulated (2.0-fold, $P < 0.0001$), whereas *CCND2* was upregulated (1.5-fold, $P < 0.0001$).

Confirmation of Microarray Data by QRT-PCR

To validate the microarray results, we performed quantitative real-time PCR (QRT)-PCR for several Wnt-related genes including *WIF1*, *WNT5A*, *DKK2*, and *CCND1*, along with keratin 16 (*KRT16*) as a positive control for upregulation in lesional psoriatic skin (Leigh *et al.*, 1995). This analysis confirmed the upregulation of *WNT5A* (2.9-fold, $P < 0.001$) and downregulation of *WIF1* (9.3-fold, $P < 0.05$), *DKK2* (8.4-fold, $P < 0.001$), and *CCND1* (4.3-fold, $P < 0.001$). As anticipated, *KRT16* showed more than 40-fold upregulation ($P < 0.001$; Figure 2).

Wnt Canonical Pathway is Suppressed in Lesional Skin

To determine the effect of gene expression changes in lesional psoriatic skin on the Wnt pathway, we used the Ingenuity Pathway Analysis software tool (www.ingenuity.com). Gene expression differences between normal control skin and lesional psoriatic skin were overlaid onto a global molecular network within the Ingenuity Pathway Knowledge Base. This revealed global downregulation of nearly all members of the canonical signaling pathway in psoriasis (Figure 3a). Decreased activity of the canonical Wnt pathway was confirmed by QRT-PCR for *AXIN2* ($P < 0.05$), a marker of canonical Wnt signaling (Jho *et al.*, 2002; Figure 3b). Consistent with these results, we found β -catenin to be decreased in lesional psoriatic skin (Figure 3c) and there was decreased nuclear localization of β -catenin in psoriatic skin compared to normal or uninvolved skin (insets in Figure 3c).

Wnt-5a is Upregulated in Lesional Skin Whereas WIF-1 is Downregulated

We examined the protein levels of Wnt-5a and WIF-1 proteins in normal, psoriatic, and symptom-free skin from psoriatic patients by semiquantitative immunohistochemical analysis. Lesional psoriatic skin showed increased Wnt-5a staining in both the epidermis and dermis compared with control and uninvolved skin (Figure 4a). These findings were confirmed using computer-assisted image quantification (Figure 4c). There were strong foci of Wnt-5a staining in the papillary dermis of lesional skin. Counterstaining with CD34 and CD31 did not show any colocalization (data not shown) indicating that the source of this staining is not from vascular structures. In addition, tissue lysates from normal, uninvolved, and lesional psoriatic skin showed increased levels of Wnt-5a protein in lesional psoriatic skin (Figure 4b). In agreement with our gene chip and QRT-PCR data, we




Figure 1. Microarray analysis reveals that *WNT5A* is strikingly upregulated, and *WIF1* downregulated, in the majority of lesional skin samples. This gene expression heat map image used transcripts from 58 paired lesional and uninvolved psoriasis and 64 normal skin samples and shows Wnt pathway members, regulators of Wnt signaling, and regulators of growth, proliferation, and transcription. In addition, genes involved in protein kinase activity, protein phosphatases, protein ubiquitination, and other genes related to Wnt signaling are shown. Color key: red, increased expression; blue, decreased expression; as indicated in the color key and histogram (inset).

observed decreased WIF-1 staining in the epidermis of lesional psoriatic skin, although, interestingly, there was a slight increase in the dermis (Figure 4a and c). There was slight nuclear staining of WIF-1 in epidermis of lesional skin (Figure 4a) but similar nuclear staining has been seen in bladder cancer (Urakami *et al.*, 2006) but not in renal cell carcinoma (Kawakami *et al.*, 2009). The significance of this nuclear staining in the psoriatic epidermis is at this time not clear.

WNT5A Expression by Keratinocytes is Induced by Several Proinflammatory Cytokines

Psoriatic skin is replete with proinflammatory cytokines and growth factors. To examine the effects of such a cytokine environment on Wnt expression by keratinocytes, we stimulated normal human keratinocytes (NHKs) with tumor necrosis factor (TNF)- α , IL-17A, IFN- γ , IL-22, transforming growth factor (TGF)- α and IL-1 α or a combination of TGF- α and IL-1 α for 24 hours. *WIF1* expression was undetectable in both proliferating and differentiated NHKs (data not shown), whereas *WNT5A* mRNA was readily detectable in both (Figure 5). We observed an approximately 1.5- to 2-fold increase in *WNT5A* expression treated with TGF- α , TNF- α , IFN- γ , or IL-1 α . The combination of both TGF- α and IL-1 α had an additive effect resulting in a 2- to 3-fold increase in

WNT5A expression (Figure 5). There was no change in *WNT5A* expression with IL-22 and IL-17A stimulation. Interestingly, the effects of these cytokines were observed only in the more differentiated keratinocytes, suggestive of a role for differentiation in the development of cytokine responsiveness. Baseline expression of *WNT5A* in both proliferating and differentiated NHKs was comparable to that of control and uninvolved skin. However, the maximum induction of *WNT5A* expression observed after cytokine or TGF- α stimulation was only about half the level of that observed in lesional psoriatic skin (data not shown), indicating that other additional mediators, or cell types, are involved in *WNT5A* mRNA induction in lesional psoriatic skin *in vivo*.

Effect of Exogenous Wnt-5a and WIF-1 on Keratinocytes

We were unable to observe any effect of recombinant Wnt-5a and Wnt-3a on NHK growth or migration (data not shown). As recombinant Wnt proteins lack posttranslational modifications that are essential for their full activity, we used secreted Wnt-5a from a transfected cell line to assess the effects of Wnt agonists and antagonists on keratinocyte growth and function. NHKs were cultured in the presence of diluted (20%, v/v) conditioned medium from Wnt-5a-transfected cell line and a control cell line in the presence of WIF-1 and anti-Wnt-5a

Table 1. Fold changes between lesional psoriatic (PP) and control (NN) skin for Wnt family members and Fz receptors present on the HU133 PLUS 2.0 microarray

Gene	Fold change PP vs. NN	P-value (FDR)	Probe ID
WNT1	1.03	NS	208570_at
WNT2	0.74	2.54E-23	205648_at
WNT2B	0.77	2.86E-20	206458_s_at
WNT3	1.08	NS	221455_s_at
WNT4	1.01	NS	208606_s_at
WNT5A	5.01	3.19E-176	205990_s_at
WNT5B	0.82	7.75E-08	221029_s_at
WNT6	1.05	NS	221609_s_at
WNT7A	1.11	9.94E-05	210248_at
WNT7B	0.774	2.80E-15	217681_at
WNT8A	1.00	NS	224259_at
WNT8B	1.04	NS	207612_at
WNT9A	0.94	NS	230643_at
WNT9B	1.03	NS	1552973_at
WNT10A	1.31	4.69E-17	223709_s_at
WNT11	0.92	NS	206737_at
WNT16	0.96	NS	221113_s_at
FZD1	0.74	1.50E-14	204451_at
FZD2	1.16	4.41E-06	210220_at
FZD3	0.95	NS	219683_at
FZD4	0.66	1.66E-19	218665_at
FZD5	1.27	4.13E-14	206136_at
FZD6	1.04	NS	203987_at
FZD7	0.58	5.19E-26	203705_s_at
FZD8	0.66	3.86E-29	227405_s_at
FZD9	1.07	NS	207639_at
FZD10	0.68	9.17E-17	219764_at

Abbreviations: FDR, false discovery rate; NS, not significant. P-value is corrected for multiple testing.

antibodies for 24, 48, 72, and 96 hours and counted at each time point. Exogenous Wnt-5a had an suppressive effect on keratinocyte growth ($P < 0.01$, Figure 6a), which was prevented by anti-Wnt-5a antibodies. There was a minimal to no effect of WIF-1 on the antiproliferative effect of Wnt-5a (Figure 6a and b). These data were confirmed by flow cytometry using Carboxyfluorescein diacetate succinimidyl ester-labeled NHK examined after 96 hours of culture (Figure 6b).

DISCUSSION

Psoriasis is a common chronic inflammatory skin disease characterized by marked changes in keratinocytes growth and differentiation. The basis of this alteration in epidermal

growth and differentiation is incompletely understood but has been shown to be dependent on the activity of the immune infiltrate within the psoriatic lesions (Valdimarsson *et al.*, 1995). Several cytokines and growth factors have been implicated in this process based on mouse models (Gudjonsson *et al.*, 2007) and *in vitro/ex vivo* studies. These include cytokines and growth factors such as interleukin (IL)-1 α (Lee *et al.*, 1997), vascular endothelial growth factor (Detmar *et al.*, 1994), the epidermal growth factors TGF- α (Elder *et al.*, 1989), amphiregulin (Cook *et al.*, 1992), HB-EGF (Stoll and Elder, 1998), and several cytokines secreted by the Th17 subset of T lymphocytes, most prominently IL-17 and IL-22 (Zaba *et al.*, 2007). As outlined earlier, the Wnt family of signaling proteins are a set of highly conserved molecules that participate and control processes such as cell proliferation, cell-fate determination, and differentiation during adult homeostasis (van Amerongen *et al.*, 2008). The changes that are observed in lesional psoriatic skin have been noted to have many similarities to wound healing (Mansbridge *et al.*, 1984; Hertle *et al.*, 1992), which is of interest as wounding has been shown to activate the Wnt-mediated signaling pathway (Ito *et al.*, 2007). Thus, based on the functions of these signaling proteins and the marked changes that occur within psoriatic lesions, it would not be unanticipated that these proteins may be important in the pathogenesis of psoriasis.

To date, very few studies have been performed to determine whether and how the Wnt pathway is activated in psoriasis. One of the reasons for this is the complexity of this signaling pathway. To date, 19 Wnt proteins and 10 Fz transmembrane receptors have been described (van Amerongen *et al.*, 2008) that require the LRP5 and LRP6 co-receptors for effective signaling (Wehrli *et al.*, 2000). Classically, Wnt signaling has been divided into the canonical pathway that leads to activation and nuclear translocation of membrane-bound β -catenin and the non-canonical pathway that is independent of β -catenin (Logan and Nusse, 2004) but can be mediated by several different signaling cascades (Logan and Nusse, 2004; Katoh, 2007). To complicate matters, it is found that several of the Wnt proteins, including Wnt-5a, can activate either pathway depending on receptor context on the surface of the responding cells (van Amerongen *et al.*, 2008). Wnt-5a is typically classified as a noncanonical Wnt as its transcriptional activation has been reported to be β -catenin independent (Sen and Ghosh, 2008). However, Wnt-5a has been shown to be able to either activate or inhibit β -catenin signaling depending on receptor context (Mikels and Nusse, 2006). The activity of the Wnt canonical pathway in psoriasis, as determined by β -catenin activation, has been controversial. A recently published study showed increased nuclear β -catenin staining high in the suprabasal layer in lesional psoriatic skin (Hampton *et al.*, 2007) whereas another study showed only membranous staining in lesional psoriatic skin (Yamazaki *et al.*, 2001), indicating lack of β -catenin activation in lesional psoriatic skin. Thus, based on these two reports it is not clear whether β -catenin activity is increased in lesional psoriatic skin.




Figure 2. Quantitative real-time PCR confirmed the altered expression of several components of the Wnt signaling pathways in psoriasis. Keratin 16 (*KRT16*) expression was used as a positive control for lesional psoriasis skin. Data are expressed as fold change relative to normal skin. Bars indicate mean \pm SD ($n = 10$). Statistical significance denoted: * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$.

To date only two studies have tried to directly address whether the Wnt pathway was involved in the pathogenesis of psoriasis (Reischl *et al.*, 2007; Romanowska *et al.*, 2009). One of these studies was based on microarray gene expression in 16 patients screening for 22,283 oligonucleotide probes (Reischl *et al.*, 2007). The authors determined that 10% of the differentially expressed genes in their study were directly or indirectly related to the canonical Wnt/ β -catenin or to the noncanonical Wnt/ Ca^{2+} pathways. Of these genes, *WNT5A* was the one most markedly changed, being fivefold upregulated compared with uninvolved psoriatic skin whereas *DKK2*, an inhibitor of Wnt signaling, was found to be downregulated (Reischl *et al.*, 2007). On the basis of decreased expression of cyclin D1 (*CCND1*), the authors argued that this indicated decreased activity of the canonical Wnt/ β -catenin pathway (Reischl *et al.*, 2007). Our study, which is based on a microarray interrogating a much larger probe set (> 55,000) and on a larger cohort (64 cases and 58 controls) has extended our knowledge on the status of the Wnt family in psoriasis. We have been able to confirm the findings of Reischl *et al.* (2007) on the upregulation of *WNT5A* and downregulation of *DKK2* and several of the Fz receptors. Furthermore, we have been able to extend these findings and show that several members of the β -catenin pathway, including β -catenin itself, are downregulated in lesional skin. Thus, there was both decreased expression of the β -catenin gene and protein in lesional skin (Figure 3). In addition there was decrease in nuclear translocation of β -catenin in psoriatic skin compared with normal and uninvolved skin (Figure 3c). These findings are consistent with the decreased *AXIN2* expression observed in lesional psoriatic skin (Figure 3b), but *AXIN2* is a reliable marker for canonical pathway activation (Jho *et al.*, 2002), and show that the activity of the canonical Wnt pathway is suppressed in lesional skin. As most, if not all, of the Wnt inhibitory genes are downregulated in lesional psoriatic skin, including *WIF1*, an interesting question is why the canonical Wnt pathway is still depressed. Of all the Wnt family members, *WNT5A* was

highly upregulated, *WNT10A* and *WNT7A* expression were moderately increased, and other Wnt members were either unchanged or showed decreased expression (Table 1). Importantly all the Fz receptors were downregulated except *FZD5*, and *FZD2*, both of which have been shown not only to be involved in non-canonical Wnt signaling (Slusarski *et al.*, 1997; Ahumada *et al.*, 2002) but also to interact with Wnt-5a (He *et al.*, 1997; Slusarski *et al.*, 1997). Thus, this pattern of gene expression in lesional psoriatic skin is consistent with a shift away from the canonical pathway toward the non-canonical signaling mediated by Wnt-5a and its cognate receptors.

The pathogenic role of Wnt-5a in psoriasis, if any, is still unclear. Our study data suggest that the main source of Wnt-5a in psoriatic lesions is the epidermis (Figure 4). Wnt-5a has been implicated both in inflammatory responses of human mononuclear cells (Blumenthal *et al.*, 2006) and vascular proliferation (Masckauchan *et al.*, 2006), processes that have been implicated in psoriasis pathogenesis (Lowe *et al.*, 2007). Vascular changes in psoriasis are characterized by capillary elongation, widening, and tortuosity predominantly in the dermal papillae (Hern and Mortimer, 2007). *WNT5A* has been described to be expressed by human endothelial cells and inhibit the canonical Wnt signaling (Masckauchan *et al.*, 2006). In addition, Wnt-5a has been shown to promote angiogenesis (Masckauchan *et al.*, 2006) and proliferation of endothelial cells in a dose-dependent manner (Cheng *et al.*, 2008). Wnt-5a can induce expression of a number of genes including Tie-2 (Masckauchan *et al.*, 2006), which is a receptor tyrosine kinase and functions as the receptor for angiopoietins 1 and 2 (Kuroda *et al.*, 2001). Tie-2 has been shown to be upregulated in psoriasis (Kuroda *et al.*, 2001) and transgenic mouse model overexpressing Tie-2 in the skin results in a psoriasis-like phenotype (Voskas *et al.*, 2005). It is well known that vascular changes reflected by redness of lesions take a longer time to resolve than the thickness and scaling. Given these data, it is enticing to speculate given the lack of normalization of Wnt-5a during anti-TNF treatment




Figure 3. Pathway analysis reveals global downregulation of nearly all members of the canonical Wnt signaling pathway in psoriasis. Global gene expression differences between normal and lesional psoriatic skin were overlaid onto a global molecular and pathway network within the Ingenuity Pathway Knowledge Base (a). This revealed near-global downregulation (green) of nearly all members of the canonical signaling pathway. However, several genes associated with the noncanonical Wnt pathway were upregulated (pink), including *WNT5A* and *FZD5*. Decreased activity of the canonical Wnt pathway was confirmed by quantitative real-time PCR for *AXIN2*, a marker of canonical Wnt signaling (b). Bars indicate mean \pm SD, * $P < 0.05$. Immunofluorescent microscopy of normal, uninvolved, and psoriatic skin (scale bar = 100 μ m) (c) (original magnification, $\times 20$, $n = 3$).




Figure 4. Immunohistochemistry of normal, uninvolved, and lesional psoriasis skin revealed increased Wnt-5a and decreased WIF-1 tissue expression. Immunohistochemical staining was performed on fresh-frozen sections of skin for Wnt-5a ($n=5$) and WIF-1 ($n=5$) (scale bar = 100 μm) (a). Western blot analyses of normal, uninvolved, and lesional psoriasis ($n=3$). Lysate from a Wnt-5a transgenic cell line was used as a positive control. The blots from normal skin were not run adjacent to lesional samples on the shown gel and have been juxtaposed for increased clarity (b). Differences in the expression of these proteins were confirmed with computer-assisted image quantification (c). Bars indicate mean \pm SD; original magnifications, $\times 100$. Isotype control antibodies were used and did not show any staining (not shown). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.




Figure 5. Expression of WNT5A by normal human keratinocytes can be induced by proinflammatory cytokines. WNT5A expression could be induced by 24-hour treatment with tumor necrosis factor (TNF)- α (10 ng ml $^{-1}$), IFN- γ (20 ng ml $^{-1}$), IL-1 α (10 ng ml $^{-1}$), or transforming growth factor (TGF- α) (24 ng ml $^{-1}$) after 24 hours. Expression could be further induced by the additive response to IL-1 α and TGF- α . Neither IL-17A (20 ng ml $^{-1}$) nor IL-22 (20 ng ml $^{-1}$) had any effect on WNT5A expression. Data are expressed as fold change relative to unstimulated normal human keratinocyte (NHK). Bars indicate mean \pm SD ($n=3$ in duplicate wells). * $P < 0.05$, ** $P < 0.01$.




Figure 6. Biological effects of Wnt-5a and WIF-1 on normal human keratinocytes. Conditioned medium containing Wnt-5a was obtained from a transfected cell line and diluted 1:4 in fresh M154 CF culture medium. Control medium was obtained from the untransfected parental cell line. Wnt5a had growth-suppressive effect on normal human keratinocyte (NHK) that could be neutralized with anti-Wnt-5a antibodies but no effect was seen with exogenous WIF-1. Bars indicate mean \pm SD ($n=3$ in duplicate wells). * $P < 0.05$, ** $P < 0.01$ (a). Carboxyfluorescein diacetate succinimidyl ester (CFSE) staining by flow cytometry correlated with cell counts (b).

(unpublished observation) that Wnt-5a may have a role in promoting and/or maintaining vascular changes in lesional skin.

WNT5A has also been shown to be expressed by human antigen-presenting cells through stimulation through Toll-like receptors and directly by TNF- α (Blumenthal *et al.*, 2006). In this context, it is of interest that we observed increased expression of Wnt-5a in lesional dermis although less than

that seen for the epidermis (Figure 4). However, we did not determine whether it was the inflammatory infiltrate or the vascular cells that were the main source of dermal Wnt-5a in psoriasis. TNF- α is a proinflammatory cytokine that has been shown to have a central role in psoriasis (Gottlieb *et al.*, 2005) and treatments directed against TNF- α are highly effective (Leonardi *et al.*, 2003; Chew *et al.*, 2004). As mentioned above and confirmed in our study, TNF- α has been shown to be able to induce the expression of Wnt-5a directly (Blumenthal *et al.*, 2006). In contrast, WNT5A expression in lesional skin does not decrease during anti-TNF- α treatment (unpublished observation), suggesting, that in psoriasis lesions, Wnt-5a is not acting downstream of TNF- α . Wnt-5a has also been shown to be induced through stimulation of Toll-like receptor signaling, in which case, neutralizing antibodies against TNF- α did not have any suppressive effect (Blumenthal *et al.*, 2006) indicating that WNT5A expression can also occur independently of TNF- α , which is consistent with the findings presented in our study. Interestingly, neutralization of Wnt-5a led to a decrease in the production of the cytokines IFN- γ and IL-12p40 (the common subunit of IL-12 and IL-23) (Blumenthal *et al.*, 2006). IFN- γ and IL-23 have been shown to be crucial in the pathogenesis of psoriasis through the maintenance and effector functions of Th1 and Th17 cells (Uyemura *et al.*, 1993; Cargill *et al.*, 2007; Krueger *et al.*, 2007; Zaba *et al.*, 2007). Recently, it was reported that Wnt-5a may synergize with type 1 IFNs such as IFN- α and IFN- β (Romanowska *et al.*, 2009), and type I IFNs have been implicated in the onset of psoriasis (Nestle *et al.*, 2005). These data support the notion that Wnt-5a has a role in amplifying and/or maintaining the inflammatory processes present in lesional skin.

Interestingly, IL-1 α , TNF- α and IFN- γ , and TGF- α , (a member of the epidermal growth factor ligand family) were able to induce expression of WNT5A by keratinocytes. Although we were not able to detect any activity on keratinocytes with recombinant Wnt-5a or Wnt-3a *in vitro* as determined by both growth assays and migration assays (data not shown), we observed growth-suppressive effect of secreted Wnt-5a on human keratinocytes *in vitro* (Figure 6). It should be noted that *in vivo* several of the Wnt proteins including Wnt-1, Wnt-3a, and Wnt-5a have significant posttranslational modifications (Kurayoshi *et al.*, 2007) that the commercially available recombinant proteins lack. The Wnts are heavily glycosylated, which is essential for folding and secretion (Kurayoshi *et al.*, 2007) and several, including Wnt-3a and Wnt-5a, are additionally conjugated to palmitate. This palmitate conjugation is essential for their biological activity and the lipid-unmodified form of Wnt-5a has been shown to be unable to bind to Fz5 and therefore unable to activate intracellular signal cascades and stimulate cell migration (Kurayoshi *et al.*, 2007). This might be one of the reasons for the observed lack of biological activity *in vitro* with either recombinant Wnt-3a or Wnt-5a. Taken together, our study data indicate that Wnt-5a may have a growth-suppressive effect on the psoriatic epidermis, a finding that has been observed in other settings (Olson *et al.*, 1998), and may therefore be a part of a regulatory mechanism keeping proliferation under control.

In conclusion, our study data indicate that there is a shift in lesional psoriatic skin away from canonical Wnt signaling toward noncanonical pathways likely mediated by the increased expression and production of Wnt-5a and its cognate receptors along with impaired homeostatic inhibition of Wnt signaling by WIF-1 and the Dkk proteins. Our study results and previously published studies indicate that Wnt-5a may have a role in inducing the marked vascular changes in lesional skin, influencing epidermal proliferation, and have a role in the amplification of inflammatory responses. Further studies are warranted to elucidate the exact role of the Wnt pathway in psoriasis pathogenesis.

MATERIALS AND METHODS

Study Subjects

In all, 58 psoriatic cases and 64 normal healthy controls were enrolled for the study. The criteria for entry as case were the manifestation of one or more well-demarcated, erythematous, scaly psoriatic plaques that were not limited to the scalp. In instances of only a single psoriatic plaque, a plaque size of at least 1% of total body surface area was required. Study subjects did not use any systemic antipsoriatic treatments for 2 weeks before biopsy. Informed consent was obtained from all subjects, under protocols approved by the institutional review board of the University of Michigan. This study was conducted in compliance with good clinical practice and according to the Declaration of Helsinki Principles. Local anesthesia was obtained with lidocaine HCl 1% with epinephrine 1:100,000 (Hospira, Lake Forest, IL). Two biopsies were taken from each patient; one 6 mm punch biopsy was obtained from lesional skin of patients and the other from uninvolved skin, taken at least 10 cm away from any active plaque. One to two biopsies were obtained from healthy controls. Immediately upon removal, biopsies were snap-frozen in liquid nitrogen and stored at -80°C .

Microarrays

The biopsy samples were prepared as previously prescribed (Gudjonsson *et al.*, 2009b). Data from the most up- or down-regulated probes were used for analysis.

Quantitation of mRNA Levels

Quantitative reverse transcription-PCR was performed on paired lesional and nonlesional samples from 10 psoriatic patients and 10 normal controls. The RNA used was from the same samples used for the gene microarrays. The reverse-transcription reaction was performed on 0.5 μg of RNA template and cDNA was synthesized using anchored-oligo(dT)18 primers as instructed by the manufacturer (Roche Diagnostics, Mannheim, Germany). QRT-PCR was carried out using a LightCycler 2.0 system (Roche Diagnostics). The reaction profile consisted of an initial denaturation at 95°C for 15 minutes followed by 40 cycles of PCR at 95°C for 10 seconds (denaturation), 58°C for 10 seconds (annealing) and 72°C for 10 seconds (extension). The fluorescence emitted was captured at the end of the extension step of each cycle at 530 nm. Primers for the genes WIF1, WNT5A, WNT3A, FZD1, FZD10, DKK2, KRT16, LRP6, RAB5A, RAB27A, GNA15, and CCND1 were obtained from Superarray Biosciences (Frederick, MD). Results were normalized to the expression of the housekeeping gene ribosomal protein, large,

P0 (*RPLP0/36B4*) (Laborda, 1991). QRT-PCR of cultured NHK was carried out using cDNA prepared as above and primer sets for *DEFB4*, *CXCL1*, *WNT5A*, and *WIF1* were obtained from Applied Biosystems and run on an Applied Biosystems 7900HT Fast Real-Time PCR System with results normalized to *RPLP0* expression.

Immunohistochemistry

Immunohistochemistry was performed on 5 μm fresh-frozen tissue sections from uninvolved, lesional psoriatic, and control skin using goat anti-Wnt5a (1 $\mu\text{g ml}^{-1}$; Santa Cruz Biotechnology, Santa Cruz, CA), mouse anti-WIF-1 (5 $\mu\text{g ml}^{-1}$; R&D Systems, Minneapolis, MN), and goat anti-Fzd5 (2 $\mu\text{g ml}^{-1}$; R&D Systems) antibodies overnight at 4 °C, followed by the appropriate biotinylated secondary antibodies (Vector Laboratories, Burlingame, CA), streptavidin-horseradish-peroxidase conjugate (Vector Laboratories) and visualized with 3-amino-9-ethyl-carbazole (BioGenex, San Ramon, CA), followed by hematoxylin counterstaining (Biocare Medical, Concord, CA). Stained sections were examined by light microscopy, and each stained tissue section was subjected to image capture in its entirety through five digital images taken with a $\times 20$ objective. All images were subsequently analyzed using Image-Pro software (Image-Pro Plus; Media Cybernetics, Silver Spring, MD), to quantify the area stained in epidermis versus dermis (calculated as density per area).

Paraffin-embedded slides from normal controls ($n=3$), uninvolved psoriatic ($n=3$), and lesional psoriatic ($n=3$) were used for β -catenin immunofluorescence staining. Antigen retrieval was performed for 20 minutes (Trizma base/EDTA) and the slides were then incubated with mouse antihuman β -catenin (Sigma, St Louis, MO) overnight at 4 °C. The slides were then washed three times and incubated with secondary antibody (AF594 chicken anti-mouse; Invitrogen, Carlsbad, CA). For Wnt5a CD31/CD34 counterstaining, paraffin-embedded slides were processed as described above and stained with goat anti-mouse Wnt5a (1:200 dilution; R&D Systems) overnight followed by incubation with biotinylated anti-goat antibodies for 30 minutes and avidin-488 (1:200 dilution; Vector Laboratories) for 10 minutes. The slides were then washed and counterstained with either mouse anti-human CD34 (ready for use; Dako North America, Carpinteria, CA) for 1 hour, or CD31 (dilution 1:20; Dako North America) for 1 hour, followed by AF594 chicken-anti-mouse antibodies (dilution 1:300; Invitrogen) for 30 minutes. The slides were washed three times and mounted with Vectashield fluorescent medium containing 46-diamidino-2-phenyl indole (Vector Laboratories).

Keratinocyte Cultures

Normal human keratinocytes were obtained from sun-protected adult skin by trypsin flotation and propagated in modified MCDB 153 medium (M154; Cascade Biologics, Portland, OR) as described previously (Stoll *et al.*, 2001), with the calcium concentration set at 0.1 mM. To determine the proliferative effects of exogenous Wnt-5a and WIF-1, we seeded NHK at a density of 2,000 cells per cm^2 on Costar 12-well polystyrene plates (Corning Life Sciences, Lowell, MA) and allowed to attach for 48 hours. Cells were treated with M154, Wnt-5a-conditioned medium (from L Wnt-5A cell line), or medium conditioned by the parental untransfected cell line (L-M (TK-1) line). The L-M(TK-1) (CCL-1.3) and L Wnt-5A cell lines (CRL-2814) were obtained from ATCC (Manassas, VA).

To condition medium for use in experiments with NHK, we grew the L Wnt-5a cell line as recommended, then passaged 1:10 and

grew in M154 without selection agents for 3 days. After 3 days culture medium was collected and stored at 4 °C whereas a second batch of conditioned medium was obtained from the same cells. On day 6, cells were discarded and the two batches of media were combined and stored at -20 °C until required,

Before use, conditioned media were diluted 1:4 in fresh M154 and subsequently added to the NHK cultures, in the presence or absence of recombinant human WIF-1 (300 ng ml^{-1} ; R&D Systems) or Wnt-5a-antibody (5 $\mu\text{g ml}^{-1}$; R&D Systems) for variable time periods (24–96 h). Cells were trypsinized and counted with a hemacytometer at the times indicated. All experiments were performed in triplicates. Evaluation of growth was also measured by flow cytometry after 96 hours using carboxyfluorescein diacetate succinimidyl ester-labeled NHK (Cell-Trace CFSE Detection kit; Invitrogen). NHK were in a similar manner exposed to recombinant Wnt-3a (200 ng ml^{-1}) and recombinant Wnt-5a (200 ng ml^{-1} ; R&D Systems) for variable time periods (24–96 hours) and were manually counted.

To examine the induction of Wnt proteins by proinflammatory cytokines, we grew NHK cultures to 40% confluency or maintained to 4 days after confluence. Cultures were then starved of growth factors in unsupplemented M154 for 24 hours before use. Cultures were stimulated with recombinant human TNF- α (10 ng ml^{-1}), IL-17A (20 ng ml^{-1}), IFN- γ (20 ng ml^{-1}), IL-22 (20 ng ml^{-1}), IL-1 α (10 ng ml^{-1}) (R&D Systems) or TGF- α (24 ng ml^{-1} ; R&D Systems) for 24 hours and processed for RNA isolation as described above.

Western Blots

One 6-mm punch biopsy was obtained from normal skin from healthy individuals ($n=3$) and paired lesional and nonlesional biopsies were obtained from psoriatic patients ($n=3$) as described above. Biopsies were snap-frozen in liquid nitrogen and stored at -80 °C. Protein lysates were obtained by pulverizing the biopsies while still frozen and the pulverized tissue was transferred to a 2 ml glass tissue grinder with 500 μl of RIPA lysis buffer. The samples were centrifuged at 4,500 r.p.m. for 4 min and supernatants collected. Protein concentrations were measured and all samples were diluted to 1 mg ml^{-1} . Samples were separated on 10% SDS-polyacrylamide gels and transferred to Immobilon-P membranes (Millipore, Billerica, MA). Western blots were blocked with Tris-buffered saline (TBS)/0.1% Tween 20 (TBS-T) containing 5% non-fat dry milk for 1 hour at room temperature and incubated with Wnt-5a antibody (dilution 1:1,000; Cell Signaling Technology, Danvers, MA) overnight at 4 °C. The blots were washed three times with TBS-T and then incubated with horseradish-peroxidase-conjugated rabbit secondary antibody (dilution 1:2,500; GE) for 1.5 hours at room temperature. Blots were washed again and detected by chemiluminescence using ECL (GE Healthcare, Piscataway, NJ) and Kodak X-Omat film (Kodak, Rochester, NY). Anti- β -actin (dilution 1:2,500; Sigma) was used as a loading control.

Statistical Analyses

Student's *t*-test was used to analyze differences. A paired *t*-test was used when uninvolved and lesional psoriatic data sets were compared, unpaired *t*-test was used for other comparisons. Quantitative immunohistochemistry data were tested for significance using Student's *t*-test assuming equal variances and *P*-values = 0.05 were considered statistically significant.

Ingenuity Pathway Analyses

Microarray data were analyzed using Ingenuity Pathway Analysis software (Ingenuity Systems, www.ingenuity.com). For network generation, a data set containing gene identifiers and corresponding expression values for WNT pathway genes was uploaded into the application. Each gene identifier was mapped to its corresponding gene object in the Ingenuity Pathways Knowledge Base. These genes, called focus genes, were overlaid onto a global molecular network developed from information contained in the Ingenuity Pathways Knowledge Base.

CONFLICT OF INTEREST

The authors state no conflict of interest.

ACKNOWLEDGMENTS

We thank all the research subjects for their participation in this study and Lynda Hodges and Kathleen McCarthy for their assistance. We also thank Dr Wang for suggestions and comments. This work was supported by grants to JTE (National Institute of Arthritis, Musculoskeletal and Skin Diseases, R01) and JEG (American Skin Association, Dermatology Foundation).

REFERENCES

- Ahumada A, Slusarski DC, Liu X *et al.* (2002) Signaling of rat Frizzled-2 through phosphodiesterase and cyclic GMP. *Science* 298:2006–10
- Belso N, Szell M, Pivarcsi A *et al.* (2008) Differential expression of D-type cyclins in HaCaT keratinocytes and in psoriasis. *J Invest Dermatol* 128:634–42
- Blumenthal A, Ehlers S, Lauber J *et al.* (2006) The Wntless homolog WNT5A and its receptor Frizzled-5 regulate inflammatory responses of human mononuclear cells induced by microbial stimulation. *Blood* 108:965–73
- Cargill M, Schrodi SJ, Chang M *et al.* (2007) A large-scale genetic association study confirms IL12B and leads to the identification of IL23R as psoriasis-risk genes. *Am J Hum Genet* 80:273–90
- Cheng CW, Yeh JC, Fan TP *et al.* (2008) Wnt5a-mediated non-canonical Wnt signalling regulates human endothelial cell proliferation and migration. *Biochem Biophys Res Commun* 365:285–90
- Chew AL, Bennett A, Smith CH *et al.* (2004) Successful treatment of severe psoriasis and psoriatic arthritis with adalimumab. *Br J Dermatol* 151:492–6
- Cook PW, Pittelkow MR, Keeble WW *et al.* (1992) Amphiregulin messenger RNA is elevated in psoriatic epidermis and gastrointestinal carcinomas. *Cancer Res* 52:3224–7
- Detmar M, Brown LF, Claffey KP *et al.* (1994) Overexpression of vascular permeability factor/vascular endothelial growth factor and its receptors in psoriasis. *J Exp Med* 180:1141–6
- Elder JT, Fisher GJ, Lindquist PB *et al.* (1989) Overexpression of transforming growth factor alpha in psoriatic epidermis. *Science* 243:811–4
- Gottlieb AB, Chamian F, Masud S *et al.* (2005) TNF inhibition rapidly down-regulates multiple proinflammatory pathways in psoriasis plaques. *J Immunol* 175:2721–9
- Gudjonsson JE, Aphale A, Grachtchouk M *et al.* (2009a) Lack of evidence for activation of the hedgehog pathway in psoriasis. *J Invest Dermatol* 129:635–40
- Gudjonsson JE, Ding J, Li X *et al.* (2009b) Global gene expression analysis reveals evidence for decreased lipid biosynthesis and increased innate immunity in uninvolved psoriatic skin. *J Invest Dermatol* 129:2795–804
- Gudjonsson JE, Johnston A, Dyson M *et al.* (2007) Mouse models of psoriasis. *J Invest Dermatol* 127:1292–308
- Hampton PJ, Ross OK, Reynolds NJ (2007) Increased nuclear beta-catenin in suprabasal involved psoriatic epidermis. *Br J Dermatol* 157:1168–77
- He X, Saint-Jeannet JP, Wang Y *et al.* (1997) A member of the Frizzled protein family mediating axis induction by Wnt-5A. *Science* 275:1652–4
- Hern S, Mortimer PS (2007) *In vivo* quantification of microvessels in clinically uninvolved psoriatic skin and in normal skin. *Br J Dermatol* 156:1224–9
- Hertle MD, Kubler MD, Leigh IM *et al.* (1992) Aberrant integrin expression during epidermal wound healing and in psoriatic epidermis. *J Clin Invest* 89:1892–901
- Hsieh JC, Kodjabachian L, Rebbert ML *et al.* (1999) A new secreted protein that binds to Wnt proteins and inhibits their activities. *Nature* 398:431–6
- Ito M, Yang Z, Andl T *et al.* (2007) Wnt-dependent *de novo* hair follicle regeneration in adult mouse skin after wounding. *Nature* 447:316–20
- Jho EH, Zhang T, Domon C *et al.* (2002) Wnt/beta-catenin/Tcf signaling induces the transcription of Axin2, a negative regulator of the signaling pathway. *Mol Cell Biol* 22:1172–83
- Katoh M (2007) WNT signaling pathway and stem cell signaling network. *Clin Cancer Res* 13:4042–5
- Kawakami K, Hirata H, Yamamura S *et al.* (2009) Functional significance of Wnt inhibitory factor-1 gene in kidney cancer. *Cancer Res* 69:8603–10
- Kawano Y, Kypta R (2003) Secreted antagonists of the Wnt signalling pathway. *J Cell Sci* 116:2627–34
- Kohn AD, Moon RT (2005) Wnt and calcium signaling: beta-catenin-independent pathways. *Cell Calcium* 38:439–46
- Krueger GG, Langley RG, Leonardi C *et al.* (2007) A human interleukin-12/23 monoclonal antibody for the treatment of psoriasis. *N Engl J Med* 356:580–92
- Kuhl M, Sheldahl LC, Park M *et al.* (2000) The Wnt/Ca²⁺ pathway: a new vertebrate Wnt signaling pathway takes shape. *Trends Genet* 16:279–83
- Kurayoshi M, Yamamoto H, Izumi S *et al.* (2007) Post-translational palmitoylation and glycosylation of Wnt-5a are necessary for its signalling. *Biochem J* 402:515–23
- Kuroda K, Sapadin A, Shoji T *et al.* (2001) Altered expression of angiopoietins and Tie2 endothelium receptor in psoriasis. *J Invest Dermatol* 116:713–20
- Laborda J (1991) 36B4 cDNA used as an estradiol-independent mRNA control is the cDNA for human acidic ribosomal phosphoprotein PO. *Nucleic Acids Res* 19:3998
- Lee RT, Briggs WH, Cheng GC *et al.* (1997) Mechanical deformation promotes secretion of IL-1 alpha and IL-1 receptor antagonist. *J Immunol* 159:5084–8
- Leigh IM, Navsaria H, Purkis PE *et al.* (1995) Keratins (K16 and K17) as markers of keratinocyte hyperproliferation in psoriasis *in vivo* and *in vitro*. *Br J Dermatol* 133:501–11
- Leigh IM, Pulford KA, Ramaekers FC *et al.* (1985) Psoriasis: maintenance of an intact monolayer basal cell differentiation compartment in spite of hyperproliferation. *Br J Dermatol* 113:53–64
- Leonardi CL, Powers JL, Matheson RT *et al.* (2003) Etanercept as monotherapy in patients with psoriasis. *N Engl J Med* 349:2014–22
- Liu G, Bañico A, Aaronson SA (2005) The mechanism of endogenous receptor activation functionally distinguishes prototype canonical and noncanonical Wnts. *Mol Cell Biol* 25:3475–82
- Logan CY, Nusse R (2004) The Wnt signaling pathway in development and disease. *Annu Rev Cell Dev Biol* 20:781–810
- Lowes MA, Bowcock AM, Krueger JG (2007) Pathogenesis and therapy of psoriasis. *Nature* 445:866–73
- Lu W, Yamamoto V, Ortega B *et al.* (2004) Mammalian Ryk is a Wnt coreceptor required for stimulation of neurite outgrowth. *Cell* 119:97–108
- Mansbridge JN, Knapp AM, Streffling AM (1984) Evidence for an alternative pathway of keratinocyte maturation in psoriasis from an antigen found in psoriatic but not normal epidermis. *J Invest Dermatol* 83:296–301

- Masckauchan TN, Agalliu D, Vorontchikhina M *et al.* (2006) Wnt5a signaling induces proliferation and survival of endothelial cells *in vitro* and expression of MMP-1 and Tie-2. *Mol Biol Cell* 17:5163–72
- Mikels AJ, Nusse R (2006) Purified Wnt5a protein activates or inhibits beta-catenin-TCF signaling depending on receptor context. *PLoS Biol* 4:e115
- Nair RP, Duffin KC, Helms C *et al.* (2009) Genome-wide scan reveals association of psoriasis with IL-23 and NF-kappaB pathways. *Nat Genet* 41:199–204
- Nestle FO, Conrad C, Tun-Kyi A *et al.* (2005) Plasmacytoid predendritic cells initiate psoriasis through interferon-alpha production. *J Exp Med* 202:135–43
- Oishi I, Suzuki H, Onishi N *et al.* (2003) The receptor tyrosine kinase Ror2 is involved in non-canonical Wnt5a/JNK signalling pathway. *Genes Cells* 8:645–54
- Olson DJ, Oshimura M, Otte AP *et al.* (1998) Ectopic expression of wnt-5a in human renal cell carcinoma cells suppresses *in vitro* growth and telomerase activity. *Tumour Biol* 19:244–52
- Prasad CP, Gupta SD, Rath G *et al.* (2007) Wnt signaling pathway in invasive ductal carcinoma of the breast: relationship between beta-catenin, dishevelled and cyclin D1 expression. *Oncology* 73:112–7
- Reischl J, Schwenke S, Beekman JM *et al.* (2007) Increased expression of Wnt5a in psoriatic plaques. *J Invest Dermatol* 127:163–9
- Romanowska M, Evans A, Kellock D *et al.* (2009) Wnt5a exhibits layer-specific expression in adult skin, is upregulated in psoriasis, and synergizes with type 1 interferon. *PLoS ONE* 4:e5354
- Sen M, Ghosh G (2008) Transcriptional outcome of Wnt-Frizzled signal transduction in inflammation: evolving concepts. *J Immunol* 181:4441–5
- Slusarski DC, Corces VG, Moon RT (1997) Interaction of Wnt and a Frizzled homologue triggers G-protein-linked phosphatidylinositol signalling. *Nature* 390:410–3
- Stoll SW, Elder JT (1998) Retinoid regulation of heparin-binding EGF-like growth factor gene expression in human keratinocytes and skin. *Exp Dermatol* 7:391–7
- Stoll SW, Kansra S, Peshick S *et al.* (2001) Differential utilization and localization of ErbB receptor tyrosine kinases in skin compared to normal and malignant keratinocytes. *Neoplasia* 3:339–50
- Tao Q, Yokota C, Puck H *et al.* (2005) Maternal wnt11 activates the canonical wnt signaling pathway required for axis formation in *Xenopus* embryos. *Cell* 120:857–71
- Urakami S, Shiina H, Enokida H *et al.* (2006) Epigenetic inactivation of Wnt inhibitory factor-1 plays an important role in bladder cancer through aberrant canonical Wnt/beta-catenin signaling pathway. *Clin Cancer Res* 12:383–91
- Uyemura K, Yamamura M, Fivenson DF *et al.* (1993) The cytokine network in lesional and lesion-free psoriatic skin is characterized by a T-helper type 1 cell-mediated response. *J Invest Dermatol* 101:701–5
- Valdimarsson H, Baker BS, Jonsdottir I *et al.* (1995) Psoriasis: a T-cell-mediated autoimmune disease induced by streptococcal superantigens? *Immunol Today* 16:145–9
- van Amerongen R, Mikels A, Nusse R (2008) Alternative wnt signaling is initiated by distinct receptors. *Sci Signal* 1:re9
- Voskas D, Jones N, Van Slyke P *et al.* (2005) A cyclosporine-sensitive psoriasis-like disease produced in Tie2 transgenic mice. *Am J Pathol* 166:843–55
- Wehrli M, Dougan ST, Caldwell K *et al.* (2000) arrow encodes an LDL-receptor-related protein essential for Wingless signalling. *Nature* 407:527–30
- Weinstein GD, McCullough JL, Ross PA (1985) Cell kinetic basis for pathophysiology of psoriasis. *J Invest Dermatol* 85:579–83
- Wright NA, Camplejohn RS (eds) (1983) *Psoriasis: Cell Proliferation*. Churchill Livingstone: Edinburgh, London, Melbourne and New York
- Yamazaki F, Aragane Y, Kawada A *et al.* (2001) Immunohistochemical detection for nuclear beta-catenin in sporadic basal cell carcinoma. *Br J Dermatol* 145:771–7
- Zaba LC, Cardinale I, Gilleaudeau P *et al.* (2007) Amelioration of epidermal hyperplasia by TNF inhibition is associated with reduced Th17 responses. *J Exp Med* 204:3183–94