



Evaluation of the combined effect of thymol, carvacrol and (*E*)-cinnamaldehyde on *Amblyomma sculptum* (Acari: Ixodidae) and *Dermacentor nitens* (Acari: Ixodidae) larvae



Tatiane Pinheiro Lopes Novato^{a,*}, Laryssa Xavier Araújo^a,
Caio Márcio Oliveira de Monteiro^a, Ralph Maturano^b, Tatiane de Oliveira Souza Senra^b,
Renata da Silva Matos^d, Geovany Amorim Gomes^c, Mario Geraldo de Carvalho^c,
Erik Daemon^a

^a Programa de Pós-graduação em Comportamento e Biologia Animal da Universidade Federal de Juiz de Fora, Rua José Lourenço Kelmer, s/n-Campus Universitário Bairro Martelos, 36036-330 Juiz de Fora, MG, Brazil

^b Programa de Pós-graduação em Ciências Veterinárias da Universidade Federal Rural do Rio de Janeiro, BR-465, Km 7, Seropédica, Rio de Janeiro, Brazil

^c Departamento de Química, Instituto de Ciências Exatas, Universidade Federal Rural do Rio de Janeiro, BR-465, Km 7, Seropédica, Rio de Janeiro, Brazil

^d Programa de Pós-graduação em Biologia Celular e Molecular da Universidade Estadual Paulista Júlio de Mesquita Filho. R. 10, 13505-470 Rio Claro, São Paulo, Brazil

ARTICLE INFO

Article history:

Received 6 January 2015

Received in revised form 12 August 2015

Accepted 19 August 2015

Keywords:

Tick

Monoterpene

Phenylpropanoid

Synergism

ABSTRACT

This study aimed at assessing the combined effect of thymol, carvacrol and (*E*)-cinnamaldehyde on *Amblyomma sculptum* and *Dermacentor nitens* larvae. The effects resulting from treatments were evaluated by means of the modified larval packet test. In order to determine the LC₅₀, components of essential oils, the monoterpenes thymol, carvacrol and phenylpropanoid (*E*)-cinnamaldehyde were individually tested at different concentrations. After determining the LC₅₀, each essential oil component was separately evaluated and then combined with another substance at a 1:1 proportion at the LC₅₀ concentration and at 1/2 and 1/4 of the LC₅₀. For *A. sculptum*, the lowest LC₅₀ value was obtained for (*E*)-cinnamaldehyde (1.40 mg/ml), followed by thymol (2.04 mg/ml) and carvacrol (3.49 mg/ml). The same order of effectiveness was observed for *D. nitens*, with values of 1.68, 2.17 and 3.33 mg/ml, respectively. In the evaluation of component associations of essential oils against *A. sculptum* larvae, only the combinations between carvacrol and thymol (LC₅₀) and carvacrol and (*E*)-cinnamaldehyde (1/4 LC₅₀) presented a moderate synergistic effect. In turn, for *D. nitens* larvae, the combinations between thymol and carvacrol (LC₅₀ and 1/2 LC₅₀) presented a synergistic effect, while the others presented an additive or antagonistic effect. Therefore, it can be concluded that the combination of thymol and carvacrol (LC₅₀) has a moderate synergistic effect against *A. sculptum* larvae, while thymol, combined with carvacrol (LC₅₀ and 1/2 LC₅₀), has a synergistic effect against *D. nitens* larvae.

© 2015 Elsevier B.V. All rights reserved.

1. Introduction

There are about 65 tick species in Brazil (Dantas-Torres et al., 2009). Among them, *Amblyomma sculptum* (Berlese, 1888), and *Dermacentor nitens* (Neumann, 1897), stand out as very important species, both for animals and public health (Nava et al., 2014; Guglielmone et al., 2006).

A. sculptum is distributed in peri-Amazonian regions in Brazil, specifically in the states of Pernambuco, Piauí, Goiás, Mato Grosso,

Mato Grosso do Sul, Espírito Santo, Minas Gerais, Rio de Janeiro, São Paulo and Paraná, as well as in Bolivia, Paraguay and northern Argentina (Nava et al., 2014). Equines and capybaras are the preferred hosts of this tick, but its low host specificity, mainly in immature stages, means it can be also found on other animals, such as deer, cattle and canids, as well as on humans (Oliveira, 2004; Guglielmone et al., 2006). Besides pain, its bite may cause inflammation, fever and stress, both in humans and animals (Oliveira, 2004). This tick is also a vector of *Rickettsia rickettsii* bacterium, the etiological agent of Brazilian Spotted Fever (Guedes et al., 2005).

D. nitens can be found from southern Florida and Texas to northern Argentina (Borges and Leite, 1993). It is known in Brazil as “carrapato-da-orelha-do-cavalo” (“horse ear tick”) because it

* Corresponding author.

E-mail address: tatinovatojf@gmail.com (T.P.L. Novato).

mainly parasitizes the ears of equines, although, in large infestations, it can also be found on other body regions (Guimarães et al., 2001; Guglielmone et al., 2006). This tick is of great veterinary importance, causing blood loss, stress, transmission of infectious agents, predisposition to myiasis, and secondary bacterial infections in its hosts (Guglielmone et al., 2006; Labruna and Machado, 2006). It is also responsible for the transmission of *Babesia caballi* (Nuttall and Strickland, 1910), the etiological agent of equine babesiosis (Labruna and Machado, 2006).

The most common method to control ticks is the application of acaricide (Borges et al., 2011). However, the indiscriminate use of these products can lead to toxicity, both in animals and humans, as well as soil and water contamination (Monteiro and Prata, 2013). For these reasons, it is of utmost importance to develop new alternatives to control these ticks, which are safer to humans, animals and the environment (Chagas, 2004; Borges et al., 2011).

Essential oils and their components have been indicated as a promising alternative to control these ectoparasites (Regnault-Roger and Philogène, 2008). Thymol and carvacrol are monoterpenes mainly found in the essential oils of Lamiaceae (Martinov, 1820) and Verbanaceae (Saint-Hilaire, 1805) plant families, which act against various tick species (Novelino et al., 2007; Coskun et al., 2008; Cetin et al., 2009, 2010; Dolan et al., 2009; Daemon et al., 2009, 2012; Monteiro et al., 2009, 2010; Mendes et al., 2011; Senra et al., 2013a). (*E*)-cinnamaldehyde is a phenylpropanoid present in the essential oil of plants of the *Cinnamomum* (Schafer, 1760) genus, which has proven activity against *Rhipicephalus microplus* (Canestrini, 1888), *D. nitens* (Senra et al., 2013a), *A. cajennense* s.l. and *Rhipicephalus sanguineus* (Latreille, 1806) (Senra et al., 2013b).

Studies have shown that the association of components of essential oils like terpenes and phenylpropanoid can have a synergetic, antagonistic or additive effect on different target organisms, such as bacteria, nematodes and insects (Didry et al., 1994; Ntalli et al., 2010; Gallardo et al., 2012). However, such research has not been carried out for ticks. Hence, the aim of this study was to assess the combined action of thymol, carvacrol and (*E*)-cinnamaldehyde on unengorged larvae of *A. sculptum* and *D. nitens*.

2. Material and methods

2.1. Experiment location and tick origin

The study was carried out in the “Laboratório de Artropódos Parasitos” (LAP) of Juiz de Fora Federal University, Minas Gerais, Brazil. *A. sculptum* and *D. nitens* larvae were obtained from the eggs laid by engorged females manually collected on naturally infested horses which had had no recent contact with acaricides, in the municipality of Descoberto, in the state of Minas Gerais. Females of *A. sculptum* were examined for species confirmation according to the new taxonomic proposal of Nava et al. (2014). The larvae of both species submitted to the tests were aged between 15 and 25 days after hatching.

2.2. Origin and dilution of the components of essential oils

The carvacrol and (*E*)-cinnamaldehyde used in this study were purchased from Sigma–Aldrich. The thymol, in the form of crystals, was obtained from Henrifarma Químicos e Farmacêuticos Ltda. All three components of the essential oils tested were 99% pure and were dissolved in 70% ethanol, a solvent with low toxicity to the unengorged larvae of these ticks (Resende et al., 2012).

2.3. Larval packet test

All the tests were carried out using the method proposed by Stone and Haydock (1962) as adapted by Monteiro et al. (2012),

in which approximately 100 larvae were placed in the center of a sheet of filter paper (6 × 6 cm). These sheets were then folded in the middle and their edges were sealed with clips to form packets, which were evenly moistened with 90 µl of the solutions to be tested. The packets (each one an experimental unit) were placed in a climate-controlled chamber (27 °C and relative humidity >80%) for 24 h, after which the mortality rate was measured (percentage of dead larvae).

In order to determine the 50% lethal concentration (LC₅₀) to *A. sculptum* larvae, thymol was tested at concentrations of 0.62, 1.25, 2.5, 5.0, 7.5 and 10.0 mg/ml, while carvacrol and (*E*)-cinnamaldehyde were tested at concentrations of 0.62, 1.25, 2.5, 3.75, 5.0 and 7.5 mg/ml, with 10 repetitions for each group. To determine the LC₅₀ to *D. nitens* larvae, thymol, carvacrol and (*E*)-cinnamaldehyde were tested at concentrations of 0.31, 0.62, 1.25, 2.5, 5.0 and 7.5 mg/ml. These concentrations were defined based on previously published results of the acaricidal activity of these components (Daemon et al., 2012; Senra et al., 2013a,b). Control groups were treated with 70% ethanol.

After determining the LC₅₀, thymol, carvacrol and (*E*)-cinnamaldehyde were evaluated separately and in combination with another in 1:1 proportion, at concentrations of LC₅₀ and 1/2 and 1/4 of LC₅₀, with 10 repetitions for each treatment. Control groups were exclusively treated with the solvent (70% ethanol). The groups were kept in different climate-controlled chambers at 27 ± 1 °C and RH of 80 ± 10%.

2.4. Analysis of the data

The lethal concentrations for 50% mortality of larvae (LC₅₀) were calculated using POLOPC[®] software. The combined action of thymol, carvacrol and (*E*)-cinnamaldehyde was measured by calculating the combination index (CI), using the CompuSyn[®] version 1.0 program (Chou, 2006). This method was chosen due to its sensitivity for the qualitative analysis categorization of the interactions between the substances. In present study, the original categories proposed by Chou (2006) were reduced to facilitate the interpretation of the results obtained (Table 1).

3. Results

The LC₅₀ values of the tested substances are presented in Table 2. No mortality was observed for *A. sculptum* or *D. nitens* in any of the control groups. In the tests with a single substance against the *A. sculptum* and *D. nitens* larvae, the substance with the lowest LC₅₀ value was (*E*)-cinnamaldehyde, with values of 1.40 and 1.68 mg/ml, respectively, followed by thymol (2.04 and 2.17 mg/ml) and carvacrol (3.49 and 3.43 mg/ml).

Among the nine combinations tested against *A. sculptum* larvae, only the association of carvacrol and thymol (LC₅₀) and (*E*)-cinnamaldehyde (1/4 of LC₅₀) showed moderate synergy. The other associations with thymol and carvacrol exhibited an additive effect, while (*E*)-cinnamaldehyde with thymol and carvacrol presented an antagonistic effect (Table 3).

For the combinations tested against *D. nitens*, thymol and carvacrol, at concentrations of 1/2 of LC₅₀ and LC₅₀ presented synergetic effect, while at 1/4 of LC₅₀ the effect was additive. An antagonistic effect was observed for all of the combinations and concentrations of carvacrol and (*E*)-cinnamaldehyde, as well as of thymol and (*E*)-cinnamaldehyde (Table 4).

4. Discussion

Investigations on the potential of essential oils and their components to control ticks have increased in recent years (Ellse and Wall,

Table 1

Adaptation of the qualitative categories defined by Chou (2006) for description of the combined effect of different substances and their respective combination index values.

Original description (Chou, 2006)	Range of combination index of original description	Adapted description (Combination index)
Very strong synergism	<0.1	Synergism (<0.7)
Strong synergism	0.1–0.3	
Synergism	0.3–0.7	
Moderate synergism	0.7–0.85	Moderate synergism (0.7–0.9)
Slight synergism	0.85–0.90	
Additive	0.90–1.10	Additive (0.9–1.10)
Slight antagonism	1.10–1.20	Moderate antagonism (1.10–1.45)
Moderate antagonism	1.20–1.45	
Antagonism	1.45–3.3	Antagonism (>1.45)
Strong antagonism	3.3–10	
Very strong antagonism	>10	

Table 2Lethal concentration (LC₅₀) of component of essential oils, thymol, carvacrol and (*E*)-cinnamaldehyde on unengorged larvae of *Amblyomma sculptum* and *Dermacentor nitens*.

Tick species	Components of essential oils	LC ₅₀ (mg/ml)	Confidence interval (95%)
<i>Amblyomma sculptum</i>	Carvacrol	3.49	3.18–3.82
	Thymol	2.04	1.76–2.37
	(<i>E</i>)-cinnamaldehyde	1.40	1.24–1.57
<i>Dermacentor nitens</i>	Carvacrol	3.33	2.87–3.86
	Thymol	2.17	1.92–2.45
	(<i>E</i>)-cinnamaldehyde	1.68	1.47–1.92

Table 3Evaluation of paired combinations of components of essential oils, carvacrol, thymol and (*E*)-cinnamaldehyde on *Amblyomma sculptum* larvae.

Fraction of LC ₅₀	Component A	Component B	Larval mortality (%)			CI	Effect CompuSyn modified
			A (mean ± S.D)	B (mean ± S.D)	A + B (mean ± S.D)		
1/4	Carvacrol	Thymol	1.1 ± 1.5	0.0 ± 0.0	10.5 ± 6.7	1.01	Additive
1/2	Carvacrol	Thymol	14.1 ± 6.6	2.81 ± 2.9	58.3 ± 33.5	1.09	Additive
LC ₅₀	Carvacrol	Thymol	89.2 ± 7.1	48.6 ± 18.2	99.5 ± 1.3	0.79	Moderate synergy
1/4	Carvacrol	(<i>E</i>)-cinnamaldehyde	1.1 ± 1.5	1.5 ± 1.1	14.1 ± 13.3	0.85	Moderate synergy
1/2	Carvacrol	(<i>E</i>)-cinnamaldehyde	14.1 ± 6.6	0.9 ± 0.9	19.8 ± 17.4	1.50	Antagonism
LC ₅₀	Carvacrol	(<i>E</i>)-cinnamaldehyde	89.2 ± 7.1	37.4 ± 18.9	65.4 ± 36.6	1.76	Antagonism
1/4	Thymol	(<i>E</i>)-cinnamaldehyde	0.0 ± 0.0	1.5 ± 1.1	0.0 ± 0.0	4.30	Antagonism
1/2	Thymol	(<i>E</i>)-cinnamaldehyde	2.8 ± 2.9	0.9 ± 0.9	2.1 ± 1.6	2.88	Antagonism
LC ₅₀	Thymol	(<i>E</i>)-cinnamaldehyde	48.6 ± 18.2	37.4 ± 18.9	16.7 ± 8.5	2.54	Antagonism

Combination index (CI) <0.70: synergy; 0.70–0.90: moderate synergy; 0.90–1.10: additive; 1.10–1.45: moderate antagonism; >1.45: antagonism. The control (70% ethanol) produced zero mortality.

Table 4Evaluation of paired combinations of components of essential oils, carvacrol, thymol and (*E*)-cinnamaldehyde on *Dermacentor nitens* larvae.

Fraction of LC ₅₀	Component A	Component B	Larval mortality (%)			CI	Effect CompuSyn modified
			A (mean ± S.D)	B (mean ± S.D)	A + B (mean ± S.D)		
1/4	Carvacrol	Thymol	5.9 ± 3.6	0.9 ± 1.5	9.4 ± 9.1	1.00	Additive
1/2	Carvacrol	Thymol	9.9 ± 3.5	3.7 ± 4.8	43.9 ± 13.7	0.54	Synergy
LC ₅₀	Carvacrol	Thymol	28.5 ± 10.6	34.9 ± 15.8	100 ± 0	0.07	Synergy
1/4	Carvacrol	(<i>E</i>)-cinnamaldehyde	5.9 ± 3.6	1.8 ± 3.0	3.6 ± 4.6	2.00	Antagonism
1/2	Carvacrol	(<i>E</i>)-cinnamaldehyde	9.9 ± 3.5	8.3 ± 5.2	11.7 ± 7.6	2.01	Antagonism
LC ₅₀	Carvacrol	(<i>E</i>)-cinnamaldehyde	28.5 ± 10.6	55.8 ± 19.6	52.7 ± 19.8	1.37	Antagonism
1/4	Thymol	(<i>E</i>)-cinnamaldehyde	0.9 ± 1.5	1.8 ± 3.0	2.2 ± 2.4	1.65	Antagonism
1/2	Thymol	(<i>E</i>)-cinnamaldehyde	3.7 ± 4.8	8.3 ± 5.2	3.3 ± 4.1	2.82	Antagonism
LC ₅₀	Thymol	(<i>E</i>)-cinnamaldehyde	34.9 ± 15.8	55.8 ± 19.6	47.3 ± 12.3	1.57	Antagonism

Combination index (CI) <0.70: synergy; 0.70–0.90: moderate synergy; 0.90–1.10: additive; 1.10–1.45: moderate antagonism; >1.45: antagonism. The control (70% ethanol) produced zero mortality.

2014) due to the potential advantages they have when compared to chemical products, such as low toxicity to mammals and short persistence in the environment (Miresmailli and Isman, 2006). Besides this, a number of components obtained from the essential oils of plants have been shown to be effective against various tick species, thus demonstrating the potential of developing new acaricidal formulations. However, this is the first study of the effect of

the association of monoterpenes (thymol, carvacrol) and a phenylpropanoid (*E*)-cinnamaldehyde on these two tick species.

According to the LC₅₀ values against *A. sculptum* and *D. nitens* larvae, (*E*)-cinnamaldehyde had the greatest activity, followed by thymol and carvacrol. The activity of these substances on these ticks' larvae has been reported in other studies (Daemon et al., 2012; Senra et al., 2013a,b), however, in those studies, these

components of essential oils caused high mortality (>80%) starting from the lowest concentrations tested, making it impossible to calculate the LC₅₀ values, as achieved in this study. With respect to the order of activity and efficacy, Senra et al. (2013b) also found that (*E*)-cinnamaldehyde was the most active substance against *A. cajennense* s.l. larvae, followed by thymol and carvacrol. In experiments with bacteria of the genera *Streptococcus* and *Prevotella*, (*E*)-cinnamaldehyde performed by Didry et al. (1994), was also the most active substance, followed by thymol and carvacrol. However, Friedman et al. (2002), assessing the bactericidal effect of essential oils on *Campylobacter jejuni*, *Escherichia coli*, *Listeria monocytogenes* and *Salmonella enterica*, observed that carvacrol had the strongest activity, followed by (*E*)-cinnamaldehyde and thymol. The differences between our results and those presented in the literature indicate that organisms have varied responses to different natural compounds.

Comparison of the activity of each substance alone against the two tick species showed that *A. sculptum* and *D. nitens* had similar sensitivity to (*E*)-cinnamaldehyde, thymol and carvacrol, with proximate LC₅₀ values and overlapping confidence intervals. Daemon et al. (2012) evaluated the activity of thymol on *R. sanguineus* and *D. nitens* larvae and also observed that these two tick species had similar sensitivity to this monoterpene when tested at concentrations of 10, 15 and 20 µl/ml, with respective mortality rates of 96.7, 95.9 and 98.1% against *R. sanguineus* and 90.2, 90.3, 99.5% against *D. nitens*.

With respect to the combinations of thymol and carvacrol, there was a moderate synergetic effect on *A. sculptum* larvae when using the concentration of the LC₅₀, and synergy for *D. nitens* when using the concentration of the LC₅₀ and 1/2 of LC₅₀. The synergetic action of thymol combined with carvacrol has already been reported for a variety of organisms, such as bacteria and nematodes (Didry et al., 1994; Ntalli et al., 2010) as well as for other arthropods, such as beetles (Lima et al., 2011) and lepidopterans (Pavela et al., 2010). Lima et al. (2011) observed a synergetic effect of the interaction of thymol and carvacrol against *Tenebrio molitor* (Linnaeus, 1758) and Pavela (2010) observed the same effect in tests with *Spodoptera littoralis* (Boisduval, 1833) larvae. Studies have shown that the toxicity of thymol and carvacrol to different organisms is partly related to the effect of these monoterpenes on the cells of target organisms, since they cause disorganization in the cell membrane, leading it to lose permeability (Medeiros et al., 2011). That effect has been demonstrated both for eukaryotic (Medeiros et al., 2011) and prokaryotic cells (Lambert et al., 2001). In the case of thymol, this effect has also been verified on oocytes of the cattle tick *R. sanguineus* (Matos et al., 2014). Hence, it can be inferred that the synergetic effect may be related to the joint action on the cell membrane of the ticks, with enhanced action in combination. However, histological tests are necessary to confirm this hypothesis.

In insects, thymol and carvacrol have action on the GABA receptor, acting on the neuromuscular system, deregulating motor activity (Waliwitiya et al., 2010; Tong and Coasts, 2010) and on the octopamine receptor, causing cell membrane disruption and compromising the tracheal system (Isman et al., 2007; Khanikar et al., 2013). It can be inferred that these monoterpenes also act on these receptors in ticks, although there are no studies about it in the literature.

The associations between carvacrol and (*E*)-cinnamaldehyde presented an antagonistic effect in all combinations tested against *D. nitens*, while in the case of *A. sculptum*, only the combination with a concentration of 1/4 of LC₅₀ did not present an antagonistic effect (showing moderate synergism instead). Pavela (2014), in assessing the effect of some aromatic compounds on *S. littoralis*, found an antagonistic effect of the combination of carvacrol and (*E*)-cinnamaldehyde. However, in a study with bacteria, Didry et al. (1994) observed a synergetic effect when combining car-

vacrol with (*E*)-cinnamaldehyde against *Streptococcus milleri* and *Prevotella intermedia*. The same result was reported by Ye et al. (2013) based on tests with the bacteria *E. coli*, *Staphylococcus aureus*, *S. intermedius* and *S. haemolyticus*, in which the association of carvacrol and (*E*)-cinnamaldehyde presented a synergetic effect. This fact indicates that different species can respond distinctly to the association of components of essential oils, so that an effect found against one organism cannot be extrapolated to others. These differences can also be related to the testing methods, concentrations tested and source of the substances.

The combination of thymol and (*E*)-cinnamaldehyde had an antagonistic effect at all the concentrations of each substance, for both tick species. The same effect was observed by Didry et al. (1994) in tests applying thymol together with (*E*)-cinnamaldehyde against the bacteria *Streptococcus mitis* and *Prevotella oris*. Pavela (2014) also found an antagonistic effect of thymol combined with (*E*)-cinnamaldehyde against *S. littoralis*. Therefore, it can be assumed that this antagonistic effect is related to the possible reaction of (*E*)-cinnamaldehyde (aldehyde) with thymol and carvacrol (phenolic alcohol), forming a hemiacetal or acetal, which normally does not have biological activity due to the unavailability of an active functional group (phenol and enone) in both compounds.

Thus, it is possible to conclude that the combination of carvacrol and thymol, at the LC₅₀, is more effective in controlling the two tick species, because there is a synergetic effect for both species, thus enabling the use of these monoterpenes in the development of commercial acaricidal formulations aimed at controlling mixed infestations. In turn, combinations containing (*E*)-cinnamaldehyde with thymol and carvacrol are not appropriate for controlling these ectoparasites, since we observed an antagonistic effect in the majority of the associations tested. In cases of synergetic effect, it will be possible to reduce the concentrations of the components of essential oils applied without losing efficiency. This knowledge can serve as basis for developing new acaricidal products. However, further studies are necessary on other stages of these tick species and under natural conditions, as well as toxicity evaluation in the hosts.

References

- Borges, L.M.F., Leite, R.C., 1993. Aspectos biológicos do *Dermacentor nitens* (Neuman, 1897) em condições de laboratório. Arq. Bras. Med. Vet. Zoo. 45, 586–591.
- Borges, L.M.F., Dias, S.L.A., Barbosa, C.S., 2011. Perspectives for the use of plant extracts to control the cattle tick *Rhipicephalus (Boophilus) microplus*. Rev. Bras. Parasitol. Vet. 20, 89–96.
- Cetin, H., Cilek, J.E., Aydin, L., Yanikoglu, A., 2009. Acaricidal effects of the essential oil of *Origanum minutiflorum* (Lamiaceae) against *Rhipicephalus turanicus* (Acari: Ixodidae). Vet. Parasitol. 160, 359–361.
- Cetin, H., Cilek, J.E., Oz, E., Aydin, L., Deveci, O., Yanikoglu, A., 2010. Acaricidal activity of *Satureja thymbra* L. essential oil and its major components, carvacrol and [gamma]-terpinene against adult *Hyalomma marginatum* (Acari: Ixodidae). Vet. Parasitol. 170, 287–290.
- Chagas, A.C.S., 2004. Controle de parasitos usando extrato de plantas. Rev. Bras. Parasitol. Vet. 13, 156–160.
- Chou, T.C., 2006. Theoretical basis, experimental design, and computerized simulation of synergism and antagonism in drug combination studies. Pharmacol. Ver. 58, 621–681.
- Coskun, S., Girisgin, O., Kürkcüoğlu, M., Malyer, H., Girisgin, A.O., Kirimer, N., Baser, K.H., 2008. Acaricidal efficacy of *Origanum onites* L. essential oil against *Rhipicephalus turanicus* (Ixodidae). Parasitol. Res. 103, 259–261.
- Daemon, E., Monteiro, C.M.O., Rosa, L.S., Clemente, M.A., Arcoverde, A., 2009. Evaluation of the acaricide activity of thymol on engorged and unengorged larvae of *Rhipicephalus sanguineus* (Latreille, 1808) (Acari: Ixodidae). Parasitol. Res. 105, 495–497.
- Daemon, E., Maturano, R., Monteiro, C.M.O., Goldner, M.S., Massoni, T., 2012. Acaricidal activity of hydroethanolic formulations of thymol against *Rhipicephalus sanguineus* (Acari: Ixodidae) and *Dermacentor nitens* (Acari: Ixodidae) larvae. Vet. Parasitol. 186, 542–545.
- Dantas-Torres, F., Onofrio, V.C., Barros-Battesti, D.M., 2009. The ticks (Acari: Ixodida: Argasidae, Ixodidae) of Brazil. Syst. Appl. Acarol. 14, 30–46.
- Didry, N., Dubreuil, L., Pinkas, M., 1994. Activity of thymol, carvacrol, cinnamaldehyde and eugenol on oral bacteria. Pharm. Acta Helv. 69, 25–28.

- Dolan, M.C., Jordan, R.A., Schulze, T.L., Schulze, C.J., Manning, M.C., Ruffolo, D., Schmidt, J.P., Piesman, J., Karchesy, J.J., 2009. Ability of two natural products, nootkatone and carvacrol, to suppress *Ixodes scapularis* and *Amblyomma americanum* (Acari: Ixodidae) in a Lyme disease endemic area of New Jersey. *J. Econ. Entomol.* 102, 2316–2324.
- Elise, L., Wall, R., 2014. The use of essential oils in veterinary ectoparasites control: a review. *Med. Vet. Entomol.* 28, 233–243.
- Friedman, M., Henika, P.R., Mandrell, R.E., 2002. Bactericidal activities of plant essential oils and some of their isolated constituents against *Campylobacter jejuni*, *E. coli*, *Listeria monocytogenes*, and *Salmonella enterica*. *J. Food Prot.* 65, 1545–1560.
- Gallardo, A., Picollo, M.I., González-Audino, P., Mougabure-Cueto, G., 2012. Insecticidal activity of individual and mixed monoterpenoids of geranium essential oil against *Pediculus humanus capitis* (Phthiraptera: Pediculidae). *J. Med. Entomol.* 49, 332–335.
- Guedes, E., Leite, R.C., Prata, M.C.A., Pacheco, R.C., Walker, D.H., Labruna, M.B., 2005. Detection of *Rickettsia rickettsii* in the tick *Amblyomma cajennense* in a new Brazilian spotted fever-endemic area in the state of Minas Gerais. *Mem. Inst. Oswaldo Cruz* 100, 841–845.
- Guglielmone, A.A., Szabó, M.P.J., Martins, J.R.S., Estrada-Peña, A., 2006. Diversidade e importância de carrapatos na sanidade animal. In: Barros-Battesti, D.M., Arzua, M., Bechara, G.H. (Eds.), Carrapatos de importância médica veterinária da região neotropical: um guia ilustrado para identificação de espécies. *Vox/ICTD-3/Butantan*, São Paulo, pp. 115–124.
- Guimarães, J.H., Tucci, E.C., Barros-Battesti, D.M. (Eds.), 2001. Ectoparasitos de importância veterinária. *Plêiade/FAPESP*, São Paulo, p. 213.
- Isman, M.B., Machial, C.M., Miresmailli, S., Bainard, L.D., 2007. Essential oil-based pesticides: new insights from old chemistry. In: Ohkawa, H., Miyagawa, H., Lee, P.W. (Eds.), *Pesticide chemistry: Crop protection, public health, environmental safety*. Wiley, pp. 201–210.
- Khanikar, B., Parida, P., Yadav, R.N.S., Bora, D., 2013. Comparative mode of action of some terpene compounds against octopamine receptor and acetyl cholinesterase of mosquito and human system by the help of homology modeling and docking studies. *J. Appl. Pharm. Sci.* 3, 6–12.
- Labruna, M.B., Machado, R.Z., 2006. Agentes transmitidos por carrapatos na região neotropical. In: Barros-Battesti, D.M., Arzua, M., Bechara, G.H. (Eds.), Carrapatos de importância médica veterinária da região neotropical: um guia ilustrado para identificação de espécies. *Vox/ICTD-3/Butantan*, São Paulo, pp. 155–164.
- Lambert, R.J.W., Skandamis, P.N., Coote, P., Nychas, G.J.E., 2001. A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. *J. Appl. Microbiol.* 91, 453–462.
- Lima, R.K., Cardoso, M.G., Moraes, J.C., Carvalho, S.M., Rodrigues, V.G., Guimarães, L.G.L., 2011. Chemical composition and fumigant effect of essential oil of *Lippia sidoides* Cham. and monoterpenes against *Tenebrio molitor* (L.) (COLEOPTERA: TENEBRIONIDAE). *Cienc. Agrotec.* 35, 664–671.
- Matos, R.S., Daemon, E., Camargo-Mathias, M.I., Furquim, K.C.S., Sampieri, B.R., Remédio, R.N., Araújo, L.X., Novato, T.P.L., 2014. Histopathological study of *Rhipicephalus sanguineus* (Acari: Ixodidae) exposed to different thymol concentrations. *Parasitol. Res.* 113, 4555–4565.
- Medeiros, M.G.F., Silva, A.C., Citó, A.M.G.L., Borges, A.R., Lima, S.G., Lopes, J.A.D., Figueredo, R.C.B.Q., 2011. In vitro antileishmanial activity and cytotoxicity of oil essential from *Lippia sidoides* Cham. *Parasitol. Int.* 60, 237–241.
- Mendes, A.S., Daemon, E., Monteiro, C.M.O., Maturano, R., Brito, F.C., Massoni, T., 2011. Acaricidal activity of thymol on larvae and nymphs of *Amblyomma cajennense* (Acari: Ixodidae). *Vet. Parasitol.* 183, 136–139.
- Miresmailli, S., Isman, M.B., 2006. Efficacy and persistence of rosemary oil as an acaricide against two spotted Spider Mite (Acari: Tetranychidae) on greenhouse tomato. *J. Econ. Entomol.* 99, 2015–2023.
- Monteiro, C.M.O., Prata, M.C.A., 2013. Controle biológico do carrapato dos bovinos *Rhipicephalus microplus* com a utilização de nematoides entomopatogênicos: conquistas e desafios. In: Veríssimo, C. (Ed.), *Controle de carrapatos nas pastagens*. FAPESP, Nova Odessa, pp. 46–68.
- Monteiro, C.M.O., Daemon, E., Clemente, M.A., Rosa, L.S., Maturano, R., 2009. Acaricidal efficacy of thymol on engorged nymphs and females of *Rhipicephalus sanguineus* (Latreille, 1808) (Acari: Ixodidae). *Parasitol. Res.* 105, 1093–1097.
- Monteiro, C.M.O., Daemon, E., Silva, A.M.R., Maturano, R., Amaral, C., 2010. Acaricide and ovicide activities of thymol on engorged females and eggs of *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae). *Parasitol. Res.* 106, 615–619.
- Monteiro, C.M., Maturano, R., Daemon, E., Catunda-Junior, F.E.A., Calmon, F., Senra, T.O.S., Faza, A., Carvalho, M.G., 2012. Acaricidal activity of eugenol on *Rhipicephalus microplus* (Acari: Ixodidae) and *Dermacentor nitens* (Acari: Ixodidae) larvae. *Parasitol. Res.* 111, 1295–1300.
- Nava, S., Beati, L., Labruna, M.B., Cáceres, A.G., Mangold, A.J., Guglielmone, A.A., 2014. Reassessment of the taxonomic status of *Amblyomma cajennense* (Fabricius, 1787) with the description of three new species, *Amblyomma tonelliae* n. sp., and reinstatement of *Amblyomma mixtum* Koch, 1844, and *Amblyomma sculptum* Berlese, 1888 (Ixodida: Ixodidae). *Tick-borne Dis.* 5, 252–276.
- Novelino, A.M.S., Daemon, E., Soares, G.L.G., 2007. Evaluation of the acaricide effect of thymol, menthol, salicylic acid, and methyl salicylate on *Boophilus microplus* (Canestrini 1887) (Acari: Ixodidae) larvae. *Parasitol. Res.* 101, 809–811.
- Ntalli, N.G., Ferrari, F., Giannakou, I., Menkissoglu-Spiroudi, U., 2010. Synergistic and antagonistic interactions of terpenes against *Meloidogyne incognita* and the nematocidal activity of essential oils from seven plants indigenous to Greece. *Pest Manage. Sci.* 67, 341–351.
- Oliveira, P.R., 2004. Biologia e controle de *Amblyomma cajennense*. *Rev. Bras. Parasitol. Vet.* 23, 118–122.
- Pavela, R., 2010. Acute and synergistic effects of monoterpenoid essential oil compounds on the larvae of *Spodoptera littoralis*. *J. Biopest.* 3, 573–578.
- Pavela, R., 2014. Acute, synergistic and antagonistic effects of some aromatic compounds on the *Spodoptera littoralis* Bois. (Lep. Noctuidae) larvae. *Ind. Crop Prod.* 60, 247–258.
- Regnault-Roger, C., Philogène, B.J.R., 2008. Past and current prospects for the use of botanicals and plant allelochemicals in integrated pest management. *Pharm. Biol.* 46, 41–52.
- Resende, J.D.S.A., Daemon, E., Monteiro, C.M.O., Maturano, R., Prata, M.C.A., Rodrigues, A.F.S.F., 2012. Toxicity of solvents and surfactants to *Amblyomma cajennense* (Fabricius, 1787) (Acari: Ixodidae) and *Dermacentor nitens* (Neumann, 1897) (Acari: Ixodidae) larvae. *Exp. Parasitol.* 131, 139–142.
- Senra, T.O.S., Zeringóta, V., Monteiro, C.M.O., Calmon, F., Maturano, R., Gomes, G.A., Faza, A., Carvalho, M.G., Daemon, E., 2013a. Assessment of the acaricidal activity of carvacrol, (E)-cinnamaldehyde, trans-anethole, and linalool on larvae of *Rhipicephalus microplus* and *Dermacentor nitens* (Acari: Ixodidae). *Parasitol. Res.* 112, 1461–1466.
- Senra, T.O.S., Calmon, F., Zeringóta, V., Monteiro, C.M.O., Maturano, R., Matos, R.S., Melo, D., Gomes, G.A., Carvalho, M.G., Daemon, E., 2013b. Investigation of activity of monoterpenes and phenylpropanoids against immature stages of *Amblyomma cajennense* and *Rhipicephalus sanguineus* (Acari: Ixodidae). *Parasitol. Res.* 112, 3471–3476.
- Stone, B.F., Haydock, K.P., 1962. A method for measuring the acaricide susceptibility of the cattle *B. microplus* (Canestrini). *Bull. Entomol. Res.* 53, 563–578.
- Tong, F., Coats, J.R., 2010. Effects of monoterpenoid insecticides on [³H]-TBOB binding in house fly GABA receptor and ³⁶Cl⁻ uptake in American cockroach ventral nerve cord. *Pestic. Biochem. Phys.* 98, 317–324.
- Waliwitiya, R., Belton, P., Nicholson, R.A., Lowenberger, C.A., 2010. Effects of the essential oil constituent thymol and other neuroactive chemicals on flight motor activity and wing beat frequency in the blowfly *Phaenicia sericata*. *Pest Manage. Sci.* 66, 277–289.
- Ye, H., Shen, S., Xu, J., Lin, S., Yuan, Y., Jones, G.S., 2013. Synergistic interactions of cinnamaldehyde in combination with carvacrol against food-borne bacteria. *Food Control* 34, 619–623.