



Contents lists available at SciVerse ScienceDirect

Tetrahedron Letters

journal homepage: www.elsevier.com/locate/tetlet

Highly enantioselective bioreduction of ethyl 3-oxohexanoate

Aline de Souza Ramos^a, Joyce Benzaquem Ribeiro^{b,*}, Raquel de Oliveira Lopes^b,
Selma Gomes Ferreira Leite^c, Rodrigo Octavio Mendonça Alves de Souza^b

^aFarmanguinhos, Fundação Oswaldo Cruz, Rua Sizenando Nabuco 100, Manguinhos, Rio de Janeiro, RJ 21041-250, Brazil

^bInstituto de Química, Universidade Federal do Rio de Janeiro, CT Bloco A, Cidade Universitária, Rio de Janeiro, RJ 21949-909, Brazil

^cEscola de Química, Universidade Federal do Rio de Janeiro, CT Bloco E, Cidade Universitária, Rio de Janeiro, RJ 21949-909, Brazil

ARTICLE INFO

Article history:

Received 9 August 2011

Revised 30 August 2011

Accepted 6 September 2011

Available online 10 September 2011

Keywords:

β-Ketoester

Biotransformation

Immobilization

Kluyveromyces marxianus

Chiral reduction

ABSTRACT

Seven wild-type microorganism strains were used to reduce ethyl 3-oxohexanoate to ethyl (R)-3-hydroxyhexanoate. Free cells of *Kluyveromyces marxianus* and *Aspergillus niger* led to higher than 99% of conversion with higher than 99% ee. After immobilization in calcium alginate spheres, cells of *K. marxianus* exhibited high enantioselectivity (>99% ee) and conversion level (99%) within 24 h even if substrate was added at concentration of 10 g/L (62 mM).

© 2011 Elsevier Ltd. Open access under the [Elsevier OA license](http://www.elsevier.com/locate/elsevier).

Chiral β-hydroxyesters are widely used as building blocks for the synthesis of fine chemicals, pharmaceuticals, and natural products.^{1,2} Ethyl 3-hydroxyhexanoate is a key intermediate in the synthesis of (+)-neopeltolide, a bioactive marine macrolide with potent antiproliferative activity against cancer cell lines. The cellular target for this class of substance is the inhibition of cytochrome bc1 complex.³

Many efforts have been made in the search for highly selective methods to obtain chiral β-hydroxyesters, as ethyl 3-hydroxyhexanoate. Asymmetric reduction of prochiral ketoesters is an alternative route. Bioreductions are attractive methods, mainly due to high enantioselectivity, mild and safe reaction conditions, and lower environmental impact compared to conventional reactions in organic chemistry.^{4–7}

Isolated enzymes and whole cells can be used as biocatalysts in enantioselective reductions. Reductase enzymes require reduced nicotinamide cofactors and when isolated enzymes are used, a second catalytic cycle is necessary to regenerate cofactors and sustain catalytic activity.^{4–7} So, whole cells are frequently preferred because of their own cofactor regeneration system even if competing enzymes within cells could decrease stereoselectivity. Reaction conditions and suitable microorganisms should be investigated to improve yields.^{4–9}

In recent years we have been investigating the microbial reduction of ketoesters.^{2,8–11} In the present work, seven wild-type

microorganism strains were employed in the asymmetric reduction of ethyl 3-oxohexanoate to ethyl (R)-3-hydroxyhexanoate (Fig. 1). Some yeasts were also tested after immobilization in calcium alginate spheres.

In the first step of this work, free cells of five yeasts (*Saccharomyces cerevisiae* 40, *Hansenula* sp., *Geotrichum candidum*, *Kluyveromyces marxianus*, and *Rhodotorula rubra*) and two filamentous fungi (*Trichoderma harzianum* and *Aspergillus niger*) were tested (Table 1).¹² All the strains were able to catalyze the reaction with excess of the (R)-hydroxyester (determined by the measurement of optical rotation)¹³ with an excellent conversion level (>99%) within 24 h. The enantiomeric excess varied from 9% (with *T. harzianum*) to higher than 99% (with *K. marxianus* and *A. niger*). Examples of chromatograms obtained are shown in Figure 2.

In previous studies, these microorganisms were successfully used in the reduction of methyl acetoacetate,¹⁰ ethyl acetoacetate,^{2,10} ethyl 2-methylacetoacetate,¹⁰ ethyl benzoylacetate,^{8,11} and ethyl 4-chloroacetoacetate.⁹ However, excess of the (S)-hydroxyester was obtained in almost all of the cases. Only *K. marxianus*, *T. harzianum* and *A. niger* led to the excess of the (R)-hydroxyester in the reduction of ethyl acetoacetate (67% ee, 51% ee, and 19% ee, respectively) and *A. niger* maintained the




Figure 1. Bioreduction of ethyl 3-oxohexanoate to ethyl 3-(R)-hydroxyhexanoate.

* Corresponding author. Tel.: +55 21 2562 7248.

E-mail address: joyce_benzaquem@yahoo.com.br (J.B. Ribeiro).

Table 1

Bioreduction of ethyl 3-oxohexanoate (5 g/L) to ethyl (*R*)-3-hydroxyhexanoate by free cells. Incubation: 30 °C, 150 rpm, 24 h

Microorganism	Conversion ^a (%)	ee (%)
<i>Saccharomyces cerevisiae</i> 40	>99	97.5
<i>Hansenula</i> sp.	>99	87.2
<i>Geotrichum candidum</i>	>99	85.7
<i>Kluyveromyces marxianus</i>	>99	>99
<i>Rhodotorula rubra</i>	>99	90
<i>Trichoderma harzianum</i>	>99	9
<i>Aspergillus niger</i>	>99	>99

^a Determined by GC analysis.

(*R*)-enantioselectivity in the reduction of methyl acetoacetate (45% ee). So, enantioselectivity depends critically on the length of the carbon chain, as also observed by other researchers.^{4,14}

Some authors described such enantioselectivity and conversion level employing recombinant microorganisms¹⁵ and isolated enzymes.^{6,16,17} Rodríguez et al.¹⁵ showed that it was possible to obtain both (*S*)- and (*R*)-hydroxyesters with higher than 98% ee by using genetic engineering of baker's yeast, although lower substrate concentration (10 mM) had been used in comparison with the present work (31 mM). Kaluzna et al.¹⁶ also achieved high (*S*)-enantioselectivity (>99% ee) by using isolated *S. cerevisiae* reductases overproduced in *Escherichia coli*, while Zhu et al.¹⁷ used recombinant ketoreductases to reach both isomers with higher than 99% ee. However, wild-type microorganism strains are generally preferred than recombinant strains because of their robustness.² Moreover, whole cells do not require the addition of expensive cofactors, which is an advantage over the use of isolated enzymes, besides the lower cost.¹⁸

Among the wild-type microorganisms, *S. cerevisiae* was most frequently assayed in the reduction of ethyl 3-oxohexanoate. North¹⁹ obtained higher than 98% ee of the (*S*)-hydroxyester and conversion of 55% with *S. cerevisiae* in petrol; Rotthaus et al.²⁰ employed *S. cerevisiae* in the reduction using different solvents and obtained 100% conversion and 86% ee of the (*R*)-isomer when the reaction was conducted in water. Rodríguez et al.¹⁵ achieved 98% ee of the same isomer when *S. cerevisiae* was used as the biocatalyst, substrate at 12.6 mM, and glucose or galactose as the carbon source. Dahl et al.²¹ achieved 99% ee of the (*R*)-hydroxyester using baker's yeast but they added allyl alcohol to the reaction medium, which is a carcinogenic and volatile enzymatic inhibitor not recommended for industrial processes.²² Buisson et al.²² described that the yeast *G. candidum* afforded the same isomer with 98% ee and 90% of conversion within 24 h. In the present work, as can be seen in Table 1, free cells of *K. marxianus* and *A. niger* carried out in the (*R*)-hydroxyester production with more than 99% ee and more than 99% conversion in medium containing substrate at 5 g/L or 31 mM.

In order to improve the method, some yeasts (*Hansenula* sp., *K. marxianus*, and *R. rubra*) were immobilized in calcium alginate spheres and tested for their ethyl (*R*)-3-oxohexanoate reduction ability in two cycles (Table 2).²³ This entrapment technique makes the product recovery much easier and the biocatalyst can be readily reused. However, the immobilization can influence enantiomeric excess and conversion level.^{2,8,9,11,24,25}

As shown in Table 2, the immobilized cells of *K. marxianus* exhibited the best results because they led to the high enantioselectivity (>99% ee) and conversion level (>99%) after immobilization in the two cycles, with storage of 12 days between the cycles. The yeast *Hansenula* sp. gave high conversion level in the two cycles, but with decrease in enantioselectivity. In the first use of immobilized cells of *R. rubra*, a little decrease was observed in conversion level and, in the second cycle, still lower conversion was obtained.




Figure 2. Typical chromatograms showing the selectivities achieved. Chiral GC analysis on column Beta DEX325 (30 m × 0.25 mm × 0.25 μm), at 90 °C (23 min): (A) ethyl 3-oxohexanoate (substrate); (B) ethyl 3-hydroxybutanoate (racemate obtained via NaBH₄ reduction); (C) *Trichoderma harzianum* reduction; (D) *Hansenula* sp. reduction; (E) *Kluyveromyces marxianus* reduction. Retention times (*t_R*): *t_R* (substrate) = 15.3 min, *t_R* (*S*-enantiomer) = 18.8 min, *t_R* (*R*-enantiomer) = 19.3 min.

High substrate concentration is important for an industrial biotransformation process^{24,26} but it can decrease conversion level and enantioselectivity.²⁷ This effect was previously observed in the reduction of ethyl 4-chloroacetoacetate to ethyl (*S*)-4-chloro-3-hydroxybutanoate by the immobilized cells of *K. marxianus*.⁹ In the reactions catalyzed by whole cells, substrate toxicity is an aggravating factor.²⁸ Immobilization could reduce toxic effects by a diffusion barrier that protects cells and it could allow the addition of substrate at higher concentrations.²⁴ To evaluate

Table 2

Bioreduction of ethyl 3-oxohexanoate (5 g/L) to ethyl (R)-3-hydroxyhexanoate by immobilized cells. Incubation: 30 °C, 150 rpm, 24 h

Microorganism	First cycle		Second cycle (after 12 d)	
	Conversion ^a (%)	ee (%)	Conversion ^a (%)	ee (%)
<i>K. marxianus</i>	>99	>99	>99	>99
<i>Hansenula</i> sp.	>99	75.1	>99	60.8
<i>R. rubra</i>	96.5	80.6	61.1	80.5

^a Determined by GC analysis.

whether substrate concentration could influence catalytic activity, the cells of *K. marxianus* were immobilized again and used in the reduction of ethyl 3-oxohexanoate at substrate concentrations of 7.5 g/L and 10 g/L. This biocatalyst also showed the same conversion level (>99%) and enantioselectivity (>99% ee) observed before.

In conclusion, the filamentous fungus *A. niger* and the yeast *K. marxianus* led to the high conversion of ethyl 3-oxohexanoate to ethyl (R)-3-hydroxyhexanoate (>99%) with extremely high enantioselectivity (>99% ee). To our knowledge, this is the first Letter on the use of these two microorganisms in the reduction of ethyl 3-oxohexanoate. The high conversion level, high enantiomeric excess, tolerance to substrate and the possibility of immobilization, and reuse make *K. marxianus* a promising biocatalyst for industrial applications.

Acknowledgments

Financial support from CAPES and CNPq-BRAZIL is acknowledged. Analytical support from DQO and DQI-IQ-UFRJ is also acknowledged.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.tetlet.2011.09.023.

References and notes

- (a) Mori, K. *Tetrahedron* **1989**, *45*, 3233–3298; (b) Sheldon, R. A. *Chirotechnology*; Dekker: New York, 1993.
- Ribeiro, J. B.; Ramos, M. C. K. V.; Aquino Neto, F. R.; Leite, S. G. F.; Antunes, O. A. C. *J. Mol. Catal. B* **2003**, *24–25*, 121–124.
- (a) Wright, A. E.; Cook Botelho, J.; Guzmán, E.; Harmody, D.; Linley, P.; McCarthy, P. J.; Pitts, T. P.; Pomponi, S. A.; Reed, J. K. *J. Nat. Prod.* **2007**, *70*, 412–416; (b) Paterson, I.; Miller, N. A. *Chem. Commun.* **2008**, 4708–4710; (c) Guinchard, X.; Roulland, E. *Org. Lett.* **2009**, *11*, 4700–4703; (d) Custar, D. W.; Zabawa, T. P.; Hines, J.; Crews, C. M.; Scheidt, K. A. *J. Am. Chem. Soc.* **2009**, *131*, 12406–12414.
- Nakamura, K.; Yamanaka, R.; Matsuda, T.; Harada, T. *Tetrahedron: Asymmetry* **2003**, *14*, 2659–2681.
- Kataoka, M.; Kita, K.; Wada, M.; Yasohara, Y.; Hasegawa, J.; Shimizu, S. *Appl. Microbiol. Biotechnol.* **2003**, *62*, 437–445.
- Yang, Y.; Zhu, D.; Piegat, T. J.; Hua, L. *Tetrahedron: Asymmetry* **2007**, *18*, 1799–1803.
- Houng, J. Y.; Liao, J. H.; Wu, J. Y.; Shen, S. C.; Hsu, H. F. *Process Biochem.* **2007**, *42*, 1–7.
- Ramos, A. S.; Ribeiro, J. B.; Vazquez, L.; Fiaux, S. B.; Leite, S. G. F.; Ramos, M. C. K. V.; Aquino Neto, F. R.; Antunes, O. A. C. *Tetrahedron Lett.* **2009**, *50*, 7362–7364.
- Ribeiro, J. B.; Ramos, A. S.; Fiaux, S. B.; Leite, S. G. F.; Ramos, M. C. K. V.; Aquino Neto, F. R.; Antunes, O. A. C. *Tetrahedron: Asymmetry* **2009**, *20*, 2263–2266.
- Ramos, A. S.; Ribeiro, J. B.; Vazquez, L.; Fiaux, S. B.; Leite, S. G. F.; Cruz, R. A.; Ramos, M. C. K. V.; Aquino Neto, F. R.; Antunes, O. A. C. *Tetrahedron: Asymmetry* **2009**, *20*, 559–561.

- Ribeiro, J. B.; Ramos, M. C. K. V.; Aquino Neto, F. R.; Leite, S. G. F.; Antunes, O. A. C. *Catal. Commun.* **2005**, *6*, 131–133.
- Microorganisms, media, growth conditions, and biotransformation with free cells. *Saccharomyces cerevisiae*, *Hansenula* sp., *Geotrichum candidum*, *Kluyveromyces marxianus*, *Rhodotorula rubra*, *Aspergillus niger*, and *Trichoderma harzianum* belong to the collection of the 'Departamento de Engenharia Bioquímica, Escola de Química, Universidade Federal do Rio de Janeiro (Cidade Universitária, CT Bloco E, Rio de Janeiro, Brazil, e-mail selma@eq.ufrj.br)' and are freely available upon request. Cells were allowed to grow for 48 h, under 150 rpm at 30 °C in a medium containing 1% glucose, 0.5% yeast extract, 0.5% peptone, 0.1% (NH₄)₂SO₄, and 0.1% MgSO₄·7H₂O. After that period, they were harvested by centrifugation, re-suspended in water, and used for the reaction. After centrifugation, the cells (4 g/L, dried weight) were added to the reduction medium containing: glucose (5%), MgCl₂ (0.1%) in a final volume of 100 mL. After 30 min of addition of the microorganisms, the substrate (0.5 g diluted in 1 mL of ethanol 96%) was added to the medium. The reaction was carried out in 500 mL cotton-plugged Erlenmeyer flasks for 24 h at 30 °C and 150 rpm. After 24 h, the medium was centrifuged again to separate the cells and the liquid phase was extracted with ethyl acetate. The organic phase was dried (anhydrous Na₂SO₄), filtered, and concentrated under vacuum. Conversions and enantiomeric excesses were determined by (chiral) gas chromatography (GC), on column Beta Dex325 (30 m × 0.25 mm × 0.25 μm), at 90 °C (23 min). The elution order was: ethyl (S)-3-hydroxyhexanoate (t_R = 18.8 min) followed by ethyl (R)-3-hydroxyhexanoate (t_R = 19.3 min). Substrate was eluted at 15.3 min. The reaction product was characterized by nuclear magnetic resonance (NMR) and mass spectroscopy.
- [α]_D²⁵ –30.8° (c 1 g/100 mL, CHCl₃); literature: [α]_D²⁵ –26.9° (c 1.14 g/100 mL, CHCl₃). Optical rotations were measured from CHCl₃ solutions using a JASCO DIP-370 polarimeter at the sodium D line (589 nm) operating at room temperature and compared to literature: Roche, C.; Desroy, N.; Haddad, M.; Phansavath, P.; Genet, J. P. *Org. Lett.* **2008**, *10*, 3911–3914.
- (a) Ema, T.; Moriya, H.; Kofukuda, T.; Ishida, T.; Maehara, K.; Utaka, M.; Sakai, T. *J. Org. Chem.* **2001**, *66*, 8682–8684; (b) Ishihara, K.; Yamaguchi, H.; Nakajima, N. *J. Mol. Catal. B* **2003**, *23*, 171–189.
- Rodríguez, S.; Kayser, M. M.; Stewart, J. D. *J. Am. Chem. Soc.* **2001**, *123*, 1547–1555.
- Kaluzna, I. A.; Matsuda, T.; Sewell, A. K.; Stewart, J. D. *J. Am. Chem. Soc.* **2004**, *126*, 12827–12832.
- Zhu, D.; Mukherjee, C.; Rozzell, J. D.; Kambourakis, S.; Hua, L. *Tetrahedron* **2006**, *62*, 901–905.
- Yang, W.; Xu, J. H.; Xie, Y.; Xu, Y.; Zhao, G.; Lin, G. Q. *Tetrahedron: Asymmetry* **2006**, *17*, 1769–1774.
- North, M. *Tetrahedron Lett.* **1996**, *37*, 1699–1702.
- Rotthaus, O.; Krüger, D.; Demuth, M.; Schaffner, K. *Tetrahedron* **1997**, *53*, 935–938.
- Dahl, A. C.; Fjeldberg, M.; Madsen, J. Ø. *Tetrahedron: Asymmetry* **1999**, *10*, 551–559.
- Buisson, D.; Azerad, R.; Sanner, C.; Larchevêque, M. *Tetrahedron: Asymmetry* **1991**, *2*, 987–988.
- Immobilization of cells (*Kluyveromyces marxianus*, *Rhodotorula rubra*, and *Hansenula* sp.) in calcium alginate and biotransformation. Cells grown during 48 h in the medium described before¹² were centrifuged and 0.4 g (dry weight) was re-suspended in 3 mL of distilled water to obtain a cell-suspension. A 1.5% sodium alginate aqueous solution (20 mL) was added and this mixture (cell-suspension sodium alginate aqueous solution) was dropped into a CaCl₂ aqueous solution (0.1 M) to form calcium alginate spheres. Spheres were filtered, washed with distilled water, and added to the medium containing: glucose (5%), MgCl₂ (0.1%), substrate (0.5 g diluted in 1 mL of ethanol 96%) in a final volume of 100 mL. Reduction was carried out in 500 mL cotton-plugged Erlenmeyer flasks, at 30 °C, 150 rpm during 24 h. After that period, medium was filtered to separate the biocatalyst and the liquid phase was treated as described above.¹² Immobilized cells were washed with distilled water and maintained at 4 °C in a CaCl₂ aqueous solution (0.1 M) for 12 days. Thus, the immobilized cells were incubated in growth medium¹² under 150 rpm at 30 °C for 2 h, washed with distilled water again, and reused under the same reaction's conditions.
- Milagre, C. D. F.; Milagre, H. M. S.; Moran, P. J. S.; Rodrigues, J. A. R. *J. Mol. Catal. B: Enzyme* **2009**, *56*, 55–60.
- Buque, E. M.; Chin-Joe, I.; Straathof, A. J. J.; Jongejan, J. A.; Heijnen, J. J. *Enzyme Microb. Technol.* **2002**, *31*, 656–664.
- Gröger, H.; Hummel, W.; Rollmann, C.; Chamouveau, F.; Hüsken, H.; Werner, H.; Wunderlich, C.; Abokitse, K.; Drauz, K.; Buchholz, S. *Tetrahedron* **2004**, *60*, 633–640.
- Yang, Z. H.; Yao, S. J.; Guan, Y. X. *Ind. Eng. Chem. Res.* **2005**, *44*, 5411–5416.
- Fatima, Y.; Kansal, H.; Soni, P.; Banerjee, U. C. *Process Biochem.* **2007**, *42*, 1412–1418.