Chemistry & Biology



Cysteine Reactivity Distinguishes Redox Sensing by the Heat-Inducible and Constitutive Forms of Heat Shock Protein 70

Yoshinari Miyata,^{1,3} Jennifer N. Rauch,^{1,3} Umesh K. Jinwal,² Andrea D. Thompson,¹ Sharan Srinivasan,¹ Chad A. Dickey,² and Jason E. Gestwicki^{1,*}

¹Department of Pathology, Department of Biological Chemistry, and the Life Sciences Institute, University of Michigan, Ann Arbor, MI 48109-2216, USA

²Department of Molecular Medicine, University of South Florida, 4001 E. Fletcher Avenue, MDC 36, Tampa, FL 33613, USA

³These authors contributed equally to this work

*Correspondence: gestwick@umich.edu

http://dx.doi.org/10.1016/j.chembiol.2012.07.026

SUMMARY

The heat shock protein 70 (Hsp70) family of molecular chaperones has important functions in maintaining proteostasis under stress conditions. Several Hsp70 isoforms, especially Hsp72 (HSPA1A), are dramatically upregulated in response to stress; however, it is unclear whether these family members have biochemical properties that are specifically adapted to these scenarios. The redox-active compound, methylene blue (MB), has been shown to inhibit the ATPase activity of Hsp72 in vitro, and it promotes degradation of the Hsp72 substrate, tau, in cellular and animal models. Here, we report that MB irreversibly inactivates Hsp72 but not the nearly identical, constitutively expressed isoform, heat shock cognate 70 (Hsc70; HSPA8). Mass spectrometry results show that MB oxidizes Cys306, which is not conserved in Hsc70. Molecular models suggested that oxidation of Cys306 exposes Cys267 to modification and that both events contribute to loss of ATP binding in response to MB. Consistent with this model, mutating Cys267 and Cys306 to serine made Hsp72 largely resistant to MB in vitro, and overexpression of the C306S mutant blocked MBmediated loss of tau in a cellular model. Furthermore, mutating Cys267 and Cys306 to the pseudo-oxidation mimic, aspartic acid, mirrored MB treatment: the C267D and C306D mutants had reduced ATPase activity in vitro, and overexpression of the C267/306D double mutant significantly reduced tau levels in cells. Together, these results suggest that redox sensing by specific cysteine residues in Hsp72, but not Hsc70, may be an important component of the chaperone response to oxidative stress.

INTRODUCTION

Reactive oxygen species (ROS), such as free radicals and peroxides, are produced as the result of normal metabolic and signaling processes (Forman et al., 2010; Giles et al., 2003; Paulsen and Carroll, 2010). However, an abundance of ROS is also implicated in oxidative damage to lipids, nucleic acids, and proteins, contributing to pathology in a number of diseases (Lin and Beal, 2006; Patten et al., 2010; Waris and Ahsan, 2006). Thus, to control the accumulation of ROS, organisms are equipped with several scavengers, such as glutathione and ascorbate (Apel and Hirt, 2004), and redox-sensitive transcription factors, including HIF1α (Majmundar et al., 2010), NF-κB (Morgan and Liu, 2011), and HSF1 (Zhang et al., 2011), that coordinate cellular adaptation to ROS. Another important cellular response is to protect the proteome from acute denaturation and aggregation, which could cause proteotoxicity. This type of ROS protection is often provided by a molecular chaperone that contains a reactive redox sensor (e.g., cysteine residue) linked to a system for protecting other proteins from oxidative unfolding. For example, the prokaryotic heat shock protein 33 (Hsp33) contains cysteine residues that are selectively oxidized in response to redox stress, which induces a conformation with high chaperone activity (Ilbert et al., 2007).

The heat shock protein 70 (Hsp70) family is a series of highly conserved molecular chaperones that are well known for their activity in maintaining global proteostasis. Hsp70s are involved in most steps in the life of a protein, including the folding of nascent polypeptides, protein trafficking, and degradation (Hartl et al., 2011). Members of the Hsp70 family specifically interact with unfolded substrates through a C-terminal, substratebinding domain (SBD). In addition, Hsp70s also have a conserved N-terminal nucleotide-binding domain (NBD), which binds and hydrolyzes ATP. These two domains are allosterically coupled, such that ATP hydrolysis within the NBD causes conformational changes in the SBD that enhance affinity for unfolded substrates (Mayer et al., 2000). In turn, binding of Hsp70s to unfolded proteins protects them against aggregation and assists with their refolding. However, if this process fails, Hsp70s are also involved in triage, shuttling misfolded proteins to the proteasome for turnover (Mayer and Bukau, 2005). Through these activities, Hsp70s have been linked to diseases associated with aberrant protein quality control, such as cancer and neurodegenerative disease (Evans et al., 2010; Patury et al., 2009). In addition, like Hsp33, Hsp70s have been linked to redox signaling based on its reactive cysteine residues (Liu et al., 1996; Vignols et al., 2003).

In humans, the cytosol contains at least six Hsp70 isoforms, including the constitutively expressed Hsc70 (HSPA8) and the major stress-inducible isoform, Hsp72 (HSPA1A) (Hageman et al., 2011). Hsp72 and Hsc70 have very high sequence similarity (85% identical and 94% similar). However, the levels of Hsp72 are typically low under normal conditions, and they are only highly induced in response to stress, including redox imbalance (Lindquist, 1986). Thus, Hsp72 belongs to a subfamily of Hsp70s, including HSPA6, HSPA7, and HSPA4, which is characterized by elevated expression in response to stress. One important question is whether the different family members, such as the stress-inducible forms, have any specialized biochemical functions that might distinguish them from the constitutive ones.

Previously, we identified methylene blue (MB) as an inhibitor of the ATPase activity of Hsp72 in a high throughput chemical screen (Jinwal et al., 2009). This compound has been shown to reduce the levels of some Hsp72 substrates, such as tau, polyglutamine fragments, and Akt, in cells (Congdon et al., 2012; Jinwal et al., 2009; Koren et al., 2010; Wang et al., 2010). MB also improves cognitive functions in mouse models of Alzheimer's disease (AD) (Congdon et al., 2012; O'Leary et al., 2010), and it has been explored in Phase IIb clinical trials in AD patients (Wischik et al., 2008). Although MB is a highly promiscuous compound, it has an enviable safety record and is used clinically for multiple indications (Schirmer et al., 2011). For these reasons, we decided to further explore the mechanism by which it might inactivate Hsp70s. Specifically, we hypothesized that MB might prevent ATP turnover in Hsp72 by oxidizing cysteine residues (Liu et al., 1996; Oz et al., 2009), because MB has been shown to oxidize sulfhydryls in other targets (Kelner and Alexander, 1985). Here, we report that Hsp72 is indeed oxidized by MB and that treatment with either MB or hydrogen peroxide irreversibly inactivates ATP turnover in vitro. Based on modeling studies and mutagenesis, this inhibition appears to be caused by oxidation-induced conformation changes in the NBD that block ATP binding. However, during the course of these studies, we also made the unexpected observation that the constitutive Hsp70 family member, Hsc70, is entirely resistant to irreversible inhibition by MB. Mass spectrometry and point mutants revealed that two reactive cysteines, C267S and C306S, are sufficient to distinguish the special sensitivity of Hsp72 to MB in vitro and in cells. Cys306 is highly conserved in the other stress-inducible Hsp70 isoforms, but it is absent from constitutive forms, supporting the idea that this residue is important in stress-related redox signaling. Thus, these finding suggest a major difference between closely related Hsp70 isoforms, which may be important in redox signaling and protecting cells from oxidative stress. Moreover, this work may provide insight into one mechanism by which MB reduces tau accumulation in cells, animals, and AD patients.

RESULTS

MB and H₂O₂ Irreversibly Inhibits Hsp72 but Not Hsc70

Because MB has been known to directly oxidize sulfhydryls (Kelner and Alexander, 1985), we hypothesized that the inhibition of Hsp70's ATPase activity by MB (Jinwal et al., 2009; Miyata et al., 2011) may involve oxidation of cysteines. To test this idea, Hsp72 was treated with MB (200 μ M) at 37°C for 1 hr and then dialyzed to remove any remaining compound. The remaining ATPase activity in the MB-treated Hsp72 sample was then measured using a malachite green assay (Chang et al., 2008). Because Hsp72 is a weak ATPase, the stimulatory cochaperone, DnaJ, was added to enhance nucleotide turnover in these experiments. Using this approach, we found that MBtreated Hsp72 had dramatically reduced ATPase activity (Figure 1A). Hsp72 was also inhibited by treatment with hydrogen peroxide (100 µM) under similar conditions, suggesting that the loss of activity may be due to oxidation (Figure 1A). Cysteine residues in proteins can be progressively oxidized to sulfenic, sulfinic and then sulfonic acids (Reddie and Carroll, 2008), but only sulfenic acids are readily reversible by glutathione or dithiothreitol (DTT) (Poole et al., 2004). The ATPase activity of MB-treated Hsp72 was only partially (~80%) recovered after exposure to DTT (1 mM) (Figure S1A available online), suggesting that MB oxidizes the protein to a mixture of oxidation states.

Initially as a control, the effects of MB on the ATPase activity of Hsc70 were evaluated. Unexpectedly, we found that MB had no effect on Hsc70 (Figure 1B). The chemical differences between these two well-conserved (85% identical and 94% similar) Hsp70 isoforms were then more closely examined. Sequence alignments of the human proteins showed that Hsp72 has five cysteine residues (three in the NBD and two in the SBD), whereas Hsc70 has only four (two each in the NBD and SBD). Thus, one difference between these isoforms is that Hsp72 has a unique Cys306, which is an asparagine in Hsc70 (Figure 1C). The sequences of a number of inducible or constitutive human Hsp70 family members were then examined, revealing that Cys306 is exclusively found in stress-inducible Hsp70s but not in any of the constitutively expressed family members (Figure S2A). Moreover, C267 is conserved in four of five inducible family members and only two of five constitutive forms (Figure S2A), suggesting that both of these residues may be important. Together, these results suggest that MB might selectively compromise ATPase by oxidizing at least one cysteine in Hsp72 but not Hsc70.

MB Oxidizes Cys306 of Hsp72

To map which cysteines in Hsp72 were oxidized by MB, a wellestablished mass spectrometry (MS) method using dimedone (5,5-dimethyl-1,3-cyclohexanedione) was used. Briefly, dimedone is known to react with oxidized cysteines to form stable thioethers that are readily observed in the MS spectra as a 138 Da shift in molecular mass (Figure 2A). Accordingly, MBtreated Hsp72 was treated with dimedone, and unreacted thiols were capped with carbamidomethyl groups. Subsequent trypsin digestions and analysis by liquid chromatography-tandem mass spectrometry (LC-MS/MS) revealed that the mass of the fragment containing Cys306 in Hsp72 was oxidized (Figure 2B). No other residues were modified, although it is important to note that the fragment containing Cys267 was not detected (this is likely because the nearby region is highly charged) (Figure 2C). As expected from the ATPase experiments, Hsc70 treated with MB was resistant to modification by dimedone (Figure 2C; Figure S2B). Thus, Hsp72 differs from Hsc70 in reactivity of its cysteines at positions 306 and perhaps 267.



Figure 1. The ATPase Activity of Hsp72, but Not Hsc70, Is Sensitive to Oxidation

(A) Purified Hsp72 (0.6μ M) was incubated with either MB, peroxide (H₂O₂), or a mock control and the remaining compound was removed by extensive dialysis. The stimulation of ATPase activity by the model J cochaperone, DnaJ, was then measured. Results are the average of three experiments performed in triplicate. Error bars, SEM.

(B) Human Hsc70 is resistant to oxidation. Experiments were performed as described for (A).

(C) The locations of cysteine residues (red) in Hsp72 are shown using the crystal structures of the NBD (Protein Data Bank [PDB] ID: 3JXU; shown in yellow) and the SBD (PDB ID: 1DKX; shown in green). In the insets, identical residues are shown in yellow, conserved residues are shown in gray, and the positions of cysteines are shown in red. Alignments were prepared in Vector NTI (Invitrogen). The chemical structure of MB is shown. See Figure S2 for the expanded alignment.

Hsp72 Cys to Ser Mutations Confer Resistance to MB

The MS studies suggested that MB-based oxidation of specific cysteines might be responsible for the compound's effects on Hsp72's ATPase activity. To test this idea, Cys306, Cys267, or both residues were mutated to serine by site-directed mutagenesis, and the resulting mutant proteins were purified. These substitutions did not change the global structure (Figure S3) or ATPase activity of the mutant chaperones (Figure 3A–3C). However, the ATPase activities of C267S (Figure 3A) and C306S (Figure 3B) were partially resistant to MB (50 μ M), and the double mutant (C267/306S) was completely resistant (Figure 3C). Thus, MB appeared to exert its inhibitory effect on Hsp72 ATPase activity via oxidation of these cysteines, and both residues appeared to be involved.

MB is known to reduce tau levels in cells through a mechanism dependent on Hsp72 (Jinwal et al., 2009). Thus, the overexpression of the C306S, C267S, and C267/306S mutants may desensitize cells to MB. In fact, when HeLaC3 cells were stably transfected with Hsp72 C306S, MB no longer reduced the levels of phosphorylated (pS396/404) or total tau (Figure 3D). Together, these results strongly suggest that MB acts on Hsp72 by

oxidizing Cys267 and Cys306 and that this activity is important in regulating the levels of tau.

C267D Mutation Causes a Conformational Change and Disrupts Nucleotide Binding

To gain some insight into the structural basis of these observation, the NBD of Hsp72 containing a C267D mutation was modeled using Robetta (Kim et al., 2004), with the structure of human Hsp72 NBD in the ADP form (3JXU) as a starting point (Wisniewska et al., 2010). An aspartic acid substitution was used because it sterically and electronically mimics a sulfinic acid (Permyakov et al., 2012), one of the possible oxidation states sampled by MB-treated cysteine residues. A comparison of the structures of the wild-type Hsp72 NBD and the C267D Hsp72 NBD showed that they were globally very similar, with an root-mean-square deviation (RMSD) of 1.55 Å and a template modeling score of 0.94 (Figure 4A). However, several residues that are specifically involved in nucleotide binding were predicted to be significantly shifted. For example, Gly339, which makes a hydrogen bond interaction with the alpha phosphate group of ADP, was shifted away from the nucleotide by 1.0 Å,

Α

Reactions of free thiol with iodoacetamide and oxidized cysteine with dimedone



residue	Hsp72		Hsc70		
	-	+	-	+	MB (200 µM)
C17	+57	+57	+57	+57	
C267	n.d.	n.d.	n.d .	n.d.	
C306	+57	+138	n.a.	n.a.	
C574	+57	+57	+57	+57	
C603	+57	+57	+57	+57	

Mass spectrometry results indicate oxidation of Hsp72 Cys306



Figure 2. Hsp72 Is Oxidized by MB at Specific Cysteine Residues

(A) Schematic of the specific reactions of iodoacetimide with free thiol and dimedone with sulfenic acid, producing a mass shift of 57 or 138 Da, respectively.
(B) Select region of the MS/MS spectra focused on the region including the C306 fragment (CSDLFR 306-311). This fragment is 797.39 Da in the mock-treated control (indicating iodoacetamide capping) and 878.43 Da in the MB-treated (indicating dimedone conjugation).

В

(C) Summary of the MS findings, showing that MB only oxidized C306 in Hsp72. n.d. = not detected. n.a = residue not conserved.

See Figure S1 for the DTT reversibility.

likely preventing the formation of this important bond (Figure 4B). Furthermore, Arg272, which interacts with the adenine ring by pi-stacking, is significantly pulled away by regional rotations in the backbone (Figure 4C). Collectively, these changes and others (Table S1) resulted in side-chain displacements totaling more than 25 Å in the nucleotide binding cleft. Very similar results were seen with the C306D mutant, suggesting that these residues might both contribute to conformational rearrangements (Figure S4A). It is interesting that oxidation of C306 is predicted to swivel residue C267 into the solvent-exposed cleft above nucleotide in the NBD (Figure S4B), perhaps making it more accessible to oxidation by MB. Together, these in silico observations suggest that oxidation of Cys267 and Cys306 in Hsp72 might damage nucleotide binding and inactivate ATP turnover. Furthermore, these results suggest that sequential oxidation of C306 and then C267 may reinforce and promote these conformational changes (Figure S4C).

Cys to Asp Mutations "Phenocopies" MB Treatment In Vitro and in Cells

To test this model, Cys267 and Cys306 of Hsp72 were mutated to the pseudo-oxidation residue, Asp, and the resulting mutant proteins were purified. Initial attempts to purify the mutants on ATP agarose immediately revealed that they had significantly reduced affinity for nucleotide (Figure 4D), consistent with the models. Switching to size exclusion, the mutants were purified and found to have normal circular dichroism

(CD) spectra (Figure S4A), suggesting that the global structure was not significantly disrupted by the Asp mutations. However, partial proteolysis showed that the mutants were more prone to digestion (Figure S4B), suggesting that they may be more flexible. It is interesting that MB-treated wild-type Hsp72 is also prone to proteolysis (Figure S4B), further enforcing the similarities between the Asp point mutants and the oxidized wild-type.

Next, the enzymatic activities of the Hsp72 pseudo-oxidation mutants were examined using ATPase assays, showing that the C267D and C306D mutants had dramatically decreased enzymatic activity (Figure 6A). These mutants essentially behaved like Hsp72 that had been treated with MB. We next tested the ability of Hsp72 and the mutants to refold denatured firefly luciferase. These experiments showed that Hsp72 treated with MB and the C267D, C306D, and C267/306D double mutants all had reduced refolding activity (Figure S4C). Taken together, these studies suggest that the Asp mutants phenocopy aspects of MB-treated Hsp72.

In HeLaC3 cells, overexpression of the C306D mutant produced modest (~40%) reductions in total tau levels and no significant effect on phosphorylated tau (Figure 5B). Overexpression of the double mutant (C267/306D) substantially (>70%) reduced both tau and phosphorylated tau levels (Figure 6B). These findings provide strong support for MB acting through oxidation of specific cysteine residues in Hsp72 to reduce tau levels.



Figure 3. Serine Mutants of Hsp72 Are Resistant to MB Treatment in ATPase and Cell-Based Assays

(A) C267S mutation confers partial resistance to MB (50 μ M).

(B) C306S confers partial resistance to MB (50 µM).

(C) C267, 306S double mutation confers resistance to MB (50 µM). All of the ATPase experiments were performed at least twice in triplicate and error bars are SEM.

(D) Overexpression of the C306S mutation blocks MB-mediated clearance of tau. HeLaC3 cells were transfected with vector, Flag-tagged WT Hsp72, or C306S Hsp72 mutant for 48 hr and then treated with MB for 10 min. Samples were analyzed by western blot and the results are representative of experiments performed in duplicate.

See Figure S3 for the CD spectra of the mutants.

DISCUSSION

One "arm" of the cellular response to redox stress likely involves the acute protection of proteins from oxidation-induced unfolding and aggregation (Ilbert et al., 2007). Here, we were specifically interested in understanding how the Hsp70 family of chaperones, especially the stress-inducible forms, might be linked to these types of redox responses. We found that Hsp72, but not Hsc70, was sensitive to oxidation by either MB or peroxide. A recent large-scale proteomic study positively identified Hsp72 as sensitive to oxidation by peroxide in HeLa cells (Leonard et al., 2009), consistent with this finding. Our mass spectrometry, modeling, and point mutagenesis results suggest that oxidation of Hsp72 occurs selectively at two cysteine residues, Cys267 and Cys306. It is important to note that Cys306 is uniquely conserved in the stress-inducible Hsp70 family members and is absent from the constitutive ones (see Figure S2). Although it might be initially surprising for such highly homologous proteins to have different biochemical properties, Goldfarb et al. recently demonstrated that Hsp72 and Hsc70 have opposing effects on the surface expression of murine epithelial sodium channel (Goldfarb et al., 2006), further suggesting that even highly conserved Hsp70 isoforms can sometimes have distinct functions. Together, our findings suggest that Hsp72, and possibly other stress-inducible Hsp70 family members, are specially adapted for sensing and responding to redox stress. Based on the observed effects on tau stability, this redox response may involve switching the triage decision to favor degradation of misfolded Hsp72 substrates, perhaps clearing the cytosol of folding intermediates that are particularly prone to oxidative damage.

What is the molecular mechanism linking oxidation of Hsp72 to a loss of ATPase activity? We propose a model in which oxidation of the unique and solvent exposed Cys306 leads to rearrangement of Cys267, such that this residue is now also sensitive to oxidation (see Figure S4). Thus, although Hsc70 has Cys267, it lacks the critical initiator (i.e., residue Cys306); therefore, treatment with either MB or peroxide does not inactivate it (see Figure 1). This model further suggests that oxidation of both Cys267 and Cys306 causes numerous, subtle rearrangements in residues that normally contact nucleotide (Figure 4; Figure S4). These changes destabilize a number of key contacts, including hydrogen bonds with ATP, hydrogen bonds with the phosphate, and a number of hydrophobic interactions (Table S1). Consistent with this idea, the purified mutants, C267D and C306D, lacked the ability to bind ATP in vitro

Chemistry & Biology Hsp72 Is a Redox Sensitive Chaperone



Figure 4. Modeling of Hsp72 C267D Reveals Structural Changes in Residues that Contact Nucleotide

(A) Overall alignment of the NBDs of Hsp72 (yellow: PDB ID 3JXU) and Hsp72 C267D Model (red: modeled from template 3JXU using Robetta Server).

(B and C) In the C267D model, (B) Gly339 is shifted away from the α -phosphate of nucleotide and (C) Arg272 is shifted away from the adenine ring. In total, C267D caused structural changes totaling 25 Å (see Table S1). Similar results were seen in the C306D mutant (see Figure S4).

(D) Purified C267D and C306D mutants do not bind nucleotide. Purified mutants and wild-type Hsp72 (5 μ M) were treated with 1 ml of ATP-agarose and washed with 3 ml of Buffer A (25 mM HEPES, 10 mM KCl, 5 mM MgCl2, pH 7.5), and flowthrough (FT) was collected. Following three additional washes, the remaining protein was collected in the eluent (E) by washing with 3 ml of Buffer A containing 3 mM ATP. Fractions were analyzed on 1%–20% Tris-Tricine gels using a polyclonal anti-Hsp72 antibody (Enzo). See Table S1 for details.

oxidation is one important way by which MB reduces tau accumulation in AD models. These findings may aid in the discovery of additional AD therapeutics that take advantage of this underexplored

Treatments with high levels of MB are tolerated in cells (Yaglom et al., 2007); moreover, it is relatively nontoxic in humans (Schirmer et al., 2011). On first

(Figure 4). Thus, we propose a model in which a cascade of oxidations severely damages the binding of Hsp72 to nucleotide. How this rearrangement favors degradation of bound substrates, such as tau, is not yet clear.

Another goal of this work was to better understand the mechanism by which MB reduces tau levels in cellular and mouse models of AD (Jinwal et al., 2009; O'Leary et al., 2010). Although MB is clearly a promiscuous compound that lacks a classic "drug-like" profile, it is remarkably nontoxic; thus, it is used in humans for the treatment of a variety of indications, including inherited and acute methemoglobinemia, prevention of urinary tract infections, ifosfamid-induced neurotoxicity, vasoplegic adrenaline-resistant shock, and pediatric malaria (Schirmer et al., 2011). Because of MB's particular promise as an AD therapeutic, we were interested in understanding whether any of its effects on tau accumulation may be mediated by oxidation of Hsp72. In this work, we found that mutating Hsp72 Cys267 and/or Cys306 to serine blocked the ability of MB to reduce tau levels in cells (Figure 3D). This is an important finding because it strongly links Hsp72 oxidation to effects on tau accumulation. To further enforce this idea, overexpression of the corresponding C267/306D pseudo-oxidation double mutant was a strong "dominant negative," and it dramatically reduced tau levels (Figure 5). Together, these results suggest that Hsp72 glance, the global inactivation of Hsp70s might be hypothesized to be acutely and dramatically toxic, given the proposed roles of this chaperone family in "housekeeping" activities. However, another point of view is that the stress-inducible Hsp72 isoform is specifically concerned with degrading only the substrates that have been damaged or misfolded in response to stress (Pratt and Toft, 2003), protecting the cytosol from the accumulation of proteotoxic intermediates. In normal cells, the levels of Hsp72 and such misfolded substrates may be low due to the action of other components of the protein quality controls system. Thus, MB may have low toxicity because of its unusual selectivity for the stress-inducible Hsp70 family members. This is an important finding in the continued search for Hsp70-modifying compounds with low toxicity and favorable therapeutic benefits. Together, these findings provide one potential mechanism by which MB has activity in tau-related diseases, such as AD.

mechanism.

SIGNIFICANCE

The heat shock protein 70 (Hsp70) family of molecular chaperones plays a central role in proteostasis, especially in the response to stress. These proteins have been implicated in regulating the stability and turnover of a number of important substrates, including the microtubule-associated





HE ATP SOH ME ATP Hsp72 NBD Hsp72 NBD

protein tau. Tau aberrantly accumulates in 15 neurodegenerative disorders, including Alzheimer's disease (AD). Accordingly, there is great interest in finding ways of using the Hsp70s to restore balance in this system and reduce tau accumulation. We found that the stress-inducible Hsp70, Hsp72, is sensitive to oxidation by peroxide or methylene blue (MB), while the constitutive Hsc70 is resistant. Mutation of key cysteines in Hsp72 to serine rendered the chaperone resistant to MB in vitro, and overexpression of this mutant partially suppressed MB-mediated degradation of tau in cells. Together, these results suggest that oxidation of Hsp72 is one mechanism by which MB reduces tau levels. These results are particularly interesting because this compound has recently undergone clinical evaluation as a taureducing therapy in AD. These results suggest that oxidation of Hsp72 might be one mechanism to explain the activity of this clinical candidate.

EXPERIMENTAL PROCEDURES

Hsp72 NBD

Proteins and Reagents

Unless otherwise specified, reagents were purchased from Sigma (St. Louis, MO, USA) or Fisher Scientific (Pittsburgh, PA, USA). Human Hsp72, Hsc70, and *Escherichia coli* DnaJ were purified according to published schemes (Chang et al., 2010). Site-directed mutagenesis primers were designed based

Figure 5. Homology Model of Hsp72 C306D NBD

(A) Pseudo-oxidation of residue 306 does not produce global changes in the NBD fold, similar to what was seen with the C267D mutant (see Figure 4). C306D is represented in green, and wild-type (PDB 3JXU) is represented in yellow. The C306D and C267D models are nearly identical (see Figure 4), with C306D also causing an \sim 25 Å total displacement of residues associated with nucleotide binding.

(B) Close-up that illustrates how C306D increases the solvent exposure of Cys267, potentially enhancing its oxidation. C267 moves by ${\sim}3$ Å in the C306D model.

(C) Model for initial oxidation at Cys306, leading to synergistic oxidation of C267 and reduced ATP binding. No specific oxidation state of the thiol is intended and the results suggest that a series of oxidations are possible.

on previous reports (Zheng et al., 2004), and mutagenesis of Hsp72 was carried out in accordance with the the user manual for the QuickChange site-directed mutagenesis kit (Stratagene, La Jolla, CA, USA). The Hsp72 C267S, C306S, and C267/306S mutants were expressed and purified using the same protocol as Hsp72 (Chang et al., 2010). The Hsp72 C267D and C306D mutants were expressed as previously described (Chang et al., 2010) and purified using nickel-nitrilotriacetic acid His-Bind resin (Novagen, Darmstadt, Germany), then buffer exchanged into a 25 mM HEPES buffer (10 mM KCl, 5 mM MgCl₂, pH 7.5).

Oxidation of Hsp72/Hsc70

Protein (10 μ M) and MB (5 mM) were incubated at 37°C for 1 hr. For H₂O₂ oxidation, protein sample

(10 μ M) was incubated with 1 mM H₂O₂ at 37°C for 1 hr. Treated protein samples were subsequently dialyzed against buffer A (100 mM Tris-HCl, pH 7.4, 20 mM KCl, 6 mM MgCl₂) at 4°C.

ATPase Activity

ATPase activity was measured according to the previously published method (Chang et al., 2008). Briefly, malachite green-based assays were used to measure phosphate release from purified Hsp72, Hsc70, or mutants (1 μ M). Reactions were initiated with 1 mM ATP, performed for 60 min, and quenched before measuring absorbance. Absorbance readings were converted to picomoles of ATP using a phosphate standard curve.

Preparation of Dimedone-Modified Hsp70s

MB-treated or untreated Hsp70s (10 μ M) were incubated with 5 mM dimedone (5,5-dimethyl-1,3-cyclohexanedione) in buffer A (100 mM Tris-HCl, pH 7.4, 20 mM KCl, 6 mM MgCl₂) at room temperature for 1 hr. The samples were analyzed by SDS-PAGE and stained with colloidal Coomassie blue (Invitrogen, Carlsbad, CA). Bands corresponding to Hsp72 were excised and stored at -20° C until use.

Mass Spectrometry

In-gel digestion was performed as previously described (Brady et al., 2010). After reduction (10 mM DTT) and alklylation (65 mM iodoacetamide) of the free cysteines at room temperature for 30 min, proteins were digested overnight with trypsin (Promega). Resulting peptides were resolved on a nanocapillary reverse-phase column (Picofritcolumn, New Objective) using a 1% acetic acid/acetonitrile gradient at 300 nl/min and subjected to LC-MS/MS using

Chemistry & Biology Hsp72 Is a Redox Sensitive Chaperone

B Α Hsp72 pseudo-oxidation mutants Pseudo-oxidation mutants reduce tau levels in cells have low ATPase activity Hsp72 3060 wt wt+MB C267D C306D total tau Wild type C261. p-tau (normalized) 25 omol ATP/ug 72/min рТан intensity (Total Tau .i 0.25 Hsp7 C261,306D C306D Wild WPE DnaJ (µM) GAPDH

Figure 6. Pseudo-Oxidation Mutants Phenocopy MB Treatment

(A) The DnaJ-stimulated ATPase activities of purified Hsp72 mutants were reduced, resembling MB-treated wild-type. These experiments were performed at least three times using two independently prepared samples. Error bars, SEM.

(B) Overexpression of wild-type Hsp72 had little effect on tau levels, but the C306D and C267/306D mutants reduced tau. The gels show two independent replicates, and the quantification of band intensities includes SEM.

See Figure S4 for the characterization of the mutants.

a LTQ Orbitrap XL mass spectrometer. MS/MS spectra were searched against the database considering either carbamidomethyl- or dimedone-modified cysteine

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, four figures, and one table and can be found with this article online at http:// dx.doi.org/10.1016/j.chembiol.2012.07.026.

ACKNOWLEDGMENTS

The authors thank S. Leonard for helpful conversations. This work was supported by an Alzheimer's Association grant and National Institutes of Health (NIH) Grants R00AG031291 and R01NS073899 to C.A.D. In addition. A.D.T. was supported by GM007863, S.S. was supported by GM008353, and this work was further supported by NIH Grant R01NS059690 to J.E.G.

Received: February 27, 2012 Revised: June 14, 2012 Accepted: July 6, 2012 Published: November 21, 2012

REFERENCES

Apel, K., and Hirt, H. (2004). Reactive oxygen species: metabolism, oxidative stress, and signal transduction. Annu. Rev. Plant Biol. 55, 373-399.

Brady, G.F., Galbán, S., Liu, X., Basrur, V., Gitlin, J.D., Elenitoba-Johnson, K.S., Wilson, T.E., and Duckett, C.S. (2010). Regulation of the copper chaperone CCS by XIAP-mediated ubiquitination. Mol. Cell. Biol. 30, 1923–1936.

Chang, L., Bertelsen, E.B., Wisén, S., Larsen, E.M., Zuiderweg, E.R., and Gestwicki, J.E. (2008). High-throughput screen for small molecules that modulate the ATPase activity of the molecular chaperone DnaK. Anal. Biochem. 372, 167-176

Chang, L., Thompson, A.D., Ung, P., Carlson, H.A., and Gestwicki, J.E. (2010). Mutagenesis reveals the complex relationships between ATPase rate and the chaperone activities of Escherichia coli heat shock protein 70 (Hsp70/DnaK). J. Biol. Chem. 285, 21282-21291.

Congdon, E., Wu, J., Myeku, N., Figueroa, Y., Herman, M., Marinec, P., Gestwicki, J., Dickey, C., Yu, W., and Duff, K. (2012). Methylthioninium chloride (methylene blue) induces autophagy and attenuates tauopathy in vitro and in vivo. Autophagy 8, 609-622.

Evans, C.G., Chang, L., and Gestwicki, J.E. (2010). Heat shock protein 70 (hsp70) as an emerging drug target. J. Med. Chem. 53, 4585-4602.

Forman, H.J., Majorino, M., and Ursini, F. (2010). Signaling functions of reactive oxygen species. Biochemistry 49, 835-842.

Giles, N.M., Watts, A.B., Giles, G.I., Fry, F.H., Littlechild, J.A., and Jacob, C. (2003). Metal and redox modulation of cysteine protein function. Chem. Biol. 10.677-693

Goldfarb, S.B., Kashlan, O.B., Watkins, J.N., Suaud, L., Yan, W., Kleyman, T.R., and Rubenstein, R.C. (2006). Differential effects of Hsc70 and Hsp70 on the intracellular trafficking and functional expression of epithelial sodium channels. Proc. Natl. Acad. Sci. USA 103, 5817-5822.

Hageman, J., van Waarde, M.A., Zylicz, A., Walerych, D., and Kampinga, H.H. (2011). The diverse members of the mammalian HSP70 machine show distinct chaperone-like activities. Biochem. J. 435, 127-142.

Hartl, F.U., Bracher, A., and Hayer-Hartl, M. (2011). Molecular chaperones in protein folding and proteostasis. Nature 475. 324-332.

Ilbert, M., Horst, J., Ahrens, S., Winter, J., Graf, P.C., Lilie, H., and Jakob, U. (2007). The redox-switch domain of Hsp33 functions as dual stress sensor. Nat. Struct. Mol. Biol. 14. 556-563.

Jinwal, U.K., Miyata, Y., Koren, J., 3rd, Jones, J.R., Trotter, J.H., Chang, L., O'Leary, J., Morgan, D., Lee, D.C., Shults, C.L., et al. (2009). Chemical manipulation of hsp70 ATPase activity regulates tau stability. J. Neurosci. 29, 12079-12088

Kelner, M.J., and Alexander, N.M. (1985). Methylene blue directly oxidizes glutathione without the intermediate formation of hydrogen peroxide. J. Biol. Chem. 260, 15168-15171.

Kim, D.E., Chivian, D., and Baker, D. (2004). Protein structure prediction and analysis using the Robetta server. Nucleic Acids Res. 32(Web Server issue), W526-W531.

Koren, J., 3rd, Jinwal, U.K., Jin, Y., O'Leary, J., Jones, J.R., Johnson, A.G., Blair, L.J., Abisambra, J.F., Chang, L., Miyata, Y., et al. (2010). Facilitating Akt clearance via manipulation of Hsp70 activity and levels. J. Biol. Chem. 285, 2498-2505,

Leonard, S.E., Reddie, K.G., and Carroll, K.S. (2009). Mining the thiol proteome for sulfenic acid modifications reveals new targets for oxidation in cells. ACS Chem. Biol. *4*, 783–799.

Lin, M.T., and Beal, M.F. (2006). Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. Nature *443*, 787–795.

Lindquist, S. (1986). The heat-shock response. Annu. Rev. Biochem. 55, 1151-1191.

Liu, Q., Levy, E.J., and Chirico, W.J. (1996). N-Ethylmaleimide inactivates a nucleotide-free Hsp70 molecular chaperone. J. Biol. Chem. 271, 29937–29944.

Majmundar, A.J., Wong, W.J., and Simon, M.C. (2010). Hypoxia-inducible factors and the response to hypoxic stress. Mol. Cell *40*, 294–309.

Mayer, M.P., and Bukau, B. (2005). Hsp70 chaperones: cellular functions and molecular mechanism. Cell. Mol. Life Sci. 62, 670–684.

Mayer, M.P., Schröder, H., Rüdiger, S., Paal, K., Laufen, T., and Bukau, B. (2000). Multistep mechanism of substrate binding determines chaperone activity of Hsp70. Nat. Struct. Biol. *7*, 586–593.

Miyata, Y., Koren, J., Kiray, J., Dickey, C.A., and Gestwicki, J.E. (2011). Molecular chaperones and regulation of tau quality control: strategies for drug discovery in tauopathies. Future Med Chem. *3*, 1523–1537.

Morgan, M.J., and Liu, Z.G. (2011). Crosstalk of reactive oxygen species and NF-kB signaling. Cell Res. *21*, 103–115.

O'Leary, J.C., 3rd, Li, Q., Marinec, P., Blair, L.J., Congdon, E.E., Johnson, A.G., Jinwal, U.K., Koren, J., 3rd, Jones, J.R., Kraft, C., et al. (2010). Phenothiazine-mediated rescue of cognition in tau transgenic mice requires neuroprotection and reduced soluble tau burden. Mol. Neurodegener. 5, 45.

Oz, M., Lorke, D.E., and Petroianu, G.A. (2009). Methylene blue and Alzheimer's disease. Biochem. Pharmacol. 78, 927–932.

Patten, D.A., Germain, M., Kelly, M.A., and Slack, R.S. (2010). Reactive oxygen species: stuck in the middle of neurodegeneration. J. Alzheimers Dis. 20 (*Suppl 2*), S357–S367.

Patury, S., Miyata, Y., and Gestwicki, J.E. (2009). Pharmacological targeting of the Hsp70 chaperone. Curr. Top. Med. Chem. 9, 1337–1351.

Paulsen, C.E., and Carroll, K.S. (2010). Orchestrating redox signaling networks through regulatory cysteine switches. ACS Chem. Biol. *5*, 47–62.

Permyakov, S.E., Zernii, E.Y., Knyazeva, E.L., Denesyuk, A.I., Nazipova, A.A., Kolpakova, T.V., Zinchenko, D.V., Philippov, P.P., Permyakov, E.A., and Senin, I.I. (2012). Oxidation mimicking substitution of conservative cysteine in recoverin suppresses its membrane association. Amino Acids *42*, 1435–1442.

Poole, L.B., Karplus, P.A., and Claiborne, A. (2004). Protein sulfenic acids in redox signaling. Annu. Rev. Pharmacol. Toxicol. *44*, 325–347.

Pratt, W.B., and Toft, D.O. (2003). Regulation of signaling protein function and trafficking by the hsp90/hsp70-based chaperone machinery. Exp. Biol. Med. (Maywood) 228, 111–133.

Reddie, K.G., and Carroll, K.S. (2008). Expanding the functional diversity of proteins through cysteine oxidation. Curr. Opin. Chem. Biol. *12*, 746–754.

Schirmer, R.H., Adler, H., Pickhardt, M., and Mandelkow, E. (2011). "Lest we forget you-methylene blue..." Neurobiol Aging *32*, 2325.e7–2325.e16. . Published online February 12, 2011. PMID: 22361619.

Vignols, F., Mouaheb, N., Thomas, D., and Meyer, Y. (2003). Redox control of Hsp70-Co-chaperone interaction revealed by expression of a thioredoxin-like Arabidopsis protein. J. Biol. Chem. 278, 4516–4523.

Wang, A.M., Morishima, Y., Clapp, K.M., Peng, H.M., Pratt, W.B., Gestwicki, J.E., Osawa, Y., and Lieberman, A.P. (2010). Inhibition of hsp70 by methylene blue affects signaling protein function and ubiquitination and modulates polyglutamine protein degradation. J. Biol. Chem. *285*, 15714–15723.

Waris, G., and Ahsan, H. (2006). Reactive oxygen species: role in the development of cancer and various chronic conditions. J. Carcinog. *5*, 14.

Wischik, C., Betham, P., Wischik, D., and Seng, K. (2008). Tau aggregation inhibitor (TAI) therapy with rember[™] arrests disease progression in mild and moderate Alzheimer's disease over 50 weeks. Alzheimer's & Dementia *4* (*Suppl.*), T167.

Wisniewska, M., Karlberg, T., Lehtiö, L., Johansson, I., Kotenyova, T., Moche, M., and Schüler, H. (2010). Crystal structures of the ATPase domains of four human Hsp70 isoforms: HSPA1L/Hsp70-hom, HSPA2/Hsp70-2, HSPA6/Hsp70B', and HSPA5/BiP/GRP78. PLoS ONE 5, e8625.

Yaglom, J.A., Gabai, V.L., and Sherman, M.Y. (2007). High levels of heat shock protein Hsp72 in cancer cells suppress default senescence pathways. Cancer Res. 67, 2373–2381.

Zhang, Y., Ahn, Y.H., Benjamin, I.J., Honda, T., Hicks, R.J., Calabrese, V., Cole, P.A., and Dinkova-Kostova, A.T. (2011). HSF1-dependent upregulation of Hsp70 by sulfhydryl-reactive inducers of the KEAP1/NRF2/ARE pathway. Chem. Biol. *18*, 1355–1361.

Zheng, L., Baumann, U., and Reymond, J.L. (2004). An efficient one-step site-directed and site-saturation mutagenesis protocol. Nucleic Acids Res. *32*, e115.