



Original article

Spectral estimates of the p -Laplace Neumann operator in conformal regular domains

V. Gol'dshtein*, A. Ukhlov

Department of Mathematics, Ben-Gurion University of the Negev, P. O. Box 653, Beer Sheva, 84105, Israel

Available online 9 April 2016

Abstract

In this paper we study spectral estimates of the p -Laplace Neumann operator in conformal regular domains $\Omega \subset \mathbb{R}^2$. This study is based on (weighted) Poincaré–Sobolev inequalities. The main technical tool is the theory of composition operators in relation with the Brennan's conjecture. We prove that if the Brennan's conjecture holds for any $p \in (4/3, 2)$ and $r \in (1, p/(2-p))$ then the weighted (r, p) -Poincaré–Sobolev inequality holds with the constant depending on the conformal geometry of Ω . As a consequence we obtain classical Poincaré–Sobolev inequalities and spectral estimates for the first nontrivial eigenvalue of the p -Laplace Neumann operator for conformal regular domains.

© 2016 Ivane Javakhishvili Tbilisi State University. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Conformal mappings; Sobolev spaces; Elliptic equations

1. Introduction and methodology

Let $\Omega \subset \mathbb{R}^2$ be a simply connected planar domain with a smooth boundary $\partial\Omega$. We consider the Neumann eigenvalue problem for the p -Laplace operator ($1 < p < 2$):

$$\begin{cases} -\operatorname{div}(|\nabla u|^{p-2}\nabla u) = \mu_p |u|^{p-2}u & \text{in } \Omega \\ \frac{\partial u}{\partial n} = 0 & \text{on } \partial\Omega. \end{cases} \quad (1.1)$$

The weak statement of this spectral problem is as follows: a function u solves the previous problem if and only if $u \in W^{1,p}(\Omega)$ and

$$\iint_{\Omega} (|\nabla u(x, y)|^{p-2}\nabla u(x, y)) \cdot \nabla v(x, y) \, dx dy = \mu_p \iint_{\Omega} |u|^{p-2}u(x, y)v(x, y) \, dx dy$$

for all $v \in W^{1,p}(\Omega)$.

* Corresponding author. Tel.: +972 86461620; fax: +972 86477648.

E-mail addresses: vladimir@bgu.ac.il (V. Gol'dshtein), ukhlov@math.bgu.ac.il (A. Ukhlov).

Peer review under responsibility of Journal Transactions of A. Razmadze Mathematical Institute.

The first nontrivial Neumann eigenvalue μ_p can be characterized as

$$\mu_p(\Omega) = \min \left\{ \frac{\iint_{\Omega} |\nabla u(x, y)|^p dx dy}{\iint_{\Omega} |u(x, y)|^p dx dy} : u \in W^{1,p}(\Omega) \setminus \{0\}, \iint_{\Omega} |u|^{p-2} u dx dy = 0 \right\}.$$

Moreover, $\mu_p(\Omega)^{-\frac{1}{p}}$ is the best constant $B_{p,p}(\Omega)$ (see, for example, [1,2]) in the following Poincaré–Sobolev inequality

$$\inf_{c \in \mathbb{R}} \|f - c\|_{L^p(\Omega)} \leq B_{p,p}(\Omega) \|\nabla f\|_{L^p(\Omega)}, \quad f \in W^{1,p}(\Omega).$$

We prove, that $\mu_p(\Omega)$ depends on the conformal geometry of Ω and can be estimated in terms of Sobolev norms of a conformal mapping of the unit disc \mathbb{D} onto Ω (Theorem A).

The main technical tool is existence of universal weighted Poincaré–Sobolev inequalities

$$\begin{aligned} & \inf_{c \in \mathbb{R}} \left(\iint_{\Omega} |f(x, y) - c|^r h(x, y) dx dy \right)^{\frac{1}{r}} \\ & \leq B_{r,p}(\Omega, h) \left(\iint_{\Omega} |\nabla f(x, y)|^p dx dy \right)^{\frac{1}{p}}, \quad f \in W^{1,p}(\Omega), \end{aligned} \quad (1.2)$$

in any simply connected planar domain $\Omega \neq \mathbb{R}^2$ for conformal weights $h(x, y) := J_{\varphi}(x, y) = |\varphi'(x, y)|^2$ induced by conformal homeomorphisms $\varphi : \Omega \rightarrow \mathbb{D}$.

Main results of this article can be divided onto two parts. The first part is the technical one and concerns weighted Poincaré–Sobolev inequalities in arbitrary simply connected planar domains with nonempty boundaries (Theorem C and its consequences). Results of the first part will be used for (non weighted) Poincaré–Sobolev inequalities in so-called conformal regular domains (Theorem B) that lead to lower estimates for the first nontrivial eigenvalue μ_p (Theorem A). To the best of our knowledge lower estimates were known before for convex domains only. The class of conformal regular domains is much larger. It includes, for example, bounded domains with Lipschitz boundaries and quasidisks, i.e images of discs under quasiconformal homeomorphisms of whole plane.

Brennan's conjecture [3] is that for a conformal mapping $\varphi : \Omega \rightarrow \mathbb{D}$

$$\iint_{\Omega} |\varphi'(x, y)|^{\beta} dx dy < +\infty, \quad \text{for all } \frac{4}{3} < \beta < 4. \quad (1.3)$$

For the inverse conformal mapping $\psi = \varphi^{-1} : \mathbb{D} \rightarrow \Omega$ Brennan's conjecture [3] states

$$\iint_{\mathbb{D}} |\psi'(u, v)|^{\alpha} dudv < +\infty, \quad \text{for all } -2 < \alpha < \frac{2}{3}. \quad (1.4)$$

A connection between Brennan's Conjecture and composition operators on Sobolev spaces was established in [4]:

Equivalence Theorem. *Brennan's Conjecture (1.3) holds for a number $\beta \in (4/3; 4)$ if and only if a conformal mapping $\varphi : \Omega \rightarrow \mathbb{D}$ induces a bounded composition operator*

$$\varphi^* : L^{1,p}(\mathbb{D}) \rightarrow L^{1,q(p,\beta)}(\Omega)$$

for any $p \in (2; +\infty)$ and $q(p, \beta) = p\beta/(p + \beta - 2)$.

The inverse Brennan's Conjecture states that for any conformal mapping $\psi : \mathbb{D} \rightarrow \Omega$, the derivative ψ' belongs to the Lebesgue space $L^{\alpha}(\mathbb{D})$, for $-2 < \alpha < 2/3$. The integrability of the derivative in the power greater than $2/3$ requires some restrictions on the geometry of Ω . If $\Omega \subset \mathbb{R}^2$ is a simply connected planar domain of finite area, then

$$\iint_{\mathbb{D}} |\psi'(u, v)|^2 dudv = \iint_{\mathbb{D}} J_{\psi}(u, v) dudv = |\Omega| < \infty.$$

Integrability of the derivative in the power $\alpha > 2$ is impossible without additional assumptions on the geometry of Ω . For example, for any $\alpha > 2$ the domain Ω necessarily has a finite geodesic diameter [5].

Let $\Omega \subset \mathbb{R}^2$ be a simply connected planar domain. Then Ω is called a conformal α -regular domain if there exists a conformal mapping $\varphi : \Omega \rightarrow \mathbb{D}$ such that

$$\iint_{\mathbb{D}} \left| (\varphi^{-1})'(u, v) \right|^\alpha dudv < \infty \quad \text{for some } \alpha > 2. \tag{1.5}$$

If Ω is a conformal α -regular domain for some $\alpha > 2$ we call Ω a conformal regular domain.

The property of α -regularity does not depend on choice of a conformal mapping φ and depends on the hyperbolic geometry of Ω only [6]. For connection between conformal mapping and hyperbolic geometry see, for example, [7].

Note that a boundary $\partial\Omega$ of a conformal regular domain can have any Hausdorff dimension between one and two, but cannot be equal two [8].

The next theorem gives lower estimates of the first nontrivial p -Laplace Neumann eigenvalue:

Theorem A. Let $\varphi : \Omega \rightarrow \mathbb{D}$ be a conformal homeomorphism from a conformal α -regular domain Ω to the unit disc \mathbb{D} and Brennan’s Conjecture holds. Then for every $p \in (\max\{4/3, (\alpha + 2)/\alpha\}, 2)$ the following inequality holds

$$\frac{1}{\mu_p(\Omega)} \leq \inf_{q \in [1, 2p/(4-p))} \left\{ \|(\varphi^{-1})'\|_{L^\alpha(\mathbb{D})}^2 \left(\iint_{\mathbb{D}} \left| (\varphi^{-1})' \right|^{\frac{(p-2)q}{p-q}} dudv \right)^{\frac{p-q}{q}} \cdot B_{\frac{\alpha p}{\alpha-2}, q}^p(\mathbb{D}) \right\}.$$

Here $B_{r,q}(\mathbb{D})$ is the best constant in the corresponding (r, q) -Poincaré–Sobolev inequality in the unit disc \mathbb{D} for $r = \alpha p / (\alpha - 2)$.

In the limit case $\alpha = \infty$ and $p = q$ we have

Corollary A. Suppose that Ω is a smooth bounded Jordan domain with a boundary $\partial\Omega$ of a class C^1 with a Dini continuous normal. Let $\varphi : \Omega \rightarrow \mathbb{D}$ be a conformal homeomorphism from Ω onto the unit disc \mathbb{D} . Then for every $p \in (1, 2)$ the following inequality holds

$$\frac{1}{\mu_p(\Omega)} \leq \|(\varphi^{-1})'\|_{L^\infty(\mathbb{D})}^p \frac{1}{\mu_p(\mathbb{D})}.$$

Remark 1.1. The constant $B_{r,q}(\mathbb{D})$ satisfies [9,10]:

$$B_{r,q}(\mathbb{D}) \leq \frac{2}{\pi^\delta} \left(\frac{1 - \delta}{1/2 - \delta} \right)^{1-\delta}, \quad \delta = 1/q - 1/r.$$

Remark 1.2. The Brennan’s conjecture was proved for $\alpha \in [\alpha_0, 2/3)$ when $\alpha_0 = -1.752$ [11].

In the Introduction we formulate main results under the assumptions that the Brennan’s conjecture holds true for all $-2 < \alpha < 2/3$. In the main part of the paper we prove main results for $\alpha_0 = -1.752 < \alpha < 2/3$ that was proved recently.

This difference is related to our belief that the Brennan’s conjecture is correct.

Remark 1.3. The estimates for the $\mu_p(\Omega)$ were known before only for convex domains. For example, in [12] it was proved that

$$\mu_p(\Omega) \geq \left(\frac{\pi_p}{d(\Omega)} \right)^p$$

where

$$\pi_p = 2 \int_0^{(p-1)^{\frac{1}{p}}} \frac{dt}{(1 - t^p/(p-1))^{\frac{1}{p}}} = 2\pi \frac{(p-1)^{\frac{1}{p}}}{p(\sin(\pi/p))}.$$

Theorem A has a direct connection with the spectral stability problem for the p -Laplace operator. See, the recent papers, [13–16], where one can find the history of the problem, main results in this area and appropriate references.

Theorem A is a corollary (after simple calculations) of the following version of the Poincaré–Sobolev inequality.

Theorem B. Suppose that $\Omega \subset \mathbb{R}^2$ is a conformal α -regular domain and Brennan’s Conjecture holds. Then for every $p \in (\max\{4/3, \alpha/(\alpha - 1)\}, 2)$, every $s \in (1, \frac{\alpha-2}{\alpha} \frac{p}{2-p})$ and every function $f \in W^{1,p}(\Omega)$, the inequality

$$\inf_{c \in \mathbb{R}} \left(\iint_{\Omega} |f(x, y) - c|^s dx dy \right)^{\frac{1}{s}} \leq B_{s,p}(\Omega) \left(\iint_{\Omega} |\nabla f(x, y)|^p dx dy \right)^{\frac{1}{p}} \tag{1.6}$$

holds with the constant

$$B_{s,p}(\Omega) \leq \|(\varphi^{-1})' |L^\alpha(\mathbb{D})\|^{\frac{2}{s}} B_{r,p}(\Omega, h) \leq \inf_{q \in [1, 2p/(4-p))} \left\{ B_{\frac{\alpha s}{\alpha-2}, q}(\mathbb{D}) \cdot \|(\varphi^{-1})' |L^\alpha(\mathbb{D})\|^{\frac{2}{s}} K_{p,q}(\mathbb{D}) \right\}.$$

Here $B_{r,p}(\Omega, h)$, $r = \alpha s/(\alpha - 2)$, is the best constant of the following weighted Poincaré–Sobolev inequality:

Theorem C. Suppose $\Omega \subset \mathbb{R}^2$ is a simply connected domain with non empty boundary, Brennan’s Conjecture holds and $h(z) = J(z, \varphi)$ is the conformal weight defined by a conformal homeomorphism $\varphi : \Omega \rightarrow \mathbb{D}$. Then for every $p \in (4/3, 2)$ and every function $f \in W^{1,p}(\Omega)$, the inequality

$$\inf_{c \in \mathbb{R}} \left(\iint_{\Omega} |f(x, y) - c|^r h(x, y) dx dy \right)^{\frac{1}{r}} \leq B_{r,p}(\Omega, h) \left(\iint_{\Omega} |\nabla f(x, y)|^p dx dy \right)^{\frac{1}{p}} \tag{1.7}$$

holds for any $r \in [1, p/(2 - p))$ with the constant

$$B_{r,p}(\Omega, h) \leq \inf_{q \in [1, 2p/(4-p))} \left\{ K_{p,q}(\mathbb{D}) \cdot B_{r,q}(\mathbb{D}) \right\}.$$

Here $B_{r,q}(\mathbb{D})$ is the best constant in the (non-weighted) (r,q) -Poincaré–Sobolev inequality in the unit disc $\mathbb{D} \subset \mathbb{R}^2$ and $K_{p,q}(\Omega)$ is the norm of composition operator

$$(\varphi^{-1})^* : L^{1,p}(\Omega) \rightarrow L^{1,q}(\mathbb{D})$$

generated by the inverse conformal mapping $\varphi^{-1} : \mathbb{D} \rightarrow \Omega$:

$$K_{p,q}(\Omega) \leq \left(\iint_{\mathbb{D}} \left| (\varphi^{-1})' \right|^{\frac{(p-2)q}{p-q}} dudv \right)^{\frac{p-q}{pq}}.$$

Remark 1.4. Theorem C holds (without referring the Brennan’s Conjecture) for

$$1 \leq r \leq \frac{2p}{2-p} \cdot \frac{|\alpha_0|}{2+|\alpha_0|} < \frac{p}{2-p} \tag{1.8}$$

and $p \in ((|\alpha_0| + 2)/(|\alpha_0| + 1), 2)$, where $\alpha_0 = -1.752$ represents the best result for which Brennan’s conjecture was proved.

Remark 1.5. Let $\Omega \subset \mathbb{R}^2$ be a simply connected smooth domain. Then $\varphi^{-1} \in L^\alpha(\mathbb{D})$ for all $\alpha \in \mathbb{R}$ and we have the weighted Poincaré–Sobolev inequality (1.2) for all $p \in [1, 2)$ and all $r \in [1, 2p/(2 - p)]$.

In the case, when we have an embedding of a weighted Lebesgue space into a non-weighted one, the weighted Poincaré–Sobolev inequality (1.7) implies the standard Poincaré–Sobolev inequality (1.6).

Let us give some historical remarks about the notion of conformal regular domains. This notion was introduced in [16] and was applied to the stability problem for eigenvalues of the Dirichlet–Laplace operator. In [10] the lower estimates for the first non-trivial eigenvalues of the Neumann–Laplace operator in conformal regular domains were obtained. In [5] we proved but did not formulated the following important fact about conformal regular domains and the Poincaré–Sobolev inequality:

Theorem 1.6. Let $\Omega \subset \mathbb{R}^2$ be a simply connected domain of finite area such that the $(s, 2)$ -Poincaré–Sobolev inequality

$$\inf_{c \in \mathbb{R}} \left(\iint_{\Omega} |f(x, y) - c|^s dx dy \right)^{\frac{1}{s}} \leq B_{s,2}(\Omega) \left(\iint_{\Omega} |\nabla f(x, y)|^2 dx dy \right)^{\frac{1}{2}}$$

does not hold for some $s \geq 2$. Then Ω is not a conformal regular domain.

In the present work we suggest for the conformal regular domains a new method based on the theory of composition operators [17,18] and its applications to the Sobolev type embedding theorems [19,20].

The following diagram illustrates this idea:

$$\begin{array}{ccc} W^{1,p}(\Omega) & \xrightarrow{\varphi^*} & W^{1,q}(\mathbb{D}) \\ \downarrow & & \downarrow \\ L^s(\Omega) & \xleftarrow{(\varphi^{-1})^*} & L^r(\mathbb{D}). \end{array}$$

Here the operator φ^* defined by the composition rule $\varphi^*(f) = f \circ \varphi$ is a bounded composition operator on Sobolev spaces induced by a homeomorphism φ of Ω and \mathbb{D} and the operator $(\varphi^{-1})^*$ defined by the composition rule $(\varphi^{-1})^*(f) = f \circ \varphi^{-1}$ is a bounded composition operator on Lebesgue spaces. This method allows to transfer Poincaré–Sobolev inequalities from regular domains (for example, from the unit disc \mathbb{D}) to Ω .

In recent works we studied composition operators on Sobolev spaces in connection with the conformal mappings theory [21]. This connection leads to weighted Sobolev embeddings [22,4] with the universal conformal weights. Another application of conformal composition operators was given in [16] where the spectral stability problem for conformal regular domains was considered.

2. Composition operators

Since all composition operators that will be used in this paper are induced by conformal homeomorphisms we formulate results about composition operators for diffeomorphisms only.

2.1. Composition operators on Lebesgue spaces

For any domain $\Omega \subset \mathbb{R}^2$ and any $1 \leq p < \infty$ we consider the Lebesgue space $L^p(\Omega)$ of measurable functions $f : \Omega \rightarrow \mathbb{R}$ equipped with the following norm:

$$\|f\|_{L^p(\Omega)} := \left(\iint_{\Omega} |f(x, y)|^p dx dy \right)^{1/p} < \infty.$$

The following theorem about composition operators on Lebesgue spaces is well known (see, for example [18]):

Theorem 2.1. Let $\varphi : \Omega \rightarrow \Omega'$ be a diffeomorphism between two planar domains Ω and Ω' . Then the composition operator

$$\varphi^* : L^r(\Omega') \rightarrow L^s(\Omega), \quad 1 \leq s \leq r < \infty,$$

is bounded, if and only if

$$\left(\iint_{\Omega'} (J_{\varphi^{-1}}(u, v))^{\frac{r}{r-s}} dudv \right)^{\frac{r-s}{rs}} = K < \infty, \quad 1 \leq s < r < \infty,$$

$$\sup_{(u,v) \in \Omega'} (J_{\varphi^{-1}}(u, v))^{\frac{1}{s}} = K < \infty, \quad 1 \leq s = r < \infty.$$

The norm $\|\varphi^*\|$ of the composition operator φ^* equals K .

2.2. Composition operators on Sobolev spaces

We define the Sobolev space $W^{1,p}(\Omega)$, $1 \leq p < \infty$ as a Banach space of locally integrable weakly differentiable functions $f : \Omega \rightarrow \mathbb{R}$ equipped with the following norm:

$$\|f\|_{W^{1,p}(\Omega)} = \left(\iint_{\Omega} |f(x, y)|^p dx dy \right)^{\frac{1}{p}} + \left(\iint_{\Omega} |\nabla f(x, y)|^p dx dy \right)^{\frac{1}{p}}.$$

We define also the homogeneous seminormed Sobolev space $L^{1,p}(\Omega)$ of locally integrable weakly differentiable functions $f : \Omega \rightarrow \mathbb{R}$ equipped with the following seminorm:

$$\|f\|_{L^{1,p}(\Omega)} = \left(\iint_{\Omega} |\nabla f(x, y)|^p dx dy \right)^{\frac{1}{p}}.$$

Recall that the embedding operator $i : L^{1,p}(\Omega) \rightarrow L^1_{\text{loc}}(\Omega)$ is bounded.

Let Ω and Ω' be domains in \mathbb{R}^2 . We say that a diffeomorphism $\varphi : \Omega \rightarrow \Omega'$ induces a bounded composition operator

$$\varphi^* : L^{1,p}(\Omega') \rightarrow L^{1,q}(\Omega), \quad 1 \leq q \leq p \leq \infty,$$

by the composition rule $\varphi^*(f) = f \circ \varphi$ if $\varphi^*(f) \in L^{1,q}(\Omega)$ and there exists a constant $K < \infty$ such that

$$\|\varphi^*(f)\|_{L^{1,q}(\Omega)} \leq K \|f\|_{L^{1,p}(\Omega')} \quad \text{for all } f \in L^{1,p}(\Omega').$$

The main result of [17] gives the analytic description of composition operators on Sobolev spaces (see also [18]) and asserts (in the case of diffeomorphisms) that

Theorem 2.2 ([17]). *A diffeomorphism $\varphi : \Omega \rightarrow \Omega'$ between two domains Ω and Ω' induces a bounded composition operator*

$$\varphi^* : L^{1,p}(\Omega') \rightarrow L^{1,q}(\Omega), \quad 1 \leq q < p < \infty,$$

if and only if

$$K_{p,q}(\Omega) = \left(\iint_{\Omega} \left(\frac{|\varphi'(x, y)|^p}{|J_{\varphi}(x, y)|} \right)^{\frac{q}{p-q}} dx dy \right)^{\frac{p-q}{pq}} < \infty.$$

Definition 2.3. We call a bounded domain $\Omega \subset \mathbb{R}^2$ as a (r, q) -embedding domain, $1 \leq q, r \leq \infty$, if the embedding operator

$$i_{\Omega} : W^{1,q}(\Omega) \hookrightarrow L^r(\Omega)$$

is bounded. The unit disc $\mathbb{D} \subset \mathbb{R}^2$ is an example of the $(r, 2)$ -embedding domain for all $r \geq 1$.

The following theorem gives a characterization of composition operators in the normed Sobolev spaces [10]. For readers convenience we reproduce here the proof of the theorem.

Theorem 2.4. *Let Ω be an (r, q) -embedding domain for some $1 \leq q \leq r < \infty$ and $|\Omega'| < \infty$. Suppose that a diffeomorphism $\varphi : \Omega \rightarrow \Omega'$ induces a bounded composition operator*

$$\varphi^* : L^{1,p}(\Omega') \rightarrow L^{1,q}(\Omega), \quad 1 \leq q \leq p < \infty,$$

and the inverse diffeomorphism $\varphi^{-1} : \Omega' \rightarrow \Omega$ induces a bounded composition operator

$$(\varphi^{-1})^* : L^r(\Omega) \rightarrow L^s(\Omega')$$

for some $p \leq s \leq r$.

Then $\varphi : \Omega \rightarrow \Omega'$ induces a bounded composition operator

$$\varphi^* : W^{1,p}(\Omega') \rightarrow W^{1,q}(\Omega), \quad 1 \leq q \leq p < \infty.$$

Proof. Because the composition operator $(\varphi^{-1})^* : L^r(\Omega) \rightarrow L^s(\Omega')$ is bounded, then the following inequality

$$\|(\varphi^{-1})^* g \mid L^s(\Omega')\| \leq A_{r,s}(\Omega) \|g \mid L^r(\Omega)\|$$

holds. Here $A_{r,s}(\Omega)$ is a positive constant.

If a domain Ω is an embedding domain and the composition operators

$$(\varphi^{-1})^* : L^r(\Omega) \rightarrow L^s(\Omega'), \quad \varphi^* : L^{1,p}(\Omega') \rightarrow L^{1,q}(\Omega)$$

are bounded, then for a function $f = g \circ \varphi^{-1}$ the following inequalities

$$\begin{aligned} \inf_{c \in \mathbb{R}} \|f - c \mid L^s(\Omega')\| &\leq A_{r,s}(\Omega) \inf_{c \in \mathbb{R}} \|g - c \mid L^r(\Omega)\| \\ &\leq A_{r,s}(\Omega) M \|g \mid L^{1,q}(\Omega)\| \leq A_{r,s}(\Omega) K_{p,q}(\Omega) M \|f \mid L^{1,p}(\Omega')\| \end{aligned}$$

hold. Here M and $K_{p,q}(\Omega)$ are positive constants.

The Hölder inequality implies the following estimate

$$\begin{aligned} |c| &= |\Omega'|^{-\frac{1}{p}} \|c \mid L^p(\Omega')\| \leq |\Omega'|^{-\frac{1}{p}} (\|f \mid L^p(\Omega')\| + \|f - c \mid L^p(\Omega')\|) \\ &\leq |\Omega'|^{-\frac{1}{p}} \|f \mid L^p(\Omega')\| + |\Omega'|^{-\frac{1}{s}} \|f - c \mid L^s(\Omega')\|. \end{aligned}$$

Since $q \leq r$ we have

$$\begin{aligned} \|g \mid L^q(\Omega)\| &\leq \|c \mid L^q(\Omega)\| + \|g - c \mid L^q(\Omega)\| \leq |c| |\Omega|^{\frac{1}{q}} + |\Omega|^{\frac{r-q}{r}} \|g - c \mid L^r(\Omega)\| \\ &\leq \left(|\Omega'|^{-\frac{1}{p}} \|f \mid L^p(\Omega')\| + |\Omega'|^{-\frac{1}{s}} \|f - c \mid L^s(\Omega')\| \right) |\Omega|^{\frac{1}{q}} + |\Omega|^{\frac{r-q}{r}} \|g - c \mid L^r(\Omega)\|. \end{aligned}$$

From previous inequalities we obtain for $\varphi^*(f) = g$ finally

$$\begin{aligned} \|g \mid L^q(\Omega)\| &\leq |\Omega|^{\frac{1}{q}} |\Omega'|^{-\frac{1}{p}} \|f \mid L^p(\Omega')\| + A_{r,s}(\Omega) K_{p,q}(\Omega) M |\Omega|^{\frac{1}{q}} |\Omega'|^{-\frac{1}{p}} \|f \mid L^{1,p}(\Omega')\| \\ &\quad + K_{p,q}(\Omega) M |\Omega|^{\frac{r-q}{r}} \|f \mid L^{1,p}(\Omega')\|. \end{aligned}$$

Therefore the composition operator

$$\varphi^* : W^{1,p}(\Omega') \rightarrow W^{1,q}(\Omega)$$

is bounded. \square

3. Poincaré–Sobolev inequalities

3.1. Weighted Poincaré–Sobolev inequalities

Let $\Omega \subset \mathbb{R}^2$ be a planar domain and let $v : \Omega \rightarrow \mathbb{R}$ be a smooth positive real valued function in Ω . For $1 \leq p < \infty$ consider the weighted Lebesgue space $L^p(\Omega, v)$ of measurable functions $f : \Omega \rightarrow \mathbb{R}$ with the finite norm

$$\|f \mid L^p(\Omega, v)\| := \left(\iint_{\Omega} |f(x, y)|^p v(x, y) \, dx dy \right)^{1/p} < \infty.$$

It is a Banach space for the norm $\|f \mid L^p(\Omega, v)\|$.

Applications of the conformal mappings theory to the Poincaré–Sobolev inequalities in planar domains are based on the following result (Theorem 3.3, Proposition 3.4 [4]) which connected the classical mappings theory and the Sobolev spaces theory.

Theorem 3.1. *Let $\Omega \subset \mathbb{R}^2$ be a simply connected domain with non-empty boundary and $\varphi : \Omega \rightarrow \mathbb{D}$ be a conformal homeomorphism. Suppose that the (Inverse) Brennan’s Conjecture holds for the interval $[\alpha_0, 2/3)$ where $\alpha_0 \in (-2, 0)$ and $p \in \left(\frac{|\alpha_0|+2}{|\alpha_0|+1}, 2\right)$.*

Then the inverse mapping φ^{-1} induces a bounded composition operator

$$(\varphi^{-1})^* : L^1_p(\Omega) \rightarrow L^1_q(\mathbb{D})$$

for any q such that

$$1 \leq q \leq \frac{p|\alpha_0|}{2 + |\alpha_0| - p} < \frac{2p}{4 - p}$$

and for any function $f \in L^1_p(\Omega)$ the inequality

$$\|(\varphi^{-1})^* f \mid L^{1,q}(\mathbb{D})\| \leq \left(\iint_{\mathbb{D}} |(\varphi^{-1})'| \frac{(p-2)q}{p-q} dudv \right)^{\frac{p-q}{pq}} \|f \mid L^{1,p}(\Omega)\|$$

holds.

Remark 3.2. Let us remark that $\frac{|\alpha_0|+2}{|\alpha_0|+1} > \frac{4}{3}$ for any $\alpha_0 \in (-2, 0)$.

Using this theorem we prove

Theorem C'. Suppose that $\Omega \subset \mathbb{C}$ is a simply connected domain with non empty boundary, the Brennan's Conjecture holds for the interval $[\alpha_0, 2/3)$, where $\alpha_0 \in (-2, 0)$ and $h(z) = J_\varphi(z)$ is the conformal weight defined by a conformal homeomorphism $\varphi : \Omega \rightarrow \mathbb{D}$. Then for every $p \in ((|\alpha_0| + 2)/(|\alpha_0| + 1), 2)$ and every function $f \in W^{1,p}(\Omega)$, the inequality

$$\inf_{c \in \mathbb{R}} \left(\iint_{\Omega} |f(x, y) - c|^r h(z) dx dy \right)^{\frac{1}{r}} \leq B_{r,p}(\Omega, h) \left(\iint_{\Omega} |\nabla f(x, y)|^p dx dy \right)^{\frac{1}{p}}$$

holds for any r such that

$$1 \leq r \leq \frac{2p}{2 - p} \cdot \frac{|\alpha_0|}{2 + |\alpha_0|} < \frac{p}{2 - p}$$

with the constant

$$B_{r,p}(\Omega, h) \leq \inf_{q \in [1, 2p/(4-p))} \{K_{p,q}(\mathbb{D}) \cdot B_{r,q}(\mathbb{D})\}.$$

Here $B_{r,q}(\mathbb{D})$ is the best constant in the (non-weighted) Poincaré–Sobolev inequality in the unit disc $\mathbb{D} \subset \mathbb{C}$ and $K_{p,q}(\Omega)$ is the norm of composition operator

$$(\varphi^{-1})^* : L^{1,p}(\Omega) \rightarrow L^{1,q}(\mathbb{D})$$

generated by the inverse conformal mapping $\varphi^{-1} : \mathbb{D} \rightarrow \Omega$:

$$K_{p,q}(\Omega) \leq \left(\iint_{\mathbb{D}} \left| (\varphi^{-1})' \right|^{\frac{(p-2)q}{p-q}} dudv \right)^{\frac{p-q}{pq}}.$$

Proof. By the Riemann Mapping Theorem, there exists a conformal mapping $\varphi : \Omega \rightarrow \mathbb{D}$, and by the (Inverse) Brennan's Conjecture,

$$\int_{\mathbb{D}} |(\varphi^{-1})'(u, v)|^\alpha dudv < +\infty, \quad \text{for all } -2 < \alpha_0 < \alpha < 2/3.$$

Hence, by Theorem 3.1, the inequality

$$\|\nabla(f \circ \varphi^{-1}) \mid L^q(\mathbb{D})\| \leq K_{p,q}(\mathbb{D}) \|\nabla f \mid L^p(\Omega)\|$$

holds for every function $f \in L^{1,p}(\Omega)$ and for any q such that

$$1 \leq q \leq \frac{p|\alpha_0|}{2 + |\alpha_0| - p} < \frac{2p}{4 - p}. \tag{3.1}$$

Choose arbitrarily $f \in C^1(\Omega)$. Then $g = f \circ \varphi^{-1} \in C^1(\mathbb{D})$ and, by the classical Poincaré–Sobolev inequality,

$$\inf_{c \in \mathbb{R}} \|f \circ \varphi^{-1} - c\|_{L^r(\mathbb{D})} \leq B_{q,r}(\mathbb{D}) \|\nabla(f \circ \varphi^{-1})\|_{L^q(\mathbb{D})} \tag{3.2}$$

for any r such that

$$1 \leq r \leq \frac{2q}{2-q}.$$

By elementary calculations from the inequality (3.1), it follows that

$$\frac{2q}{2-q} \leq \frac{2p}{2-p} \cdot \frac{|\alpha_0|}{2+|\alpha_0|} < \frac{p}{2-p}.$$

Combining inequalities for $2q/(2-q)$ and r we conclude that the inequality (3.2) holds for any r such that

$$1 \leq r \leq \frac{2p}{2-p} \cdot \frac{|\alpha_0|}{2+|\alpha_0|} < \frac{p}{2-p}.$$

Using the change of variable formula, the classical Poincaré–Sobolev inequality for the unit disc

$$\inf_{c \in \mathbb{R}} \left(\iint_{\mathbb{D}} |g(u, v) - c|^r \, dudv \right)^{\frac{1}{r}} \leq B_{r,q}(\mathbb{D}) \left(\iint_{\mathbb{D}} |\nabla g(u, v)|^q \, dudv \right)^{\frac{1}{q}}$$

and Theorem 3.1, we finally infer

$$\begin{aligned} & \inf_{c \in \mathbb{R}} \left(\iint_{\Omega} |f(x, y) - c|^r h(x, y) \, dxdy \right)^{\frac{1}{r}} \\ &= \inf_{c \in \mathbb{R}} \left(\iint_{\Omega} |f(x, y) - c|^r J_{\varphi}(x, y) \, dxdy \right)^{\frac{1}{r}} = \inf_{c \in \mathbb{R}} \left(\iint_{\mathbb{D}} |g(u, v) - c|^r \, dudv \right)^{\frac{1}{r}} \\ &\leq B_{r,q}(\mathbb{D}) \left(\iint_{\mathbb{D}} |\nabla g(u, v)|^q \, dudv \right)^{\frac{1}{q}} \leq K_{p,q}(\mathbb{D}) \cdot B_{r,q}(\mathbb{D}) \left(\iint_{\Omega} |\nabla f(x, y)|^p \, dxdy \right)^{\frac{1}{p}}. \end{aligned}$$

Approximating an arbitrary function $f \in W^{1,p}(\Omega)$ by smooth functions we obtain

$$\inf_{c \in \mathbb{R}} \left(\iint_{\Omega} |f(x, y) - c|^r h(z) \, dxdy \right)^{\frac{1}{r}} \leq B_{r,p}(\Omega, h) \left(\iint_{\Omega} |\nabla f(x, y)|^p \, dxdy \right)^{\frac{1}{p}}$$

with the constant

$$B_{r,p}(\Omega, h) \leq \inf_{q:q \in [1, 2p/(4-p))} \{K_{p,q}(\mathbb{D}) \cdot B_{r,q}(\mathbb{D})\}. \quad \square$$

The property of the conformal α -regularity implies the integrability of a Jacobian of conformal mappings (conformal weights) and therefore for any conformal α -regular domain we have the embedding of weighted Lebesgue spaces $L^r(\Omega, h)$ into non-weighted Lebesgue spaces $L^s(\Omega)$ for $s = \frac{\alpha-2}{\alpha}r$.

Lemma 3.3. *Let Ω be a conformal α -regular domain. Then for any function $f \in L^r(\Omega, h)$, $\alpha/(\alpha - 2) \leq r < \infty$, the inequality*

$$\|f\|_{L^s(\Omega)} \leq \left(\iint_{\mathbb{D}} |(\varphi^{-1})'|^{\alpha} \, dudv \right)^{\frac{2}{\alpha} \cdot \frac{1}{s}} \|f\|_{L^r(\Omega, h)}$$

holds for $s = \frac{\alpha-2}{\alpha}r$.

Proof. Since Ω is a conformal α -regular domain then there exists a conformal mapping $\varphi : \Omega \rightarrow \mathbb{D}$ such that

$$\left(\iint_{\mathbb{D}} |J_{\varphi^{-1}}(u, v)|^{\frac{r}{r-s}} \, dudv \right)^{\frac{r-s}{rs}} = \left(\iint_{\mathbb{D}} |(\varphi^{-1})'|^{\alpha} \, dudv \right)^{\frac{2}{\alpha} \cdot \frac{1}{s}} < \infty,$$

for $s = \frac{\alpha-2}{\alpha}r$. Then

$$\begin{aligned} \|f \mid L^s(\Omega)\| &= \left(\iint_{\Omega} |f(x, y)|^s dx dy \right)^{\frac{1}{s}} = \left(\iint_{\Omega} |f(x, y)|^s J_{\varphi^{\frac{s}{r}}}(x, y) J_{\varphi^{-\frac{s}{r}}}(x, y) dx dy \right)^{\frac{1}{s}} \\ &\leq \left(\iint_{\Omega} |f(x, y)|^r J_{\varphi}(x, y) dx dy \right)^{\frac{1}{r}} \left(\iint_{\Omega} J_{\varphi^{-\frac{s}{r-s}}}(x, y) dx dy \right)^{\frac{r-s}{rs}} \\ &= \left(\iint_{\Omega} |f(x, y)|^r h(x, y) dx dy \right)^{\frac{1}{r}} \left(\iint_{\mathbb{D}} J_{\varphi^{-1}}^{\frac{r}{r-s}}(u, v) dudv \right)^{\frac{r-s}{rs}} \\ &= \left(\iint_{\Omega} |f(x, y)|^r h(x, y) dx dy \right)^{\frac{1}{r}} \left(\iint_{\mathbb{D}} |(\varphi^{-1})'(u, v)|^{\alpha} dudv \right)^{\frac{2}{\alpha} \cdot \frac{1}{s}}. \end{aligned} \tag{3.3}$$

The Lemma is proved. \square

From Theorem C' and Lemma 3.3 follows Theorem B':

Theorem B'. Suppose that $\Omega \subset \mathbb{C}$ is a simply connected domain with non empty boundary, the Brennan's Conjecture holds for the interval $[\alpha_0, 2/3)$, where $\alpha_0 \in (-2, 0)$.

Then for every

$$p \in \left(\max \left\{ \frac{4}{3}, \frac{2\alpha(|\alpha_0| + 2)}{(2\alpha + 3\alpha|\alpha_0| - 4|\alpha_0|)}, 2 \right\}, 2 \right),$$

every $s \in [1, \frac{\alpha-2}{\alpha} \frac{p}{2-p} \frac{|\alpha_0|}{2+|\alpha_0|}]$ and every function $f \in W^{1,p}(\Omega)$, the inequality

$$\inf_{c \in \mathbb{R}} \left(\iint_{\Omega} |f(x, y) - c|^s dx dy \right)^{\frac{1}{s}} \leq B_{s,p}(\Omega) \left(\iint_{\Omega} |\nabla f(x, y)|^p dx dy \right)^{\frac{1}{p}} \tag{3.4}$$

holds with the constant

$$B_{s,p}(\Omega) \leq \|(\varphi^{-1})' \mid L^{\alpha}(\mathbb{D})\|^{\frac{2}{s}} B_{r,p}(\Omega, h) \leq \inf_{q \in [1, 2p/(4-p))} \left\{ B_{\frac{\alpha s}{\alpha-2}, q}(\mathbb{D}) \cdot \|(\varphi^{-1})' \mid L^{\alpha}(\mathbb{D})\|^{\frac{2}{s}} K_{p,q}(\mathbb{D}) \right\}.$$

Proof. The inequality (3.4) immediately follows from the main inequality of Theorem C' and the main inequality of Lemma 3.3. The last part of this inequality used known estimates for the constant of the Poincaré–Sobolev inequality in the unit disc.

It is necessary to clarify the restrictions for parameters p, r, s , because these restrictions do not follow directly from Theorem A' and Lemma 3.3.

By Lemma 3.3 $s = \frac{\alpha-2}{\alpha}r$. By Theorem C'

$$1 \leq r \leq \frac{2p}{2-p} \cdot \frac{|\alpha_0|}{2+|\alpha_0|} < \frac{p}{2-p}.$$

Hence

$$1 \leq s \leq \frac{\alpha-2}{\alpha} \cdot \frac{2p}{2-p} \cdot \frac{|\alpha_0|}{2+|\alpha_0|} < \frac{\alpha-2}{\alpha} \cdot \frac{p}{2-p}.$$

Since $1 \leq s$ we have from this inequality that

$$\frac{\alpha}{\alpha-2} \leq \frac{2p}{2-p} \cdot \frac{|\alpha_0|}{2+|\alpha_0|} < \frac{p}{2-p}.$$

By elementary calculations

$$p \geq \frac{2\alpha(2 + |\alpha_0|)}{2\alpha + 3\alpha|\alpha_0| - 4|\alpha_0|} > \frac{\alpha}{\alpha - 1}.$$

The last inequality is correct by factor that Brennan’s conjecture is correct for all $\alpha : -2 < \alpha < 2/3$. \square

Theorem A’ follows from **Theorem B’**, using $s = p$, that is necessary for coincidence of the first nontrivial Neumann–Laplace eigenvalue and the constant in the Poincaré–Sobolev inequality of **Theorem B’**.

Theorem A’. Let $\varphi : \Omega \rightarrow \mathbb{D}$ be a conformal homeomorphism from a conformal α -regular domain Ω to the unit disc \mathbb{D} and the Brennan’s Conjecture holds for the interval $[\alpha_0, 2/3)$, where $\alpha_0 \in (-2, 0)$.

Then for every

$$p \in \left(\max \left\{ 4/3, \frac{4(\alpha + |\alpha_0|)}{\alpha(2 + |\alpha_0|)} \right\}, 2 \right)$$

the following inequality holds

$$\frac{1}{\mu_p(\Omega)} \leq \inf_{q \in [1, 2p/(4-p))} \left\{ B_{\frac{\alpha p}{\alpha-2}, q}^p(\mathbb{D}) \cdot \|(\varphi^{-1})'\|_{L^\alpha(\mathbb{D})}^2 \left(\iint_{\mathbb{D}} \left| (\varphi^{-1})' \right|^{\frac{(p-2)q}{p-q}} dudv \right)^{\frac{p-q}{q}} \right\}.$$

Proof. By **Lemma 3.3** $p = \frac{\alpha-2}{\alpha}r$. By **Theorem C’**

$$1 \leq r \leq \frac{2p}{2-p} \cdot \frac{|\alpha_0|}{2 + |\alpha_0|} < \frac{p}{2-p}.$$

Hence

$$\frac{\alpha}{\alpha - 2} \leq \frac{1}{2 - p} \cdot \frac{2|\alpha_0|}{2 + |\alpha_0|}.$$

By elementary calculations

$$p \geq 2 - \frac{2|\alpha_0|}{2 + |\alpha_0|} \frac{\alpha - 2}{\alpha} = \frac{4(\alpha + |\alpha_0|)}{\alpha(2 + |\alpha_0|)} > \frac{\alpha + 2}{\alpha}.$$

The last inequality holds provided that the Brennan’s conjecture holds true all $\alpha : -2 < \alpha < 2/3$. \square

Corollary A’. Suppose that Ω is smoothly bounded Jordan domain with a boundary $\partial\Omega$ of a class C^1 with a Dini continuous normal. Let $\varphi : \Omega \rightarrow \mathbb{D}$ be a conformal homeomorphism from Ω onto the unit disc \mathbb{D} . Then for every $p \in (1, 2)$ the following inequality is correct

$$\frac{1}{\mu_p(\Omega)} \leq \|(\varphi^{-1})'\|_{L^\infty(\mathbb{D})}^p \frac{1}{\mu_p(\mathbb{D})}.$$

Proof. If Ω is smoothly bounded Jordan domain with a boundary $\partial\Omega$ of a class C^1 with a Dini continuous normal, then for a conformal mapping $\varphi : \Omega \rightarrow \mathbb{D}$, the derivative φ' is bounded away from 0 and ∞ [23]. Hence, we can apply **Theorem A’** in the limit case $\alpha = \infty$ and $p = q$. Then

$$\frac{1}{\mu_p(\Omega)} \leq B_{p,p}^p(\mathbb{D}) \cdot \|(\varphi^{-1})'\|_{L^\infty(\mathbb{D})}^2 \|(\varphi^{-1})'\|_{L^\infty(\mathbb{D})}^{p-2} = \|(\varphi^{-1})'\|_{L^\infty(\mathbb{D})}^p \frac{1}{\mu_p(\mathbb{D})}.$$

The corollary is proved. \square

As an application we obtain the lower estimate of the first non-trivial eigenvalue on the Neumann eigenvalue problem for the p -Laplace operator in the interior of the cardioid (which is a non-convex domain with a non-smooth boundary).

Example 3.4. Let Ω_c be the interior of the cardioid $\rho = 2(1 + \cos \theta)$. The diffeomorphism

$$z = \psi(w) = (w + 1)^2, \quad w = u + iv,$$

is conformal and maps the unit disc \mathbb{D} onto Ω_c . Then by [Theorem A'](#):

$$\begin{aligned} \frac{1}{\mu_p(\Omega_c)} &\leq \inf_{1 \leq q \leq \frac{2p}{4-p}} \left(\frac{2}{\pi^\delta} \left(\frac{1-\delta}{1/2-\delta} \right)^{1-\delta} \right)^p \left(\iint_{\mathbb{D}} (2|w+1|)^\alpha \, dudv \right)^{\frac{2}{\alpha}} \\ &\quad \times \left(\iint_{\mathbb{D}} (2|w+1|)^{\frac{(p-2)q}{p-q}} \, dudv \right)^{\frac{p-q}{q}}. \end{aligned} \quad (3.5)$$

Here $\delta = 1/q - (\alpha - 2)/\alpha p$.

Acknowledgements

The first author was supported by the United States–Israel Binational Science Foundation (BSF Grant No. 2014055).

The authors thank the reviewer for careful reading of the paper and useful comments.

References

- [1] B. Brandolini, F. Chiacchio, C. Trombetti, Optimal lower bounds for eigenvalues of linear and nonlinear Neumann problems, *Proc. Roy. Soc. Edinburgh Sect. A* 145 (1) (2015) 31–45.
- [2] V. Maz'ya, *Sobolev Spaces: With Applications to Elliptic Partial Differential Equations*, Springer, Berlin/Heidelberg, 2010.
- [3] J. Brennan, The integrability of the derivative in conformal mapping, *J. Lond. Math. Soc.* (2) 18 (2) (1978) 261–272.
- [4] V. Gol'dshtein, A. Ukhlov, Brennan's conjecture and universal Sobolev inequalities, *Bull. Sci. Math.* 138 (2) (2014) 253–269.
- [5] V. Gol'dshtein, A. Ukhlov, Sobolev homeomorphisms and Brennan's conjecture, *Comput. Methods Funct. Theory* 14 (2–3) (2014) 247–256.
- [6] J. Becker, C. Pommerenke, Hölder continuity of conformal maps with quasiconformal extension, *Complex Variables Theory Appl.* 10 (4) (1988) 267–272.
- [7] A.F. Beardon, D. Minda, *The hyperbolic metric and geometric function theory. Quasiconformal mappings and their applications*, Narosa, New Delhi, 2007, pp. 9–56.
- [8] S. Hencl, P. Koskela, T. Nieminen, Dimension gap under conformal mappings, *Adv. Math.* 230 (3) (2012) 1423–1441.
- [9] D. Gilbarg, N.S. Trudinger, *Elliptic partial differential equations of second order*, in: *Grundlehren der Mathematischen Wissenschaften*, vol. 224, Springer-Verlag, Berlin-New York, 1977.
- [10] V. Gol'dshtein, A. Ukhlov, On the first eigenvalues of free vibrating membranes in conformal regular domains, *Arch. Rational Mech. Anal.* <http://dx.doi.org/10.1007/s00205-016-0988-9>.
- [11] H. Hedenmalm, S. Shimorin, Weighted Bergman spaces and the integral means spectrum of conformal mappings, *Duke Math. J.* 127 (2) (2005) 341–393.
- [12] L. Esposito, C. Nitsch, C. Trombetti, Best constants in Poincaré inequalities for convex domains, *J. Convex Anal.* 20 (1) (2013) 253–264.
- [13] V.I. Burenkov, P.D. Lamberti, Spectral stability of the p -Laplacian, *Nonlinear Anal.* 71 (5–6) (2009) 2227–2235.
- [14] V.I. Burenkov, P.D. Lamberti, M. Lanza de Cristoforis, Spectral stability of nonnegative selfadjoint operators, *Sovrem. Mat. Fundam. Naprav.* 15 (2006) 76–111 (in Russian); *J. Math. Sci. (N. Y.)* 149 (4) (2008) 1417–1452. (transl.).
- [15] P.D. Lamberti, A differentiability result for the first eigenvalue of the p -Laplacian upon domain perturbation, in: *Nonlinear Analysis and Applications: to V. Lakshmikantham on his 80th birthday*, Vol. 1, 2, Kluwer Acad. Publ., Dordrecht, 2003, pp. 741–754.
- [16] V.I. Burenkov, V. Gol'dshtein, A. Ukhlov, Conformal spectral stability estimates for the Dirichlet Laplacian, *Math. Nachr.* 288 (16) (2015) 1822–1833.
- [17] A. Ukhlov, On mappings, which induce embeddings of Sobolev spaces, *Siberian Math. J.* 34 (1993) 185–192.
- [18] S.K. Vodop'yanov, A.D. Ukhlov, Superposition operators in Sobolev spaces, *Izv. Vyssh. Uchebn. Zaved. Mat.* (10) (2002) 11–33 (in Russian); *Math. (Iz. VUZ)* 46 (10) (2002) 9–31. (transl. Russian) 2003.
- [19] V. Gol'dshtein, L. Gurov, Applications of change of variables operators for exact embedding theorems, *Integral Equations Operator Theory* 19 (1) (1994) 1–24.
- [20] V. Gol'dshtein, A. Ukhlov, Weighted Sobolev spaces and embedding theorems, *Trans. Amer. Math. Soc.* 361 (7) (2009) 3829–3850.
- [21] V. Gol'dshtein, A. Ukhlov, Brennan's conjecture for composition operators on Sobolev spaces, *Eurasian Math. J.* 3 (4) (2012) 35–43.
- [22] V. Gol'dshtein, A. Ukhlov, Conformal weights and Sobolev embeddings, in: *Problems in Mathematical Analysis*, no. 71, *J. Math. Sci. (N. Y.)* 193 (2) (2013) 202–210.
- [23] C. Pommerenke, Univalent functions. With a chapter on quadratic differentials by Gerd Jensen, in: *Studia Mathematica/Mathematische Lehrbücher*, Band XXV, Vandenhoeck & Ruprecht, Göttingen, 1975.