



ELSEVIER

Journal de
Pediatria

www.jpmed.com.br



SOCIEDADE BRASILEIRA
DE PEDIATRIA

REVIEW ARTICLE

Exacerbation of asthma and airway infection: is the virus the villain?☆☆☆



Lusmaia D.C. Costa^{a,*}, Paulo Sucasas Costa^a, Paulo A.M. Camargos^b

^a Department of Pediatrics, School of Medicine, Universidade Federal de Goiás (UFG), Goiânia, GO, Brazil

^b Department of Pediatrics, School of Medicine, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, MG, Brazil

Received 24 April 2014; accepted 1 July 2014

Available online 26 September 2014

KEYWORDS

Asthma;
Exacerbation;
Virus;
Child;
Respiratory tract
infection

Abstract

Objective: To review the available literature on the association between acute viral respiratory tract infection and the onset of asthma exacerbations, identifying the most prevalent viruses, detection methods, as well as preventive and therapeutic aspects.

Sources: A search was conducted in PubMed, Lilacs, and SciELO databases, between the years 2002 and 2013, using the following descriptors: asthma exacerbation, virus, child, and acute respiratory infection.

Summary of the findings: A total of 42 original articles addressing the identification of respiratory viruses during episodes of asthma exacerbation were selected, mostly cross-sectional studies. There was a wide variation in the methodology of the assessed studies, particularly in relation to the children's age and methods of collection and viral detection. The results indicate that, in up to 92.2% of exacerbations, a viral agent was potentially the main triggering factor, and human rhinovirus was the most frequently identified factor. The pattern of viral circulation may have been responsible for the seasonality of exacerbations. The association between viral infections and allergic inflammation appears to be crucial for the clinical and functional uncontrolled asthma, but few studies have evaluated other triggering factors in association with viral infection.

Conclusions: Respiratory viruses are present in the majority of asthmatic children during episodes of exacerbation. The involved physiopathological mechanisms are yet to be fully established, and the synergism between allergic inflammation and viral infection appears to determine uncontrolled disease. The role of other triggering and protective agents is yet to be clearly determined.

☆ Please cite this article as: Costa LD, Costa PS, Camargos PA. Exacerbation of asthma and airway infection: is the virus the villain? J Pediatr (Rio J). 2014;90:542–55.

☆☆ Study conducted at the Department of Pediatrics, Universidade Federal de Goiás (UFG), Goiânia, GO, Brazil.

* Corresponding author.

E-mail: lusmaiapneumoped@gmail.com, lusmaiacosta@uol.com.br (L.D.C. Costa).

PALAVRAS-CHAVE

Asma;
Exacerbação;
Vírus;
Criança;
Infecção do trato
respiratório

Exacerbação da asma e infecção das vias aéreas: o vírus é o vilão?**Resumo**

Objetivo: Rever a literatura disponível sobre a relação entre infecção viral aguda do trato respiratório e o desencadeamento de exacerbações da asma, identificando os vírus mais prevalentes, os métodos de detecção, bem como os aspectos preventivos e terapêuticos.

Fonte dos dados: Foi realizada uma busca nas bases de dados PubMed, SciELO e Lilacs utilizando os descritores: asma, exacerbação, vírus, criança e infecção respiratória aguda, entre os anos de 2002 e 2013.

Síntese dos dados: Foram selecionados 42 artigos originais que tratavam da identificação de vírus respiratórios durante episódios de exacerbação da asma, em sua maioria estudos transversais. Houve ampla variação na metodologia dos trabalhos avaliados, principalmente em relação à idade das crianças e métodos de coleta e detecção viral. Os resultados apontam que, em até 92,2% das exacerbações, um agente viral foi potencialmente o principal fator desencadeante, sendo o rinovírus humano o mais identificado. O padrão de circulação viral pode ter sido responsável pela sazonalidade das exacerbações. A associação entre infecção viral e inflamação alérgica parece ser determinante para levar ao descontrole clínico-funcional da asma, porém poucos estudos avaliaram outros fatores desencadeantes em associação com a infecção viral.

Conclusões: Os vírus respiratórios estão presentes na maioria das crianças asmáticas durante os episódios de exacerbação. Os mecanismos fisiopatológicos envolvidos ainda não estão totalmente estabelecidos e o sinergismo entre a inflamação alérgica e infecção viral parece determinar o descontrole da doença. O papel dos outros agentes desencadeantes e protetores não estão claramente determinados.

© 2014 Sociedade Brasileira de Pediatria. Publicado por Elsevier Editora Ltda.

Este é um artigo Open Access sob a licença de [CC BY-NC-ND](#)

Introduction

Asthma is a chronic, genetically-determined disease, whose prevalence in the pediatric population ranges between 19.0% and 24.3% among Brazilian adolescents and schoolchildren, respectively.¹ From the physiopathological viewpoint, it is characterized by chronic inflammation with the involvement of several cell types, associated with airway hyperresponsiveness, with episodes of reversible airflow limitation. It is clinically manifested by recurrent exacerbations, also called "asthma attacks" or, more appropriately, acute asthma, characterized by progressive worsening of dyspnea, coughing, wheezing, chest tightness, or a combination of these.²

The loss of clinical and functional asthma control usually occurs gradually, but it can occur abruptly in a subgroup of patients.² It is one of the main causes of emergency consultations, having been responsible, in 2007, for 195 deaths in children younger than 19 years in Brazil.³ Public policies have been developed to promote both scientific knowledge about the disease and its management, as well as to organize assistance programs in public health, which include, among others, the dispensing of medications. However, exacerbations continue to represent a significant number in statistics, with great impact on public and private healthcare systems.²

The multifactorial origin of the clinical-functional lack of disease control is well known; since the early 1970s, respiratory viruses have been associated with the triggering of asthma exacerbations in adults and children.³ In the 1990s, the development of more sensitive and specific molecular techniques allowed for the increase in respiratory virus detection and therefore, ways to better explain this

association. Studies using reverse transcriptase polymerase chain reaction (RT-PCR) as the detection technique, isolated or combined with traditional methods, observed positivity for respiratory viruses in up to 92.2% of episodes of acute asthma exacerbation in children.⁴

Considering the possibility of a causal relationship between respiratory virus infection and the triggering of asthma attacks in children, the implications of this association, as well as the possibility of specific prophylaxis and therapy for these agents, special attention to this subject is justified. Therefore, this literature review aimed to analyze articles, published between 2002 and 2013, assessing the association between asthma exacerbation and acute viral airway infection.

Methods

A search was conducted in the PubMed, Lilacs and SciELO databases, using the descriptors: "Asthma Exacerbation", "Viral Infection", and "Child", resulting in a total of 283 references for that period. After selecting the articles published in Portuguese, English, Spanish, or French, 195 articles remained. After reading the titles and abstracts, 42 original articles that assessed respiratory tract viral infection in asthmatic children during exacerbation were selected. Some articles of historical importance or review articles that included the three descriptors were added to generate the bibliography of this review. The list of references was inserted into Endnote X6 (Thompson Corp., CA, USA), a bibliographic citation management software.

Results

Respiratory viruses and immune response

The most frequently identified respiratory viruses in association with asthma exacerbation were human rhinovirus (HRV), respiratory syncytial virus (RSV), human adenovirus (hAdV), influenza (Flu), *Parainfluenza* (PFlu), human metapneumovirus (hMPV), and human coronavirus (hCoV). Of the listed viruses, most have RNA as the nucleic acid; their biological characteristics and taxonomy⁵ are described in Table 1.

The main transmission methods for these viruses are through contaminated fomites, droplets, aerosols, or direct contamination. By infecting the nasal epithelium cells, these agents trigger an immune response, which involves mainly the dendritic cells and natural killer (NK) cells, inducing the release of a number of pro-inflammatory cytokines and chemokines by the infected epithelial cells, such as interferon (IFN-1) and tumor necrosis factor alpha (TNF α), among others (Fig. 1).

In asthma patients, the viral infection causes an imbalance in the immune homeostasis of the respiratory system. Several mechanisms related to viral infection and allergic inflammation, as well as their role in triggering acute asthma, have been proposed; among them, the deficient function of the epithelial barrier caused by the virus, which has been implicated as a predisposing factor by some.⁶ However, both asthma and atopy are associated with epithelial damage, which may contribute to increased susceptibility to infections, including viral diseases and sensitization by aeroallergens.⁷ Another evaluated factor was mucus production as an airway defense mechanism; in mice studies, it was demonstrated that allergic inflammation and viral infection act synergistically increasing mucus production, which can lead to airway impaction and obstruction in asthma patients.⁸

Virus-induced alterations in interferon production have also been observed. For instance, *in vivo* and *in vitro* studies in epithelial cells from healthy adults and asthma patients infected with HRV demonstrated a decreased production of type I interferon (α and β) in the latter, making them more susceptible to infection associated with viral exacerbation.^{6,9} Similar results were obtained in studies performed with children, where the production of interferon and Th2 cytokines by bronchial epithelial cells was assessed after HRV-16 infection.

Lower interferon production and higher concentrations of viral RNA have been demonstrated in children with asthma, regardless of their atopic status, and in atopic children without asthma, suggesting that an impaired immune response to viral infection occurs not only in asthma patients, but in children with other disorders associated with Th2 lymphocytes.¹⁰ However, other studies failed to demonstrate the same reduction in interferon production; others even found an increase in its production in exacerbated asthma.^{11,12}

The bronchial epithelium produces some cytokines, including interleukin 25 and 33, as well as thymic stromal lymphopoietin, which promotes the differentiation of innate lymphoid cells into Th2. The latter can be induced by viral infection, and its production can be increased by

interleukin-4 (IL-4), suggesting that the interaction between viruses and allergic airway inflammation may enhance the inflammatory Th2 response and potentially reduce the antiviral response.^{11,13}

Collection and viral detection methods

Viral detection is highly dependent on the quality of the collected sample, on the time of symptom onset to the time of collection (ideally within 72 hours), and on transportation and storage of the sample before testing. The analysis for respiratory viruses should be performed in material from the airways. Upper airway secretion is used in most cases, and several methods are employed for this collection, such as nasopharyngeal aspirate (NPA), nasopharyngeal swab (NPS), nasopharyngeal lavage, and the combined nasal-oral swab; the first technique is considered the gold standard.¹⁴

Recently, new flocked swabs (Copan, Brescia, Italy) were developed, and presented better performance during data collection. Recent studies using this type of swab presented a sensitivity comparable to that of NPA when the detection is performed by PCR, suggesting that this swab can be used in epidemiological research and surveillance studies, due to its greater technical simplicity.^{15,16}

There are few data to support the use of combined oral-nasal swabs for virus detection, and its sensitivity is lower than that of the nasopharyngeal swab or aspirate, which can be explained by the lower viral load in the oropharynx than in the nasopharynx. The collection can also be performed on material from the lower airways, such as induced sputum and bronchial lavage.¹⁷

The methods for detection of respiratory viruses are varied and include rapid tests for antigen detection, culture, direct and indirect immunofluorescence, and nucleic acid amplification reactions, such as RT-PCR, which can detect a single agent (monoplex) or perform multiple detections (multiplex). The sensitivity of the latter is higher, and it was used in most recent studies.^{18,19} Immunofluorescence reactions have lower cost, are faster to perform, and are also able to detect multiple viruses. A panel of seven viruses (IFlu A and B, PFlu 1 to 3, hAdV, and RSV) is generally used. Some viruses, such as HRV and *Bocavirus*, can only be detected through nucleic acid amplification reactions.¹⁷

Virus and exacerbation

Several authors have performed studies aiming to detect viruses in respiratory secretions of exacerbated asthma patients, showing a prevalence of viral identification that varies with several factors, such as patient age, time of the year, method of sample collection, and method of viral detection. Table 2 presents the original articles that demonstrate results of viral testing in children with exacerbated asthma. Most studies found prevalence rates between 36.0%²⁰ and 92.2%;⁵ in these cases, the most frequently identified virus was HRV.

An investigation of respiratory viruses in 209 children aged between three and 16 years hospitalized for asthma exacerbation was performed over a period of 12 months in Buenos Aires, Argentina. Due to the possibility of other diagnoses for wheezing episodes in infants, the definition

Table 1 Biological characteristics and taxonomy of the major respiratory viruses.

	Influenza	Parainfluenza	Syncytial	Adenovirus	Rhinovirus	Coronavirus	Metapneumovirus	Bocavirus
Family	Orthomyxoviridae	Paramyxoviridae	Paramyxoviridae	Adenoviridae	Picornaviridae	Coronaviridae	Paramyxoviridae	Parvoviridae
Core of nucleic acid	RNA	RNA	RNA	DNA	RNA	RNA	RNA	DNA
Symmetry of the capsid	Helical	Helical	Helical	Icosahedral	Icosahedral	Pleomorphic	Pleomorphic	Icosahedral
Viral envelope	Enveloped	Enveloped	Enveloped	Naked	Naked	Enveloped	Enveloped	Naked
Sensitivity to ether	Sensitive	Sensitive	Sensitive	Resistant	Resistant	Sensitive	Sensitive	Sensitive
Type of nucleic acid	Single strand negative	Single negative	Single negative	Double	Single positive	Single positive	Single negative	Single negative
Nucleic acid size (kb/kbp)	10 to 13.6	15	15	40	7.2	27 to 31.5	13	5
Particle size (nm)	80 to 120nm	150 to 300	150 to 300	70 to 90	28 to 30	100 to 160	150 to 600	22 to 28

Table 2 Studies that investigated viral infection in exacerbated asthmatic children included in the literature review.

Author, year, country	n	Age	Control	Collection time	Collection and identification methods	Total detection	More frequently identified virus	Other identified viruses	Observations
Khetsuriani et al., 2007, ²⁸ USA	65 cases 77 controls	≥ 2 years	Stable asthma	March/2003 to February/2004	Nasopharyngeal swab PCR (RT-PCR) for hRV	37%	hRV (37% of total, with 60% in cases and 18% in controls)	Not researched	Association of asthma exacerbation with hRV infection genogroup C
Lopez Perez et al., 2009, ²² Mexico	100 cases 68 controls	2 to 17 years	Wheezers	Winter (April to June)	Swab DIF	Identification in 75% of cases and 44% of controls	FLUVA and hRSV in wheezers; hAdV, FLUVB and PF in asthma patients	hRSV, hAdV, FLUV	Co-detection in 17%
Olenec et al., 2010, ³⁰ Korea	58 cases	6 to 8 years	No	April and September/2006, 2007 and 2008 (501 collected samples)	NPA multiplex PCR; hRV sequencing		hRV (72 to 99%) hRV C in 16 cases	hRSV A and B, PF 1-4, hAdV, hCoV, enterovirus and hMPV, hBoV.	Virus associated with greater severity and duration; interaction between allergic sensitization and virus
Bizzintino et al., 2011, ⁴ Australia	128 cases	2 to 16 years	No	April/2003 to February/2010	NPA or swab DIF and multiplex PCR (18 primers) - nine viruses Characterization of hRV (types)	92.2%	hRV (87.5%)	hRSV, hAdV, FLUVA and B, PIV1-4, hMPV, hEV, hCoV and hBoV	hRV C was detected in most children with acute asthma (59.4%) and associated to severity
Miller et al., 2009, ²⁹ USA	1,052 cases	≤ 5 years	No	October/2001 to September/2003	NPA and swab PCR and sequencing for hRV		hRV (15.9%)	Not studied	hRV type C associated to asthma exacerbation
Chang et al., 2009, ³² Austrália	201 cases	2 to 15 years	No	March/2005 to February/2007	NPA PCR multiplex	53.8%	hRV (41%)	hRSV, hAdV, FLUV, PIV, hMPV	Co-detection in ten cases No difference in evolution for deferment viruses.

Table 2 (Continued)

Author, year, country	n	Age	Control	Collection time	Collection and identification methods	Total detection	More frequently identified virus	Other identified viruses	Observations
Ozcan et al., 2011, ²⁵ Turkey	104 cases 31 controls	3 to 17 yeras	Stable Asthma patients	12 months: September/2009 to September/2010	Swab Real-time PCR	53.8%	hRV (35.6%)	FLUVA e B, PIV 1-4, hCoV, hRSV, hRV, hMPV and hAdV	
Camara et al., 2004, ⁴³ Brazil	132 cases 65 controls	0 to 12 years	Non-asthma patients	October/1998 to June/2000	NPA Immunofluorescence (IIF); culture for hAdV and RT-PCR for hRV and hCoV	Cases (60.8%) Controls (13.3%)	RSV in those younger than 2 years hRV and hAdV in those older than 2 years	hRSV, FLUV, PIV hAdV, hCoV	There was no association between virus detection and wheezing. Allergic sensitization more associated with exacerbation. IL1, 5, 6, 10 and ECP were higher in exacerbation than in controls
Kato et al., 2011, ²⁶ Japan	174 cases 93 controls	Median 4.7 years	Stable Asthma patients Non-asthma patients	November/2003 to October/2003	NPA Kit detection of antigens and RT-PCR	79%	hRV (33.9%)	hRSV, hAdV, FLUV and PIV hEV	IL1, 5, 6, 10 and ECP were higher in exacerbation than in controls
Rawlinson et al., 2003, ¹⁹ Australia	179 cases 79 controls	1 month to 16 years	Stable Asthma patients Non-asthma patients with URTI	Winter, spring and summer of 2000 to 2002	NPA DIF PCR for hRV and hMPV Culture for virus		hRV (50% in total: 79% in cases and 17% in controls)	hAdV, FLUVA e B, hRSV, PIV3	Co-infection was common especially in winter and by hRSV
Fujitsuka et al., 2011, ²³ Japan	115	Mean 20.8 months	No	November/2007 to March/2009	Swab PCR	86.1%	hRV (31.3%)	hRSV, hAdV, PIV, FLUV, hMPV, hRV, hBoV, hEV	-
Dawood et al., 2011, ²⁷ USA	701 asthma patients in 2,165 cases of influenza	2 to 17 years	No	2003 to 2009	Swab FLUVA, B, H1N1	-	Only Influenza was screened	Not screened	Of the children hospitalized with influenza in 2003, 32% and in 2009, 44% were asthmatics. More complications with Influenza A

Table 2 (Continued)

Author, year, country	n	Age	Control	Collection time	Collection and identification methods	Total detection	More frequently identified virus	Other identified viruses	Observations
Leung et al., 2010, ³¹ China	209 cases 77 controls	3 to 18 years	Stable Asthma patients	12 months January/2007 to February/2008	NPA or swab multiplex PCR	51%	hRV	hRSV, hAdV, PIV, FluV, hMPV, hRV, hBoV, hEV	No agent was correlated with severity
Thumerelle et al., 2003, ²⁴ France	82 cases 27 controls	2 a 16 years	Stable Asthma patients	9 months October/1998 to June/1999	Swab IIF and PCR	Asthma patients (45%) and controls (3.7%)	hRV (12%)	FLUVA and B; PIV1-3; hAdV;RSV; hCoV <i>Mycoplasma</i> and chlamydia	Serology 7.4% Atypical 10%
Maffey et al., 2010, ²¹ Argentina	209 cases	3 months to 16 years	No	12 months January/2006 to December/2006	NPA or swab IIF and PCR	78%	RSV and hRV	hRSV, hAdV PIV1-3, FLUVA e B) hEV, hMPV, hBoV, mycoplasma and chlamydia	Co-detection in 20%, Seasonal circulation: three peaks: start of October (hRV); winter (hRSV) and spring (hSV, hRV and hMPV)
Mandelcwajg et al., 2010, ²⁰ France	232 cases 107 controls	1.5 to 15 years	Exacerbated asthma patients Non-hospitalized	2005 to 2009 November to March	NPA DIF and PCR	47% (hospitalized) 31% (non-hospitalized)	hBoV and RSV	FLUVA and B, hAdV, PIV-3, hMPV	hRV not screened

NPA, nasopharyngeal aspirate; IIF, indirect immunofluorescence; DIF, direct immunofluorescence; PCR, polymerase chain reaction; RT-PCR, real-time polymerase chain reaction; hRV, rhinovirus; hAdV, adenovirus; hRSV, respiratory syncytial virus; FLUV, *Influenza*; FLUVA, *Influenza A*; FLUVB, *Influenza B*; PIV1, *Parainfluenza 1*; PIV2, *Parainfluenza 2*; PIV3, *Parainfluenza 3*; hCoV, coronavirus; hMPV, metapneumovirus; hBoV, Human Bocavirus-infection; hEV, Hepatitis E virus; ECP, eosinophil cationic protein; URTI, upper respiratory tract infection.




Figure 1 Immune process involved in response to respiratory viruses and their association to allergic inflammation. Respiratory viruses infect bronchial epithelial cells (BECs) through toll-like receptors (TLRs). During replication, they trigger an inflammatory process with induction of cytokine and chemokine production by BECs, among them interferon (IFN-1), tumor necrosis factor alpha (TNF- α), interleukins (IL-33, IL-25), and thymic stromal lymphopoietin (TSLP). The dendritic cells (DCs), components of the innate immunity, are directed to secondary lymphoid organs after capturing viral antigens, where they stimulate the lymphoid cells, the protagonists of the specific immune response. In asthma patients, the production of IFNs is reduced, allowing for greater viral replication and under stimulation of TSLP, there is a deviation from lymphoid profile to helper T lymphocyte 2 (Th2), promoting lower antiviral response and increased allergic inflammation, with bronchial hyperreactivity and increased production of mucus, causing bronchial obstruction and asthma exacerbation.

of asthma was based on the criteria of Castro-Rodriguez for children younger than 3 years, and on the Global Initiative for Asthma (GINA) criteria for those older than 3. Immunofluorescence and PCR were performed in nasopharyngeal secretions of the children and showed a positive rate of 78.0% for viruses in general; hRV and hRSV were the most frequently identified types. There was also 20.0% of dual detection, with the involvement of all analyzed viruses.²¹

In México, the frequency of the viral positivity at the immunofluorescence was higher in children with asthma (75.0%) than in a control group of wheezing children without asthma (44.0%). hRV was not included in that study, and IFlu, PFlu, and hAdV were the most frequently identified virus in the group of asthma patients.²² In Japan, respiratory viruses were detected by multiplex PCR in 86.1% of 115 children with exacerbated asthma, with a mean age of 20.8 months. The hRSV was related to a single episode of wheezing ($p < 0.05$).²³ hRV was more frequently observed in patients with a history of asthma ($p < 0.05$).

A group of 82 French children with exacerbation treated at home was compared to 27 stable asthmatic children. Immunofluorescence, PCR, and serology for viruses (*Mycoplasma pneumoniae* and *Chlamydomphila pneumoniae*) detected a pathogen in 45.0% of samples, with significantly higher frequency in cases than in controls (3.7%). Viral detection tests were positive in 38% of cases, and hRV was

the most common (12.0%). In 10.0% of cases, the serologic tests were positive for both atypical pathogens.²⁴

Another series of 104 children with exacerbation, compared to 31 stable children, was studied by Turkish authors and showed positivity of 53.8% in the cases and 22.6% in controls, through RT-PCR reaction. hRV was the most commonly found virus in 35.6% of the samples.²⁵ In Japan, 174 children with acute asthma were compared to 79 stable asthmatic children and 14 children without asthma. Using an antigen detection kit and RT-PCR, respiratory viruses were detected in 79.0% of nasal aspirate samples in exacerbated asthmatic children, and hRV was the most common (33.9%). In parallel, the assessment of inflammatory markers showed a significant elevation ($p < 0.01$) of interleukins IL-1, 5, 6, and 10 in serum and in the nasal aspirates of patients in exacerbation, as well as an increase in serum eosinophilic cationic protein (ECP) levels ($p < 0.01$).²⁶

Flu, although less frequently associated with these episodes, appears to be responsible for increased morbidity in patients with an underlying chronic disease, including asthma. Of 2,165 children aged 2 to 17 years admitted with a diagnosis of infections by FLUV-A and B between 2003 and 2009 in the United States, 44.0% were asthma patients, and complications were more significantly associated with FLUV-A ($p < 0.01$). Other viruses were not assessed in that population.²⁷

Another study compared exacerbated children treated in hospitals (n=232) with those treated at home (n=107). Immunofluorescence for Flu, hAdV, hRSV, and PIV was performed, as well as PCR for *Bocavirus*. A 36.0% rate of viral detection was obtained, but no difference was observed regarding the viral profile between inpatients and outpatients. The most frequently observed viruses were RSV (15.0%) and *Bocavirus* (12.0%), but hRV was not included in the viral panel of this study.²⁰

A group of 179 Australian children aged up to 16 years had their nasal secretions collected in three periods between 2000 and 2002, and were compared with a control group of non-asthmatic children with upper respiratory tract infection (URTI) in the same period and another group of 28 children with controlled asthma, assessed during routine consultations. hRV and hMPV were screened by RT-PCR and a panel of seven viruses (FLUV- A and B, PIV- 1 to 3, hAdV, and hRSV) was studied by immunofluorescence. hRV infection accounted for 50.0% of the URTI of non-asthmatic children, and co-infection was common, especially with the hRSV, especially in children younger than 2 years.

Children with symptomatic asthma had the highest rates of hRV infection (79.0% vs. 52.0% among all children). Finally, children with controlled asthma had the lowest rates of hRV identification (17.0% vs. 79.0%).¹⁹

Studies conducted in 2007²⁸ and 2009²⁹ aimed to the identification of hRV in exacerbated asthmatic patients through RT-PCR, found an overall frequency of viral identification of 37.0% and 15.9%, respectively. The first study used a group of comparison consisting of stable asthma patients, in which the identification rate was lower (18.0%) than in the case group (60.0%). Both studies found a greater association between exacerbations and the presence of hRV C.

Virus and exacerbation severity

Another important issue in the complex association between viruses and asthma is related to the intensity of the association of exacerbations with viral infection. In this sense, several studies^{4,30-34} presented inconclusive results, although hRV was associated with increased severity or worse response to treatment.^{30,33,34} The association between viral infection and acute asthma severity was evaluated in 128 children aged 2 to 16 years. A positivity rate of 92.2% for the presence of virus was observed by direct immunofluorescence (DIF) and multiplex PCR; hRV was detected in 87.5% of cases, and type C was observed in half the cases and was associated with greater severity.⁴

Fifty-eight asthmatic children aged 6 to 8 years were monitored for a period of five weeks between April and September of 2009. They had nasal lavage samples collected weekly for multiplex PCR analysis, in addition to a symptom diary, peak expiratory flow, and notes on rescue medication use. A virus was detected in 36.0% to 50.0% of the specimens; hRV was identified in 72.0% to 99.0% of the positive samples, and was associated with greater symptom severity.³⁰

Nonetheless, viral testing by multiplex PCR for 20 pathogens in 209 children with exacerbated asthma compared with 77 controlled asthma patients, performed in Hong Kong between 2007 and 2008, showed no association between the presence of the virus and exacerbation

severity. One virus was identified in 51.0% of cases, and this detection was, in general, more associated with exacerbations (OR 2.77; 95% CI: 1:51 to 5:11; $p < 0.01$). When analyzed individually, no virus was associated with exacerbation, although hRV was the most frequent, being identified in 26.2% of exacerbated and in 13.0% of controlled asthma patients, but with no significant difference ($p = 0.27$).³¹

Nasopharyngeal aspirate samples of 201 asthmatic children aged between 2 and 15 years collected during episodes of exacerbation were referred for viral identification by PCR. The positivity rate was 53.8%; the most frequently observed were hRV (41.0%), followed by hRSV (9.0%). There was no association with exacerbation severity.³²

A study compared the response to treatment with bronchodilators between exacerbated children with viral respiratory infection symptoms (n=168) and a group without such symptoms (n=50). The mean age was 6.6 years, and exacerbation severity did not differ between groups. Children with viral symptoms had poor response to bronchodilators, requiring more doses of beta-agonists after 6, 12, and 24 hours. The viral screening was conducted in 77.0% of cases; hRV was the most frequently found virus (61.4%).³³

In another study, 78 exacerbated children were treated at the hospital and compared to 78 asymptomatic adults. Multiplex PCR reactions for eight respiratory viruses and monoplex PCT for *Enterovirus*, hRV, and *Bocavirus* detected the presence of respiratory viruses in 65.4% of cases; hRV was once again the most frequently observed virus (52.6%). Genotyping showed a higher frequency (56.0%) of type C hRV and association with type A showed a worse clinical outcome.³⁴

Seasonality of exacerbations

Asthma exacerbations have seasonal distribution, occurring cyclically in both adults and children, and can be explained by the viral circulation pattern or change in the level of pollutants and aeroallergens. One example is what occurs in temperate countries, where a higher rate of occurrence is more likely to be observed in the fall and spring among schoolchildren.³⁵ A combination of factors may explain this phenomenon, such as increased circulation of hRV in late summer and early autumn, increased circulation of pollutants and aeroallergens, and the return to school after the summer vacations. The influence of the return to school activities may be explained by lower adherence to maintenance treatment during the vacation period. The circulation of other viruses has been reported in other countries in the northern hemisphere, especially hRSV during autumn-winter, Flu in winter, PIV- 1 and 2 in the fall, and PIV- 3 in the spring.^{36,37}

In Brazil, data on viral circulation were collected from the Brazilian system of epidemiological surveillance on Flu viruses and their counterparts in the period between 2000 and 2010. Samples obtained from nasopharyngeal swabs of patients in different sentinel units distributed throughout the country were analyzed by immunofluorescence. They showed a predominance of FLUV and hRSV, with circulation throughout the year, with peaks for the latter between March and June, and between May and August for FLUV.³⁸ No data were located concerning the movement of hRV in Brazil.

Few published data regarding the seasonality of exacerbations were found. To make a parallel to virus circulation and the occurrence of exacerbations, the authors analyzed data obtained in some studies, such as the study conducted in the Federal District, which observed a higher frequency in the month of March.³⁹ Still in the Midwest region, in the state of Goiás, an increased frequency of respiratory symptoms, not specified as asthma, was observed in winter.⁴⁰ An observation regarding the distribution of the occurrence of asthma in the state of Minas Gerais also showed higher concentrations in fall-winter, between May and July,⁴¹ indicating a predominance of respiratory and/or asthma symptoms in the Brazilian fall-winter seasons.

Viral infection and allergic sensitization

In addition to the seasonal variation of the virus, other factors involved in the genesis of asthma exacerbation may explain this variation, such as aeroallergens and pollutants, which also vary throughout the different seasons. It is likely that the combination of these and other factors result in the observed seasonal peaks in exacerbations. In the month of April of the years 2006 and 2008, a study was conducted in Korea aiming to monitor viral infection and to identify sensitization to aeroallergens in 58 children with acute asthma or diagnosis of a cold, whose mean age was 6.5 years. Children with allergic sensitization presented the same number of viral infections, but with more symptoms than those non-sensitized.³⁰

In another study, conducted in Manchester, England, 84 children hospitalized for exacerbation were compared to children with stable asthma and children hospitalized for non-respiratory disease. The authors concluded that the association between viral infection and allergen exposure increased the risk of hospital admission by 19.4-fold.⁴²

In Brazil, Camara et al.⁴³ investigated the role of viral infections, sensitization, and exposure to aeroallergens as risk factors for wheezing in children aged up to 12 years. In those younger than 2 years, the frequency of viral positivity was significantly higher in cases (60.8%) than in controls (13.3%). In older children, there was no significant difference: 69.7% of cases and 43.4% of the positive controls. They concluded that in children younger than 2 years, the risk factors associated with wheezing were viral infection and a family history of atopy; among older children, sensitization to inhalant allergens was the most important event for the onset of crises.

Pollutants and aeroallergens

The effect of air pollutants is usually disregarded in the presence of viruses or allergens. However, there is evidence that acute exposure to specific pollutants may contribute to the symptoms and severity of exacerbations. For instance, cigarette smoke induces a model of non-eosinophilic inflammation with relative resistance to corticosteroids.⁴⁴ Passive smoking is quite common in homes of asthmatic children, causing a negative impact on disease control. In Scotland, the 2006 legislation that banned smoking in public places reduced hospitalizations for asthma by 18.2%.⁴⁵

Other pollutants appear to contribute to asthma exacerbations, such as those resulting from the combustion of natural gas and engine oil, such as nitrogen dioxide (NO₂). Children spend most of their time outside and breathe in a greater amount of pollutants per kilogram of weight when compared to adults, and the increased levels of NO₂ are associated with the severity of virus-induced exacerbations. This emphasizes a potential synergism between these two inflammatory stimuli.⁴⁴

Moreover, controlled exposure in asthma patients demonstrated that NO₂ increases the response to inhaled allergens. A cohort of 114 asthmatic children aged between 8 and 11 years were monitored for symptoms, measurement of peak expiratory flow, measurement of exposure to NO₂, and presence of virus in nasal secretion during a period of 13 months. One or more viruses were detected in 78% of the reported episodes of respiratory symptoms; it was demonstrated that exposure to high concentrations of NO₂ in the week before the onset of a viral respiratory infection was related to the exacerbation severity.⁴⁶

A longitudinal study conducted in the United States measured exposure to cigarette smoke in 1,444 children with asthma and NO₂ in a subset of 663 of them, over a period of nine months. They demonstrated increased symptoms in those exposed to NO₂, but only among non-atopic children, with a relative risk of 1.8 (95% CI: 1.1 to 2.8). There was no association between symptoms and increased cigarette smoke exposure.⁴⁷ Two cross-sectional studies compared children exposed to different levels of cigarette smoke and showed that those exposed to high levels had higher symptom scores ($p < 0.01$), nocturnal symptoms (OR 3.4; 95% CI: 1.3 to 8.8), and need for relief ($p = 0.03$) and control ($p = 0.02$) medications.⁴⁸

A study in which 937 children aged between 5 and 11 years were randomized to intervention with environmental education guidelines aimed at reducing exposure, showed a reduction in exposure in the group that received instructions for a period of 14 months. The intervention group had fewer days with symptoms ($p < 0.01$) after one year of follow-up, in addition to a decrease in the levels of dust mites (*Dermatophagoides pteronyssinus* and *Dermatophagoides farinae*) and cockroach antigens in the home environment.⁴⁹

Fungal sensitization is prevalent in children with asthma, although few studies have addressed this issue, compared to studies related to dust mites. One study demonstrated that children with a positive skin test for fungi had more days of symptoms when compared to those with negative tests (6.3 vs. 5.7 for two weeks, $p = 0.04$). During the study period, fungi were grown from the intra- and extra-domestic environment; increased exposure to fungi was associated with increased days of symptoms and unscheduled physician visits for asthma.⁴⁴

Preventive and therapeutic aspects

The preponderance of virus participation among the infectious agents in exacerbations makes the indiscriminate prescription of antibiotics in this situation pointless in most cases. Previous studies suggest that chest radiography is improperly and unnecessarily used in children and adults with acute asthma treated in emergency rooms.⁵⁰ Due to

the alterations that are usually found in patients during asthmatic crises, such as hyperinsufflation, fluid extravasation, and atelectasis associated with hypoxemia,⁵¹ the misinterpretation of these findings as a sign of pneumonia is common and, consequently, unnecessary prescription of antibiotics.

A multicenter study of 734 asthmatic patients treated in emergency rooms evaluated the request for additional tests, in this case, chest radiography and blood tests. Severely ill patients, those under 1 year, and those with a comorbidity were excluded. A total of 302 (41.0%) children underwent additional tests, such as chest radiography (27.0%) and blood tests (14.0%). After excluding febrile or hypoxic patients, 32.0% were still subjected to at least one of the exams.⁵⁰ Despite the lack of Brazilian data, the routine of pediatric emergency care services in the country appears to adhere to this rule.

In order to prevent the dissemination of viral agents, due to the high capacity of viral spread through droplets and fomites, hand washing, and the use of respiratory masks are simple strategies that have been proven to be effective.⁵² Staying away from situations that favor clusters of people during periods of increased viral circulation has been recommended, although there are no studies that proved the effectiveness of this strategy.^{52,53} The use of substances such as herbal preparations including *Echinacea* and vitamin C has been evaluated, but double-blind, placebo-controlled studies failed to demonstrate their benefit.⁵⁴

The prevention of viral infections through vaccines has been the most effective way to control diseases caused by viruses. In the case of respiratory viruses, the only vaccine available is for Flu, although there are ongoing studies for the development of vaccines for other respiratory viruses, especially HRV. However, their great antigenic diversity hinders research success; recent studies have tried to establish a more adequate antigenic target in the viral structure.^{54,55} Specific RSV immunoglobulin has been successfully used in reducing hospitalizations for viral bronchiolitis, and new perspectives for the treatment of exacerbations triggered by viral infections have emerged from studies directed to synthetic agonists of TLR3 receptor, IFN- β agonist, and IL33- and IL25-antagonists, among others.^{56,57}

There is no specific treatment for most respiratory viruses. Some antivirals have been successfully used, as in the case of Flu infection, such as amantadine, rimantadine, oseltamivir phosphate, and zanamivir; the latter is not indicated for patients with asthma. Ribavirin is indicated for the treatment of severe infections caused by RSV. Other antiviral agents are being studied and have not yet been approved for clinical use, such as pleconaril, vapendavir, pirodavir, and rupintrivir.⁵⁴

Glucocorticoids have potent anti-inflammatory effects and have been successfully used in maintenance treatment in patients with persistent asthma, controlling inflammation and preventing exacerbations. Some studies have assessed its effect on virus-induced asthma. The suppression of the release of pro-inflammatory mediators induced by HRV infection *in vitro* in bronchial epithelial cells, such as CCL5, CCL10, CXCL8, and IL6, as well as the reduction of factors associated with remodeling, was achieved after the use of budesonide.⁵⁸ Other *in vitro* studies documented the action of other corticosteroids alone or in combination with bronchodilators or leukotriene antagonists in reducing

the release of several inflammatory molecules, with potential modulation of the deleterious effects of viruses on the asthmatic population.^{51,59} Despite the proven benefits of inhaled corticosteroids in the control of asthma triggered by multiple factors, their action on virus-induced exacerbations is unclear. The use of low-to-moderate doses of inhaled corticosteroids as maintenance therapy cannot prevent intermittent viral-induced wheezing.^{60,61}

However, better results have been obtained with the intermittent use of inhaled corticosteroids at high doses.⁶²

Discussion

The use of viral detection techniques with high sensitivity and specificity has increased the identification of some respiratory viruses in children with asthma exacerbation. The direct or indirect immunofluorescence reactions still have great practical importance, as they can detect a panel of seven viruses (FLUV- A and B, PIV- 1-3, hAdV and hRSV), being an affordable and fast method, with good sensitivity, especially in children.¹⁸ It is currently used by the Brazilian Ministry of Health for the screening of respiratory viruses in the diagnosis of severe acute flu-like illness and Flu-like illness in sentinel units.

The techniques for nucleic acid amplification (RT-PCR) are more expensive, but more sensitive; thus, they are used in research and by the Brazilian Ministry of Health for the identification and genotyping of Flu.³⁸ Furthermore, it allows for the identification of some viruses with significant clinical and epidemiological importance, such as hRV and *Bocavirus*, not identified by immunofluorescence.^{17,54}

As for the method used to obtain the sample, it is worth mentioning the controversial issue of nasopharyngeal swab in viral research. Although its use has been consolidated for bacterial infections (*S. pneumoniae* and *S. aureus*), its role in viral infections still deserves some consideration. The authors agree that, from the practical point of view, it is more feasible, eliminating the use of suction systems, probes, and more specialized training, when compared to aspirate or lavage samples. However, only those with more advanced technology (flocked swab), which provides best capture and release of cells and, therefore, of the virus, are equivalent to the aspirate in terms of sample quality. Nevertheless, this swab is not routinely used in services and researches in Brazil.³⁸

Regarding the association between viral infection and asthma exacerbation, a wide variation was observed concerning the methods of studies that assessed viral infection in exacerbated asthmatic children in the studies included in this review. For instance, sample size varied from 58 to 1,052 cases and the age ranged from 0 to 17 years. This finding is important, given the difficulty in defining asthma in children younger than 3 years, which should be considered in case selection.^{2,21} Moreover, it is known that there is a considerable difference between the age groups and the most prevalent viruses, such as the hRSV.⁶⁰ Regarding the methods of respiratory secretion collection in the included studies, there was no uniform means of collection; the aspirate was used in 43.8% of studies, the swab in 31.6%, both in 25%, and the flocked swab was not used. There was a wide

variation regarding the detection methods and in relation to some outcomes.

In addition to the differences in sample collections, all these studies were cross-sectional, which does not allow for the establishment of a cause-effect association between viral infection and the onset of exacerbation, but suggest such an association. In relation to other factors known to be associated with uncontrolled asthma, such as allergens and irritants, most studies did not include these variables in the evaluation. When the inflammatory process typical of asthma is associated with a viral respiratory infection, there is a tendency to greater severity and duration, as well as a poorer response to conventional treatment of the acute episode.^{32,33} The involved mechanisms still need to be fully elucidated, evidencing the synergistic effect between viral infection and allergic airway inflammation in the pathogenesis of exacerbations.^{30,43}

Another pertinent issue is the role of inhaled corticosteroids in attenuating the inflammation triggered by the virus, also seldom mentioned in these studies. Its action in the control and reduction of morbidity associated with asthma is well established,² but it is still a controversial subject regarding the prevention of viral-induced wheezing. Its effectiveness in the inflammatory process triggered by a virus has been demonstrated in *in vitro* studies,^{51,58,59} but studies evaluating its clinical benefit have yet to reach conclusive results.^{61,62}

Regardless of the direction of virus-allergen interaction, the present findings strongly suggest that an adequate strategy to prevent virus-induced exacerbations should focus on two courses, namely the improvement of antiviral response and the reduction of allergic sensitization or inflammation. The latter can be achieved with appropriate treatment of the asthma patient at risk with medications that reduce airway inflammation. Conversely, the preventive measures for viral infection acquisition and its timely diagnosis allow for a proper management of exacerbations, and reduction of the number of hospitalizations and unnecessary additional tests, especially in children who are febrile at the time of assessment.

The association between viral infection and asthma in childhood still has several points that need clarification, especially the actual role of viruses in triggering exacerbations and that of inhaled corticosteroids in its attenuation.

Funding

Fundação de Amparo à Pesquisa do Estado de Goiás (N° 20120267001128).

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

To the Laboratory of Virology of the Instituto de Patologia Tropical e Saúde Pública of the Universidade Federal de Goiás.

References

1. Solé D, Wandalsen GF, Camelo-Nunes IC, Naspitz CK. Prevalência de sintomas de asma, rinite e eczema atópico entre crianças e adolescentes brasileiros identificados pelo International Study of Asthma and Allergies (ISAAC): fase 3. *J Pediatr* (Rio J). 2006;82:341–6.
2. Global Strategy for Asthma Management and Prevention. Global Initiative for Asthma (GINA) 2012. Available from: <http://www.ginasthma.org/>
3. Gregg I, Horn M, Brain E, Inglis JM, Yealland SJ. Proceedings: The role of viral infection in asthma and bronchitis. *Tubercle*. 1975;56:171.
4. Bizzintino J, Lee WM, Laing IA, Vang F, Pappas T, Zhang G, et al. Association between human rhinovirus C and severity of acute asthma in children. *Eur Respir J*. 2011;37:1037–42.
5. Butel JS. Virologia. In: Brooks GF, Carroll KC, Butel JS, Morse SA, Mietzner TA, editors. *Microbiologia médica de Jawetz, Melnick e Adelberg*. 25th ed. Porto Alegre: McGraw-Hill; 2010. p. 373–93.
6. Contoli M, Message S, Laza-Stanca V, Edwards MR, Wark PA, Bartlett NW, et al. Role of deficient type III interferon-lambda production in asthma exacerbations. *Nat Med*. 2006;12:1023–6.
7. Holgate ST. Epithelium dysfunction in asthma. *J Allergy Clin Immunol*. 2007;120:1233–44.
8. Hashimoto K, Graham BS, Ho SB, Adler KB, Collins RD, Olson SJ, et al. Respiratory syncytial virus in allergic lung inflammation increases Muc5ac and Gob-5. *Am J Respir Crit Care Med*. 2004;170:306–12.
9. Papadopoulos NG, Christodoulou I, Rohde G, Agache I, Almqvist C, Bruno A, et al. Viruses and bacteria in acute asthma exacerbations—a GA(2) LEN-DARE systematic review. *Allergy*. 2011;66:458–68.
10. Baraldo S, Contoli M, Bazzan E, Turato G, Padovani A, Marku B, et al. Deficient antiviral immune responses in childhood: distinct roles of atopy and asthma. *J Allergy Clin Immunol*. 2012;130:1307–14.
11. Gavala M, Bashir H, Gern J. Virus/allergen interactions in asthma. *Curr Allergy Asthma Rep*. 2013;13:298–307.
12. Holt P, Sly P. Interaction between adaptive and innate immune pathways in the pathogenesis of atopic asthma: operation of a lung/bone marrow axis. *Chest*. 2011;139:1165–71.
13. Hams E, Fallon PG. Innate type 2 cells and asthma. *Curr Opin Pharmacol*. 2012;12:503–9.
14. Chan KH, Peiris JS, Lim W, Nicholls JM, Chiu SS. Comparison of nasopharyngeal flocked swabs and aspirates for rapid diagnosis of respiratory viruses in children. *J Clin Virol*. 2008;42:65–9.
15. Sung RYT, Chan PKS, Choi KC, Yeung AC, Li AM, Tang JW, et al. Comparative study of nasopharyngeal aspirate and nasal swab specimens for diagnosis of acute viral respiratory infection. *J Clin Microbiol*. 2008;46:3073–6.
16. Blaschke AJ, Allison MA, Meyers L, Rogatcheva M, Heyrend C, Mallin B, et al. Non-invasive sample collection for respiratory virus testing by multiplex PCR. *J Clin Virol*. 2011;52:210–4.
17. Ginocchio CC, McAdam AJ. Current best practices for respiratory virus testing. *J Clin Microbiol*. 2011;49:544–8.
18. She RC, Polage CR, Caram LB, Taggart EW, Hymas WC, Woods CW, et al. Performance of diagnostic tests to detect respiratory viruses in older adults. *Diagn Microbiol Infect Dis*. 2010;67:246–50.
19. Rawlinson WD, Waliuzzaman Z, Carter IW, Belessis YC, Gilbert KM, Morton JR. Asthma exacerbations in children associated with *Rhinovirus* but not human metapneumovirus infection. *J Infect Dis*. 2003;187:1314–8.
20. Mandelcwaig A, Moulin F, Menager C, Rozenberg F, Lebon P, Gendrel D. Underestimation of influenza viral infection in childhood asthma exacerbations. *J Pediatr*. 2010;157:505–6.

21. Maffey AF, Barrero PR, Venialgo C, Fernández F, Fuse VA, Saia M, et al. Viruses and atypical bacteria associated with asthma exacerbations in hospitalized children. *Pediatr Pulmonol.* 2010;45:619–25.
22. Lopez Perez G, Morfin Maciel BM, Navarrete N, Aguirre A. Identification of influenza, parainfluenza, adenovirus and respiratory syncytial virus during rhinopharyngitis in a group of Mexican children with asthma and wheezing. *Rev Alerg Mex.* 2009;56:86–91.
23. Fujitsuka A, Tsukagoshi H, Arakawa M, Goto-Sugai K, Ryo A, Okayama Y, et al. A molecular epidemiological study of respiratory viruses detected in Japanese children with acute wheezing illness. *BMC Infect Dis.* 2011;11:168.
24. Thumerelle C, Deschildre A, Bouquillon C, Santos C, Sardet A, Scalbert M, et al. Role of viruses and atypical bacteria in exacerbations of asthma in hospitalized children: a prospective study in the Nord-Pas de Calais region (France). *Pediatr Pulmonol.* 2003;35:75–82.
25. Ozcan C, Toyran M, Civelek E, Erkoçoğlu M, Altaş AB, Albayrak N, et al. Evaluation of respiratory viral pathogens in acute asthma exacerbations during childhood. *J Asthma.* 2011;48:888–93.
26. Kato M, Tsukagoshi H, Yoshizumi M, Saitoh M, Kozawa K, Yamada Y, et al. Different cytokine profile and eosinophil activation are involved in rhinovirus- and RS virus-induced acute exacerbation of childhood wheezing. *Pediatr Allergy Immunol.* 2011;22:e87–94.
27. Dawood FS, Kamimoto L, D’Mello TA, Reingold A, Gershman K, Meek J, et al. Children with asthma hospitalized with seasonal or pandemic *Influenza*, 2003-2009. *Pediatrics.* 2011;128:e27–32.
28. Khetsuriani N, Kazerouni NN, Erdman DD, Lu X, Redd SC, Anderson LJ, et al. Prevalence of viral respiratory tract infections in children with asthma. *J Allergy Clin Immunol.* 2007;119:314–21.
29. Miller EK, Edwards KM, Weinberg GA, Iwane MK, Griffin MR, Hall CB, et al. A novel group of rhinoviruses is associated with asthma hospitalizations. *J Allergy Clin Immunol.* 2009;123:98–104.
30. Olenec JP, Kim WK, Lee W-M, Vang F, Pappas TE, Salazar LE, et al. Weekly monitoring of children with asthma for infections and illness during common cold seasons. *J Allergy Clin Immunol.* 2010;125:1001–6, e1.
31. Leung TF, To MY, Yeung ACM, Wong YS, Wong GWK, Chan PKS. Multiplex molecular detection of respiratory pathogens in children with asthma exacerbation. *Chest.* 2010;137:348–54.
32. Chang AB, Clark R, Acworth JP, Petsky HL, Sloots TP. The impact of viral respiratory infection on the severity and recovery from an asthma exacerbation. *Pediatr Infect Dis J.* 2009;28:290–4.
33. Rueter K, Bizzintino J, Martin AC, Zhang G, Hayden CM, Geelhoed GC, et al. Symptomatic viral infection is associated with impaired response to treatment in children with acute asthma. *J Pediatr.* 2012;160:82–7.
34. Arden KE, Chang AB, Lambert SB, Nissen MD, Sloots TP, Mackay IM. Newly identified respiratory viruses in children with asthma exacerbation not requiring admission to hospital. *J Med Virol.* 2010;82:1458–61.
35. Sears MR, Johnston NW. Understanding the September asthma epidemic. *J Allergy Clin Immunol.* 2007;120:526–9.
36. Litwin CM, Bosley JG. Seasonality and prevalence of respiratory pathogens detected by multiplex PCR at a tertiary care medical center. *Arch Virol.* 2014;65–72.
37. Johnston SL. Innate immunity in the pathogenesis of virus-induced asthma exacerbations. *Proc Am Thorac Soc.* 2007;4:267–70.
38. de Mello Freitas FT. Sentinel surveillance of influenza and other respiratory viruses, Brazil, 2000-2010. *Braz J Infect Dis.* 2013;17:62–8.
39. Valença LM, Restivo PCN, Nunes MS. Variação sazonal nos atendimentos de emergência por asma em Gama. *Brasil J Bras Pneumol.* 2006;32:284–9.
40. Silva Junior JL, Padilha TF, Rezende JE, Rabelo EC, Ferreira AC, Rabahi MF. Effect of seasonality on the occurrence of respiratory symptoms in a Brazilian city with a tropical climate. *J Bras Pneumol.* 2011;37:759–67.
41. Ezequiel Oda S, Gazeta GS, Freire NM. Prevalência dos atendimentos por crises de asma nos serviços públicos do Município de Juiz de Fora (MG). *J Bras Pneumol.* 2007;33:20–7.
42. Murray CS, Poletti G, Kebabdzé T, Morris J, Woodcock A, Johnston SL, et al. Study of modifiable risk factors for asthma exacerbations: virus infection and allergen exposure increase the risk of asthma hospital admissions in children. *Thorax.* 2006;61:376–82.
43. Camara AA, Silva JM, Ferriani VPL, Tobias KR, Macedo IS, Padovani MA, et al. Risk factors for wheezing in a subtropical environment: Role of respiratory viruses and allergen sensitization. *J Allergy Clin Immunol.* 2004;113:551–7.
44. Dick S, Doust E, Cowie H, Ayres JG, Turner S. Associations between environmental exposures and asthma control and exacerbations in young children: a systematic review. *BMJ open.* 2014;4:e003827.
45. Mackay D, Haw S, Ayres JG, Fischbacher C, Pell JP. Smoke-free legislation and hospitalizations for childhood asthma. *N Engl J Med.* 2010;363:1139–45.
46. Chauhan AJ, Inskip HM, Linaker CH, Smith S, Schreiber J, Johnston SL, et al. Personal exposure to nitrogen dioxide (NO₂) and the severity of virus-induced asthma in children. *Lancet.* 2003;361:1939–44.
47. Kattan M, Gergen PJ, Eggleston P, Visness CM, Mitchell HE. Health effects of indoor nitrogen dioxide and passive smoking on urban asthmatic children. *J Allergy Clin Immunol.* 2007;20:618–24.
48. Morkjaroenpong V, Rand CS, Butz AM, Huss K, Eggleston P, Malveaux FJ, et al. Environmental tobacco smoke exposure and nocturnal symptoms among inner-city children with asthma. *J Allergy Clin Immunol.* 2002;110:147–53.
49. Morgan WJ, Crain EF, Gruchalla RS, O’Connor GT, Kattan M, Evans R 3rd, et al. Results of a home-based environmental intervention among urban children with asthma. *N Engl J Med.* 2004;351:1068–80.
50. Gentile NT, Ufberg J, Barnum M, McHugh M, Karras D. Guidelines reduce X-ray and blood gas utilization in acute asthma. *Am J Emerg Med.* 2003;21:451–3.
51. Matsukura S, Kurokawa M, Homma T, Watanabe S, Suzuki S, Ieki K, et al. Basic Research on virus-induced asthma exacerbation: inhibition of inflammatory chemokine expression by fluticasone propionate. *Int Arch Allergy Immunol.* 2013;161:84–92.
52. Ahanchian H, Jones C, Chen Y-s, Sly P. Respiratory viral infections in children with asthma: do they matter and can we prevent them? *BMC Pediatr.* 2012;12:147.
53. Weedon KM, Rupp AH, Hefron AC, Kelly SF, Zheng X, Shulman ST, et al. The impact of infection control upon hospital-acquired influenza and respiratory syncytial virus. *Scand J Infect Dis.* 2013;45:297–303.
54. Jacobs SE, Lamson DM, Kirsten S, Walsh TJ. Human rhinoviruses. *Clin Microbiol Rev.* 2013;26:135–62.
55. Papi A, Contoli M. Rhinovirus vaccination: the case against. *Eur Respir J.* 2011;37:5–7.
56. Jackson DJ, Sykes A, Mallia P, Johnston SL. Asthma exacerbations: origin, effect, and prevention. *J Allergy Clin Immunol.* 2011;128:1165–74.
57. Christopher ME, Wong JP. Use of toll-like receptor 3 agonists against respiratory viral infections. *Antiinflamm Antiallergy Agents Med Chem.* 2011;10:327–38.
58. Skevaki CL, Christodoulou I, Spyridaki IS, Tiniakou I, Georgiou V, Xepapadaki P, et al. Budesonide and formoterol inhibit

- inflammatory mediator production by bronchial epithelial cells infected with rhinovirus. *Clin Exp Allergy*. 2009;39:1700–10.
59. Bochkov Y, Busse W, Brockman-Schneider R, Evans MD, Jarjour NN, McCrae C, et al. Budesonide and formoterol effects on rhinovirus replication and epithelial cell cytokine responses. *Respir Res*. 2013;14:98.
 60. Alvarez AE, Marson FAL, Bertuzzo CS, Arns CW, Ribeiro JD. Epidemiological and genetic characteristics associated with the severity of acute viral bronchiolitis by respiratory syncytial virus. *J Pediatr (Rio J)*. 2013;89:531–43.
 61. Wilson N, Sloper K, Silverman M. Effect of continuous treatment with topical corticosteroid on episodic viral wheeze in preschool children. *Arch Dis Child*. 1995;72:317–20.
 62. Ducharme FM, Lemire C, Noya FJD, Davis GM, Alos N, Leblond H, et al. Preemptive use of high-dose fluticasone for virus-induced wheezing in young children. *N Engl J Med*. 2009;360:339–53.