

Session 5 – Cardiovascular Multimodality Imaging Techniques

Friday May 29 – 10.00 – 11.00

27

Non invasive ultrasonic chordal cutting

Olivier Villemain^{*1}, M. Pernot², W. Kwiecinski², A. Bel¹, M. Tanter¹, A. Hagege¹, E. Messas¹

¹ Hôpital Européen Georges-Pompidou, Université Paris-Descartes, Cardiology, Paris, France,

² Waves and acoustics laboratory Institute Langevin, Paris, France.

* Auteur correspondant : olovillemain@gmail.com

Objective Chordal cutting targeting leaflet tethering has been described to improve the efficacy of annuloplasty during ischemic mitral regurgitation surgery. Histotripsy is a novel ultrasound based technique for tissue fragmentation through the cavitation generated at the focus of a very intense ultrasonic pulse. We investigate the feasibility of using histotripsy for chordal cutting to avoid cardiopulmonary bypass and invasive surgery in infarcted heart.

Methods Experiments were performed in vitro in explanted sheep heart (N=10) and in vivo in sheep beating heart (N=5, 40±4kg). In vitro, the mitral valve basal chordae was fixed. The ultrasound pulses were emitted from the therapeutic device (1-MHz focused transducer) placed at a distance of 64mm. In vivo, we performed sternotomy and the device was applied on the thorax cavity. We analysed MV coaptation and chordae by real time 3D echocardiography before and after chordal cutting. A postmortem anatomical exploration have been realized to confirm the section of the chordae.

Results In vitro, all the basal chordae were completely cut. The mean procedure time was 6 (±3) minutes. The thickness of the chordae was the main criteria affecting the duration of procedure. In the sheep, central basal chordae of anterior leaflet were completely cut. The mean procedure time was 19 (±9) minutes. By echography, the sectioned chordae was visible and no mitral valve prolapse was found. All the postmortem anatomical exploration of hearts confirmed the section of the basal chordae. No additional lesions were objectified.

Conclusions Noninvasive ultrasound histotripsy succeed to cut mitral valve basal chordae in vitro and in vivo in beating heart. Future investigation will be needed to test this technique on its ability to decrease ischemic or functional MR secondary to leaflet tethering. If positive, this will open the door of completely noninvasive technique for MV repair especially in case of functional MR.

28

Monitoring of LVAD implantation and set-up with miniaturized transesophageal echocardiography: Initial experience at La Pitié-Salpêtrière Hospital and possible application fields

Nadia Bouabdallaoui^{*}, Guillaume Lebreton, Pierre Demondion, Pascal Leprince

Department of thoracic and cardiovascular surgery; Pierre-et-Marie Curie University, Paris VI, Paris, France.

* Auteur correspondant : nadia.bouabdallaoui@gmail.com

Background Left ventricular assist devices (LVAD) are valuable options for mechanical circulatory support in patients with severe heart failure (HF) as a bridge to cardiac transplantation, to recovery, or as destination therapy. Transesophageal echocardiography (TEE) plays a key role in the management of LVAD patients; either in guiding the device implantation or in the post-operative hemodynamic assessment. The aim of this observational study is to investigate the safety and usefulness of a miniaturized TEE probe designed for continuous hemodynamic monitoring (hTEE, hemodynamic transesophageal echocardiography; ImaCor, Garden City, NY) in hemodynamic assessment of patients undergoing LVAD implantation.

Methods Between June 2014 and September 2014, 7 consecutive patients underwent LVAD implantation for severe HF. hTEE probe was used for continuous hemodynamic monitoring in the pre-, intra- and postoperative periods. All procedures were performed by hTEE-trained physicians.

Results All data were recorded and used without need of further hemodynamic device. The average number of hemodynamic assessment in each patient was of 5.3. The average number of hemodynamic assessment leading to therapeutic changes in each patient was 2.6. The mean duration of hTEE probe insertion was approximately of 20 hours (9 hours, 72 hours). No device complications were noticed during this period.

Conclusions This observation suggest that hemodynamic monitoring using the hTEE system in LVAD patients is feasible and provides valid data after a brief training period. The overall impact of hTEE use in these patients is yet to be determined and further studies are still required.

29

Comparison of cardiac magnetic resonance and echocardiography for the assessment of aortic valve area in aortic stenosis

Dania Mohty^{*2}, Marc-Antoine Isorni¹, Stéphan Chassaing¹, Arnaud Maudière¹, Christophe Barbey¹, Olivier Bar¹, Didier Bruchère¹, Didier Brunchar¹, Etienne Puymirat³

¹ Clinique Saint-Gatien, Cardiology Department, Tours, France,

² Centre hospitalier Universitaire de Limoges, Cardiology Department, Limoges, France,

³ HEGP, cardiology department, Paris, France.

* Auteur correspondant : dania.mohty@chu-limoges.fr

Background Aortic valve area (AVA) calculated by the continuity equation by echocardiography (CE-TTE) is the method of reference to assess aortic valve stenosis severity (AS). The Hakki's formula (simplified Gorlin formula) is another method, sometimes used during cardiac catheterization to calculate AVA. It can also be adapted to magnetic resonance imaging (CMR) to assess the stenotic AVA as previously demonstrated.

The aim of our study is to investigate using a large number of patients with AS, the accuracy of the "Hakki-CMR" method to determine the severity of AS as compared to CE-TTE and to anatomic AVA by planimetry performed by CMR.

Methods and results Between 2007 and 2014, 390 consecutive patients with AS (mean age 81±10 years, men 55%, mean LVEF= 60±13%, underwent clinically indicated TTE (Philips IE 33.) and CMR (Philips ACHIEVA 1.5 Tesla) within 30 days.

The mean aortic pressure gradient was 43±18 mmHg; the AVA was 0.63±0.25cm² by CE-TTE, 0.75±0.30cm² using Hakki-CMR and 0.65±0.18cm² by planimetry-CMR.

The mean differences in AVA between different methods were respectively:

1) 0.04cm² (95% limits of agreement -0.15 to 0.21, p<0.0001) between Hakki-CMR and CE-TTE, 2) 0.04cm² (95% limits of agreement -0.17 to 0.23, p=0.0002), between planimetry-CMR and the CE-TTE, 3) 0.01cm² (95% limits of agreement -0.16 to 0.14, p<0.0001) between Hakki-CMR and planimetry.

Conclusion In a large cohort of patients with AS, assessment of AVA using the Hakki's formula by CMR, yielded more reliable results than those obtained using planimetry by CMR, and similar to those obtained using the CE-TTE. Despite the time and cost issue of CMR, the Hakki formula used during flow quantification, is a valuable method that is reliable easy, and fast to apply. It may be highly useful especially in cases of suboptimal echogenicity by TTE and regardless of the valve's degree of calcification.

30

Pocket-size ultrasound does not alter the diagnosis of the cardiologist in patients admitted for chest pain in a cardiac unit

L. Soulat-Dufour, S. Lang, A. Pasteur Rousseau, C. Poulizac, C. Hermand, C. El Asri, L. Boyer-Chatenet, S. Adavane, S. Ederhy, F. Boccarda, D. Pateron, Ariel Cohen^{*}

* Hôpital Saint-Antoine, Paris, France.

* Auteur correspondant : ariel.cohen@sat.aphp.fr

Purpose Chest pain is a very common cause of hospitalization in cardiology departments and is associated with a large variety of cardiac disease.