

Lin28A and Lin28B Inhibit let-7 MicroRNA Biogenesis by Distinct Mechanisms

Elena Piskounova,^{1,2,4} Christos Polytaichou,^{3,5} James E. Thornton,^{1,2,4} Robert J. LaPierre,^{1,2,4} Charalabos Pothoulakis,⁶ John P. Hagan,^{1,2,4,7} Dimitrios Iliopoulos,^{3,5,*} and Richard I. Gregory^{1,2,4,*}

¹Stem Cell Program, Children's Hospital, Boston, MA 02115, USA

²Department of Biological Chemistry and Molecular Pharmacology

³Department of Microbiology and Immunobiology

Harvard Medical School, Boston, MA 02115, USA

⁴Harvard Stem Cell Institute, Boston, MA 02115, USA

⁵Department of Cancer Immunology and AIDS, Dana-Farber Cancer Institute, Boston, MA 02115, USA

⁶Inflammatory Bowel Disease Center, Division of Digestive Diseases, David Geffen School of Medicine at UCLA, Los Angeles, CA 90095, USA

⁷Department of Molecular Virology, Immunology, and Medical Genetics, The Ohio State University Medical Center, Columbus, OH 43210, USA

*Correspondence: dimitrios_iliopoulos@dfci.harvard.edu (D.I.), rgregory@enders.tch.harvard.edu (R.I.G.)

DOI 10.1016/j.cell.2011.10.039

SUMMARY

Lin28A and Lin28B selectively block the expression of let-7 microRNAs and function as oncogenes in a variety of human cancers. Lin28A recruits a TUTase (Zcchc11/TUT4) to let-7 precursors to block processing by Dicer in the cell cytoplasm. Here we find that unlike Lin28A, Lin28B represses let-7 processing through a Zcchc11-independent mechanism. Lin28B functions in the nucleus by sequestering primary let-7 transcripts and inhibiting their processing by the Microprocessor. The inhibitory effects of Zcchc11 depletion on the tumorigenic capacity and metastatic potential of human cancer cells and xenografts are restricted to Lin28A-expressing tumors. Furthermore, the majority of human colon and breast tumors analyzed exclusively express either Lin28A or Lin28B. Lin28A is expressed in HER2-overexpressing breast tumors, whereas Lin28B expression characterizes triple-negative breast tumors. Overall our results illuminate the distinct mechanisms by which Lin28A and Lin28B function and have implications for the development of new strategies for cancer therapy.

INTRODUCTION

Control of gene expression by microRNAs (miRNAs) is important for normal development. Altered miRNA expression is linked with various diseases including cancer (Small and Olson, 2011). miRNA biogenesis begins with transcription of primary transcripts (pri-miRNAs) that contain a stem-loop structure. In the cell nucleus, pri-miRNAs are processed by the Microprocessor, containing the ribonuclease Drosha and its essential cofactor DGCR8 (Denli et al., 2004; Gregory et al., 2004). The Microprocessor cleaves the double-stranded RNA toward the base of

the stem loop to release a 60–80 nucleotide (nt) precursor (pre-miRNA) that is exported to the cell cytoplasm and cleaved by Dicer to generate a 22 nt duplex (Hutvagner et al., 2001; Krol et al., 2010). One RNA strand is bound by Argonaute and incorporated into the RNA-induced silencing complex (RISC) (Gregory et al., 2005; Liu et al., 2004). Basepairing between the miRNA and target mRNA guides RISC to complementary transcripts, leading to gene repression through mRNA degradation and/or translational repression (Krol et al., 2010).

Altered miRNA expression is directly associated with cancer initiation, progression, and metastasis and is observed in a wide variety of human malignancies (Di Leva and Croce, 2010). The let-7 miRNA family members act as tumor suppressors by inhibiting expression of oncogenes and key regulators of mitogenic pathways including RAS, MYC, and HMGA2 (Büssing et al., 2008). let-7 is downregulated in numerous different cancers, and low let-7 correlates with poor prognosis (Boyerinas et al., 2010; Shell et al., 2007; Takamizawa et al., 2004). Restoration of let-7 expression effectively inhibits cancer growth in mouse models of lung and breast cancers (Barh et al., 2010; Esquela-Kerscher et al., 2008; Slack, 2009; Trang et al., 2010; Yu et al., 2007a). In humans, there are 12 let-7 family members (let-7a-1, -2, -3; let-7b; let-7c; let-7d; let-7e; let-7f-1, -2; let-7g; let-7i; miR-98) located at eight different chromosomal loci. Of note, many tumors are characterized by the coordinate downregulation of multiple let-7 miRNAs (Shell et al., 2007).

The developmentally regulated RNA-binding protein Lin28 was found to selectively repress expression of let-7 miRNAs (Heo et al., 2008; Newman et al., 2008; Rybak et al., 2008; Viswanathan et al., 2008). This posttranscriptional regulation of let-7 by Lin28 is required for normal development and contributes to the pluripotent state by preventing let-7-mediated differentiation of embryonic stem cells (ESCs). Lin28 overexpression or let-7 inhibition with antisense RNAs promotes reprogramming of human and mouse fibroblasts to induced pluripotent stem cells (iPSCs) (Martinez and Gregory, 2010; Melton et al., 2010; Yu et al., 2007b). Unlike in *C. elegans* where a single *Lin28* gene is responsible for repression of let-7 expression and control of

developmental timing, the mammalian genome encodes two Lin28 paralogs, Lin28 (hereafter Lin28A) and Lin28B (Guo et al., 2006; Lehrbach et al., 2009; Moss et al., 1997; Van Wynsberghe et al., 2011; Viswanathan and Daley, 2010). Lin28B also represses expression of multiple let-7 members, and genome-wide association studies (GWAS) have linked *Lin28B* with the determination of human height and control of the age of onset of puberty and menopause, phenotypes that are recapitulated in a mouse model (Zhu et al., 2010). Activation of Lin28A/Lin28B occurs in several different primary human tumors, and these tumors display low levels of let-7 expression (Iliopoulos et al., 2009; Viswanathan et al., 2009). Indeed, Lin28A/Lin28B function as oncogenes that promote cellular transformation when ectopically expressed (Iliopoulos et al., 2009; Viswanathan et al., 2009; West et al., 2009). Importantly, this effect is abrogated when let-7 is reintroduced into these cells (Iliopoulos et al., 2009; Viswanathan et al., 2009). Therefore, Lin28-mediated cellular transformation is directly dependent on let-7 levels. Conversely, depletion of Lin28A or Lin28B in human cancer cells results in decreased cell proliferation (Chang et al., 2009; Iliopoulos et al., 2009; Viswanathan et al., 2009). Lin28A/Lin28B may contribute to the development of aggressive, poorly differentiated tumors because their expression is associated with advanced disease in hepatocellular carcinoma (HCC), chronic myeloid leukemia (CML), Wilms' tumor, ovarian carcinoma, colon adenocarcinoma, and germ cell tumors (Dangi-Garimella et al., 2009; Guo et al., 2006; Iliopoulos et al., 2009; Ji and Wang, 2010; King et al., 2011; Liang et al., 2010; Lu et al., 2009; Oh et al., 2010; Peng et al., 2010; Viswanathan et al., 2009; Wang et al., 2010; West et al., 2009; Yang et al., 2010) and is associated with poor clinical outcome and patient survival in HCC, colon, and ovarian cancer (King et al., 2011; Lu et al., 2009; Viswanathan et al., 2009). In the case of Lin28B, rare amplification or translocation events might explain activation in some cases (Viswanathan et al., 2009). A more common mechanism might be transcriptional activation by upstream factors. For example, c-Myc binds to both *Lin28A* and *Lin28B* loci and activates expression of these genes (Chang et al., 2009). In a breast cancer model, transient expression of Src oncoprotein results in a transformed cell line that forms self-renewing mammospheres harboring tumor-initiating cells (Iliopoulos et al., 2009). The transformation process involves NF- κ B activation leading to direct transcriptional upregulation of Lin28B, consequent let-7 loss, and derepression of the let-7 target gene IL-6. Because IL-6 activates NF- κ B, this regulatory circuit represents a positive feedback loop, providing a molecular link between inflammation and cancer.

Selective regulation of let-7 expression involves Lin28A binding to the terminal loop of let-7 precursors, a molecular recognition that requires both the cold shock domain (CSD) and CCHC-type zinc finger RNA-binding domains of the Lin28A protein (Piskounova et al., 2008). Lin28A recruits the activity of a terminal uridylyltransferase (TUTase), Zcchc11 (also known as TUTase4 or TUT4), that inhibits pre-let-7 processing by Dicer and leads to the rapid decay of oligouridylylated pre-let-7 RNAs (Hagan et al., 2009; Heo et al., 2009). Although both Lin28A and Lin28B can recruit Zcchc11/TUT4 to uridylylate pre-let-7 in vitro, the molecular mechanism of the Lin28B-mediated

blockade of let-7 expression has yet to be determined (Heo et al., 2008, 2009). Here we investigate the regulation of let-7 expression by Lin28B. Surprisingly, we find that despite their high degree of homology, Lin28A and Lin28B function through distinct mechanisms. Depletion of Zcchc11 affects let-7 expression only in Lin28A-expressing cancer cells, whereas Lin28B functions through a Zcchc11-independent mechanism. We find that Lin28A and Lin28B are differentially localized in cells with predominantly cytoplasmic Lin28A, whereas due to its functional nuclear localization signals, Lin28B accumulates in the nucleus, where it binds pri-let-7 miRNAs to block processing by the Microprocessor. In contrast, Lin28A functions in the cytoplasm by blocking at the Dicer step and recruiting the TUTase to uridylylate pre-let-7. Our findings identify Zcchc11 as a possible therapeutic target in Lin28A-expressing cancers. Accordingly, we demonstrate that Zcchc11 depletion selectively inhibits the tumorigenic capacity and metastatic potential of Lin28A- but not Lin28B-expressing human cancer cells and xenografts. Our results illuminate the distinct mechanisms by which Lin28A and Lin28B function and have broad implications for the development of new strategies for cancer therapy.

RESULTS

Lin28B Regulates let-7 Expression through a Zcchc11-Independent Mechanism

The paralogous RNA-binding proteins Lin28A and Lin28B have a high degree of sequence identity and conserved domain organization (Figure 1A), and both proteins selectively block let-7 expression (Newman et al., 2008; Viswanathan et al., 2008). We screened several human cancer cell lines and found that some express Lin28A, whereas others express Lin28B (Figure 1B). We did not observe coexpression of both Lin28A and Lin28B in any cell line, suggesting that their expression may be mutually exclusive. We found ubiquitous Zcchc11 expression. HeLa cells express Zcchc11 but neither Lin28A nor Lin28B. Because Lin28A-mediated repression of let-7 in mouse ESCs (mESCs) involves the TUTase Zcchc11, we next asked whether Lin28A and Lin28B function through the same mechanism to block let-7 processing. Previous reports have used recombinant Lin28A and Lin28B interchangeably in biochemical assays, demonstrating that Lin28B is capable of enhancing Zcchc11 activity in vitro; however, the physiological relevance of these observations remains unknown (Heo et al., 2009).

To begin to investigate whether both Lin28A and Lin28B function through a Zcchc11 TUTase-dependent mechanism, we performed coimmunoprecipitation (co-IP) experiments. Myc-tagged Lin28A, Lin28B, or Ago2 were coexpressed with either Flag-tagged Zcchc11 or Flag-EIF6 control (Figure 1C). Because the Lin28A-Zcchc11 interaction has been shown to be RNA dependent, we also coexpressed pri-let-7g (Heo et al., 2009). Consistent with earlier reports, myc-Lin28A was found to be associated with affinity-purified Flag-Zcchc11 (Heo et al., 2009). However, we were unable to detect a physical interaction between myc-Lin28B and Flag-Zcchc11. We performed additional co-IP experiments in which we titrated the amount of exogenously expressed Flag-Zcchc11. These experiments confirmed the specific physical interaction of Zcchc11 and Lin28A,




Figure 1. Lin28B Regulates let-7 Biogenesis through a Zcchc11-Independent Mechanism

(A) Schematic representation of human Lin28A and Lin28B.

(B) Western blot analysis of Zcchc11, Lin28A, and Lin28B in extracts prepared from human cancer cell lines.

(C) Coimmunoprecipitation (co-IP): HeLa cells were cotransfected with human myc-Lin28A, myc-Lin28B, or myc-Ago2 with either Flag-Zcchc11 or Flag-EIF6. Flag-IP and Flag- and Myc-western blots were performed to detect expression and interaction, respectively. See also Figure S1.

(D) Stable knockdown of Zcchc11 leads to upregulation of mature let-7g levels in Lin28A-expressing cells but not Lin28B-expressing cell lines. miRNA levels were measured by qRT-PCR. Error bars represent SEM (n = 3). Protein knockdown was monitored by western blot.

whereas myc-Lin28B was not detected in any of the Flag-Zcchc11 IPs (Figure S1A available online). This was additionally confirmed by the co-IP of endogenous Lin28A in Igrv1 cells (Figure S1B). Together, these results indicate that unlike for

Lin28A, we could not detect any physical interaction between Lin28B and Zcchc11.

Next, to address the functional requirement of Zcchc11 in the Lin28A- and Lin28B-mediated repression of let-7 expression, we

performed a series of knockdown experiments to deplete *Zcchc11* in a panel of human cancer cell lines. We used shRNAs to deplete *Lin28A* or *Zcchc11* expression in Igrov1 cells and measured the effect on *let-7* expression by quantitative reverse transcription PCR (qRT-PCR). As expected, depletion of *Lin28A* led to an ~10-fold increase in *let-7* levels. Knockdown of *Zcchc11* with three independent shRNAs also led to elevated mature *let-7* levels (Figure 1D). Therefore, *Zcchc11* is involved in the repression of *let-7* expression in this *Lin28A*-expressing human cancer cell line as has been previously reported in mESCs and embryonal carcinoma cells (Hagan et al., 2009; Heo et al., 2009). We performed analogous experiments in three different *Lin28B*-expressing cancer cell lines, HepG2, K562, and H1299 (Figure 1D), and found no significant effect on mature *let-7* levels in any of the cell lines when *Zcchc11* was depleted. In contrast, knockdown of *Lin28B* consistently led to the expected increase in mature *let-7*. Overall our results indicate that *Zcchc11* negatively regulates *let-7* expression in *Lin28A*- but not *Lin28B*-expressing cell lines, suggesting that *Lin28B* employs a *Zcchc11*-independent mechanism to block *let-7* processing.

Lin28B Localizes to the Nucleus

We sought potential explanations for the functional differences between *Lin28A* and *Lin28B*. We used immunofluorescence assays to examine the subcellular localization of the endogenous *Lin28A* and *Lin28B* proteins (Figure 2A). *Lin28A* was mostly localized to the cytoplasm of Igrov1 cells, whereas *Lin28B* localized to specific foci in the nuclei of H1299 cells where it colocalized with the nucleolar marker Fibrillarin. To further confirm the localization of *Lin28B* in the nucleoli, we performed immunofluorescence assays on H1299 cells in which *Lin28B* expression (or control) was stably knocked down by shRNA and showed that the observed nucleolar staining pattern is specific to *Lin28B* (Figure 2B). These data were further confirmed by biochemical fractionation and western blot of both cell lines (Figure 2C). Consistent with published data, we found *Zcchc11* only in the cytoplasmic fraction in both the *Lin28A*- and *Lin28B*-expressing cell lines (Figure 2C). These data suggest that the divergence in the mechanisms by which *Lin28A* and *Lin28B* block *let-7* biogenesis derives from their differential subcellular localization. The lack of physical and functional interactions between *Zcchc11* and *Lin28B* is therefore likely due to their localization to distinct cellular compartments, even though recombinant *Lin28B* has the capacity to enhance *Zcchc11* activity *in vitro* (Heo et al., 2009).

Lin28B Contains Functional Nuclear Localization Signals

Lin28B protein has an extended C terminus compared to *Lin28A*, which upon closer inspection contains a putative bipartite nuclear localization signal (NLS), KK[GPSVQ]KRKK. Another potential NLS, RRPK[GKTLQ]KRKPK, was identified in the linker region that connects the two functional RNA-binding domains (Figure 2D). To test the function of these putative NLS, we generated constructs for the expression of a series of GFP fusion proteins. We transiently transfected HeLa cells with these constructs and analyzed the subcellular localization of the GFP

fusion proteins by microscopy (Figure 2E). Consistent with the localization of endogenous *Lin28A* in Igrov1 cells, we found that *Lin28A*-GFP localized mainly to the cytoplasm. *Lin28B*-GFP predominantly localized to specific foci in the nucleus, again recapitulating the nucleolar localization of endogenous *Lin28B* observed in H1299 cells. When we exogenously expressed the *Lin28B* Δ NLS#1 truncation, we observed increased signal in the cytoplasm; however, some nucleolar localization still remained consistent with the presence of a second NLS. Indeed, the double-mutant *Lin28B*-GFP protein lacking both NLS showed cellular localization similar to that of GFP alone, suggesting that both NLS elements are important for nuclear and nucleolar localization of *Lin28B* (Figure 2E). To further determine whether these sequences represent functional NLS, we examined the localization of the NLS#1-GFP and NLS#2-GFP (Figure 2E). When exogenously expressed in HeLa cells, NLS#1-GFP localized nearly entirely throughout the nucleus including the nucleoli. This is in contrast to the control GFP construct that is broadly distributed throughout the cell. NLS#2-GFP was nearly entirely localized to the nucleoli (Figure 2E). Together these results identify that NLS#1 amino acid sequence represents a bona fide NLS, and that NLS#2 is a functional nucleolar localization signal (NoLS). NoLS properties are less well known and have only recently been studied at the amino acid sequence level (Scott et al., 2010). Several reports suggest that proteins with a NoLS also contain an NLS that allows them to cross the nuclear membrane before localizing to the nucleoli. These GFP-fusion results were confirmed by biochemical fractionation of HeLa cells transiently expressing either Flag-*Lin28A*, full-length, truncated, or double-mutant Flag-*Lin28B*. As expected Flag-*Lin28A* was predominantly present in the cytoplasmic fraction, Flag-*Lin28B* was mostly nuclear, NLS#1 mutant only showed marginal increase of signal in the cytoplasmic fraction, whereas the double-mutant Flag-*Lin28B* showed a more significant increase in cytoplasmic signal (Figure 2F).

Lin28B Localizes to Nucleoli where the Microprocessor Is Absent

We next compared the specific localization of *Lin28B* with that of the nuclear miRNA processing machinery. We found that the Microprocessor components DGCR8 and Drosha colocalize in the nucleoplasm but are excluded from nucleoli (Figure 3A). To further confirm that localization of *Lin28B* is distinct and nonoverlapping with the Microprocessor, we examined the colocalization of mCherry-DGCR8 and GFP-*Lin28A/B* proteins in transfected cells (Figure 3B). *Lin28A* localized mostly to the cytoplasm and therefore showed no overlap with the nuclear DGCR8. *Lin28B* localized to nucleoli and did not overlap with DGCR8 either. In contrast, the localization of the *Lin28B* NLS/NoLS mutant showed a broadly dispersed localization throughout the nucleus and cytoplasm (similar to GFP control) and displayed colocalization with DGCR8 in the nucleoplasm. We confirmed this finding that *Lin28B* and the Microprocessor normally occupy distinct compartments in the nucleus by performing large-scale biochemical fractionation and western blot of a stable HeLa cell line expressing Flag-*Lin28B*. *Lin28B* was specifically present in the nucleolar-enriched fractions whereas DGCR8 was only detectable in the nuclear fraction and not in the nucleolar




Figure 2. Lin28A and Lin28B Are Differentially Localized within the Cell

(A) Immunofluorescence detection of endogenous Lin28A in Igrov1 and Lin28B in H1299 cell lines. Fibrillarin, a known nucleolar protein, was used as a positive control.

(B) Immunofluorescence analysis of control and Lin28B knockdown H1299 cell lines.

(C) Biochemical fractionation of Igrov1 and H1299 cell lines. Endogenous Lin28A, Lin28B, and Zcchc11 in each fraction were detected by western blot. Fibrillarin was used as a nuclear marker; Tubulin was used as a cytoplasmic marker.

(D) Schematic of NLS in the Lin28B protein. An arginine as well as several lysines that were replaced by glycines are underlined and italicized.

(E) Localization of GFP-Lin28 fusion proteins in HeLa cells.

(F) Fractionation of Flag-Lin28 proteins, exogenously expressed in HeLa cells. Proteins were detected by Flag-western blot.




Figure 3. Lin28B Localizes to Nucleoli where the Microprocessor Is Absent

(A) Colocalization of the Microprocessor components GFP-Drosha and mCherry-DGCR8 in HeLa cells reveals their distribution throughout the nucleus and exclusion from nucleoli.

(B) Localization of GFP-Lin28A, Lin28B, or mutant Lin28B proteins with mCherry-DGCR8 in HeLa cells reveals nonoverlapping localization of Lin28B and DGCR8.

(C) Fractionation of a Flag-Lin28B HeLa stable cell line. Flag-Lin28B and endogenous DGCR8 were detected by western blot and show a nonoverlapping subcellular localization of Lin28B and the Microprocessor. Fibrillarin was used as a control for nucleolar localization.

fractions (Figure 3C). Overall, these findings suggest a possible mechanism by which Lin28B blocks let-7 processing in the nucleus by sequestering pri-let-7 miRNAs in the nucleoli away from the Microprocessor.

Lin28B Directly Binds and Sequesters Pri-let-7

To further dissect the mechanism of the Lin28B-mediated let-7 processing block, we first compared the relative abilities of recombinant human Lin28A and Lin28B proteins to bind pre-let-7 (Figures 4A, 4B, and S2). We performed electromobility shift assay (EMSA) with pre-let-7g to analyze the relative binding affinities of the two recombinant proteins. We found that Lin28A and Lin28B have apparent dissociation constants (K_D) of approximately 0.6 nM and 0.5 nM, respectively (Figure 4B).

Both these estimated K_D are much lower than previously reported for recombinant mouse Lin28A. This difference is likely due to the omission here of nonspecific yeast transfer RNA (tRNA) competitor used previously in the binding buffer (Piskounova et al., 2008). We also performed EMSA with pri-let-7g and demonstrated that both recombinant Lin28A and Lin28B are able to bind pri-let-7g with similar affinities (Figure 4C). Collectively, these assays reveal that both Lin28 proteins can directly bind to let-7 precursors with high affinity in vitro.

To gain further support for our model in which Lin28B binds and sequesters pri-let-7 in the nucleus to inhibit the Microprocessor, we next examined the RNA associated with Lin28B. We individually expressed and purified Flag-Lin28A and Flag-Lin28B, extracted the associated RNA, and analyzed relative levels of pri-let-7g by qRT-PCR. This RNA immunoprecipitation (RIP) analysis revealed that Lin28B directly associates with pri-let-7g RNA (Figure 4D). We detected an \sim 18-fold enrichment of pri-let-7 associated with Lin28B. Furthermore, we found substantially more pri-let-7 associated with Lin28B than with Lin28A, which is consistent with the differential subcellular localization of these proteins. Taken together, these results indicate that this preferential association of pri-let-7g with Lin28B likely reflects the distinct mechanism by which Lin28B represses let-7 expression rather than any possible intrinsic differences in the relative RNA-binding affinities of Lin28A and Lin28B proteins. Next, we examined by qRT-PCR the effect of transient Lin28B overexpression on pri-let-7g levels. Transient Lin28B overexpression led to \sim 12-fold accumulation of pri-let-7g levels (Figure 4E). In contrast, overexpression of Lin28A only had a more modest effect on pri-let-7g levels, consistent with its predominantly cytoplasmic localization. In order to further assess the effects of Lin28B overexpression on both pri- and mature miRNA levels, we utilized a Flag-Lin28B expressing a HeLa stable cell line. Analysis of several pri-miRNAs by qRT-PCR in this cell line demonstrated a substantial accumulation of pri-let-7 miRNAs, >10 -fold for pri-let-7g and >3 -fold for pri-let-7a-1. There was, however, no effect on levels of pri-miR-21 (Figure 4F). We detected a corresponding decrease in the levels of mature let-7, with $>90\%$ decrease for mature let-7g and $\sim 40\%$ decrease for mature let-7a. Again no effect was observed on levels of mature miR-21 (Figure 4F). Together, these data support our model whereby nuclear Lin28B directly associates with pri-let-7, sequestering it from cleavage by the Microprocessor to selectively inhibit let-7 maturation, and underscore our findings that the paralogous RNA-binding proteins Lin28A and Lin28B operate by distinct mechanisms to selectively repress let-7 expression.

Zcchc11 Inhibition Blocks the Tumorigenicity and Invasiveness of Lin28A- but Not Lin28B-Expressing Breast Cancer Cells In Vitro and In Vivo

We were motivated to further explore the relevance of our findings that Lin28A and Lin28B block let-7 processing through distinct mechanisms and to examine the effect of Zcchc11 inhibition on the tumorigenicity and invasiveness of human Lin28A/B-expressing cancer cells. Initially, we tested the consequences of Zcchc11 inhibition in the MCF10A ER-*Src*-inducible model of cellular transformation, where MCF10A-immortalized




Figure 4. Lin28B Directly Binds and Sequesters Pri-let-7

(A) Colloidal blue staining of purified recombinant His-Lin28A and His-Lin28B proteins.

(B) Binding of r.Lin28A and r.Lin28B to pre-let-7g was assessed by EMSA performed with 0.5 nM 5'-end labeled pre-let-7g RNA and the indicated concentration of recombinant protein. Band intensities were quantitated from three independent experiments and represented as the fraction of bound pre-let-7g RNA in the plots. Values are given as average \pm SEM (n = 3). See also Figure S2.

(C) EMSA performed indicated concentration of r.Lin28A and r.Lin28B with in vitro transcribed uniformly labeled pri-let-7g.

(D) RIP analysis of RNA associated with immunopurified Flag-Lin28A and Flag-Lin28B from HeLa cells. RNA was extracted from IP material and analyzed by qRT-PCR. Error bars \pm SEM (n = 3). Lower panel indicates relative Lin28A and Lin28B expression levels by Flag-western blot.

(E) Accumulation of pri-let-7 by transient Lin28B expression in HeLa cell detected by qRT-PCR. Error bars \pm SEM (n = 3). Lower panel indicates relative expression levels of Lin28A and Lin28B proteins detected by Flag-western blot in transfected cells.

(F) pri-let-7 accumulates (top panel) and mature let-7 levels decrease (bottom panel) in HeLa cells stably overexpressing Lin28B. Error bars \pm SEM (n = 3).

breast epithelial cells become transformed 36 hr post-tamoxifen (TAM) treatment (Iliopoulos et al., 2009). We previously reported that Lin28B expression is activated and required for induction

and maintenance of the transformed phenotype in this model (Figures S3A and S3B). In contrast, Zcchc11 depletion did not affect transformation or colony formation (Figures S3B–S3D)




Figure 5. Zcchc11 Inhibition Blocks Tumorigenicity and Invasiveness of Lin28A-Expressing Breast Cancer Cells

(A) qRT-PCR analysis of Zcchc11 knockdown in MDA-MB-231 and T47D breast cancer cells. Error bars \pm SEM ($n = 3$).

(B) Inhibition of Zcchc11 expression does not affect let-7a expression in Lin28B-expressing cells (MDA-MB-231), whereas it upregulates let-7a expression in Lin28A-expressing cells (T47D). Let-7a expression levels were measured by qRT-PCR in cells treated with siRNAs for 48 hr. Error bars \pm SEM ($n = 3$).

(C) Inhibition of Zcchc11 expression did not affect the colony formation ability of MDA-MB-231 cells but suppressed the colony formation ability of T47D cells. The number of colonies was evaluated 20 days post-plating in soft agar. The experiment was repeated thrice, and the statistical significance was calculated using Student's *t* test.

(D) Inhibition of Zcchc11 expression did not affect the invasiveness of MDA-MB-231 cells but suppressed the invasive ability of T47D cells. The number of invasive cells was measured 16 hr post-transfection with indicated siRNAs. In all assays, 10 fields per insert were scored, and SD was measured. The experiment was repeated thrice, and the statistical significance was calculated using Student's *t* test.

(E) Inhibition of Zcchc11 expression did not suppress tumor growth of MDA-MB-231 cells in xenografts, but it inhibited tumor growth of T47D cells in xenografts. The treatments with indicated siRNA were performed i.p. for five cycles starting on day 15. Each treatment group consisted of five mice.

and did not affect the expression of let-7a and its direct downstream target IL-6 (Figures S3D–S3F). Accordingly, Zcchc11 inhibition did not affect the tumorigenicity of MCF10A ER-*Src*-transformed cells in xenograft assays (Figure S3G). Overall, these data suggest that inhibition of Zcchc11 does not have an inhibitory effect on the tumorigenicity and invasiveness of Lin28B-expressing MCF10A ER-*Src* cells.

To further explore the distinct requirements for Zcchc11 in Lin28A- and Lin28B-expressing cancer, we compared the effects of Zcchc11 inhibition on the tumorigenicity and invasiveness of MDA-MB-231 breast cancer cells (Lin28B-expressing cells) relative to T47D breast cancer cells (Lin28A-expressing cells). We found that suppression of Zcchc11 expression did not affect let-7a expression in MDA-MB-231 cells but led to a 7-fold increase in mature let-7a levels in T47D cells (Figures 5A and 5B). Furthermore, Zcchc11 inhibition did not affect the tumorigenicity and invasiveness of MDA-MB-231 cells, although it suppressed both the colony formation ability and invasiveness of T47D cells (Figures 5C and 5D). Zcchc11 inhibition had similar effects on the tumor growth of these cell lines in xenografts. Specifically, Zcchc11 knockdown did not affect tumor growth of MDA-MB-231, whereas it suppressed T47D tumor growth (Figure 5E). Synthetic let-7a miRNA suppressed both MDA-MB-231 and T47D tumor growth (Figure 5E). Also, in the tumors derived from MDA-MB-231 xenografts (day 30), let-7a expression was not affected by inhibition of Zcchc11, whereas Lin28B suppression increased let-7a levels about 5-fold (Figures 5F and 5G). On the other hand, both Zcchc11 and Lin28A inhibition resulted in upregulation of let-7a expression to similar levels in T47D-derived tumors (day 30) (Figures 5F and 5G).

In addition to the breast cancer cells, we tested the effects of Zcchc11 inhibition on tumor growth of several other (liver, lung, ovarian, melanoma, colon) cancer cell types (Figure 6A). As above, we found that Zcchc11 inhibition (Figure 6B) blocked the growth of Lin28A-expressing tumors (Igrov1) and did not affect the growth of Lin28B-expressing tumors (HepG2, H1299, SK_MEL_28, CaCO2) (Figure 6A). Lin28A and Lin28B inhibition suppressed the growth of the corresponding tumors (Figures 6C and 6D). Taken together, these data suggest that Zcchc11 plays a role in the tumorigenicity and invasiveness of Lin28A-expressing cancer cells, but depletion of Zcchc11 in Lin28B-expressing cancer cell lines has no effect on cancer growth.

Lin28A and Lin28B Expression Levels in Human Colon and Breast Tissues

To further study the disease relevance of our findings, we measured Lin28A and Lin28B expression in human colon and breast tissues. We found that Lin28A or Lin28B is upregulated whereas let-7a is downregulated in colon adenocarcinomas relative to normal colon tissues (Figure 7A). Specifically, we

identified that Lin28A was upregulated in 19/45 colon adenocarcinomas, whereas Lin28B was upregulated in 14/45 colon adenocarcinomas. We found that the colon tumors with Lin28A overexpression had very low levels of Lin28B and vice versa. Furthermore, immunohistochemistry and in situ hybridization analyses in normal and colon cancer tissues revealed that Lin28A or Lin28B proteins are upregulated whereas let-7a is downregulated in colon carcinomas relative to normal colon tissues (Figures 7B and S4). Similar to the mRNA data, immunohistochemistry revealed that tumors that expressed high Lin28A protein levels had low levels for Lin28B and vice versa. This is consistent with our analysis of human cancer cell lines where we did not find cells that express both Lin28A and Lin28B (Figure 1B and data not shown).

We also found that Lin28A or Lin28B was upregulated whereas let-7a was downregulated in breast cancer relative to normal breast tissues (Figure 7C). Specifically, we identified that Lin28A was upregulated in 9/33 breast carcinomas, whereas Lin28B was upregulated in 10/33 breast carcinomas. Similar to the colon tissues, we detected that the breast tumors that overexpressed Lin28A had very low levels of Lin28B and vice versa. In addition, we detected that Lin28A was significantly upregulated in HER2-overexpressing breast tumors, whereas Lin28B was significantly upregulated in triple-negative (ER⁻, PR⁻, HER2⁻) breast tumors (Figure 7D). Furthermore, according to our previous studies, Lin28B expression is a part of an inflammatory circuit and is regulated by NF- κ B transcription factor (Iliopoulos et al., 2009). Thus, we tested how NF- κ B activity correlates with Lin28A or Lin28B mRNA levels in human breast tumors. We found a statistically significant correlation between NF- κ B nuclear levels and Lin28B but not Lin28A expression levels (Figure 7E). These data suggest that NF- κ B regulates the Lin28B but not Lin28A pathway. Also, in order to have a broader view of Lin28A and Lin28B expression levels in human cancer tissues, we tested their expression levels in kidney, liver, lung, ovarian, prostate, and thyroid cancer (Figure 7F). These data reveal that Lin28B is upregulated in liver, ovarian, and thyroid carcinomas.

DISCUSSION

Lin28A and Lin28B inhibit let-7 miRNA biogenesis in ESCs and cancer, and it has been assumed that both proteins block let-7 expression through the same mechanism, by recruiting the TUTase Zcchc11 (TUT4) in the cell cytoplasm to uridylate pre-let-7 and target it for degradation. Moreover, these paralogous proteins have been used interchangeably in several in vitro assays (Heo et al., 2009). Our results provide evidence that despite their high degree of homology, Lin28A and Lin28B function through distinct mechanisms to block let-7 processing, a finding that has broad implications for the development

(F) qRT-PCR analysis of siRNA inhibition of Lin28B, Lin28A, and Zcchc11 in xenograft tumors (day 30) derived from MDA-MB-231 and T47D cells. Error bars \pm SEM (n = 3).

(G) Inhibition of Lin28B but not of Zcchc11 allows upregulation of let-7a expression levels in MDA-MB-231 xenograft tumors (day 30). However, inhibition of Lin28A or Zcchc11 results in let-7a upregulation in T47D xenograft tumors. Let-7a expression levels were measured by qRT-PCR on tumors untreated or treated with indicated siRNA. Error bars \pm SEM (n = 3). See also Figure S3.




Figure 6. Inhibition of Zcchc11 Expression Suppresses Tumor Growth of Lin28A- but Not Lin28B-Expressing Xenografts

(A) Xenograft experiments were performed with a variety of different human cancer cell lines. Mice were treated with the indicated siRNA for five cycles starting on day 15. For all cells lines tested, each treatment group consisted of five mice. Although inhibition of Lin28A or Lin28B suppressed tumor growth in the relevant xenografts, inhibition of Zcchc11 inhibited growth only of Lin28A- but not Lin28B-expressing tumors. Error bars \pm SEM (n = 3).

(B) Analysis of siRNA inhibition of Zcchc11 in xenograft tumors (day 30) derived from the indicated cells.

(C) Analysis of siRNA inhibition of Lin28B in xenograft tumors (day 30) derived from the indicated cells.

(D) Analysis of siRNA inhibition of Lin28A in xenograft tumors (day 30) derived from IGRV1 cells. mRNA expression levels were measured by qRT-PCR on tumors untreated or treated with the indicated siRNA. Error bars \pm SEM (n = 3).

of new cancer therapies and the potential use of Zcchc11 inhibitors in Lin28A-expressing tumors but not in Lin28B-expressing tumors. This distinction derives from the differential subcellular localization of these two proteins: Lin28A localizes primarily to the cytoplasm, whereas Lin28B contains func-

tional nuclear localization signals and specifically localizes to nucleoli.

Due to the differential subcellular localization of the two proteins, we find that in human cancer cell lines, Lin28A and Lin28B block let-7 processing at different steps of the




Figure 7. Lin28A and Lin28B Expression in Primary Human Cancers

(A) qRT-PCR analysis of Lin28A, Lin28B, and let-7a expression levels in normal and colon cancer tissues. Tumor samples were further classified into two groups expressing either high Lin28A or Lin28B. Data expressed as mean \pm SEM (n = 3).

(B) Immunohistochemistry for Lin28A and Lin28B and in situ hybridization for let-7a and U6 in normal colon tissues and colon adenocarcinomas. See also Figure S4.

miRNA-processing pathway. However, the steps at which let-7 processing is blocked by Lin28 in various different organisms are controversial. A recently published report by Van Wynsberghe et al. proposes that Lin28 binds pri-let-7 and blocks let-7 expression cotranscriptionally in *C. elegans* and disputes earlier conclusions that Lin28 functions at steps further downstream in the let-7 biogenesis pathway (Lehrbach et al., 2009; Van Wynsberghe et al., 2011). They also report that a very small fraction of Lin28A in human ESCs (hESCs) localizes to the nucleus and binds pri-let-7 miRNAs, although pre-let-7 is bound abundantly (Van Wynsberghe et al., 2011). This is consistent with our results that demonstrate that a small fraction of Lin28A localizes to the nucleus in an Igrv1 cell line. We also find that Lin28A binds pri-let-7 miRNA, but not as much as Lin28B, and earlier reports have demonstrated that purified Lin28A can inhibit the Microprocessor in vitro (Newman et al., 2008; Viswanathan et al., 2008). Also, although our data demonstrate that Lin28B-mediated repression of let-7 expression is Zcchc11 (TUT4) independent in multiple different cell types, it remains possible that in certain contexts or cell types, Lin28B may localize to the cytoplasm and utilize Zcchc11/TUT4 to repress let-7 biogenesis. For example, uridylylated pre-let-7 was previously detected in Huh7 cells, and Lin28B is reportedly localized to the cytoplasm in Huh7 cells (Guo et al., 2006; Heo et al., 2008).

Our demonstration that the Microprocessor is excluded from nucleoli raises the possibility that sequestration of certain pri-miRNAs to nucleoli by specific RNA-binding proteins may prove to be a more general strategy for the posttranscriptional control of miRNA biogenesis. Previous reports have demonstrated that nucleoli contain machinery responsible for modifying small nucleolar RNAs, for example through RNA methylation or 3' uridylation. Whether these nucleolar mechanisms play a role in pri-miRNA regulation remains to be determined (Boisvert et al., 2007; Matera et al., 2007). It is possible that additional new factors may be involved in sequestering and possibly degrading and/or modifying pri-let-7 miRNAs in the nucleoli in a Lin28B-dependent manner. The identification of such factors could reveal new potential therapeutic targets aimed at restoring let-7 expression in Lin28B-expressing cancers.

Our proof-of-concept studies with human breast and ovarian cancer cell lines demonstrate that inhibition of Zcchc11 may represent a possible new therapeutic target for cancer. Indeed knockdown of Zcchc11 effectively inhibits the tumorigenic capacity and metastatic potential of human breast and ovarian cancer cells and xenografts. Importantly, however, our data also predict that the therapeutic potential of Zcchc11 inhibition will be limited to Lin28A-expressing cancers. Although Lin28A expression is relatively uncommon in several human cancer cell lines, our analysis of primary human colon and breast cancer indicates that upregulation of Lin28A or Lin28B occurs in a large proportion of tumors with approximately equal frequency for

each protein. Furthermore, expression of Lin28A or Lin28B seems to distinguish different classes of breast cancers (Figure 7). Therefore, the identification of small-molecule inhibitors of Zcchc11 may lead to the development of novel chemotherapies for Lin28A-expressing cancers.

EXPERIMENTAL PROCEDURES

Cloning

Myc-Lin28A and -Lin28B were cloned into pBK-EF1. Lin28A, Lin28B, and Lin28B Δ NLS#1 were cloned into pFLAG-CMV2 vector (Sigma). Lin28B Δ NoLS Δ NLS#1 was generated by site-directed mutagenesis using the Quick Change kit (Stratagene). Lin28A, Lin28B, Lin28B Δ NLS#1, and Lin28B Δ NoLS Δ NLS#1 were cloned into CT-GFP-Topo (Invitrogen). NLS#1 and NLS#2 oligos were annealed before ligating into CT-GFP-Topo. N-terminal Cherry-DGCR8 fusion construct was generated by subcloning Cherry cDNA into p3xFLAG-CMV14-DGCR8 (Gregory et al., 2004). Lin28A and Lin28B were subcloned into Pet21 for His-tagged recombinant protein expression. Pri-let-7g was previously reported (Viswanathan et al., 2008). Cloning primers are listed in Table S1.

Transfection, Knockdowns, and Stable Cell Line Generation

All transfections were performed with Lipofectamine (Invitrogen) per manufacturer's instructions. shRNA constructs were generated using pLKO.1 Puro (Addgene #8453) and pLKO.1 Hygro (Addgene #24150) vectors. The sequences of the shRNA hairpins and siRNAs are listed in Table S2. Lentivirus production, infection, and stable cell line selection are as described (Viswanathan et al., 2009). siRNAs used in this study were as follows: siRNA negative control (100 nM), (cat no. AM4611, Ambion Inc); siRNA Zcchc11#1 (100 nM) (cat no. s23551, Ambion Inc); siRNA Zcchc11#2 (100 nM) (cat no. s23553, Ambion Inc); siRNA Lin28B (100 nM) (cat no. s52477, Ambion Inc); siRNA Lin28A (100 nM) (cat no. s36195, Ambion Inc).

Western Blotting

The following antibodies were used: Lin28A (Cell Signaling, 3978), Lin28B (Cell Signaling, 4196), Zcchc11 (Protein Tech Group, 18980-1-AP), Zcchc11 (Imgenex, IMX-3587), DGCR8 (Protein Tech Group, 10996-1-AP), Fibrillarin (Abcam, ab18380), and β -Tubulin (Abcam AB6046).

Subcellular Fractionation

Cellular fractionation was done using NE-PER Nuclear and Cytoplasmic Extraction Kits (Pierce) per manufacturer's instructions. Large-scale fractionation was performed as described in the Extended Experimental Procedures.

EMSA

EMSA with purified His-Lin28A/B was performed as described but without competitor yeast tRNA (Piskounova et al., 2008). Complexes were resolved on native 3.5% or 5% polyacrylamide gels and visualized by autoradiography. Band intensities of scanned gels were quantified using ImageJ software and used to calculate percentage of probe bound. Graph-Pad Prism was used to plot data. Percent active protein was determined using stoichiometric binding reactions as described in Ryder et al. (2008). Hills equation for specific binding with one site, $Y = B_{max} * X^n / (K_D^n + X^n)$, was used to calculate K_D .

Colony Formation Assay

MDA-MB-231 cells and T47D cells were transfected with different siRNAs for 48 hr. Triplicate samples of 10^5 cells from each cell line were mixed 4:1 (v/v) with 2.0% agarose in growth medium for a final concentration of 0.4% agarose. The cell mixture was plated on top of a solidified layer of 0.5% agarose in

(C) qRT-PCR analysis of Lin28A, Lin28B, and let-7a in human normal and breast cancer tissues. Tumor samples were further classified into two groups expressing either high Lin28A or Lin28B. Data expressed as mean \pm SEM (n = 3).

(D) Lin28A, Lin28B, and let-7a expression levels in different breast cancer subtypes.

(E) Correlation between Lin28A and Lin28B mRNA levels assessed by qRT-PCR with NF- κ B phosphorylation status assessed by ELISA assay.

(F) Heatmap representation of Lin28A and Lin28B in carcinomas of different origin measured by qRT-PCR.

growth medium. Cells were fed every 6 to 7 days with growth medium containing 0.4% agarose. The number of colonies was counted after 20 days.

Invasion Assays

We performed invasion assays in MDA-MB-231, and T47D breast cancer cells were transfected with different siRNAs for 16 hr. Invasion of matrigel was conducted by using standardized conditions with BDBioCoat growth factor-reduced MATRIGEL invasion chambers (PharMingen). Assays were conducted per manufacturer's protocol, using 10% FBS as chemoattractant. Noninvading cells on the top side of the membrane were removed, whereas invading cells were fixed and stained with 4'-6-diamidino-2-phenylindole (DAPI, Vector Laboratories Inc.), 16 hr post-seeding.

Mouse Experiments

Xenograft experiments with human cancer cell lines are described in detail the [Extended Experimental Procedures](#). Briefly, cells were injected subcutaneously in the right flank of athymic nude mice. Tumor growth was monitored every 5 days. In Vivo Ready siRNAs (Ambion Inc.) were mixed with Invivofermine 2.0 liposomes (Ambion Inc) and injected intraperitoneally (i.p.) in a volume of 100 μ l at a dose of 5 mg/kg.

Human Tissues and RNAs

Thirty normal colon tissues and 45 colon adenocarcinomas were collected from the translational pathology core laboratory of the Department of Pathology at UCLA. All subjects gave informed consent, and the study was approved by the UCLA Institutional Review Board. RNAs from 12 normal mammary tissues and 33 breast cancer tissues were purchased from Origene Inc. The ER, PR, and HER2 status for each of these breast carcinomas was known. Additional RNAs were purchased from Origene (8 renal cell carcinomas, 8 hepatocellular carcinomas, 8 squamous cell lung carcinomas, 8 ovarian adenocarcinomas, 8 prostate adenocarcinomas, 8 papillary thyroid carcinomas).

In Situ Hybridization and Immunohistochemistry

Sections of the colon adenocarcinomas and adjacent uninvolved tissues were analyzed by in situ hybridization as described ([Iliopoulos et al., 2009](#)) with modification ([Extended Experimental Procedures](#)). Double-DIG labeled Mircury LNA Detection probe for the detection of hsa-let-7 (1800-15, Exiqon) was used. Protocol for immunohistochemistry is described in detail the [Extended Experimental Procedures](#), with the following antibodies: Lin28A (3978, Cell Signaling Technology) and Lin28B (LS-B3423, LSBio).

SUPPLEMENTAL INFORMATION

Supplemental Information includes [Extended Experimental Procedures](#), four figures, and two tables and can be found with this article online at [doi:10.1016/j.cell.2011.10.039](https://doi.org/10.1016/j.cell.2011.10.039).

ACKNOWLEDGMENTS

We thank Dr. Akihiko Yoshimura, Keio University, Japan, for the human Flag-Zcchc11 plasmid; Dr. Na Liu, Memorial Sloan Kettering Cancer Center, for the Myc-Ago2 plasmid; Dr. Bharat Ramratnam, Brown University, for the GFP-Drosha plasmid; and Dr. Ramin Shiekhattar, Wistar Institute, for the Flag-Eif6 expression plasmid. We thank Chiara Giacomelli, Dr. Saurabh Patel, and Dr. Maria Hatziaepostolou for technical assistance. R.I.G. was supported by grants from the US National Institute of General Medical Sciences (NIGMS) (1R01GM086386-01A1), The Harvard Stem Cell Institute, and The American Cancer Society (121635-RSG-11-175-01-RMC). R.I.G. is a Pew Research Scholar. D.I. was supported by start-up funds from the Department of Cancer Immunology & AIDS at Dana-Farber Cancer Institute.

Received: March 8, 2011

Revised: July 25, 2011

Accepted: October 24, 2011

Published: November 23, 2011

REFERENCES

- Barh, D., Malhotra, R., Ravi, B., and Sindhurani, P. (2010). MicroRNA let-7: an emerging next-generation cancer therapeutic. *Curr. Oncol.* *17*, 70–80.
- Boisvert, F.M., van Koningsbruggen, S., Navascués, J., and Lamond, A.I. (2007). The multifunctional nucleolus. *Nat. Rev. Mol. Cell Biol.* *8*, 574–585.
- Boyerinas, B., Park, S.M., Hau, A., Murmann, A.E., and Peter, M.E. (2010). The role of let-7 in cell differentiation and cancer. *Endocr. Relat. Cancer* *17*, F19–F36.
- Büssing, I., Slack, F.J., and Grosshans, H. (2008). let-7 microRNAs in development, stem cells and cancer. *Trends Mol. Med.* *14*, 400–409.
- Chang, T.C., Zeitels, L.R., Hwang, H.W., Chivukula, R.R., Wentzel, E.A., Dewes, M., Jung, J., Gao, P., Dang, C.V., Beer, M.A., et al. (2009). Lin-28B transactivation is necessary for Myc-mediated let-7 repression and proliferation. *Proc. Natl. Acad. Sci. USA* *106*, 3384–3389.
- Dangi-Garimella, S., Yun, J., Eves, E.M., Newman, M., Erkeland, S.J., Hammond, S.M., Minn, A.J., and Rosner, M.R. (2009). Raf kinase inhibitory protein suppresses a metastasis signalling cascade involving LIN28 and let-7. *EMBO J.* *28*, 347–358.
- Denli, A.M., Tops, B.B., Plasterk, R.H., Ketting, R.F., and Hannon, G.J. (2004). Processing of primary microRNAs by the Microprocessor complex. *Nature* *432*, 231–235.
- Di Leva, G., and Croce, C.M. (2010). Roles of small RNAs in tumor formation. *Trends Mol. Med.* *16*, 257–267.
- Esquela-Kerscher, A., Trang, P., Wiggins, J.F., Patrawala, L., Cheng, A., Ford, L., Weidhaas, J.B., Brown, D., Bader, A.G., and Slack, F.J. (2008). The let-7 microRNA reduces tumor growth in mouse models of lung cancer. *Cell Cycle* *7*, 759–764.
- Gregory, R.I., Yan, K.P., Amuthan, G., Chendrimada, T., Doratotaj, B., Cooch, N., and Shiekhattar, R. (2004). The Microprocessor complex mediates the genesis of microRNAs. *Nature* *432*, 235–240.
- Gregory, R.I., Chendrimada, T.P., Cooch, N., and Shiekhattar, R. (2005). Human RISC couples microRNA biogenesis and posttranscriptional gene silencing. *Cell* *123*, 631–640.
- Guo, Y., Chen, Y., Ito, H., Watanabe, A., Ge, X., Kodama, T., and Aburatani, H. (2006). Identification and characterization of lin-28 homolog B (LIN28B) in human hepatocellular carcinoma. *Gene* *384*, 51–61.
- Hagan, J.P., Piskounova, E., and Gregory, R.I. (2009). Lin28 recruits the TUTase Zcchc11 to inhibit let-7 maturation in mouse embryonic stem cells. *Nat. Struct. Mol. Biol.* *16*, 1021–1025.
- Heo, I., Joo, C., Cho, J., Ha, M., Han, J., and Kim, V.N. (2008). Lin28 mediates the terminal uridylation of let-7 precursor MicroRNA. *Mol. Cell* *32*, 276–284.
- Heo, I., Joo, C., Kim, Y.K., Ha, M., Yoon, M.J., Cho, J., Yeom, K.H., Han, J., and Kim, V.N. (2009). TUT4 in concert with Lin28 suppresses microRNA biogenesis through pre-microRNA uridylation. *Cell* *138*, 696–708.
- Hutvagner, G., McLachlan, J., Pasquinelli, A.E., Bálint, E., Tuschl, T., and Zamore, P.D. (2001). A cellular function for the RNA-interference enzyme Dicer in the maturation of the let-7 small temporal RNA. *Science* *293*, 834–838.
- Iliopoulos, D., Hirsch, H.A., and Struhl, K. (2009). An epigenetic switch involving NF- κ B, Lin28, Let-7 MicroRNA, and IL6 links inflammation to cell transformation. *Cell* *139*, 693–706.
- Ji, J., and Wang, X.W. (2010). A Yin-Yang balancing act of the lin28/let-7 link in tumorigenesis. *J. Hepatol.* *53*, 974–975.
- King, C.E., Cuatrecasas, M., Castells, A., Sepulveda, A.R., Lee, J.S., and Rustgi, A.K. (2011). LIN28B promotes colon cancer progression and metastasis. *Cancer Res.* *71*, 4260–4268.
- Krol, J., Loedige, I., and Filipowicz, W. (2010). The widespread regulation of microRNA biogenesis, function and decay. *Nat. Rev. Genet.* *11*, 597–610.
- Lehrbach, N.J., Armisen, J., Lightfoot, H.L., Murfitt, K.J., Bugaut, A., Balasubramanian, S., and Miska, E.A. (2009). LIN-28 and the poly(U) polymerase PUP-2 regulate let-7 microRNA processing in *Caenorhabditis elegans*. *Nat. Struct. Mol. Biol.* *16*, 1016–1020.

- Liang, L., Wong, C.M., Ying, Q., Fan, D.N., Huang, S., Ding, J., Yao, J., Yan, M., Li, J., Yao, M., et al. (2010). MicroRNA-125b suppressed human liver cancer cell proliferation and metastasis by directly targeting oncogene LIN28B2. *Hepatology* 52, 1731–1740.
- Liu, J., Carmell, M.A., Rivas, F.V., Marsden, C.G., Thomson, J.M., Song, J.J., Hammond, S.M., Joshua-Tor, L., and Hannon, G.J. (2004). Argonaute2 is the catalytic engine of mammalian RNAi. *Science* 305, 1437–1441.
- Lu, L., Katsaros, D., Shaverdashvili, K., Qian, B., Wu, Y., de la Longrais, I.A., Preti, M., Menato, G., and Yu, H. (2009). Pluripotent factor lin-28 and its homologue lin-28b in epithelial ovarian cancer and their associations with disease outcomes and expression of let-7a and IGF-II. *Eur. J. Cancer* 45, 2212–2218.
- Martinez, N.J., and Gregory, R.I. (2010). MicroRNA gene regulatory pathways in the establishment and maintenance of ESC identity. *Cell Stem Cell* 7, 31–35.
- Matera, A.G., Terns, R.M., and Terns, M.P. (2007). Non-coding RNAs: lessons from the small nuclear and small nucleolar RNAs. *Nat. Rev. Mol. Cell Biol.* 8, 209–220.
- Melton, C., Judson, R.L., and Belloch, R. (2010). Opposing microRNA families regulate self-renewal in mouse embryonic stem cells. *Nature* 463, 621–626.
- Moss, E.G., Lee, R.C., and Ambros, V. (1997). The cold shock domain protein LIN-28 controls developmental timing in *C. elegans* and is regulated by the lin-4 RNA. *Cell* 88, 637–646.
- Newman, M.A., Thomson, J.M., and Hammond, S.M. (2008). Lin-28 interaction with the Let-7 precursor loop mediates regulated microRNA processing. *RNA* 14, 1539–1549.
- Oh, J.S., Kim, J.J., Byun, J.Y., and Kim, I.A. (2010). Lin28-let7 modulates radio-sensitivity of human cancer cells with activation of K-Ras. *Int. J. Radiat. Oncol. Biol. Physiol.* 76, 5–8.
- Peng, S., Maihle, N.J., and Huang, Y. (2010). Pluripotency factors Lin28 and Oct4 identify a sub-population of stem cell-like cells in ovarian cancer. *Oncogene* 29, 2153–2159.
- Piskounova, E., Viswanathan, S.R., Janas, M., LaPierre, R.J., Daley, G.Q., Sliz, P., and Gregory, R.I. (2008). Determinants of microRNA processing inhibition by the developmentally regulated RNA-binding protein Lin28. *J. Biol. Chem.* 283, 21310–21314.
- Rybak, A., Fuchs, H., Smirnova, L., Brandt, C., Pohl, E.E., Nitsch, R., and Wulczyn, F.G. (2008). A feedback loop comprising lin-28 and let-7 controls pre-let-7 maturation during neural stem-cell commitment. *Nat. Cell Biol.* 10, 987–993.
- Ryder, S.P., Recht, M.I., and Williamson, J.R. (2008). Quantitative analysis of protein-RNA interactions by gel mobility shift. *Methods Mol. Biol.* 488, 99–115.
- Scott, M.S., Boisvert, F.M., McDowall, M.D., Lamond, A.I., and Barton, G.J. (2010). Characterization and prediction of protein nucleolar localization sequences. *Nucleic Acids Res.* 38, 7388–7399.
- Shell, S., Park, S.M., Radjabi, A.R., Schickel, R., Kistner, E.O., Jewell, D.A., Feig, C., Lengyel, E., and Peter, M.E. (2007). Let-7 expression defines two differentiation stages of cancer. *Proc. Natl. Acad. Sci. USA* 104, 11400–11405.
- Slack, F. (2009). let-7 microRNA reduces tumor growth. *Cell Cycle* 8, 1823.
- Small, E.M., and Olson, E.N. (2011). Pervasive roles of microRNAs in cardiovascular biology. *Nature* 469, 336–342.
- Takamizawa, J., Konishi, H., Yanagisawa, K., Tomida, S., Osada, H., Endoh, H., Harano, T., Yatabe, Y., Nagino, M., Nimura, Y., et al. (2004). Reduced expression of the let-7 microRNAs in human lung cancers in association with shortened postoperative survival. *Cancer Res.* 64, 3753–3756.
- Trang, P., Medina, P.P., Wiggins, J.F., Ruffino, L., Kelnar, K., Omotola, M., Homer, R., Brown, D., Bader, A.G., Weidhaas, J.B., and Slack, F.J. (2010). Regression of murine lung tumors by the let-7 microRNA. *Oncogene* 29, 1580–1587.
- Van Wynsberghe, P.M., Kai, Z.S., Massirer, K.B., Burton, V.H., Yeo, G.W., and Pasquinelli, A.E. (2011). LIN-28 co-transcriptionally binds primary let-7 to regulate miRNA maturation in *Caenorhabditis elegans*. *Nat. Struct. Mol. Biol.* 18, 302–308.
- Viswanathan, S.R., and Daley, G.Q. (2010). Lin28: A microRNA regulator with a macro role. *Cell* 140, 445–449.
- Viswanathan, S.R., Daley, G.Q., and Gregory, R.I. (2008). Selective blockade of microRNA processing by Lin28. *Science* 320, 97–100.
- Viswanathan, S.R., Powers, J.T., Einhorn, W., Hoshida, Y., Ng, T.L., Toffanin, S., O'Sullivan, M., Lu, J., Phillips, L.A., Lockhart, V.L., et al. (2009). Lin28 promotes transformation and is associated with advanced human malignancies. *Nat. Genet.* 41, 843–848.
- Wang, Y.C., Chen, Y.L., Yuan, R.H., Pan, H.W., Yang, W.C., Hsu, H.C., and Jeng, Y.M. (2010). Lin-28B expression promotes transformation and invasion in human hepatocellular carcinoma. *Carcinogenesis* 31, 1516–1522.
- West, J.A., Viswanathan, S.R., Yabuuchi, A., Cunniff, K., Takeuchi, A., Park, I.H., Sero, J.E., Zhu, H., Perez-Atayde, A., Frazier, A.L., et al. (2009). A role for Lin28 in primordial germ-cell development and germ-cell malignancy. *Nature* 460, 909–913.
- Yang, X., Lin, X., Zhong, X., Kaur, S., Li, N., Liang, S., Lassus, H., Wang, L., Katsaros, D., Montone, K., et al. (2010). Double-negative feedback loop between reprogramming factor LIN28 and microRNA let-7 regulates aldehyde dehydrogenase 1-positive cancer stem cells. *Cancer Res.* 70, 9463–9472.
- Yu, F., Yao, H., Zhu, P., Zhang, X., Pan, Q., Gong, C., Huang, Y., Hu, X., Su, F., Lieberman, J., and Song, E. (2007a). let-7 regulates self renewal and tumorigenicity of breast cancer cells. *Cell* 131, 1109–1123.
- Yu, J., Vodyanik, M.A., Smuga-Otto, K., Antosiewicz-Bourget, J., Frane, J.L., Tian, S., Nie, J., Jonsdottir, G.A., Ruotti, V., Stewart, R., et al. (2007b). Induced pluripotent stem cell lines derived from human somatic cells. *Science* 318, 1917–1920.
- Zhu, H., Shah, S., Shyh-Chang, N., Shinoda, G., Einhorn, W.S., Viswanathan, S.R., Takeuchi, A., Grasemann, C., Rinn, J.L., Lopez, M.F., et al. (2010). Lin28a transgenic mice manifest size and puberty phenotypes identified in human genetic association studies. *Nat. Genet.* 42, 626–630.