



Contents lists available at ScienceDirect

Physics Letters B

www.elsevier.com/locate/physletb



Propagation peculiarities of mean field massive gravity

S. Deser^{a,b}, A. Waldron^c, G. Zahariade^d^a Walter Burke Institute for Theoretical Physics, California Institute of Technology, Pasadena, CA 91125, USA^b Physics Department, Brandeis University, Waltham, MA 02454, USA^c Department of Mathematics, University of California, Davis, CA 95616, USA^d Department of Physics, University of California, Davis, CA 95616, USA

ARTICLE INFO

Article history:

Received 18 April 2015

Accepted 16 July 2015

Available online 28 July 2015

Editor: M. Cvetič

ABSTRACT

Massive gravity (mGR) describes a dynamical “metric” on a fiducial, background one. We investigate fluctuations of the dynamics about mGR solutions, that is about its “mean field theory”. Analyzing mean field massive gravity (\bar{m} GR) propagation characteristics is not only equivalent to studying those of the full non-linear theory, but also in direct correspondence with earlier analyses of charged higher spin systems, the oldest example being the charged, massive spin 3/2 Rarita–Schwinger (RS) theory. The fiducial and mGR mean field background metrics in the \bar{m} GR model correspond to the RS Minkowski metric and external EM field. The common implications in both systems are that hyperbolicity holds only in a weak background-mean-field limit, immediately ruling both theories out as fundamental theories; a situation in stark contrast with general relativity (GR) which is at least a consistent classical theory. Moreover, even though both \bar{m} GR and RS theories can still in principle be considered as predictive effective models in the weak regime, their lower helicities then exhibit superluminal behavior: lower helicity gravitons are superluminal as compared to photons propagating on either the fiducial or background metric. Thus our approach has uncovered a novel, dispersive, “crystal-like” phenomenon of differing helicities having differing propagation speeds. This applies both to \bar{m} GR and mGR, and is a *peculiar* feature that is also problematic for consistent coupling to matter.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

Consistency is a powerful tool for studying field theories. Already classically, there are stringent conditions that are extremely difficult to fulfill for systems with spin $s > 1$, the most important exception being ($s = 2$, $m = 0$) general relativity. Key consistency requirements are

- (i) Correct degree of freedom (DoF) counts.
- (ii) Non-ghost kinetic terms.
- (iii) Predictability.
- (iv) (Sub)luminal propagation.

Requirements (i) and (ii) are closely related (as are (iii) and (iv)). Models whose constraints do not single out the correct propagating DoF suffer from relatively ghost kinetic terms: the relevant

E-mail addresses: deser@brandeis.edu (S. Deser), wally@math.ucdavis.edu (A. Waldron), zahariad@ucdavis.edu (G. Zahariade).

<http://dx.doi.org/10.1016/j.physletb.2015.07.055>

0370-2693/© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

example here is the sixth ghost excitation that plagues generic massive gravity (mGR) theories [1]. The discovery that a class of mGR models satisfied requirements (i) and (ii) generated a revival of interest in massive spin 2 theories [2–7] even though failure of the propagation requirements (iii) and (iv) were long known to be devil higher spin theories [8,9].

The predictability requirement is that initial data can be propagated to the future of spacetime hypersurfaces. In PDE terms, this means that the underlying equations must be hyperbolic [10]. The final requirement, that signals cannot propagate faster than light, can be imposed once the hyperbolicity requirement is satisfied. The classic example of a model that obeys requirements (i) and (ii) as well as (iii) but only in a weak field region, is the charged, massive, $s = 3/2$ RS theory. Curiously enough, the propagation problems of this model were first discovered in a quantum setting by Johnson and Sudarshan [11] who studied the model's canonical field commutators (this is easy to understand in retrospect, because field commutators and propagators are directly related [12]). The first detailed analysis of the model's propagation characteristics was carried out by Velo and Zwanziger; our aim is

to reproduce their RS results in $\bar{m}GR$, so we quote their 1971 abstract verbatim [8]:

The Rarita–Schwinger equation in an external electromagnetic potential is shown to be equivalent to a hyperbolic system of partial differential equations supplemented by initial conditions. The wave fronts of the classical solutions are calculated and are found to propagate faster than light. Nevertheless, for sufficiently weak external potentials, a consistent quantum mechanics and quantum field theory may be established. These, however, violate the postulates of special relativity.

In previous works we and other authors have shown that similar conclusions hold for the full non-linear mGR models [5,6,13,14,7,15]. These investigations rely on the method of characteristics, which amounts to studying leading kinetic terms and is thus essentially equivalent to an analysis of linear fluctuations around a mean field background. Since this mean field massive gravity ($\bar{m}GR$) fluctuation model depends both on a background and a fiducial metric, it is in direct correspondence with the charged RS model. Hence, without any computation at all, one can readily predict that: (a) mGR loses hyperbolicity in some strong field regime and (b) in the weak field hyperbolic regime where predictability is restored, lower helicity modes have propagation characteristics differing from maximal helicity ± 2 ; thus superluminality with respect to (luminal) photons is inevitable. Apart from confirming earlier conclusions in a very simple setting, our results give a precise description of mGR's effective, weak field, regime.

2. Massive gravity

At its genesis, the first known non-linear mGR model of [16] was originally formulated in terms of dynamical and fiducial vierbeine e^m and f^m . It took some forty years for researchers – independently in an effective field theory-inspired metric formulation – to discover that this model was one of a three-parameter family [2] that avoided the sixth, ghost-like excitation of [1]. The action describing these fiducial mGR models is given by¹

$$S_{\text{mGR}}[e, \omega; f] = - \int \epsilon_{mnr s} e^m \left\{ \frac{1}{4} e^n [d\omega^{rs} + \omega^r_t \omega^{ts}] - m^2 \left[\frac{\beta_0}{4} e^n e^r e^s + \frac{\beta_1}{3} e^n e^r f^s + \frac{\beta_2}{2} e^n f^r f^s + \beta_3 f^n f^r f^s \right] \right\}.$$

The parameter β_0 governs a standard cosmological term; this is required to obtain the Fierz–Pauli (FP) linearized limit when both the fiducial and mGR backgrounds are Minkowski. When both the fiducial and mGR backgrounds are Einstein with cosmological constant $\bar{\Lambda}$, the model's parameters must obey $\frac{\bar{\Lambda}}{3!} = m^2(\beta_0 + \beta_1 + \beta_2 + \beta_3)$ and the linearized theory is FP with mass $m_{\text{FP}}^2 := m^2(\beta_1 + 2\beta_2 + 3\beta_3)$.

Varying the model's dynamical fields (e^m, ω^{mn}) gives equations of motion

$$\nabla e^m \approx 0 \approx G_m - m^2 t_m, \quad (1)$$

where $t_m := \epsilon_{mnr s} [\beta_0 e^n e^r e^s + \beta_1 e^n e^r f^s + \beta_2 e^n f^r f^s + \beta_3 f^n f^r f^s]$. Also, the Einstein three-form is defined by $G_m := \frac{1}{2} \epsilon_{mnr s} e^n R^{rs}$ and

$R^{mn} := d\omega^{mn} + \omega^m_r \omega^{rn}$ is the Riemann curvature; ∇ is the connection of ω^{mn} . The forty equations above are subject to thirty constraints that are spelled out in detail in [7]. In particular, these include the covariant algebraic relations²

$$e^m f_m \approx 0 \approx K_{mn} e^m f^n \approx \epsilon_{mnr s} M^{mn} K^{rs},$$

where the tensor $K^{mn} := \omega^{mn} - \chi^{mn}$ denotes the contorsion and $M^{mn} := \beta_1 e^m e^n + 2\beta_2 e^{[m} f^{n]} + 3\beta_3 f^m f^n$.

3. Mean field massive gravity

Consider mGR propagating in an arbitrary fiducial (pseudo-) Riemannian manifold $(M, \bar{g}_{\mu\nu})$ with corresponding vierbeine and spin connections (f^m, χ^{mn}) . Now let (e^m, ω^{mn}) be a solution to the mGR equations of motion (1). We wish to study fluctuations $(\varepsilon^m, \lambda^{mn})$ about this configuration:

$$\tilde{e}^m = e^m + \varepsilon^m, \quad \tilde{\omega}^{mn} = \omega^{mn} + \lambda^{mn}.$$

The action governing these is the quadratic part of $S_{\text{mGR}}[\tilde{e}, \tilde{\omega}; f] - S_{\text{mGR}}[e, \omega; f]$, namely

$$S[h, \lambda; e, f] := -\frac{1}{2} \int \epsilon_{mnr s} \left[e^m \varepsilon^n \nabla \lambda^{rs} + \frac{1}{2} (e^m e^n \lambda^r_t \lambda^{ts} + R^{mn} \varepsilon^r \varepsilon^s) - m^2 (3\beta_0 e^m e^n \varepsilon^r \varepsilon^s + 2\beta_1 e^m f^n \varepsilon^r \varepsilon^s + \beta_2 f^m f^n \varepsilon^r \varepsilon^s) \right].$$

The mean field model is a theory of forty dynamical fields $(\varepsilon^m, \lambda^{mn})$. In the above, ∇ is the Levi-Civita connection of e^m , and R^{mn} its Riemann tensor; we stress that henceforth the fiducial field $(f^m, \chi^{mn}(f), \bar{g}_{\mu\nu}(f))$ and mGR background fields $(e^m, \omega^{mn}(e), g_{\mu\nu}(e))$ are *non-dynamical*; all index manipulations will be carried out using the mGR background metric and vierbein.

The $\bar{m}GR$ equations of motion are

$$\begin{aligned} \mathcal{T}^m &:= \nabla \varepsilon^m + \lambda^{mn} e_n \approx 0, \\ \mathcal{G}_m &:= \frac{1}{2} \epsilon_{mnr s} [e^n \nabla \lambda^{rs} + \varepsilon^n R^{rs}] - m^2 \tau_m \approx 0, \end{aligned} \quad (2)$$

where $\tau_m := \epsilon_{mnr s} [3\beta_0 e^n e^r \varepsilon^s + 2\beta_1 e^n f^r \varepsilon^s + \beta_2 f^n f^r \varepsilon^s]$.

4. Mean field degrees of freedom

In principle, since we are describing the linearization of a model whose constraints have been completely analyzed in [7], we know *a priori* that $\bar{m}GR$ describes five propagating degrees of freedom. However, for completeness and our causality study, we reanalyze its constraints.

The first step is to introduce a putative choice of time coordinate t , which for now need not rely in any way on either the fiducial or background metric, and use this to decompose any p -form θ (with $p < 4$) as

$$\theta := \theta + \dot{\theta}, \quad (3)$$

where $\dot{\theta} \wedge dt = 0$. Thus θ is the purely spatial part of the form θ . Hence for any on-shell relation $\mathcal{P} \approx 0$ polynomial in $(\nabla, \varepsilon, \lambda)$,

¹ Here d is the exterior derivative and the dynamical vierbeine and spin connection (e, ω) , are one-forms. We suppress wedge products unless necessary for clarity.

² The first of these assumes invertibility of the operator M^{mn} as a map from two-forms to antisymmetric Lorentz tensors; we shall always work on the model's branch where this holds.

its spatial $\mathcal{P} \approx 0$ part is a constraint because it contains no t -derivatives.

Thus we immediately find *sixteen* primary constraints:

$$\begin{aligned} \mathcal{T}^m &= \nabla \mathbf{e}^m + \lambda^{mn} \mathbf{e}_n \approx 0 \approx \mathcal{G}_m \\ &= \frac{1}{2} \epsilon_{mnr s} [\mathbf{e}^n \nabla \lambda^{rs} + \mathbf{e}^n \mathbf{R}^{rs}] - m^2 \boldsymbol{\tau}_m. \end{aligned}$$

There are *ten* secondary constraints in total. The first six of these follow from the integrability condition $e^{[m} \nabla \mathcal{T}^{n]} \approx 0$ which yields the so-called *symmetry constraint*

$$e_{[m} \boldsymbol{\tau}_{n]} + \varepsilon_{[m} \dot{t}_{n]} \approx 0.$$

As mentioned above, we assume that the set of two forms $\{M^{mn}\}$ is a basis for the space of two-forms, so the symmetry constraint yields

$$\varepsilon^m f_m \approx 0.$$

The remaining four secondary constraints come from the covariant curl $\nabla \mathcal{G}_m \approx 0$ and give the *vector constraint*

$$\nabla \boldsymbol{\tau}_m + \lambda_m{}^n \dot{t}_n \approx 0.$$

Employing the equation of motion $\mathcal{T}^m \approx 0$, this implies

$$\epsilon_{mnr s} [M^{mn} \lambda^{rs} + 2(\beta_1 \mathbf{e}^m + \beta_2 f^m) \varepsilon^n K^{rs}] \approx 0.$$

Finally there are *four* tertiary constraints stemming from covariant curls of the secondaries: the temporal part of the *curled symmetry constraint* $K_{mn} \varepsilon^m f^n + \lambda_{mn} \varepsilon^m f^n \approx 0$, i.e.

$$\begin{aligned} \dot{K}_{mn} \mathbf{e}^m f^n + \mathbf{K}_{mn} \dot{\varepsilon}^m f^n + \mathbf{K}_{mn} \varepsilon^m \dot{f}^n + \dot{\lambda}_{mn} \mathbf{e}^m f^n \\ + \lambda_{mn} \dot{\varepsilon}^m f^n + \lambda_{mn} \varepsilon^m \dot{f}^n \approx 0, \end{aligned}$$

and the *scalar constraint*

$$\begin{aligned} \epsilon_{mnr s} \left(\beta_1 (\varepsilon^m \dot{e}^t + \varepsilon^m \varepsilon^t) - 2\beta_2 \varepsilon^{(m} f^{t)} \right) K^{nr} K^s{}_t \\ + \epsilon_{mnr s} \left(\beta_1 \mathbf{e}^m \dot{e}^t - 2\beta_2 \mathbf{e}^{(m} f^{t)} - 3\beta_3 f^m f^t \right) (\lambda^{nr} K^s{}_t + K^{nr} \lambda^s{}_t) \\ + 2m^2 \beta_1 \varepsilon^m \dot{t}_m + 2m^2 (\beta_1 \varepsilon^m + 2\beta_2 f^m) \boldsymbol{\tau}_m \\ + 3\epsilon_{mnr s} \beta_3 f^m f^n \nabla \lambda^{rs} - 4\beta_2 \varepsilon^m \bar{G}_m - 2\epsilon_{mnr s} \beta_1 \varepsilon^m \mathbf{e}^n \bar{R}^{rs} \approx 0. \end{aligned}$$

The $\nabla \lambda^{rs}$ term seems to indicate that the above display is not a constraint for $\beta_3 \neq 0$, however, as shown in [7], this quantity (weakly) equals one without time derivatives of fields. In summary, the model describes forty fields subject to thirty constraints and thus propagates five³ DoF.

5. Characteristic matrix

We now study whether $\bar{m}GR$ can propagate initial data off a given hypersurface Σ . This amounts to asking if derivatives normal to Σ are determined by the equations of motion (2). For that, we simply replace all derivatives in the equations of motion and gradients of their constraints by the normal covector ξ_μ to Σ multiplied by the normal derivative of the corresponding field:

³ Although this conclusion for $\bar{m}GR$ is guaranteed by previous studies [3,4,7] of the non-linear model's DoF count, it verifies the linearized mGR study [17]. In that work, the fiducial metric is eliminated in terms of the mean field in order to argue that linearized spin 2 fields can propagate consistently in any gravitational background. This result is consistent with earlier work in [18] which relies on a $1/m^2$ expansion to study leading DoF and causality properties of gravitating, massive spin 2 models.

$$\partial_\mu \varepsilon^m \Big|_\Sigma = \xi_\mu \partial_n \varepsilon^m \quad \text{and} \quad \partial_\mu \lambda^{mn} \Big|_\Sigma = \xi_\mu \partial_n \lambda^{mn}.$$

We will also, for simplicity alone, restrict to the parameter choices $\beta_2 = \beta_3 = 0$ (the model's characteristic matrix for its entire parameter range has been computed in [7]). In particular, we must focus on the question whether the linear system of equations for the normal derivatives $\partial_n \varphi := (\partial_n \varepsilon, \partial_n \lambda)$ implied by the equations of motion along Σ is invertible. This amounts to a matrix problem encoded by the theory's characteristic matrix \mathcal{C} . In what follows we compute the system of equations given by the homogeneous linear system $\mathcal{C} \cdot \partial_n \varphi = 0$. Starting with the equations of motion we find

$$\begin{aligned} \xi \wedge \partial_n \varepsilon^m &= 0, \\ \epsilon_{mnr s} \mathbf{e}^n \wedge \xi \wedge \partial_n \lambda^{rs} &= 0. \end{aligned} \quad (4)$$

The gradients of the secondary and tertiary constraints then imply

$$\begin{aligned} \xi_\mu f_m \wedge \partial_n \varepsilon^m &= 0 = \xi_\mu \epsilon_{mnr s} \mathbf{e}^m \wedge [\mathbf{e}^n \wedge \partial_n \lambda^{rs} - 2K^{nr} \wedge \partial_n \varepsilon^s], \\ \xi_\mu f^m \wedge [K_{mn} \wedge \partial_n \varepsilon^n + \mathbf{e}^n \wedge \partial_n \lambda_{mn}] &= 0, \\ \xi_\mu \epsilon_{mnr s} \mathbf{e}^m \wedge [\mathbf{e}^n \wedge K^{rt} \wedge \partial_n \lambda_t^s \\ + (K^{nt} \wedge K_t^r - \bar{R}^{nr} + m^2(4\beta_0 \mathbf{e}^n \wedge \mathbf{e}^r + 3\beta_1 \mathbf{e}^n \wedge f^r)) \wedge \partial_n \varepsilon^s] &= 0. \end{aligned} \quad (5)$$

In the above the prefactor ξ_μ was included to indicate the origin of these equations but can be removed with impunity. To handle Equation (4) we decompose form-valued normal derivatives as earlier in Equation (3), and find⁴

$$\partial_n \mathbf{e}^m = 0 = \partial_n \lambda^{mn}.$$

Supposing that the one-form $\xi = dt$, for some evolution coordinate t , we now use a shorthand notation $\partial_n \varepsilon^m = dt \dot{\varepsilon}_t^m$ and $\partial_n \lambda^{mn} = dt \dot{\lambda}_t^{mn}$. We thus have the *reduced characteristic system*

$$\begin{pmatrix} \mathbf{f}_m & 0 \\ 2\epsilon_{mnr s} \mathbf{e}^n \times \mathbf{K}^{rs} & \epsilon_{mnr s} \mathbf{e}^r \times \mathbf{e}^s \\ \mathbf{f}^n \times \mathbf{K}_{nm} & \mathbf{f}_m \times \mathbf{e}_n \\ \mathcal{R}_m & \mathcal{K}_{mn} \end{pmatrix} \begin{pmatrix} \dot{\varepsilon}_t^m \\ \dot{\lambda}_t^{mn} \end{pmatrix} = 0. \quad (6)$$

In the above square matrix, \times denotes the standard three-dimensional cross product while the spatial densities on its last line can be read off from (5) and are simple for flat fiducial metrics. Vanishing of the determinant of the above 10×10 matrix completely characterizes the boundary of the model's predictive hyperbolic regime (modulo the restriction explained in footnote 4). As we shall see, the reduced characteristic system describes the propagation of superluminal lower helicity modes: In the next section, we specialize to flat fiducial spaces and show how to analyze this determinant in direct analogy with the RS system.

6. Analogy with Rarita–Schwinger

The charged, spin 3/2, Rarita–Schwinger (RS) equation of motion reads

$$\gamma^{\mu\nu\rho} (\nabla_\nu + ieA_\nu + \frac{m}{2} \gamma_\nu) \psi_\rho = 0.$$

Here ∇ is the Levi-Civita connection of the fiducial spacetime and A is the background EM potential. These are analogous to the

⁴ Here we assumed that the pullback of the mGR background vierbeine to the hypersurface Σ is invertible and ignore likely pathologies when this fails.

$\bar{m}GR$ fiducial and background-mean-field metrics. The RS characteristic matrix was computed in [8] for flat fiducial metrics and spacelike hypersurfaces Σ and found to have zero determinant when the magnetic field \mathbf{B} obeyed⁵

$$1 - \left(\frac{2e}{3m^2}\right)^2 \mathbf{B}^2 = 0. \quad (7)$$

The condition $\frac{2e|\mathbf{B}|}{3m^2} < 1$ thus determines the weak field, hyperbolic, regime. Our aim now is to develop the analogous statement for $\bar{m}GR$ and inherit the conclusions of [8].

We begin with a short calculation. Consider now a flat fiducial metric so $f^m = \delta_{\mu}^m dx^{\mu}$ and $d\bar{s}^2 = -dt^2 + d\mathbf{x}^2$. This simplifies the reduced characteristic system considerably. Firstly the equation $\mathbf{f}_m \dot{\epsilon}_t^m = 0$ of (6) implies $\dot{\epsilon}_t^a = 0$, where we have decomposed the Lorentz index $m = (0, a)$. Let us introduce an EM-like notation

$$\epsilon_{abc} \mathbf{K}^{bc} =: \mathbf{B}_a, \quad \mathbf{K}^{0a} =: \mathbf{E}^a.$$

For simplicity only, we now restrict to the case where $\mathbf{e}^0 = \mathbf{E}^a = 0$. Thus using $\epsilon^0_{abc} = \epsilon_{abc}$ the second equation of (6) implies

$$\dot{\lambda}_t^{0a} = -\frac{1}{2} \epsilon^{abc} \mathbf{B}_b \cdot \tilde{\mathbf{e}}_c \dot{\epsilon}_t^0.$$

Here the 3-vectors $\tilde{\mathbf{e}}_a$ form the 3-inverse of \mathbf{e}^a so that $\tilde{\mathbf{e}}_a \cdot \mathbf{e}^b = \delta_a^b$. The third equation of (6) then gives

$$\mathbf{f}_{[a} \times \mathbf{e}_{b]} \dot{\lambda}_t^{ab} = 0.$$

This equation generically allows $\dot{\lambda}_t^{ab}$ to be expressed as a function of $\dot{\epsilon}_t^0$ but this requires a non-trivial condition on \mathbf{e}^a . Under the hypothesis that the eigenvalue spectra of the matrices $\mathbf{f}^a \cdot \tilde{\mathbf{e}}_b$ and $-\mathbf{f}^a \cdot \tilde{\mathbf{e}}_b$ do not intersect, the above equation implies that $\dot{\lambda}_t^{ab} = 0$. In this framework, the last equation of (6) gives the single VZ-type condition

$$\left[m_{\text{FP}}^2 (4 - (\mathbf{f}^a \cdot \tilde{\mathbf{e}}_a)) - \frac{1}{6} \left((\mathbf{B}^a \cdot \tilde{\mathbf{e}}^b) (\mathbf{B}_{[a} \cdot \tilde{\mathbf{e}}_{b]}) - \frac{1}{2} (\mathbf{B}^a \cdot \tilde{\mathbf{e}}_{[a}) (\mathbf{B}^b \cdot \tilde{\mathbf{e}}_{b]}) \right) \right] \dot{\epsilon}_t^0 = 0. \quad (8)$$

In the weak field limit where the mean field approaches Minkowski space, the coefficient of m_{FP}^2 approaches unity but can change sign in a strong-field, large $\tilde{\mathbf{e}}$ limit. Hence there are certainly strong field configurations where the model loses hyperbolicity [10] and closed causal curves are unavoidable. (This signals the onset of strong coupling in an effective field theory.) Now, comparing Equations (7) and (8), we see that we have reduced $\bar{m}GR$'s weak field propagation analysis to a previous – well understood – case. Finally, we note that the same analogy and characteristic method can also be applied to the bimetric theory by treating the two background metrics as a (fiducial, background) pair [20], and clearly will lead to similar findings as exhibited here for mGR.

7. Conclusions

mGR is not a fundamental theory but rather an effective one with a range of validity determined by requiring hyperbolicity in a weak field regime. Excepting the further caveats explained in the text, the reduced characteristic matrix of Equation (6) completely determines this allowed regime. Even in the weak regime, modes exhibit a crystal structure with differing helicities' maximal propagation speeds. For mGR to give a useful effective theory for

physical applications, one must couple to matter (or at least photons) and require consistent causal cones for all modes. Once these couplings are decided upon, the characteristic method will determine their effective range of validity, if any. There is, in principle, also the logical possibility that there exists a causal, luminal, UV completion of mGR analogous to that for QED in curved space [21]. The general discussion of UV completions [22] supports the conclusion that quantizing these models is unlikely to improve their causal consistency properties – indeed quite the opposite.

Acknowledgements

We thank C. Deffayet, S. Dubovsky, K. Hinterbichler, K. Izumi, M. Porrati and Y.C. Ong for discussions. S.D. was supported in part by grants NSF PHY-1266107 and DOE # de-sc0011632. A.W. was supported in part by a Simons Foundation Collaboration Grant for Mathematicians, ID 317562. G.Z. was supported in part by DOE Grant DE-FG03-91ER40674. This research was supported in part by Perimeter Institute for Theoretical Physics. This research was supported in part by the Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Economic Development and Innovation. A.W. and G.Z. thank the Perimeter Institute for an illuminating ‘‘Superluminality in Effective Field Theories for Cosmology’’ workshop.

References

- [1] D. Boulware, S. Deser, Phys. Rev. D 6 (1972) 3368; D. Boulware, S. Deser, Phys. Lett. B 40 (2) (1972) 227.
- [2] C. de Rham, G. Gabadadze, Phys. Rev. D 82 (2010) 044020, arXiv:1007.0443v2 [hep-th]; C. de Rham, G. Gabadadze, A.J. Tolley, Phys. Rev. Lett. 106 (2011) 231101, arXiv:1011.1232v2 [hep-th]; C. de Rham, G. Gabadadze, A.J. Tolley, Phys. Lett. B 711 (2012) 190, arXiv:1107.3820v1 [hep-th]; C. de Rham, G. Gabadadze, A.J. Tolley, J. High Energy Phys. 1111 (2011) 093, arXiv:1108.4521v2 [hep-th].
- [3] S.F. Hassan, R.A. Rosen, Phys. Rev. Lett. 108 (2012) 041101, arXiv:1106.3344v3 [hep-th]; S.F. Hassan, R.A. Rosen, J. High Energy Phys. 1204 (2012) 123, arXiv:1111.2070v1 [hep-th]; S.F. Hassan, R.A. Rosen, A. Schmidt-May, J. High Energy Phys. 1202 (2012) 026, arXiv:1109.3230v2 [hep-th]; S. Hassan, A. Schmidt-May, M. von Strauss, Phys. Lett. B 715 (2012) 335, arXiv:1203.5283v2 [hep-th]; See, however, L. Albarte, A.H. Chamseddine, V. Mukhanov, J. High Energy Phys. 1104 (2011) 004, arXiv:1011.0183 [hep-th]; A.H. Chamseddine, V. Mukhanov, J. High Energy Phys. 1108 (2011) 091, arXiv:1106.5868 [hep-th].
- [4] C. Deffayet, J. Mourad, G. Zahariade, J. Cosmol. Astropart. Phys. 1301 (2013) 032, arXiv:1207.6338 [hep-th].
- [5] S. Deser, A. Waldron, Phys. Rev. Lett. 110 (2013) 111101, arXiv:1212.5835v3 [hep-th].
- [6] S. Deser, M. Sandora, A. Waldron, Phys. Rev. D 87 (2013) 101501, arXiv:1301.5621 [hep-th].
- [7] S. Deser, M. Sandora, A. Waldron, G. Zahariade, Phys. Rev. D 90 (2014) 104043, arXiv:1408.0561 [hep-th].
- [8] G. Velo, D. Zwanziger, Phys. Rev. 186 (1969) 1337; G. Velo, D. Zwanziger, Phys. Rev. 188 (1969) 2218.
- [9] M. Kobayashi, A. Shamaly, Phys. Rev. D 17 (1978) 2179; M. Kobayashi, A. Shamaly, Prog. Theor. Phys. 61 (1979) 656; S. Deser, V. Pascalutsa, A. Waldron, Phys. Rev. D 62 (2000) 105031, arXiv:hep-th/0003011; I.L. Buchbinder, D.M. Gitman, V.D. Pershin, Phys. Lett. B 492 (2000) 161, arXiv:hep-th/0006144; S. Deser, A. Waldron, Nucl. Phys. B 631 (2002) 369, arXiv:hep-th/0112182.
- [10] See, for example, R. Courant, D. Hilbert, Methods of Mathematical Physics, Differential Equations, Wiley, 2008.
- [11] K. Johnson, E.C.G. Sudarshan, Ann. Phys. 13 (1961) 126.
- [12] See, for example, J. Bjorken, S. Drell, Relativistic Quantum Fields, McGraw–Hill, 1965.
- [13] K. Izumi, Y.C. Ong, Class. Quantum Gravity 30 (2013) 184008, arXiv:1304.0211 [hep-th].

⁵ See [19] for models designed to cure this pathology by adding higher background derivative, string-inspired terms to the RS action.

- [14] S. Deser, K. Izumi, Y.C. Ong, A. Waldron, *Phys. Lett. B* 726 (2013) 544, arXiv:1306.5457 [hep-th];
S. Deser, K. Izumi, Y.C. Ong, A. Waldron, Superluminal propagation and acausality of nonlinear massive gravity, in: *Proceedings, Conference in Honor of the 90th Birthday of Freeman Dyson*, Singapore, 2013, World Scientific, 2014, arXiv:1312.1115 [hep-th].
- [15] C. Burrage, C. de Rham, L. Heisenberg, A.J. Tolley, *J. Cosmol. Astropart. Phys.* 1207 (2012) 004, arXiv:1111.5549 [hep-th];
P. de Fromont, C. de Rham, L. Heisenberg, A. Matas, Superluminality in the bi- and multi-Galileon, arXiv:1303.0274 [hep-th];
A. Gruzinov, All Fierz–Paulian massive gravity theories have ghosts or superluminal modes, arXiv:1106.3972 [hep-th].
- [16] B. Zumino, Effective Lagrangians and broken symmetries, in: S. Deser, M. Grisaru, H. Pendleton (Eds.), *Brandeis Univ. Lectures on Elementary Particles and Quantum Field Theory*, vol. 2, MIT Press, Cambridge, MA, 1970, p. 437.
- [17] L. Bernard, C. Deffayet, M. von Strauss, arXiv:1410.8302 [hep-th].
- [18] I.L. Buchbinder, D.M. Gitman, V.A. Krykhtin, V.D. Pershin, *Nucl. Phys. B* 584 (2000) 615, arXiv:hep-th/9910188;
I.L. Buchbinder, D.M. Gitman, V.D. Pershin, *Phys. Lett. B* 492 (2000) 161.
- [19] M. Porrati, R. Rahman, A. Sagnotti, *Nucl. Phys. B* 846 (2011) 250, arXiv:1011.6411 [hep-th];
M. Porrati, R. Rahman, *Phys. Rev. D* 80 (2009) 025009, arXiv:0906.1432 [hep-th];
M. Porrati, R. Rahman, *Phys. Rev. D* 84 (2011) 045013, arXiv:1103.6027 [hep-th].
- [20] S. Deser, A. Waldron, G. Zahariade, in preparation.
- [21] T.J. Hollowood, G.M. Shore, *Nucl. Phys. B* 795 (2008) 138, arXiv:0707.2303 [hep-th].
- [22] A. Adams, N. Arkani-Hamed, S. Dubovsky, A. Nicolis, R. Rattazzi, *J. High Energy Phys.* 0610 (2006) 014, arXiv:hep-th/0602178.