



# A proof of the strong no loop conjecture

Kiyoshi Igusa<sup>a</sup>, Shiping Liu<sup>b,\*</sup>, Charles Paquette<sup>b</sup>

<sup>a</sup> Department of Mathematics, Brandeis University, Waltham, MA 02453, United States

<sup>b</sup> Département de Mathématiques, Université de Sherbrooke, Sherbrooke, Québec, Canada, J1K 2R1

Received 28 March 2011; accepted 28 June 2011

Available online 12 August 2011

Communicated by Henning Krause

---

## Abstract

The strong no loop conjecture states that a simple module of finite projective dimension over an artin algebra has no non-zero self-extension. The main result of this paper establishes this well known conjecture for finite dimensional algebras over an algebraically closed field.

© 2011 Elsevier Inc. All rights reserved.

*Keywords:* Artin algebra; Module; Projective dimension; Hochschild homology; Quiver; Strong no loop conjecture

---

## Introduction

Let  $A$  be an artin algebra, and denote by  $\text{mod}A$  the category of finitely generated right  $A$ -modules. It is an important problem in the representation theory of algebras to determine whether  $A$  has finite or infinite global dimension, and more specifically, whether a simple  $A$ -module has finite or infinite projective dimension. For instance, the derived category  $D^b(\text{mod}A)$  has Auslander–Reiten triangles if and only if  $A$  has finite global dimension; see [7,8]. One approach to this problem is to consider the extension quiver of  $A$ , which has vertices given by a complete set of non-isomorphic simple  $A$ -modules and single arrows  $S \rightarrow T$ , where  $S$  and  $T$  are vertices such that  $\text{Ext}_A^1(S, T)$  does not vanish. Then the *no loop conjecture* affirms that the extension quiver of  $A$  contains no loop if  $A$  is of finite global dimension, while the *strong no*

---

\* Corresponding author.

*E-mail addresses:* [igusa@brandeis.edu](mailto:igusa@brandeis.edu) (K. Igusa), [shiping.liu@usherbrooke.ca](mailto:shiping.liu@usherbrooke.ca) (S. Liu), [charles.paquette@usherbrooke.ca](mailto:charles.paquette@usherbrooke.ca) (C. Paquette).

*loop conjecture*, which is due to Zacharia, strengthens this to state that a vertex in the extension quiver admits no loop if it has finite projective dimension; see [1,10].

The no loop conjecture was first explicitly established for artin algebras of global dimension two; see [5]. For finite dimensional elementary algebras (that is, the simple modules are all one dimensional), as shown in [10], this can be easily derived from an earlier result of Lenzing on Hochschild homology in [13]. Lenzing's technique was to extend the notion of the trace of endomorphisms of projective modules, defined by Hattori and Stallings in [9,18], to endomorphisms of modules over a noetherian ring with finite global dimension, and apply it to a particular kind of filtration for the regular module.

In contrast, up to now, the strong no loop conjecture has only been verified for some special classes of algebras such as monomial algebras; see [2,10], special biserial algebras; see [14], and algebras with at most two simple modules and radical cubed zero; see [12]. Many other partial results can be found in [3,4,6,15,16,20]. Most recently, Skorodumov generalized and localized Lenzing's filtration to indecomposable projective modules. This allowed him to prove this conjecture for finite dimensional elementary algebras of finite representation type; see [17].

In this paper, we shall localize Lenzing's trace function to endomorphisms of modules in  $\text{mod } \Lambda$  with an  $e$ -bounded projective resolution, where  $e$  is an idempotent in  $\Lambda$ . The key point is that every module in  $\text{mod } \Lambda$  has an  $e$ -bounded projective resolution whenever the semi-simple module supported by  $e$  is of finite injective dimension. This enables us to obtain a local version of Lenzing's result. As a consequence, we shall solve the strong no loop conjecture for a large class of artin algebras including finite dimensional elementary algebras over any field, and particularly, for finite dimensional algebras over an algebraically closed field.

## 1. Localized trace and Hochschild homology

Throughout,  $J$  will stand for the Jacobson radical of  $\Lambda$ . The additive subgroup of  $\Lambda$  generated by the elements  $ab - ba$  with  $a, b \in \Lambda$  is called the *commutator group* of  $\Lambda$  and written as  $[\Lambda, \Lambda]$ . One defines then the Hochschild homology group  $\text{HH}_0(\Lambda)$  to be  $\Lambda/[\Lambda, \Lambda]$ . We shall say that  $\text{HH}_0(\Lambda)$  is *radical-trivial* if  $J \subseteq [\Lambda, \Lambda]$ .

To start with, we recall the notion of the trace of an endomorphism  $\varphi$  of a projective module  $P$  in  $\text{mod } \Lambda$ , as defined by Hattori and Stallings in [9,18]; see also [10,13]. Write  $P = e_1 \Lambda \oplus \cdots \oplus e_r \Lambda$ , where the  $e_i$  are primitive idempotents in  $\Lambda$ . Then  $\varphi = (a_{ij})_{r \times r}$ , where  $a_{ij} \in e_i \Lambda e_j$ . The *trace* of  $\varphi$  is defined to be

$$\text{tr}(\varphi) = \sum_{i=1}^r a_{ii} + [\Lambda, \Lambda] \in \text{HH}_0(\Lambda).$$

We shall collect some well known properties of this trace function in the following proposition, in which statement (2) is precisely the reason for defining the trace to be an element in  $\text{HH}_0(\Lambda)$ .

**1.1. Proposition** (*Hattori–Stallings*). *Let  $P, P'$  be projective modules in  $\text{mod } \Lambda$ .*

- (1) *If  $\varphi, \psi \in \text{End}_\Lambda(P)$ , then  $\text{tr}(\varphi + \psi) = \text{tr}(\varphi) + \text{tr}(\psi)$ .*
- (2) *If  $\varphi : P \rightarrow P'$  and  $\psi : P' \rightarrow P$  are  $\Lambda$ -linear, then  $\text{tr}(\varphi\psi) = \text{tr}(\psi\varphi)$ .*
- (3) *If  $\varphi = (\varphi_{ij})_{2 \times 2} : P \oplus P' \rightarrow P \oplus P'$ , then  $\text{tr}(\varphi) = \text{tr}(\varphi_{11}) + \text{tr}(\varphi_{22})$ .*

- (4) If  $\psi : P \rightarrow P'$  is an isomorphism and  $\varphi \in \text{End}_\Lambda(P)$ , then  $\text{tr}(\psi\varphi\psi^{-1}) = \text{tr}(\varphi)$ .
- (5) If  $\varphi : \Lambda \rightarrow \Lambda$  is the left multiplication by  $a \in \Lambda$ , then  $\text{tr}(\varphi) = a + [\Lambda, \Lambda]$ .

Next, we recall Lenzing’s extension of this function to endomorphisms of modules of finite projective dimension. For  $M \in \text{mod}\Lambda$ , let  $\mathcal{P}_M$  denote a projective resolution of  $M$  in  $\text{mod}\Lambda$  as follows:

$$\cdots \longrightarrow P_i \xrightarrow{d_i} P_{i-1} \longrightarrow \cdots \longrightarrow P_0 \xrightarrow{d_0} M \longrightarrow 0.$$

For each  $\varphi \in \text{End}_\Lambda(M)$ , one can construct a commutative diagram

$$\begin{array}{ccccccccccc} \cdots & \longrightarrow & P_i & \xrightarrow{d_i} & P_{i-1} & \longrightarrow & \cdots & \longrightarrow & P_0 & \xrightarrow{d_0} & M & \longrightarrow & 0 \\ & & \downarrow \varphi_i & & \downarrow \varphi_{i-1} & & & & \downarrow \varphi_0 & & \downarrow \varphi & & \\ \cdots & \longrightarrow & P_i & \xrightarrow{d_i} & P_{i-1} & \longrightarrow & \cdots & \longrightarrow & P_0 & \xrightarrow{d_0} & M & \longrightarrow & 0 \end{array}$$

in  $\text{mod}\Lambda$ . We shall call  $\{\varphi_i\}_{i \geq 0}$  a *lifting* of  $\varphi$  to  $\mathcal{P}_M$ . If  $M$  is of finite projective dimension, then one may assume that  $\mathcal{P}_M$  is bounded and define the *trace* of  $\varphi$  by

$$\text{tr}(\varphi) = \sum_{i=0}^{\infty} (-1)^i \text{tr}(\varphi_i) \in \text{HH}_0(\Lambda),$$

which is independent of the choice of  $\mathcal{P}_M$  and  $\{\varphi_i\}$ ; see [13], and also [10].

Our strategy is to localize this construction. Let  $e$  be an idempotent in  $\Lambda$ . Set

$$\Lambda_e = \Lambda/\Lambda(1 - e)\Lambda.$$

The canonical algebra projection  $\Lambda \rightarrow \Lambda_e$  induces a group homomorphism

$$H_e : \text{HH}_0(\Lambda) \rightarrow \text{HH}_0(\Lambda_e).$$

For an endomorphism  $\varphi$  of a projective module in  $\text{mod}\Lambda$ , we define its *e-trace* by

$$\text{tr}_e(\varphi) = H_e(\text{tr}(\varphi)) \in \text{HH}_0(\Lambda_e).$$

It is evident that this *e-trace* function has the properties (1) to (4) stated in Proposition 1.1. We shall state another important property in the following result. For doing so, we recall that the *top* of a module in  $\text{mod}\Lambda$  is the quotient of the module by its Jacobson radical.

**1.2. Lemma.** *Let  $e$  be an idempotent in  $\Lambda$ , and let  $P$  be a projective module in  $\text{mod}\Lambda$  whose top is annihilated by  $e$ . If  $\varphi \in \text{End}_\Lambda(P)$ , then  $\text{tr}_e(\varphi) = 0$ .*

**Proof.** We may assume that  $P$  is non-zero. Then  $1 - e = e_1 + \cdots + e_r$ , where the  $e_i$  are pairwise orthogonal primitive idempotents in  $\Lambda$ . Let  $\varphi \in \text{End}_\Lambda(P)$ . By Proposition 1.1(3), we may assume

that  $P$  is indecomposable. Then  $P \cong e_s \Lambda$  for some  $1 \leq s \leq r$ . By Proposition 1.1(4), we may assume that  $P = e_s \Lambda$ . Then  $\varphi$  is the left multiplication by some  $a \in e_s \Lambda e_s$ . By Proposition 1.1(5),

$$\text{tr}_e(\varphi) = H_e(a + [\Lambda, \Lambda]) = \bar{a} + [\Lambda_e, \Lambda_e],$$

where  $\bar{a} = a + \Lambda(1 - e)\Lambda$ . Since  $a = e_s a e_s = (1 - e)a(1 - e) \in \Lambda(1 - e)\Lambda$ , we get  $\text{tr}_e(\varphi) = 0$ .  $\square$

In order to extend the  $e$ -trace function, we shall say that a projective resolution  $\mathcal{P}_M$  of a module  $M$  in  $\text{mod}\Lambda$  is  $e$ -bounded if all but finitely many tops of the terms in  $\mathcal{P}_M$  are annihilated by  $e$ . In this case, if  $\varphi$  is an endomorphism of  $M$  with a lifting  $\{\varphi_i\}_{i \geq 0}$  to  $\mathcal{P}_M$ , then it follows from Lemma 1.2 that  $\text{tr}_e(\varphi_i) = 0$  for all but finitely many  $i$ . This allows us to define the  $e$ -trace of  $\varphi$  by

$$\text{tr}_e(\varphi) = \sum_{i=0}^{\infty} (-1)^i \text{tr}_e(\varphi_i) \in \text{HH}_0(\Lambda_e).$$

**1.3. Lemma.** *Let  $e$  be an idempotent in  $\Lambda$ . Then the  $e$ -trace is well defined for endomorphisms of modules in  $\text{mod}\Lambda$  having an  $e$ -bounded projective resolution.*

**Proof.** Let  $M$  be a module in  $\text{mod}\Lambda$  having an  $e$ -bounded projective resolution

$$\mathcal{P}_M: \quad \cdots \longrightarrow P_i \xrightarrow{d_i} P_{i-1} \longrightarrow \cdots \longrightarrow P_0 \xrightarrow{d_0} M \longrightarrow 0.$$

Fix  $\varphi \in \text{End}_\Lambda(M)$ . We first show that  $\text{tr}_e(\varphi)$  is independent of the choice of a lifting of  $\varphi$  to  $\mathcal{P}_M$ . By Proposition 1.1(1), it suffices to prove, for any lifting  $\{\psi_i\}_{i \geq 0}$  of the zero endomorphism of  $M$ , that  $\sum_{i=0}^{\infty} (-1)^i \text{tr}_e(\psi_i) = 0$ . Indeed, let  $h_i : P_i \rightarrow P_{i+1}$  be morphisms such that  $\psi_0 = d_1 h_0$  and  $\psi_i = d_{i+1} h_i + h_{i-1} d_i$ . By Proposition 1.1,  $\text{tr}_e(\psi_i) = \text{tr}_e(d_{i+1} h_i) + \text{tr}_e(h_{i-1} d_i) = \text{tr}_e(d_{i+1} h_i) + \text{tr}_e(d_i h_{i-1})$ , for all  $i \geq 1$ .

On the other hand, by assumption, there exists some  $m \geq 0$  such that the top of  $P_i$  is annihilated by  $e$ , for every  $i \geq m$ . By Lemma 1.2,  $\text{tr}_e(d_{m+1} h_m) = 0$  and  $\text{tr}_e(\psi_i) = 0$  for all  $i \geq m$ . This yields

$$\begin{aligned} \sum_{i=0}^{\infty} (-1)^i \text{tr}_e(\psi_i) &= \text{tr}_e(\psi_0) + \sum_{i=1}^m (-1)^i \text{tr}_e(\psi_i) \\ &= \text{tr}_e(d_1 h_0) + \sum_{i=1}^m (-1)^i (\text{tr}_e(d_{i+1} h_i) + \text{tr}_e(d_i h_{i-1})) \\ &= (-1)^m \text{tr}_e(d_{m+1} h_m) \\ &= 0. \end{aligned}$$

Next, we verify that  $\text{tr}_e(\varphi)$  is independent of the choice of the  $e$ -bounded projective resolution  $\mathcal{P}_M$ . Suppose that  $M$  has another  $e$ -bounded projective resolution

$$\mathcal{P}'_M : \quad \cdots \longrightarrow P'_i \xrightarrow{d'_i} P'_{i-1} \longrightarrow \cdots \longrightarrow P'_0 \xrightarrow{d'_0} M \longrightarrow 0.$$

Considering  $\varphi$ , we get morphisms  $u_i : P_i \rightarrow P'_i$  with  $i \geq 0$  such that  $d'_0 u_0 = \varphi d_0$  and  $d'_i u_i = u_{i-1} d_i$  for  $i \geq 1$ . Similarly, considering the identity map  $1_M$ , we obtain maps  $v_i : P'_i \rightarrow P_i$  with  $i \geq 0$  such that  $d_0 v_0 = d'_0$  and  $d_i v_i = v_{i-1} d'_i$  for  $i \geq 1$ . Observe that  $\{v_i u_i\}_{i \geq 0}$  and  $\{u_i v_i\}_{i \geq 0}$  are liftings of  $\varphi$  to  $\mathcal{P}_M$  and  $\mathcal{P}'_M$ , respectively. By Proposition 1.1(2),

$$\sum_{i=0}^{\infty} (-1)^i \operatorname{tr}_e(u_i v_i) = \sum_{i=0}^{\infty} (-1)^i \operatorname{tr}_e(v_i u_i). \quad \square$$

The following result says that the  $e$ -trace function is additive in some generalized Grothendieck group defined in [10].

**1.4. Proposition.** *Let  $e$  be an idempotent in  $\Lambda$ . Consider a commutative diagram*

$$\begin{array}{ccccccccc} 0 & \longrightarrow & L & \xrightarrow{u} & M & \xrightarrow{v} & N & \longrightarrow & 0 \\ & & \downarrow \varphi_L & & \downarrow \varphi_M & & \downarrow \varphi_N & & \\ 0 & \longrightarrow & L & \xrightarrow{u} & M & \xrightarrow{v} & N & \longrightarrow & 0 \end{array}$$

in  $\operatorname{mod} \Lambda$  with exact rows. If  $L, N$  have  $e$ -bounded projective resolutions, then so does  $M$  and  $\operatorname{tr}_e(\varphi_M) = \operatorname{tr}_e(\varphi_L) + \operatorname{tr}_e(\varphi_N)$ .

**Proof.** Assume that  $L$  and  $N$  have  $e$ -bounded projective resolutions as follows:

$$\mathcal{P}_L : \quad \cdots \longrightarrow P_i \xrightarrow{d_i} P_{i-1} \longrightarrow \cdots \longrightarrow P_0 \xrightarrow{d_0} L \longrightarrow 0$$

and

$$\mathcal{P}_N : \quad \cdots \longrightarrow P'_i \xrightarrow{d'_i} P'_{i-1} \longrightarrow \cdots \longrightarrow P'_0 \xrightarrow{d'_0} N \longrightarrow 0.$$

By the Horseshoe lemma, there exists in  $\operatorname{mod} \Lambda$  a commutative diagram

$$\begin{array}{ccccccccc} \cdots & \longrightarrow & P_i & \xrightarrow{d_i} & P_{i-1} & \longrightarrow & \cdots & \longrightarrow & P_0 & \xrightarrow{d_0} & L & \longrightarrow & 0 \\ & & \downarrow q_i & & \downarrow q_{i-1} & & & & \downarrow q_0 & & \downarrow u & & \\ \cdots & \longrightarrow & P_i \oplus P'_i & \xrightarrow{d''_i} & P_{i-1} \oplus P'_{i-1} & \longrightarrow & \cdots & \longrightarrow & P_0 \oplus P'_0 & \xrightarrow{d''_0} & M & \longrightarrow & 0 \\ & & \downarrow p_i & & \downarrow p_{i-1} & & & & \downarrow p_0 & & \downarrow v & & \\ \cdots & \longrightarrow & P'_i & \xrightarrow{d'_i} & P'_{i-1} & \longrightarrow & \cdots & \longrightarrow & P'_0 & \xrightarrow{d'_0} & N & \longrightarrow & 0 \end{array}$$

with exact rows, where  $q_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ,  $p_i = (0, 1)$  for all  $i \geq 0$ . In particular, the middle row is an  $e$ -bounded projective resolution of  $M$  which we denote by  $\mathcal{P}_M$ . Choose a lifting  $\{f_i\}_{i \geq 0}$  of  $\varphi_L$

to  $\mathcal{P}_L$  and a lifting  $\{g_i\}_{i \geq 0}$  of  $\varphi_N$  to  $\mathcal{P}_N$ . It is well known; see, for example, [19, p. 46] that there exists a lifting  $\{h_i\}_{i \geq 0}$  of  $\varphi_M$  to  $\mathcal{P}_M$  such that

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & P_i & \xrightarrow{q_i} & P_i \oplus P'_i & \xrightarrow{p_i} & P'_i & \longrightarrow & 0 \\
 & & \downarrow f_i & & \downarrow h_i & & \downarrow g_i & & \\
 0 & \longrightarrow & P_i & \xrightarrow{q_i} & P_i \oplus P'_i & \xrightarrow{p_i} & P'_i & \longrightarrow & 0
 \end{array}$$

is commutative, for every  $i \geq 0$ . Since  $h_i q_i = q_i f_i$  and  $g_i p_i = p_i h_i$ , we may choose to write  $h_i$  as a  $(2 \times 2)$ -matrix whose diagonal entries are  $f_i$  and  $g_i$ . By Proposition 1.1(3),  $\text{tr}_e(h_i) = \text{tr}_e(f_i) + \text{tr}_e(g_i)$ . Hence,  $\text{tr}_e(\varphi_M) = \text{tr}_e(\varphi_N) + \text{tr}_e(\varphi_L)$ .  $\square$

In the sequel,  $S_e$  will stand for the semi-simple  $\Lambda$ -module  $e\Lambda/eJ$ . The following observation is essential in our investigation.

**1.5. Lemma.** *Let  $e$  be an idempotent in  $\Lambda$ . If  $S_e$  is of finite injective dimension, then the  $e$ -trace is defined for every endomorphism in  $\text{mod}\Lambda$ .*

**Proof.** Suppose that  $S_e$  is of finite injective dimension  $n$ . Let  $M$  be a module in  $\text{mod}\Lambda$  with a minimal projective resolution

$$\mathcal{P}_M: \quad \cdots \longrightarrow P_i \longrightarrow P_{i-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow M \longrightarrow 0.$$

For each  $i > n$ , we have  $\text{Hom}_\Lambda(P_i, S_e) = \text{Ext}_\Lambda^i(M, S_e) = 0$ , and hence  $e$  annihilates the top of  $P_i$ . This shows that  $\mathcal{P}_M$  is  $e$ -bounded. Therefore,  $\text{tr}_e(\varphi)$  is defined for every endomorphism  $\varphi$  of  $M$ .  $\square$

**Remark.** If  $\Lambda$  is of finite global dimension, then we recover Lenzing’s trace function by taking  $e$  to be the identity of  $\Lambda$ .

Now, we are able to describe the Hochschild homology group  $\text{HH}_0(\Lambda_e)$  in case  $S_e$  is of finite injective dimension.

**1.6. Theorem.** *Let  $\Lambda$  be an artin algebra, and let  $e$  be an idempotent in  $\Lambda$ . If  $S_e$  is of finite injective dimension, then  $\text{HH}_0(\Lambda_e)$  is radical-trivial.*

**Proof.** Suppose that  $S_e$  is of finite injective dimension. By Lemma 1.5, the  $e$ -trace is defined for every endomorphism in  $\text{mod}\Lambda$ . Let  $x \in \Lambda$  be such that  $\bar{x} = x + \Lambda(1 - e)\Lambda$  lies in the radical of  $\Lambda_e$ , which is  $(J + \Lambda(1 - e)\Lambda)/\Lambda(1 - e)\Lambda$ . Hence,  $\bar{x} = \bar{a}$  for some  $a \in J$ . Let  $r > 0$  be such that  $a^r = 0$ , and consider the chain

$$0 = M_r \subseteq M_{r-1} \subseteq \cdots \subseteq M_1 \subseteq M_0 = \Lambda,$$

of submodules of  $\Lambda$ , where  $M_i = a^i \Lambda$ ,  $i = 0, \dots, r$ . Let  $\varphi_0 : \Lambda \rightarrow \Lambda$  be the left multiplication by  $a$ . Since  $\varphi_0(M_i) \subseteq M_{i+1}$ , we see that  $\varphi_0$  induces morphisms  $\varphi_i : M_i \rightarrow M_i$ ,  $i = 1, \dots, r$ , such that

$$\begin{array}{ccccccc}
 0 & \longrightarrow & M_{i+1} & \longrightarrow & M_i & \longrightarrow & M_i/M_{i+1} \longrightarrow 0 \\
 & & \downarrow \varphi_{i+1} & & \downarrow \varphi_i & & \downarrow 0 \\
 0 & \longrightarrow & M_{i+1} & \longrightarrow & M_i & \longrightarrow & M_i/M_{i+1} \longrightarrow 0
 \end{array}$$

commutes. By Proposition 1.4,  $\text{tr}_e(\varphi_i) = \text{tr}_e(\varphi_{i+1})$ , for  $i = 0, 1, \dots, r - 1$ . Applying Proposition 1.1(5), we get

$$\bar{a} + [\Lambda_e, \Lambda_e] = H_e(a + [\Lambda, \Lambda]) = H_e(\text{tr}(\varphi_0)) = \text{tr}_e(\varphi_0) = \text{tr}_e(\varphi_r) = 0,$$

that is,  $\bar{x} = \bar{a} \in [\Lambda_e, \Lambda_e]$ .  $\square$

Taking  $e$  to be the identity of  $\Lambda$ , we recover the following well known result; see, for example, [13].

**1.7. Corollary.** *If  $\Lambda$  is an artin algebra of finite global dimension, then  $\text{HH}_0(\Lambda)$  is radical-trivial.*

Let  $\Lambda$  be a finite dimensional algebra over a field of characteristic zero. If  $\Lambda$  is of finite global dimension, then all the Hochschild homology groups  $\text{HH}_i(\Lambda)$  with  $i \geq 1$  vanish; see [13]. However, in the situation as in Theorem 1.6, even if  $\Lambda$  is of finite global dimension,  $\Lambda_e$  may be of infinite global dimension with non-vanishing higher Hochschild homology groups.

**Example.** Let  $\Lambda = kQ/I$ , where  $k$  is a field,  $Q$  is the quiver

$$\begin{array}{ccc}
 1 & \xrightarrow{\alpha} & 2 \\
 \gamma \downarrow & \swarrow \varepsilon & \downarrow \beta \\
 4 & \xrightarrow{\delta} & 3
 \end{array}$$

and  $I$  is the ideal in the path algebra  $kQ$  generated by  $\alpha\beta - \gamma\delta, \beta\varepsilon, \delta\varepsilon, \varepsilon\alpha$ . It is easy to see that  $\Lambda$  is of finite global dimension. Let  $e$  be the sum of the primitive idempotents in  $\Lambda$  associated to the vertices 1, 2, 3. By Theorem 1.6,  $\text{HH}_0(\Lambda_e)$  is radical-trivial. On the other hand,  $\Lambda_e$  is a Nakayama algebra of infinite global dimension, and a direct computation shows that  $\text{HH}_2(\Lambda_e)$  does not vanish; see [11].

## 2. Main results

The main objective of this section is to apply the previously obtained result to solve the strong no loop conjecture for finite dimensional algebras over an algebraically closed field.

We start with an artin algebra  $\Lambda$  and a primitive idempotent  $e$  in  $\Lambda$ . We shall say that  $\Lambda$  is *locally commutative at  $e$*  if  $e\Lambda e$  is commutative and that  $\Lambda$  is *locally commutative* if it is locally commutative at every primitive idempotent. Moreover,  $e$  is called *basic* if  $e\Lambda$  is not isomorphic to any direct summand of  $(1 - e)\Lambda$ . In this terminology,  $\Lambda$  is basic if and only if its primitive idempotents are all basic.

**2.1. Proposition.** *Let  $\Lambda$  be an artin algebra, and let  $e$  be a basic primitive idempotent in  $\Lambda$  such that  $\Lambda/J^2$  is locally commutative at  $e + J^2$ . If  $S_e$  is of finite projective or injective dimension, then  $\text{Ext}^1_\Lambda(S_e, S_e) = 0$ .*

**Proof.** Firstly, assume that  $S_e$  is of finite injective dimension. For proving that  $\text{Ext}_A^1(S_e, S_e) = 0$ , it suffices to show that  $eJe/eJ^2e = 0$ . Let  $a \in eJe$ . Then  $a + \Lambda(1 - e)\Lambda \in [\Lambda_e, \Lambda_e]$  by Theorem 1.6. Since  $e$  is basic,  $e\Lambda(1 - e)\Lambda e \subseteq eJ^2e$ . This yields an algebra homomorphism

$$f : \Lambda_e \rightarrow e\Lambda e/eJ^2e : x + \Lambda(1 - e)\Lambda \mapsto exe + eJ^2e.$$

Thus,  $a + eJ^2e = f(a + \Lambda(1 - e)\Lambda)$  lies in the commutator group of  $e\Lambda e/eJ^2e$ . On the other hand,  $e\Lambda e/eJ^2e \cong (e + J^2)(\Lambda/J^2)(e + J^2)$ , which is commutative. Therefore,  $a + eJ^2e = 0$ , that is,  $a \in eJ^2e$ . The result follows in this case.

Next, assume that  $S_e$  is of finite projective dimension. Let  $D$  be the standard duality between  $\text{mod } \Lambda$  and  $\text{mod } \Lambda^{\text{op}}$ . Then  $D(S_e)$  is a simple  $\Lambda^{\text{op}}$ -module of finite injective dimension, which is supported by the primitive idempotent  $e^o$  corresponding to  $e$ . Since  $e^o$  is basic such that the quotient of  $\Lambda^{\text{op}}$  modulo the square of its radical is locally commutative at the class of  $e^o$ , we have  $\text{Ext}_{\Lambda^{\text{op}}}^1(D(S_e), D(S_e)) = 0$ , and consequently,  $\text{Ext}_A^1(S_e, S_e) = 0$ .  $\square$

**Remark.** In particular, Proposition 2.1 establishes the strong no loop conjecture for basic artin algebras  $\Lambda$  such that  $\Lambda/J^2$  is locally commutative.

Now we specialize the preceding result to finite dimensional algebras over a field.

**2.2. Theorem.** *Let  $\Lambda$  be a finite dimensional algebra over a field  $k$ , and let  $S$  be a simple  $\Lambda$ -module of  $k$ -dimension one. If  $S$  is of finite projective or injective dimension, then  $\text{Ext}_A^1(S, S) = 0$ .*

**Proof.** Let  $e$  be a primitive idempotent in  $\Lambda$  which does not annihilate  $S$ . Then  $\Lambda$  admits a complete set  $\{e_1, \dots, e_n\}$  of orthogonal primitive idempotents with  $e = e_1$ . We may assume that  $e_1\Lambda, \dots, e_r\Lambda$ , with  $1 \leq r \leq n$ , are the non-isomorphic indecomposable projective modules in  $\text{mod } \Lambda$ . Then

$$\Lambda/J \cong M_{n_1}(D_1) \times \dots \times M_{n_r}(D_r),$$

where  $D_i = \text{End}_\Lambda(e_i\Lambda/e_iJ)$  and  $n_i$  is the number of indices  $j$  with  $1 \leq j \leq n$  such that  $e_j\Lambda \cong e_i\Lambda$ , for  $i = 1, \dots, r$ . Observe that  $S$  is a simple  $M_{n_1}(D_1)$ -module, and hence  $S \cong D_1^{n_1}$ . Since  $S$  is one dimensional over  $k$ , it is one dimensional over  $D_1$ . In particular,  $n_1 = 1$ . That is,  $e$  is a basic primitive idempotent. Moreover,  $e\Lambda e/eJe \cong Se \cong k$ . Thus, for  $x_1, x_2 \in e\Lambda e$ , we can write  $x_i = \lambda_i e + a_i$ , where  $\lambda_i \in k$  and  $a_i \in eJe$ ,  $i = 1, 2$ . This yields  $x_1x_2 - x_2x_1 = a_1a_2 - a_2a_1 \in eJ^2e$ . Therefore,  $e\Lambda e/eJ^2e$  is commutative, and so is  $(e + J^2)(\Lambda/J^2)(e + J^2)$ . Now the result follows immediately from Proposition 2.1.  $\square$

**Remark.** A finite dimensional algebra over a field is called *elementary* if its simple modules are all one dimensional over the base field, or equivalently, it is isomorphic to an algebra given by a quiver with relations; see [1]. It is well known that a finite dimensional algebra over an algebraically closed field is Morita equivalent to an elementary algebra. In this connection, Theorem 2.3 establishes the strong no loop conjecture for finite dimensional elementary algebras over any field, and consequently, for finite dimensional algebras over an algebraically closed field.



We shall extend our results further. For this purpose, some more terminology is needed. From now on, let  $\Lambda$  stand for a finite dimensional elementary algebra over a field  $k$ , which is isomorphic to an algebra given by a quiver with relations. To simplify the notation, assume that  $\Lambda = kQ/I$ , where  $Q$  is a finite quiver,  $kQ$  is the path algebra of  $Q$  over  $k$ , and  $I$  is an admissible ideal in  $kQ$ . Recall that  $I$  is *admissible* if  $(kQ^+)^n \subseteq I \subseteq (kQ^+)^2$  for some  $n \geq 2$ , where  $kQ^+$  is the ideal in  $kQ$  generated by the arrows. Consider  $\rho = \lambda_1 p_1 + \dots + \lambda_r p_r \in I$ , where the  $p_i$  are distinct paths in  $Q$  from one fixed vertex to another, and the  $\lambda_i$  are non-zero scalars in  $k$ . We say that  $\rho$  is a *minimal relation* for  $\Lambda$  if  $\sum_{i \in \Omega} \lambda_i p_i \notin I$  for any  $\Omega \subset \{1, \dots, r\}$ . Observe that a minimal relation in this sense does not necessarily lie in a minimal set of generators of  $I$ . For instance, a path  $p$  in  $Q$  is a minimal relation for  $\Lambda$  if and only if  $p \in I$ . Moreover, a path  $p$  in  $Q$  is said to be *non-zero* in  $\Lambda$  if  $p \notin I$ ; and *free* in  $\Lambda$  if  $p$  is not a summand of any minimal relation for  $\Lambda$ .

Now, let  $\sigma = \alpha_1 \alpha_2 \dots \alpha_r$  be an oriented cycle in  $Q$ , where the  $\alpha_i$  are arrows. We denote by  $\text{supp}(\sigma)$  the set of vertices occurring as starting points of  $\alpha_1, \dots, \alpha_r$ , and define the *idempotent supporting*  $\sigma$  to be the sum of all primitive idempotents in  $\Lambda$  associated to the vertices in  $\text{supp}(\sigma)$ . Furthermore, the *cyclic permutations* of  $\sigma$  are the oriented cycles  $\sigma_1 = \sigma, \sigma_2 = \alpha_2 \dots \alpha_r \alpha_1, \dots$ , and  $\sigma_r = \alpha_r \alpha_1 \dots \alpha_{r-1}$ . Now, we say that  $\sigma$  is *cyclically non-zero* (respectively, *cyclically free*) in  $\Lambda$  if each of  $\sigma_1, \dots, \sigma_r$  is non-zero (respectively, free) in  $\Lambda$ . For example, a loop in  $Q$  is always cyclically free in  $\Lambda$ .

**2.3. Theorem.** *Let  $\Lambda = kQ/I$  with  $Q$  a finite quiver and  $I$  an admissible ideal in  $kQ$ , and let  $\sigma$  be an oriented cycle in  $Q$  with supporting idempotent  $e$  in  $\Lambda$ . If  $\sigma$  is cyclically free in  $\Lambda$ , then  $S_e$  is of infinite projective and injective dimensions.*

**Proof.** Suppose that  $\sigma$  is cyclically free in  $\Lambda$ . If  $\sigma$  is a power of a shorter oriented cycle  $\delta$ , then it is easy to see that  $\delta$  is also cyclically free in  $\Lambda$  and  $\text{supp}(\delta) = \text{supp}(\sigma)$ . Hence, we may assume that  $\sigma$  is not a power of any shorter oriented cycle. It is then well known that the cyclic permutations  $\sigma_1, \dots, \sigma_r$  of  $\sigma$ , where  $\sigma_1 = \sigma$ , are pairwise distinct.

For any  $x \in kQ$ , denote by  $\bar{x}$  its class in  $\Lambda$  and by  $\tilde{x}$  the class of  $\bar{x}$  in  $\Lambda_e$ . Let  $W$  be the vector subspace of  $\Lambda_e$  spanned by the classes  $\tilde{p}$ , where  $p$  ranges over the paths in  $Q$  different from  $\sigma_1, \dots, \sigma_r$ . Then, there exist paths  $p_1, \dots, p_m$  in  $Q$  different from  $\sigma_1, \dots, \sigma_r$  such that  $\{\tilde{p}_1, \dots, \tilde{p}_m\}$  is a  $k$ -basis of  $W$ . We claim that  $\{\tilde{\sigma}_1, \dots, \tilde{\sigma}_r, \tilde{p}_1, \dots, \tilde{p}_m\}$  is a  $k$ -basis of  $\Lambda_e$ . Indeed, it clearly spans  $\Lambda_e$ . Assume that

$$\sum_{i=1}^r \lambda_i \tilde{\sigma}_i + \sum_{j=1}^m \nu_j \tilde{p}_j = \tilde{0}, \quad \lambda_i, \nu_j \in k.$$

That is,  $\sum_{i=1}^r \lambda_i \tilde{\sigma}_i + \sum_{j=1}^m \nu_j \tilde{p}_j \in \Lambda(1 - e)\Lambda$ . Then

$$\sum_{i=1}^r \lambda_i \tilde{\sigma}_i + \sum_{j=1}^m \nu_j \tilde{p}_j = \sum_{l=1}^s \mu_l \tilde{q}_l, \quad \mu_l \in k,$$

where  $q_1, \dots, q_s$  are some distinct paths in  $Q$  passing through a vertex not in  $\text{supp}(\sigma)$ . Fix some  $t$  with  $1 \leq t \leq r$ . Letting  $\varepsilon_t$  be the trivial path in  $Q$  associated to the starting point  $a_t$  of  $\sigma_t$ , we get

$$\rho = \sum_{i=1}^r \lambda_i(\varepsilon_i \sigma_i \varepsilon_i) + \sum_{j=1}^m v_j(\varepsilon_j p_j \varepsilon_j) - \sum_{l=1}^s \mu_l(\varepsilon_l q_l \varepsilon_l) \in I.$$

Note that the non-zero elements of the  $\varepsilon_t \sigma_i \varepsilon_t, \varepsilon_t p_j \varepsilon_t, \varepsilon_t q_l \varepsilon_t \in kQ$  are distinct oriented cycles from  $a_t$  to  $a_t$ . If  $\lambda_t \neq 0$ , then  $\lambda_t(\varepsilon_t \sigma_i \varepsilon_t)$ , that is  $\lambda_t \sigma_i$ , would be a summand of a minimal non-zero summand  $\rho'$  of  $\rho$  with  $\rho' \in I$ . By definition,  $\rho'$  is a minimal relation for  $\Lambda$ , contrary to the hypothesis that  $\sigma$  is cyclically free in  $\Lambda$ . Therefore,  $\lambda_t = 0$ . This shows that  $\lambda_1, \dots, \lambda_r$  are all zero, and consequently, so are  $v_1, \dots, v_m$ . Our claim is established. Suppose now that  $\tilde{\sigma} \in [\Lambda_e, \Lambda_e]$ . Then


$$\tilde{\sigma} = \sum_{i=1}^n \eta_i(\tilde{u}_i \tilde{v}_i - \tilde{v}_i \tilde{u}_i), \tag{1}$$

where  $\eta_i \in k$  and  $u_i, v_i \in \{\sigma_1, \dots, \sigma_r, p_1, \dots, p_m\}$ . For each  $1 \leq i \leq n$ , we see easily that  $u_i v_i \notin \{\sigma_1, \dots, \sigma_r\}$  if and only if  $v_i u_i \notin \{\sigma_1, \dots, \sigma_r\}$ , and in this case,  $\tilde{u}_i \tilde{v}_i - \tilde{v}_i \tilde{u}_i \in W$ . Therefore, Eq. (1) becomes

$$\tilde{\sigma} = \sum \eta_{ij}(\tilde{\sigma}_i - \tilde{\sigma}_j) + w, \tag{2}$$

where  $\eta_{ij} \in k$  and  $w \in W$ . Let  $L$  be the linear form on  $\Lambda_e$ , which sends each of  $\tilde{\sigma}_1, \dots, \tilde{\sigma}_r$  to 1 and vanishes on  $W$ . Since  $\sigma = \sigma_1$ , applying  $L$  to Eq. (2) yields  $1 = 0$ , a contradiction. Therefore, the class of  $\tilde{\sigma}$  in  $\text{HH}_0(\Lambda_e)$  is non-zero. Since  $\tilde{\sigma}$  lies in the radical of  $\Lambda_e$ , by Theorem 1.6,  $S_e$  is of infinite projective and injective dimensions.  $\square$

**Example.** Let  $\Lambda = kQ/I$ , where  $Q$  is the following quiver



and  $I$  is the ideal in  $kQ$  generated by  $\alpha\beta, \delta\gamma, \beta\varepsilon, \varepsilon\beta, \nu\delta, \nu\mu, \mu\nu, \gamma\mu, \alpha\gamma\delta\beta\alpha\gamma - \varepsilon\gamma$ . It is easy to see that the oriented cycle  $\beta\alpha\gamma\delta$  is cyclically free in  $\Lambda$ . By Theorem 2.3, one of the simple modules  $S_1, S_2, S_3$  is of infinite projective dimension, and one is of infinite injective dimension.

**2.4. Corollary.** *Let  $\Lambda = kQ/I$ , where  $Q$  is a finite quiver and  $I$  is an admissible ideal in  $kQ$ . If  $Q$  contains an oriented cycle which is cyclically free in  $\Lambda$ , then  $\Lambda$  is of infinite global dimension.*

An admissible ideal  $I$  in  $kQ$  is called *monomial* if it is generated by some paths. In this case, every minimal relation for  $\Lambda$  is a multiple of a path in  $Q$ . Therefore, an oriented cycle in  $Q$  is cyclically free in  $\Lambda$  if and only if it is cyclically non-zero in  $\Lambda$ . This yields the following consequence, which can also be derived from some results stated in [11].

**2.5. Corollary.** *Let  $\Lambda = kQ/I$ , where  $Q$  is a finite quiver and  $I$  is a monomial ideal in  $kQ$ . If  $Q$  contains an oriented cycle which is cyclically non-zero in  $\Lambda$ , then  $\Lambda$  is of infinite global dimension.*

**Example.** Consider  $\Lambda = kQ/I$ , where  $Q$  is the quiver

$$\begin{array}{ccc} & \alpha & \\ & \curvearrowright & \\ 1 & & 2 \\ & \curvearrowleft & \\ & \beta & \end{array}$$

and  $I$  is the ideal in  $kQ$  generated by  $\alpha\beta$ . It is easy to see that  $\Lambda$  is of global dimension two. Observe that  $Q$  contains an oriented cycle  $\beta\alpha$  which is non-zero but not cyclically non-zero in  $\Lambda$ .

To conclude, we would like to draw the reader's attention to the following even stronger version of the strong no loop conjecture.

**2.6. Extension Conjecture.** *Let  $S$  be a simple module over an artin algebra. If  $\text{Ext}^1(S, S)$  is non-zero, then  $\text{Ext}^i(S, S)$  is non-zero for infinitely many integers  $i$ .*

This conjecture was originally posed in [14] under the name of *extreme no loop conjecture*. It remains open except for monomial algebras and special biserial algebras; see [6,14].

## Acknowledgments

This research was supported in part by the Natural Sciences and Engineering Research Council of Canada and was carried out during the first author's visit to the Université de Sherbrooke.

## References

- [1] M. Auslander, I. Reiten, S.O. Smalø, Representation Theory of Artin Algebras, Cambridge Stud. Adv. Math., vol. 36, Cambridge University Press, Cambridge, 1995.
- [2] W.D. Burgess, K.R. Fuller, E.L. Green, D. Zacharia, Left monomial rings – a generalization of monomial algebras, Osaka J. Math. 30 (3) (1993) 543–558.
- [3] W.D. Burgess, M. Saorín, Homological aspects of semigroup gradings on rings and algebras, Canad. J. Math. 51 (3) (1999) 488–505.
- [4] L. Diracca, S. Koenig, Cohomological reduction by split pairs, J. Pure Appl. Algebra 212 (3) (2008) 471–485.
- [5] E.L. Green, W.H. Gustafson, D. Zacharia, Artin rings of global dimension two, J. Algebra 92 (1985) 375–379.
- [6] E.L. Green, Ø. Solberg, D. Zacharia, Minimal projective resolutions, Trans. Amer. Math. Soc. 353 (7) (2001) 2915–2939.
- [7] D. Happel, On the derived category of a finite dimensional algebra, Comment. Math. Helv. 62 (1987) 339–389.
- [8] D. Happel, Auslander–Reiten triangles in derived categories of finite-dimensional algebras, Proc. Amer. Math. Soc. 112 (1991) 641–648.
- [9] A. Hattori, Rank element of a projective module, Nagoya Math. J. 25 (1965) 113–120.
- [10] K. Igusa, Notes on the no loops conjecture, J. Pure Appl. Algebra 69 (2) (1990) 161–176.
- [11] K. Igusa, D. Zacharia, On the cyclic homology of monomial relation algebras, J. Algebra 151 (2) (1992) 502–521.
- [12] B.T. Jensen, Strong no-loop conjecture for algebras with two simples and radical cube zero, Colloq. Math. 102 (1) (2005) 1–7.
- [13] H. Lenzing, Nilpotente Elemente in Ringen von endlicher globaler Dimension, Math. Z. 108 (1969) 313–324.
- [14] S. Liu, J.P. Morin, The strong no loop conjecture for special biserial algebras, Proc. Amer. Math. Soc. 132 (12) (2004) 3513–3523.
- [15] N. Marmaridis, A. Papistas, Extensions of abelian categories and the strong no-loops conjecture, J. Algebra 178 (1) (1995) 1–20.
- [16] C. Paquette, Strictly stratified algebras revisited, Comm. Algebra 37 (8) (2009) 2591–2600.
- [17] D. Skorodumov, The strong no loop conjecture is true for mild algebras, arXiv:1011.1143v2.
- [18] J. Stallings, Centerless groups – an algebraic formulation of Gottlieb's theorem, Topology 4 (1965) 129–134.

- [19] C. Weibel, *An Introduction to Homological Algebra*, Cambridge Stud. Adv. Math., vol. 38, Cambridge University Press, Cambridge, 1994.
- [20] D. Zacharia, Special monomial algebras of finite global dimension, in: *NATO Adv. Sci. Inst.*, vol. 233, Kluwer, Dordrecht, 1988, pp. 375–378.