




Contents lists available at SciVerse ScienceDirect

Annals of Pure and Applied Logic

www.elsevier.com/locate/apal

Abstract elementary classes and accessible categories

T. Beke^{a,1}, J. Rosický^{b,*,2}^a Department of Mathematics, University of Massachusetts Lowell, One University Avenue, Lowell, MA 01854, USA^b Department of Mathematics and Statistics, Masaryk University, Faculty of Sciences, Kotlářská 2, 611 37 Brno, Czech Republic

ARTICLE INFO

Article history:

Received 18 November 2011

Received in revised form 15 May 2012

Accepted 29 June 2012

Available online 9 August 2012

Communicated by I. Moerdijk

MSC:

03C48

18C35

Keywords:

Abstract elementary class

Accessible category

ABSTRACT

We investigate properties of accessible categories with directed colimits and their relationship with categories arising from Shelah's Abstract Elementary Classes. We also investigate ranks of objects in accessible categories, and the effect of accessible functors on ranks.

© 2012 Elsevier B.V. All rights reserved.

1. Introduction

M. Makkai and R. Paré [9] introduced accessible categories as categories sharing two typical properties of categories of structures described by infinitary first-order theories – the existence of sufficiently many directed colimits and the existence of a set of objects generating all objects by means of distinguished colimits. Their (purely category-theoretic) definition has since then found applications in various branches of mathematics. Very often, accessible categories have all directed colimits. These arise as categories of models and elementary embeddings of infinitary theories with finitary quantifiers. In model theory, S. Shelah went in a similar direction and introduced abstract elementary classes as a formalization of properties of models of generalized logics with finitary quantifiers. Our aim is to relate these two approaches. In Section 5, we introduce a hierarchy of accessible categories with directed colimits. The main result of that section, Corollary 5.7, sandwiches Shelah's Abstract Elementary Classes between two natural families of accessible categories.

Unlike abstract elementary classes, accessible categories are not equipped with canonical 'underlying sets'. Nonetheless, there exists a good substitute for 'size of the model', namely, the *presentability rank* of an object, that can be expressed purely in the language of category theory, i.e. in terms of objects and morphisms. In this sense, accessible categories take an extreme 'signature-free' and 'elements-free' view of abstract elementary classes. From this point of view, Shelah's Categoricity Conjecture, the driving force of abstract elementary classes (see [3]), turns on the subtle interaction between ranks of objects and directed colimits in accessible categories.

* Corresponding author.

E-mail addresses: tibor_beke@uml.edu (T. Beke), rosicky@math.muni.cz (J. Rosický).¹ Supported by the NSA under Grant H98230-11-1-0172.² Supported by MSM 0021622409.

The connection between abstract elementary classes and accessible categories was discovered, independently, by M.J. Lieberman as well; see [7] and [8]. Our Corollary 5.7 simplifies Lieberman’s description; cf. Proposition 4.8 and Claim 4.9 in [8].

2. Accessible categories

In order to define accessible categories one just needs the concept of a λ -directed colimit where λ is a regular cardinal number. This is a colimit over a diagram $D : \mathcal{D} \rightarrow \mathcal{K}$ where \mathcal{D} is a λ -directed poset, considered as a category. An object K of a category \mathcal{K} is called λ -presentable if its hom-functor $\text{hom}(K, -) : \mathcal{K} \rightarrow \mathbf{Set}$ preserves λ -directed colimits; here \mathbf{Set} is the category of sets.

A category \mathcal{K} is called λ -accessible, where λ is a regular cardinal, provided that

- (1) \mathcal{K} has λ -directed colimits,
- (2) \mathcal{K} has a set \mathcal{A} of λ -presentable objects such that every object of \mathcal{K} is a λ -directed colimit of objects from \mathcal{A} .

A category is *accessible* if it is λ -accessible for some regular cardinal λ .

A *signature* Σ is a set of (infinitary) operation and relation symbols. These symbols are S -sorted where S is a set of sorts. It is advantageous to work with many-sorted signatures but it is easy to reduce them to single-sorted ones. One just replaces sorts by unary relation symbols and adds axioms saying that there are disjoint and cover the underlying set of a model. Thus the underlying set of an S -sorted structure A is the disjoint union of underlying sets A_s over all sorts $s \in S$. $|A|$ will denote the cardinality of the underlying set of the Σ -structure A . The category of all Σ -structures and homomorphisms (i.e., mappings preserving all operations and relations) is denoted by $\mathbf{Str}(\Sigma)$. A homomorphism is called a *substructure embedding* if it is injective and reflects all relations. Any inclusion of a substructure is a substructure embedding. Conversely, if $h : A \rightarrow B$ is a substructure embedding then A is isomorphic to the substructure $h(A)$ of B . The category of all Σ -structures and substructure embeddings is denoted by $\mathbf{Emb}(\Sigma)$. Both $\mathbf{Str}(\Sigma)$ and $\mathbf{Emb}(\Sigma)$ are accessible categories, cf. [2, 5.30 and 1.70].

A signature Σ is *finitary* if all relation and function symbols are finitary. For a finitary signature, the category $\mathbf{Str}(\Sigma)$ is locally finitely presentable and $\mathbf{Emb}(\Sigma)$ is finitely accessible. In both cases, there is a cardinal κ such that, for each regular cardinal $\kappa \leq \mu$, a Σ -structure K is μ -presentable if and only if $|K| < \mu$. This follows from the downward Löwenheim-Skolem theorem; see [9, 3.3.1].

Let λ be a cardinal and Σ be a λ -ary signature, i.e., all relation and function symbols have arity smaller than λ . Given a cardinal κ , the language $L_{\kappa\lambda}(\Sigma)$ allows less than κ -ary conjunctions and disjunctions and less than λ -ary quantifications. A substructure embedding of Σ -structures is called $L_{\kappa\lambda}$ -*elementary* if it preserves all $L_{\kappa\lambda}$ -formulas. A *theory* T is a set of sentences of $L_{\kappa\lambda}(\Sigma)$. $\mathbf{Mod}(T)$ denotes the category of T -models and homomorphisms, $\mathbf{Emb}(T)$ the category of T -models and substructure embeddings while $\mathbf{Elem}(T)$ will denote the category of T -models and $L_{\kappa\lambda}$ -elementary embeddings. The category $\mathbf{Elem}(T)$ is accessible (see [2, 5.42]). For certain theories T , the category $\mathbf{Mod}(T)$ does not have μ -directed colimits for any regular cardinal μ and thus fails to be accessible.

A theory T is called *basic* if it consists of sentences

$$(\forall x)(\varphi(x) \Rightarrow \psi(x))$$

where φ and ψ are positive-existential formulas and x is a string of variables. For a basic theory T , the category $\mathbf{Mod}(T)$ is accessible. Conversely, every accessible category is equivalent to the category of models and homomorphisms of a basic theory. All these facts can be found in [9] or [2].

Locally presentable categories are defined as cocomplete accessible categories. Following [2, 1.20], each locally λ -presentable category is locally μ -presentable for each regular cardinal $\mu > \lambda$. Let λ be an uncountable regular cardinal. The category \mathbf{Pos}_λ of λ -directed posets and substructure embeddings is λ -accessible but it is not μ -accessible for all regular cardinals $\mu > \lambda$. Following [9, 2.3], let us write $\lambda \triangleleft \mu$ whenever \mathbf{Pos}_λ is μ -accessible. There are arbitrarily large regular cardinals μ such that $\lambda \triangleleft \mu$ and, at the same time, arbitrarily large regular cardinals μ such that $\lambda \triangleleft \mu$ does not hold. Thus the *accessibility spectrum* of \mathbf{Pos}_λ has a proper class of gaps. Generally, if $\lambda \triangleleft \mu$ then any λ -accessible category \mathcal{K} is μ -accessible; see [9, 2.3.10] or [2, 2.11]. By [2, 2.13(1)], one has $\omega \triangleleft \lambda$ for every uncountable regular cardinal λ . Thus a finitely accessible category is μ -accessible for all uncountable regular cardinals μ .

Definition 2.1. We say that a category \mathcal{K} is *well λ -accessible* if it is μ -accessible for each regular cardinal $\lambda \leq \mu$.

\mathcal{K} is *well accessible* if it is well λ -accessible for some regular cardinal λ .

We have just seen that any locally presentable category and any finitely accessible category is well accessible.

Definition 2.2. Let λ be a regular cardinal. We say that an object K of a category \mathcal{K} has *presentability rank* (or, for brevity, *rank*) λ if it is λ -presentable but not μ -presentable for any regular cardinal $\mu < \lambda$. We will write $\text{rank}(K) = \lambda$.

This concept was introduced by Makkai and Paré under the name of *presentability* of an object. See the very last line of p. 29 of [9].

Remark 2.3. (1) We have mentioned that in $\mathbf{Emb}(\Sigma)$, Σ finitary, there is a cardinal κ such that, for each cardinal $\kappa \leq \mu$, a Σ -structure A is μ -presentable if and only if $|A| < \mu$. Therefore for all large enough structures, $\text{rank}(A) = |A|^+$, i.e., presentability ranks play the role of cardinalities.

(2) Let T be a basic theory in a language $L_{\kappa\omega}(\Sigma)$. Since $\mathbf{Emb}(T)$ is closed under directed colimits in $\mathbf{Emb}(\Sigma)$, $\text{rank}(X) \leq |X|^+$ for each T -model X . By the downward Löwenheim–Skolem theorem, cf. [9, 3.3.1], there is a cardinal μ such that, for $\mu \leq |X|$, we have $\text{rank}(X) = |X|^+$. Moreover, either the presentability ranks of T -models in $\mathbf{Emb}(T)$ form a set or T has models of presentability rank ν^+ for all $\mu \leq \nu$. This means that the *presentability spectrum* of $\mathbf{Emb}(T)$ does not have arbitrarily large gaps.

(3) If T is a basic theory in a language $L_{\kappa\omega}(\Sigma)$ then $\mathbf{Mod}(T)$ does not need to be closed under directed colimits in $\mathbf{Str}(\Sigma)$. The simplest example would involve countably many unary relation symbols and the axiom that their conjunction is false. Clearly, this property is not preserved by directed colimits. The situation changes for basic theories T in a language $L_{\kappa\omega}^*(\Sigma)$ allowing only finitary conjunctions. In this case, $\mathbf{Mod}(T)$ is closed under directed colimits in $\mathbf{Str}(\Sigma)$. [9] calls categories equivalent to such $\mathbf{Mod}(T)$ ∞, ω -elementary. Every finitely accessible category is ∞, ω -elementary; see [9, 4.3.2]. The downward Löwenheim–Skolem theorem applies to ∞, ω -elementary $\mathbf{Mod}(T)$, and implies, similarly to (2), that $\text{rank}(X)$ eventually coincides with $|X|^+$ and the presentability spectrum of $\mathbf{Mod}(T)$ does not have arbitrarily large gaps.

Even if T is only a basic theory in a language $L_{\kappa\omega}(\Sigma)$, then $\mathbf{Mod}(T)$ is closed under λ -directed colimits in $\mathbf{Str}(\Sigma)$ for some regular cardinal λ . Then the downward Löwenheim–Skolem theorem still implies that $\text{rank}(X)$ eventually coincides with $|X|^+$ and the presentability spectrum of $\mathbf{Mod}(T)$ does not have arbitrarily large gaps.

(4) Let T be an arbitrary theory in a language $L_{\kappa\omega}(\Sigma)$. We know that $\mathbf{Elem}(T)$ is an accessible category and, following [2, 5.39], $\mathbf{Elem}(T)$ is closed under directed colimits in $\mathbf{Str}(\Sigma)$. Thus $\mathbf{Elem}(T)$ is an accessible category with directed colimits. Moreover, $\text{rank}(X) = |X|^+$ in $\mathbf{Elem}(T)$ for all sufficiently large T -models X and the presentability spectrum of $\mathbf{Elem}(T)$ does not have arbitrarily large gaps.

Following [9, 3.2.8], there is a basic theory T' in another language $L_{\kappa\omega}^*(\Sigma')$ such that the categories $\mathbf{Elem}(T)$ and $\mathbf{Mod}(T')$ are equivalent. Thus $\mathbf{Elem}(T)$ is ∞, ω -elementary.

(5) Assuming GCH, in any accessible category \mathcal{K} , $\text{rank}(K)$ is a successor cardinal for all objects K (with the possible exception of a set of isomorphism types). Indeed, let \mathcal{K} be a κ -accessible category and $\text{rank}(K) = \lambda$ with $\kappa < \lambda$. λ is a regular cardinal by definition; if λ was a limit cardinal then (as it is uncountable) it would be a weakly inaccessible cardinal. Since GCH is assumed, λ is inaccessible. Thus, given $\alpha < \kappa < \lambda$ and $\beta < \lambda$, we have $\beta^\alpha < \lambda$. Following [2, 2.13(4)], $\kappa \triangleleft \lambda$. Following [9, 2.3.11], the object K can be exhibited as a κ -directed colimit of κ -presentable objects along a diagram of size less than λ . Let the size of that diagram be ν ; then K is ν^+ -presentable. That would mean $\nu^+ = \lambda$, contradicting that λ is a limit cardinal.

By ‘category’ we always mean a locally small one, i.e., having a set of morphisms between any two objects. A category is called *small* if it has a set of objects. We say that a category is *large* if it is not equivalent to a small category. This means that it has a proper class of non-isomorphic objects. In Remark 2.3 we saw several families of large accessible categories that, starting from some cardinal, possess objects of every possible presentation rank. Since in those examples this property followed from the downward Löwenheim–Skolem theorem, we will call such categories LS-accessible.

Definition 2.4. An accessible category \mathcal{K} will be called λ -LS-accessible if \mathcal{K} has an object of presentability rank μ^+ for each cardinal $\mu \geq \lambda$.

\mathcal{K} is LS-accessible if it is λ -LS-accessible for some cardinal λ .

We will deal with this concept later. We should note that we have not been able to find any large accessible category which is not LS-accessible.

3. Accessible functors

A functor $F : \mathcal{K} \rightarrow \mathcal{L}$ is called λ -accessible if \mathcal{K} and \mathcal{L} are λ -accessible categories and F preserves λ -directed colimits. F is called accessible if it is λ -accessible for some regular cardinal λ . By the Uniformization theorem (see [9, 2.4.9] or [2, 2.19]), for each accessible functor F there is a regular cardinal λ such that F is λ -accessible and preserves λ -presentable objects; it means that if K is λ -presentable then $F(K)$ is λ -presentable. The same is then true for each $\mu \triangleright \lambda$.

Definition 3.1. A functor $F : \mathcal{K} \rightarrow \mathcal{L}$ will be called well λ -accessible if \mathcal{K} and \mathcal{L} are well λ -accessible categories and F preserves μ -directed colimits and μ -presentable objects for each $\lambda \leq \mu$. F is called well accessible if it is well λ -accessible for some regular cardinal λ .

Remark 3.2. (1) Of course, whenever F preserves λ -directed colimits then it preserves μ -directed ones for $\mu \geq \lambda$.

(2) Every colimit preserving functor between locally λ -presentable categories is well λ -accessible. This immediately follows from the fact that, for regular cardinals $\lambda \leq \mu$, an object of a locally λ -presentable category is μ -presentable if and only if it is a μ -small colimit of λ -presentable objects. Recall that a category is μ -small if it has less than μ morphisms.

(3) Every finitely accessible functor is well finitely accessible. This follows from [2, 2.13(1)] and the fact that, for regular cardinals $\lambda \triangleleft \mu$, an object of a λ -presentable category is μ -presentable if and only if it is a λ -directed μ -small colimit of λ -presentable objects (see [9, 2.3.11]).

(4) Let I be an infinite set and consider the functor $\mathbf{Set} \rightarrow \mathbf{Set}$ that sends X to X^I . This functor is accessible but not well accessible since there are arbitrarily large $|X|$ such that $|X^I| > |X|$, hence the functor does not preserve μ -presentable objects for all arbitrarily large enough μ . On the other hand, for I finite, the functor is well accessible since it is finitely accessible.

A well accessible functor, by definition, will take μ -presentable objects to μ -presentable ones for all large enough μ ; however, it can lower presentability ranks.

Example 3.3. Let $F : \mathbf{Gr} \rightarrow \mathbf{Ab}$ be the abelianization functor, i.e., the reflector from groups to abelian groups. Since F is a left adjoint, it is well finitely accessible by Remark 3.2(2). There exist simple groups G of arbitrarily large infinite cardinalities. Since $F(G) = 0$ for such a G , and since 0 is finitely presentable in \mathbf{Ab} and $\text{rank}(G) = |G|^+$, the functor F can lower presentability ranks from κ^+ to ω for arbitrarily large κ .

We say that a functor $F : \mathcal{K} \rightarrow \mathcal{L}$ reflects λ -presentable objects if $F(K)$ λ -presentable implies that K is λ -presentable.

Recall that a morphism $g : B \rightarrow A$ is a split epimorphism if there exists $f : A \rightarrow B$ with $gf = \text{id}_A$. Since, in this case, g is a coequalizer of fg and id_B , A is λ -presentable whenever B is λ -presentable (see [2, 1.16]).

Definition 3.4. We say that a functor $F : \mathcal{K} \rightarrow \mathcal{L}$ reflects split epimorphisms if f is a split epimorphism whenever $F(f)$ is a split epimorphism.

Remark 3.5. Any functor $F : \mathcal{K} \rightarrow \mathcal{L}$ reflecting split epimorphisms is conservative, i.e., it reflects isomorphisms. If all morphisms of \mathcal{K} are monomorphisms then F reflects split epimorphisms if and only if it is conservative.

Lemma 3.6. Let $F : \mathcal{K} \rightarrow \mathcal{L}$ be a λ -accessible functor which reflects split epimorphisms. Then F reflects λ -presentable objects.

Proof. Let $F(K)$ be λ -presentable in \mathcal{L} . Since \mathcal{K} is λ -accessible, K is a λ -directed colimit $(k_i : K_i \rightarrow K)_{i \in I}$ of λ -presentable objects. Since F preserves λ -directed colimits and $F(K)$ is λ -presentable, there is $i \in I$ and $f : F(K) \rightarrow F(K_i)$ such that $F(k_i)f = \text{id}_{F(K)}$. Since F reflects split epimorphisms, this $k_i : K_i \rightarrow K$ is a split epimorphism. Thus K is λ -presentable. \square

Proposition 3.7. Let $F : \mathcal{K} \rightarrow \mathcal{L}$ be a well λ -accessible functor which reflects split epimorphisms. Then F preserves presentability ranks μ for $\lambda < \mu$.

Proof. Let $\text{rank}(K) = \mu$, $\lambda < \mu$. Then $F(K)$ is μ -presentable. Assume that $F(K)$ is ν -presentable for some $\nu < \mu$. Without loss of generality, we can assume that $\lambda \leq \nu$. Following Lemma 3.6, K is ν -presentable, which is a contradiction. Thus $\text{rank}(F(K)) = \mu$. \square

4. Accessible categories with directed colimits

An important class of accessible categories consists of accessible categories having directed colimits. It includes both ∞, ω -elementary categories and locally presentable ones. On the other hand, the basic $L_{\omega_1 \omega_1}$ theory T of well-ordered sets has both $\mathbf{Mod}(T)$ and $\mathbf{Emb}(T)$ ω_1 -accessible without having directed colimits.

Proposition 4.1. Any accessible category with directed colimits is well accessible.

Proof. Let \mathcal{K} be a λ -accessible category with directed colimits and consider a regular cardinal $\mu > \lambda$. Given an object K of \mathcal{K} , there is a λ -directed colimit $(a_i : A_i \rightarrow K)_{i \in I}$ of λ -presentable objects A_i . Let \hat{I} be the poset of all directed subsets of I of cardinalities less than μ (ordered by inclusion). Since every subset of I having less than μ elements is contained in a directed subset of I having less than μ elements (cf. [2, 2.11]), clearly, \hat{I} is μ -directed. For each $M \in \hat{I}$, let B_M be a colimit of the subdiagram indexed by M . Then B_M is μ -presentable. K is a μ -directed colimit of the B_M , $M \in \hat{I}$. Thus \mathcal{K} is μ -accessible. \square

Lemma 4.2. Let \mathcal{K} be a λ -accessible category with directed colimits and K an object of \mathcal{K} which is not λ -presentable. Then $\text{rank}(K)$ is a successor cardinal.

Proof. Let $\lambda < \mu = \text{rank}(K)$. Following the proof of Proposition 4.1, K is a μ -directed colimit of objects K_i which are directed colimits of size $v_i < \mu$ of λ -presentable objects. Since K is μ -presentable, it is a retract of some K_i . If $v_i < \lambda$ then K_i is λ -presentable and thus K is λ -presentable as well. Let $\lambda \leq v_i$. Then K_i is v_i^+ -presentable and thus K is v_i^+ -presentable too. Hence $\mu = v_i^+$. \square

Proposition 4.3. Let \mathcal{K} and \mathcal{L} be accessible categories with directed colimits and $F : \mathcal{K} \rightarrow \mathcal{L}$ a functor preserving directed colimits. Then F is well accessible.

Proof. There is a regular cardinal λ such that both \mathcal{K} and \mathcal{L} are λ -accessible and F preserves λ -presentable objects (see [2, 2.19]). Consider a regular cardinal $\mu > \lambda$ and let K be a μ -presentable object of \mathcal{K} . Following the proof of Proposition 4.1, K is a μ -directed colimit of objects B_M where each B_M is a μ -small directed colimit of λ -presentable objects. Since K is μ -presentable, it is a retract of some B_M . Since $F(B_M)$ is a μ -small directed colimit of λ -presentable objects, $F(B_M)$ is μ -presentable in \mathcal{L} . Thus $F(K)$ is μ -presentable in \mathcal{L} as a retract of $F(B_M)$. We have proved that F is well λ -accessible. \square

Remark 4.4. Let **Lin** be the category of linearly ordered sets and order preserving injective mappings (they coincide with substructure embeddings). It is a finitely accessible category which is a “minimal” ∞, ω -elementary category in the sense that for every large ∞, ω -elementary category \mathcal{K} there is a faithful functor $E : \mathbf{Lin} \rightarrow \mathcal{K}$ preserving directed colimits; see [9, 3.4.1]. Its construction is based on Ehrenfeucht–Mostowski models. Following Propositions 4.1 and 4.3, E is well accessible. Since it is faithful, it reflects epimorphisms. Epimorphisms in **Lin** are isomorphisms and thus E reflects split epimorphisms. Following Proposition 3.7, E preserves presentability ranks μ starting from some cardinal λ .

Recall that a full subcategory \mathcal{K} of a category \mathcal{L} is called *accessibly embedded* if there is a regular cardinal λ such that \mathcal{K} is closed under λ -directed colimits in \mathcal{L} .

Theorem 4.5. Accessible categories with directed colimits are precisely the reflective and accessibly embedded subcategories of finitely accessible categories.

Proof. Every reflective and accessibly embedded subcategory of a finitely accessible category \mathcal{L} is accessible (see [2, 2.53]) and has directed colimits calculated as reflections of directed colimits in \mathcal{L} . Conversely, let \mathcal{K} be a λ -accessible category with directed colimits. Then \mathcal{K} is equivalent to the full subcategory $\text{Ind}_\lambda(\mathcal{C})$ of $\mathbf{Set}^{\mathcal{C}^{\text{op}}}$ consisting of all λ -directed colimits of hom-functors (see [2, 2.26]); \mathcal{C} is a small category. Let $\text{Ind}(\mathcal{C})$ be the full subcategory of $\mathbf{Set}^{\mathcal{C}^{\text{op}}}$ consisting of all directed colimits of hom-functors. Then $\text{Ind}_\lambda(\mathcal{C})$ is closed in $\text{Ind}(\mathcal{C})$ under λ -directed colimits. Given an object X in $\text{Ind}(\mathcal{C})$, we express it as a directed colimit of hom-functors and take their colimit $F(X)$ in $\text{Ind}_\lambda(\mathcal{C})$ (recall that \mathcal{K} has directed colimits). Clearly, $F(X)$ is a reflection of X in $\text{Ind}_\lambda(\mathcal{C})$. Thus $\text{Ind}_\lambda(\mathcal{C})$ is a reflective subcategory of $\text{Ind}(\mathcal{C})$. \square

Theorem 4.6. Each large locally presentable category is LS-accessible.

Proof. Let \mathcal{K} be a locally presentable category. Then \mathcal{K} is locally λ -presentable for some regular cardinal λ and, following [2, 5.30], \mathcal{K} is equivalent to $\mathbf{Mod}(T)$ for a limit theory of $L_{\lambda\lambda}(\Sigma)$ where Σ is an S -sorted signature. Thus it suffices to prove that $\mathbf{Mod}(T)$ is LS-accessible.

Let \mathbf{Set}^S denote the category of S -sorted sets. The category \mathbf{Set}^S is locally finitely presentable and, given an S -sorted set $X = (X_s)_{s \in S}$, $\text{rank}(X) = |X|^+$ where $|X|$ is defined as the cardinality of the disjoint union of X_s , $s \in S$. Let $U : \mathbf{Mod}(T) \rightarrow \mathbf{Set}^S$ denote the forgetful functor. Since $\mathbf{Mod}(T)$ is assumed large, there is $t \in S$ such that the sets $U(K)_t$ are arbitrarily large, as K ranges over objects of $\mathbf{Mod}(T)$. Let $V : \mathbf{Mod}(T) \rightarrow \mathbf{Set}$ be the composition of U with the functor $\mathbf{Set}^S \rightarrow \mathbf{Set}$ sending (X_s) to X_t . The functor V preserves limits and λ -directed colimits (see the proof of [2, 5.9]). By [2, 1.66], V has a left adjoint F which preserves μ -presentable objects for each $\mu \geq \lambda$, cf. [2, Example 1.s(1)]. Thus $\text{rank}(FX) \leq \text{rank}(X)$ for each set X with $\lambda \leq \text{rank}(X)$.

Assume that $\mathbf{Mod}(T)$ is not LS-accessible. Then there is a set X with $\lambda \leq \text{rank}(X)$ and $\mu = \text{rank}(FX) < \text{rank}(X)$. Since \mathbf{Set} is locally μ -presentable, X is a μ -directed colimit of μ -presentable objects X_i , $i \in I$. Let

$$(u_i : X_i \rightarrow X)_{i \in I}$$

denote a colimit cocone. Since F preserves colimits,

$$(Fu_i : FX_i \rightarrow FX)_{i \in I}$$

is a colimit cocone. Since FX is μ -presentable, there is $j \in I$ and $r : FX \rightarrow FX_j$ such that $F(u_j)r = \text{id}_{FX}$. Hence Fu_j is a split epimorphism. There is a mapping $f : X \rightarrow X$ such that $fu_j = u_j$ and $f \neq \text{id}_X$. Since $F(f)F(u_j) = F(u_j)$, we have $F(f) = \text{id}_{FX}$.

Let $\eta : \text{Id} \rightarrow VF$ be the adjunction unit. Since $\eta_X f = VF(f)\eta_X = \eta_X$, η_X is not a monomorphism. Since the sets VK are arbitrarily large, there is K in \mathcal{K} such that $|X| < |VK|$. Thus there is a monomorphism $X \rightarrow VK$, which implies that η_X is a monomorphism. We get a contradiction. \square

Remark 4.7. (1) We have proved the stronger result that the functor $F : \mathbf{Set} \rightarrow \mathcal{K}$ preserves presentability ranks $\geq \lambda$.

(2) The argument even works for weakly locally presentable categories, i.e., for accessible categories with products. Such categories are equivalent to $\mathbf{Mod}(T)$ where T is a regular theory of $L_{\lambda\lambda}(\Sigma)$ (see [2, Example 5.e]). The forgetful functor $U : \mathcal{K} \rightarrow \mathbf{Set}^S$ has a weak left adjoint $F : \mathbf{Set}^S \rightarrow \mathcal{K}$ equipped with a natural transformation $\eta : \text{Id} \rightarrow UF$. This follows from the fact that \mathcal{K} is a small-injectivity class in a locally presentable category \mathcal{L} (see [2, 4.8]) and thus it is naturally weakly reflective in \mathcal{L} (cf. [1]). This suffices for the argument.

(3) Let \mathcal{K} be a large accessible category with coproducts. Assume that \mathcal{K} is λ -accessible and take the coproduct K of a (representative) set of all λ -presentable objects. Then the functor

$$U = \mathcal{K}(K, -) : \mathcal{K} \rightarrow \mathbf{Set}$$

has arbitrarily large values. U has a left adjoint F given by

$$FX = \coprod_X K$$

By applying the proof of Theorem 4.6 to F , we get that \mathcal{K} is LS-accessible.

Example 4.8. Consider a one-sorted signature Σ given by a sort s and an ω -ary function symbol $f : s^\omega \rightarrow s$. Let T be the $L_{\omega_1\omega_1}(\Sigma)$ theory saying that f is a bijection. Thus T consists of the formula

$$(\forall y)(\exists!x)(f(x) = y)$$

where x is an ω -string of variables, and $\exists!$ denotes unique existence. By general facts about categories of models of $\exists!$ -sentences (called *limit theories*), the category $\mathbf{Mod}(T)$ is locally ω_1 -presentable; see [2, 5.30]. Hence, in particular, it is accessible and has directed colimits.

If μ has cofinality ω then

$$\mu^\omega > \mu$$

cf. [4, Corollary 4 of Theorem 17]. Thus T does not have models of cardinalities of cofinality ω . Since $\mu^\omega = \mu$ whenever the cardinal μ is of the form ν^ω , $\mathbf{Mod}(T)$ is a large category. Thus there is a proper class of cardinalities in which T has a model and a proper class of cardinalities in which T does not have a model. But, following Theorem 4.6, $\mathbf{Mod}(T)$ is LS-accessible. The point is that presentation ranks and cardinalities will never start to coincide.

Without changing the category of models, one can change the theory T to an equational $L_{\omega_1\omega_1}$ theory T' in the signature Σ' consisting of f and ω unary function symbols $g_i : s \rightarrow s$. Equations of T' state that $f : X^\omega \rightarrow X$ and $\langle g_i \rangle_{i \in \omega} : X \rightarrow X^\omega$ are inverse maps (thus, bijections). The functor V of the proof of Theorem 4.6 is the underlying set functor $\mathbf{Mod}(T') \rightarrow \mathbf{Set}$ and the functor F is the free algebra functor $\mathbf{Set} \rightarrow \mathbf{Mod}(T')$. Following the proof of Theorem 4.6, $\text{rank}(FX) = |X|^+$ for any uncountable set X . For free T' -algebras FX , the difference between $\text{rank}(FX)$ and the size of the underlying set $|VFX|$ can become arbitrarily large.

Definition 4.9. A functor $H : \mathcal{K} \rightarrow \mathcal{L}$ will be called *iso-full* if for any isomorphism $f : HA \rightarrow HB$ there is an isomorphism $\bar{f} : A \rightarrow B$ such that $H(\bar{f}) = f$.

Remark 4.10. Any iso-full functor H is *essentially injective on objects* in the sense that $HA \cong HB$ implies that $A \cong B$. Any iso-full and faithful functor is conservative, i.e. reflects isomorphisms.

A functor H is called *transportable* if for an isomorphism $f : HA \rightarrow B$ there is a unique isomorphism $\bar{f} : A \rightarrow \bar{B}$ such that $H(\bar{f}) = f$ (this includes $H\bar{B} = B$).

Theorem 4.11. Let \mathcal{K} be a large accessible category with directed colimits admitting an iso-full and faithful functor into a finitely accessible category preserving directed colimits. Then \mathcal{K} is LS-accessible.

Proof. Following Remark 2.3(3), every finitely accessible category admits a finitely accessible full embedding into $\mathbf{Str}(\Sigma)$ for some finitary signature Σ . Thus we can assume that there is an iso-full and faithful functor preserving directed colimits $H : \mathcal{K} \rightarrow \mathbf{Str}(\Sigma)$. Consider the pullback

$$\begin{array}{ccc}
 \mathcal{K} & \xrightarrow{H} & \mathbf{Str}(\Sigma) \\
 \bar{G} \uparrow & & \uparrow G \\
 \mathcal{L} & \xrightarrow{\bar{H}} & \mathbf{Emb}(\Sigma)
 \end{array}$$


where G is the inclusion. Since G is transportable, the pullback is equivalent to a pseudopullback (see [9, 5.1.1]). Hence \mathcal{L} is accessible (see [9, 5.1.6]) and clearly has directed colimits. The functors \bar{H} and \bar{G} preserve directed colimits because H and G have this property. Since the pullback of an iso-full, resp. faithful functor is iso-full, resp. faithful, the functors \bar{H} and \bar{G} are iso-full and faithful. Consequently, both \bar{H} and \bar{G} are essentially injective on objects (see Remark 4.10) and \bar{G} is surjective on objects. Following Propositions 4.1 and 4.3, there is a regular cardinal λ such that both \bar{G} and \bar{H} are well λ -accessible. Since \bar{G} is iso-full and faithful, it is conservative and thus reflects split epimorphisms (see Remarks 4.10 and 3.5).

Now, following Proposition 3.7, \bar{G} preserves presentability ranks ν for $\nu > \lambda$. Thus it suffices to prove that \mathcal{L} is λ -LS-accessible. Consider a cardinal $\mu \geq \lambda$. Since \mathcal{K} is large and \bar{G} is essentially injective on objects, \mathcal{L} is large. Since \bar{H} is essentially injective on objects, there is an object L in \mathcal{L} such that $\mu \leq |\bar{H}L|$. Since \mathcal{L} is μ^+ -accessible, L is a μ^+ -directed colimit of μ^+ -presentable objects $L_i, i \in I$. Let $(l_i : L_i \rightarrow L)_{i \in I}$ denote the colimit cocone. There is an embedding $f : A \rightarrow \bar{H}L$ with $|A| = \mu$. Since A is μ^+ -presentable, $f = \bar{H}(l_j)g$ for some $g : A \rightarrow \bar{H}L_j$. Hence $\mu \leq |\bar{H}L_j|$. Since \bar{H} preserves presentability ranks ν for $\lambda < \nu$ (following Proposition 3.7), $\bar{H}L_j$ is μ^+ -presentable, i.e., $|\bar{H}L_j| \leq \mu$. Hence $|\bar{H}L_j| = \mu$, i.e., $\text{rank}(\bar{H}L_j) = \mu^+$. Therefore $\text{rank}(L_j) = \mu^+$. We have proved that \mathcal{L} is λ -LS-accessible. \square

Using the Ehrenfeucht–Mostowski models, we can prove a stronger result. (Note that any finitely accessible category has a faithful, directed colimit preserving functor into **Set** and thus any category \mathcal{K} from Theorem 4.11 has this property.)

Theorem 4.12. *Let \mathcal{K} be a large accessible category with directed colimits equipped with a faithful functor $H : \mathcal{K} \rightarrow \mathbf{Set}$ preserving directed colimits. Then \mathcal{K} is LS-accessible.*

Proof. Consider a pullback



analogous to that in the proof of Theorem 4.11. Again, \mathcal{L} is a λ -accessible category with directed colimits. Following Theorem 4.5, $\mathcal{L} = \text{Ind}_\lambda \mathcal{C}$ is a full reflective subcategory of the finitely accessible category $\text{Ind} \mathcal{C}$. Let $F : \text{Ind} \mathcal{C} \rightarrow \mathcal{L}$ be a left adjoint to the inclusion $\mathcal{L} \rightarrow \text{Ind} \mathcal{C}$. The composition $H\bar{G}F : \text{Ind} \mathcal{C} \rightarrow \mathbf{Set}$ preserves directed colimits and thus it is uniquely determined by its domain restriction on \mathcal{C} . Since F is the identity functor on its domain restriction on \mathcal{C} , the domain restriction of $H\bar{G}F$ on \mathcal{C} is faithful. We will prove that $H\bar{G}F$ is faithful. Consider two distinct morphisms $f, g : X \rightarrow Y$ in $\text{Ind} \mathcal{C}$. There are objects $C, D \in \mathcal{C}$, morphisms $u : C \rightarrow X, v : D \rightarrow Y$ and distinct morphisms $f', g' : C \rightarrow D$ such that $fu = vf'$ and $gu = vg'$. Then $H\bar{G}F(f')$ and $H\bar{G}F(g')$ are distinct and, since $H\bar{G}F(v) = G\bar{H}F(v)$ is a monomorphism, $H\bar{G}F(fu)$ and $H\bar{G}F(gu)$ are distinct. Therefore $H\bar{G}F(f)$ and $H\bar{G}F(g)$ are distinct. We have proved that $H\bar{G}F$ is faithful and thus $\bar{G}F$ is faithful.

Since every finitely accessible category is ∞, ω -elementary (see [9, 5.2.6]), Remark 4.4 provides a faithful functor $E : \mathbf{Lin} \rightarrow \text{Ind} \mathcal{C}$ preserving directed colimits. Thus $\bar{G}FE$ is faithful and preserves directed colimits. By the same argument as at the end of Remark 4.4 (applied to $\bar{G}FE$ instead to E), this functor preserves presentability ranks starting from some cardinal. Thus \mathcal{K} is LS-accessible. \square

Remark 4.13. We have proved that each category \mathcal{K} from Theorem 4.12 admits a faithful functor $E : \mathcal{L} \rightarrow \mathcal{K}$ from a finitely accessible category \mathcal{L} which preserves directed colimits and is surjective on objects. Consequently, it admits a faithful functor $\mathbf{Lin} \rightarrow \mathcal{K}$ preserving directed colimits. Moreover, if all morphisms in \mathcal{K} are monomorphisms, we do not need a pullback from the proof of Theorem 4.12 and the functor E is then even surjective on morphisms.

We will give an example of an accessible category \mathcal{W} with directed colimits with no faithful functor $\mathcal{W} \rightarrow \mathbf{Set}$ preserving directed colimits. This shows that the two parts of the assumption on the category \mathcal{K} in Theorem 4.12 are independent. At the same time, this category \mathcal{W} is an example of an accessible category with directed colimits that is not ∞, ω -elementary. On the other hand, we do not know whether the assumptions of Theorem 4.12 are necessary for the conclusion. Possibly every accessible category with directed colimits is LS-accessible; possibly every accessible category is LS-accessible.

Example 4.14. Let \mathcal{W} be the category of well-ordered sets where morphisms are either order preserving mappings or constant mappings. Since the category \mathcal{W}_0 of well-ordered sets and substructure embeddings is ω_1 -accessible, \mathcal{W} is ω_1 -accessible as well. The category \mathcal{W}_0 does not have all directed colimits. For example, a countable chain

$$A_0 \rightarrow A_1 \rightarrow \dots \rightarrow A_n \rightarrow \dots$$

such that each A_{n+1} , $0 \leq n$ contains an element a smaller than any $x \in A_n$, does not have a colimit in \mathcal{W}_0 . In fact, it does not have any compatible cocone $A_n \rightarrow A$, $0 \leq n$. In \mathcal{W} , it does have a colimit: the one-element chain 1. One can show that \mathcal{W} has all directed colimits.

Assume that there exists a faithful functor $H : \mathcal{W} \rightarrow \mathbf{Set}$ preserving directed colimits. Following Remark 4.13, there is a faithful functor $\mathbf{Lin} \rightarrow \mathcal{W}$. This is impossible because \mathcal{W} is automorphism rigid – the only isomorphisms $X \rightarrow X$ in \mathcal{W} are identities.

5. Abstract elementary classes


Consider the following hierarchy of accessible categories with directed colimits:

- (1) finitely accessible categories,
- (2) ∞, ω -elementary categories,
- (3) accessible categories with directed colimits admitting a faithful functor preserving directed colimits into a finitely accessible category,
- (4) accessible categories with directed colimits.

Each class is contained in the next. The original definition of class (2), ∞, ω -elementary categories, is model-theoretic, as categories equivalent to the category of models and homomorphisms of theories in $L_{\infty, \omega}^*$. There exist categorical descriptions too: these are the categories equivalent to the category of points of some Grothendieck topos; and, by [9, 5.2.6], exactly the 2-categorical limits of finitely accessible categories and finitely accessible functors. Since any finitely accessible category is equipped with a faithful functor into \mathbf{Set} preserving directed colimits, class (3) coincides with the categories from Theorem 4.12. Following Example 4.14, the inclusion of (3) in (4) is proper. The inclusion of (1) in (2) is also proper. We are not aware of any example of a category belonging to (3) but not to (2). We do know that any category from (2) admits a full embedding preserving directed colimits into a finitely accessible category.

We will show that the categories of models-and-strong-embeddings coming from Shelah’s Abstract Elementary Classes are between (2) and (4). We will also introduce a class of abstract elementary categories as “abstract elementary classes” without any assumption about morphisms being embeddings.

Definition 5.1. A functor $H : \mathcal{K} \rightarrow \mathcal{L}$ will be called *nearly full* if for each commutative triangle



there is $\bar{f} : A \rightarrow B$ in \mathcal{K} such that $H(\bar{f}) = f$.

A subcategory \mathcal{K} of a category \mathcal{L} will be called *nearly full* if the embedding $\mathcal{K} \rightarrow \mathcal{L}$ is nearly full.

Remark 5.2. (1) Every full functor is nearly full. If H is faithful and nearly full, we also have $g\bar{f} = h$ in the definition above. Thus every faithful and nearly full functor reflects split epimorphisms.

(2) [5] calls nearly full subcategories coherent. Since this term is overused in category theory, we are changing the terminology.

(3) Recall that a subcategory is called *replete* if it is closed under isomorphic objects. Any subcategory \mathcal{K} of a finitely accessible category \mathcal{L} closed under directed colimits is replete and iso-full. Indeed, being closed under directed colimits means that each directed colimit in \mathcal{L} of objects from \mathcal{K} belongs to \mathcal{K} . And, directed colimits are determined up to an isomorphism.

If \mathcal{K} is a full subcategory of an accessible category \mathcal{L} such that \mathcal{K} has directed colimits and the embedding of \mathcal{K} into \mathcal{L} preserves them then the *replete closure* $\bar{\mathcal{K}}$ of \mathcal{K} is closed under directed colimits in \mathcal{L} . Moreover the categories \mathcal{K} and $\bar{\mathcal{K}}$ are equivalent. Of course, the replete closure of \mathcal{K} is the full subcategory of \mathcal{L} consisting of all objects isomorphic to an object of \mathcal{K} .

Definition 5.3. An accessible categories with directed colimits will be called an *abstract elementary category* if it admits an iso-full and nearly full embedding preserving directed colimits into a finitely accessible category.

Abstract elementary categories are closely related to Shelah’s Abstract Elementary Classes; they differ from those introduced in [5]. We recall the definition of an abstract elementary class using the language of category theory.

Definition 5.4. Let Σ be a finitary signature. A nearly full subcategory \mathcal{K} of $\mathbf{Emb}(\Sigma)$ is called an *abstract elementary class* if it is closed in $\mathbf{Emb}(\Sigma)$ under directed colimits and there is a cardinal λ such that if $f : A \rightarrow B$ is a substructure with $B \in \mathcal{K}$ then there is $h : A' \rightarrow B$ in \mathcal{K} such that f factorizes through h and $|A'| \leq |A| + \lambda$.

The standard formulation (see [3]) uses colimits of continuous chains instead of directed colimits. But it is well known that this does not change anything (see [3], or [2, 1.7]; continuous chains are called smooth there). The standard formulation also includes that \mathcal{K} is replete and iso-full but Remark 5.2(3) shows that our formulation is the same. We allow many-sorted signatures in Definition 5.4. Since each many-sorted signature can be made single-sorted, this does not change the concept of an abstract elementary class.

Theorem 5.5. *Let Σ be a finitary signature. A nearly full subcategory \mathcal{K} of $\mathbf{Emb}(\Sigma)$ is an abstract elementary class if and only if it is an accessible category closed under directed colimits in $\mathbf{Emb}(\Sigma)$.*

Proof. Let \mathcal{K} be an abstract elementary class in $\mathbf{Emb}(\Sigma)$. We know that $\mathbf{Emb}(\Sigma)$ is finitely accessible and there is a regular cardinal $\lambda < \kappa$ such that, for each regular cardinal $\kappa \leq \mu$, μ -presentable objects in $\mathbf{Emb}(\Sigma)$ are precisely Σ -structures A such that $|A| < \mu$. Since $\mathbf{Emb}(\Sigma)$ is κ -accessible, Definition 5.4 yields that each object of \mathcal{K} is a κ -directed colimit of κ -presentable objects in \mathcal{K} . Thus \mathcal{K} is κ -accessible.

Conversely, let \mathcal{K} be an accessible nearly full subcategory of $\mathbf{Emb}(\Sigma)$ closed under directed colimits. Then the embedding $\mathcal{K} \rightarrow \mathbf{Emb}(\Sigma)$ is well λ -accessible for some regular cardinal λ greater than κ above. Let $f : A \rightarrow B$ be a substructure embedding with $B \in \mathcal{K}$ and put $\mu = |A| + \lambda$. Since \mathcal{K} is μ^+ -accessible, B is a μ^+ -directed colimit of μ^+ -presentable objects B_i , $i \in I$ in \mathcal{K} . Since the embedding $\mathcal{K} \rightarrow \mathbf{Emb}(\Sigma)$ preserves μ^+ -directed colimits and μ^+ -presentable objects, B is a μ^+ -directed colimit of objects B_i which are μ^+ -presentable in $\mathbf{Emb}(\Sigma)$. Since A is μ^+ -presentable (because $\kappa < \mu^+$), the substructure embedding $f : A \rightarrow B$ factorizes through some B_j , $j \in I$. It suffices to put $A' = B_j$. \square

This result is an improvement of [7, 5.9 and 5.10] (see [8, 4.1 and 4.9], as well; there is a related work [5]). Lieberman assumes that \mathcal{K} is well accessible while we use Proposition 4.1 to prove this from the existence of directed colimits. Otherwise, our proof is the same as that of Lieberman.

Remark 5.6. (1) Categories $\mathbf{Elem}(T)$ where T is an $L_{\kappa\omega}$ theory and $\mathbf{Emb}(T)$ where T is a basic $L_{\kappa\omega}$ theory are abstract elementary classes.

(2) Each large abstract elementary class is an LS-accessible category. This follows from Theorem 4.11 but also directly from the proof of Theorem 5.5. Following this proof, we have $\text{rank}(A) = |A|^+$ for each A in \mathcal{K} with $|A| \geq \lambda$.

We can write our characterization of abstract elementary classes in the language of category theory, i.e., without using Σ -structures.

Corollary 5.7. *A category is equivalent to an abstract elementary class if and only if it is an accessible category with directed colimits whose morphisms are monomorphisms and which admits an iso-full and nearly full embedding into a finitely accessible category preserving directed colimits and monomorphisms.*

Proof. Necessity is evident because $\mathbf{Emb}(\Sigma)$ is finitely accessible. Consider a category \mathcal{K} satisfying the conditions above. Let $H : \mathcal{K} \rightarrow \mathcal{L}$ be the corresponding functor into a finitely accessible category. Let \mathcal{A} be the full subcategory of \mathcal{L} consisting of finitely presentable objects. Consider the canonical functor

$$E : \mathcal{L} \rightarrow \mathbf{Set}^{A^{\text{op}}}$$

(see [2, 1.25]). E preserves directed colimits (see [2, 1.26]). Since objects of $\mathbf{Set}^{A^{\text{op}}}$ can be viewed as many-sorted unary algebras, $\mathbf{Set}^{A^{\text{op}}}$ is a full subcategory of $\mathbf{Str}(\Sigma)$ for a finitary signature Σ containing operation symbols only. Thus embeddings in $\mathbf{Str}(\Sigma)$ coincide with monomorphisms. Hence the codomain restriction of the composition EH is the iso-full and nearly full embedding

$$\mathcal{K} \rightarrow \mathbf{Emb}(\Sigma)$$

preserving directed colimits. Following Remark 5.2(3) and Theorem 5.5, \mathcal{K} is equivalent to an abstract elementary class. \square

Remark 5.8. (1) Our definition of abstract elementary category is motivated by Corollary 5.7; it results from dropping the hypotheses on monomorphisms. Any finitely accessible category whose morphisms are not monomorphisms, like posets and isotone mappings, is an example of an abstract elementary category that is not an abstract elementary class.

(2) Any abstract elementary class is an abstract elementary category. On the other hand, let \mathcal{K} be an abstract elementary category with a functor H into a finitely accessible category from Definition 5.3. Without any loss of generality, we can assume that $H : \mathcal{K} \rightarrow \mathbf{Str}(\Sigma)$ where Σ is a finitary signature. Consider the pullback from the proof of Theorem 4.11. Then

\mathcal{L} is an accessible category with directed colimits whose morphisms are monomorphisms and \bar{H} is an iso-full and nearly full embedding preserving directed colimits and monomorphisms. Thus \mathcal{L} is equivalent to an abstract elementary class. Moreover, the functor \bar{G} is onto on objects and preserves presentability ranks starting from some regular cardinal.

(3) We are not aware of any abstract elementary category which is not ∞, ω -elementary. A simple example of an abstract elementary class \mathcal{K} in Σ which is not closed under $L_{\infty, \omega}$ -elementary equivalence (thus not axiomatizable in any $L_{\kappa\omega}(\Sigma)$) is given in [6, 2.10]. There, Σ is a single-sorted signature containing just a unary relation symbol P . Objects of \mathcal{K} are Σ -structures K such that P_K is countable and the complement of P_K is infinite. Morphisms of \mathcal{K} are substructure embeddings which are identities on P . But, \mathcal{K} is isomorphic to the category of infinite sets and monomorphisms. The latter category is axiomatizable by a basic $L_{\omega\omega}$ theory in the empty signature. Thus it is ∞, ω -elementary.

Proposition 5.9. *Let \mathcal{K} be an abstract elementary category. Then there is an ∞, ω -elementary category \mathcal{L} and a faithful functor $E : \mathcal{L} \rightarrow \mathcal{K}$ which preserves directed colimits and is surjective on objects.*

Proof. Let $\bar{G} : \mathcal{L} \rightarrow \mathcal{K}$ be the embedding of an abstract elementary class \mathcal{L} from Remark 5.8(2). Shelah's Presentation Theorem (see [3, 4.15]) yields a finitary signature $\Sigma \subseteq \Sigma'$, an $L_{\omega\omega}$ theory T' and a set Γ of quantifier free types in Σ' such that \mathcal{L} consists of Σ -reducts of T' -models omitting types from Γ . Omitting a type can be expressed as an $L_{\infty\omega}$ -sentence. By adding these sentences for types from Γ to T' , we get an $L_{\infty\omega}$ theory T'' . Since $\mathbf{Elem}(T'')$ is ∞, ω -elementary by Remark 2.3(4), the reduct functor $R : \mathbf{Elem}(T'') \rightarrow \mathcal{L}$ has the desired properties for \mathcal{L} . Thus $\bar{G}R$ has these properties for \mathcal{K} . \square

6. Categoricity

Definition 6.1. Let λ be an infinite cardinal. A category \mathcal{K} is called λ -categorical if it has, up to isomorphism, precisely one object of the presentability rank λ^+ .

Remark 6.2. Following Remark 5.6(2), Definition 6.1 (suggested in [10]) is in accordance with its model theoretic meaning in abstract elementary classes for sufficiently large cardinals λ .

Shelah's Categoricity Conjecture claims that for every abstract elementary class \mathcal{K} there is a cardinal κ such that \mathcal{K} is either λ -categorical for all cardinals $\lambda \geq \kappa$ or \mathcal{K} is not λ -categorical for any cardinal $\lambda \geq \kappa$. Following Remark 5.6(2), this is a statement about \mathcal{K} as a category, i.e., it does not depend on the signature in which the abstract elementary class is presented.

By Remark 5.8(2), Shelah's Categoricity Conjecture is equivalent to the Categoricity Conjecture for abstract elementary categories. It is natural to ask about the status of the Categoricity Conjecture for accessible categories belonging to other levels of our hierarchy. At present, we can offer little information other than this easy observation.

Example 6.3. Suppose \mathcal{K} is an accessible category which is not LS-accessible. Let $\mathcal{L} = \mathcal{K} \sqcup \mathbf{Set}$ be the disjoint union of \mathcal{K} and the category of sets. Then \mathcal{L} is an accessible category and there is a proper class of cardinals λ such that \mathcal{L} is λ -categorical and, at the same time, a proper class of cardinals λ such that \mathcal{L} is not λ -categorical.

If this \mathcal{K} has directed colimits then the categoricity conjecture fails for the class (4). By Theorem 4.12, however, this simple trick does not help for categories in class (3) of our hierarchy.

Remark 6.4. Shelah's Conjecture seems to be unknown even for finitely accessible categories, lying at level (1) of the hierarchy. It would be interesting to extend the exquisite Galois-theoretic machinery of [3] to this setting.

References

- [1] J. Adámek, H. Herrlich, J. Rosický, W. Tholen, On a generalized small-object argument for the injectivity subcategory problem, Cah. Topol. Géom. Différ. Catég. XLIII (2002) 83–106.
- [2] J. Adámek, J. Rosický, Locally Presentable and Accessible Categories, Cambridge University Press, 1994.
- [3] J.T. Baldwin, Categoricity, Amer. Math. Soc., 2009.
- [4] T. Jech, Set Theory, Academic Press, 1978.
- [5] J. Kirby, Abstract elementary categories, preprint, 2008.
- [6] D.W. Kueker, Abstract elementary classes and infinitary logic, Ann. Pure Appl. Logic 156 (2008) 274–286.
- [7] M.J. Lieberman, Topological and category-theoretic aspects of abstract elementary classes, Thesis, The University of Michigan, 2009.
- [8] M.J. Lieberman, Category theoretic aspects of abstract elementary classes, Ann. Pure Appl. Logic 162 (2011) 903–915.
- [9] M. Makkai, R. Paré, Accessible Categories: The Foundation of Categorical Model Theory, Contemp. Math., vol. 104, Amer. Math. Soc., 1989.
- [10] J. Rosický, Accessible categories, saturation and categoricity, J. Symbolic Logic 62 (1997) 891–901.