



Egyptian Mathematical Society  
**Journal of the Egyptian Mathematical Society**

[www.etms-eg.org](http://www.etms-eg.org)  
[www.elsevier.com/locate/joems](http://www.elsevier.com/locate/joems)



## Original Article

# Picard, Adomian and predictor–corrector methods for an initial value problem of arbitrary (fractional) orders differential equation



Adel S. Mohamed, R.A. Mahmoud\*

Zagazig University, Faculty of Science, Egypt

Received 24 November 2014; revised 27 December 2014; accepted 1 January 2015  
 Available online 10 February 2015

### Keywords

Fractional differential equation;  
 Picard method;  
 Adomian method;  
 Predictor corrector method;  
 Convergence analysis;  
 Error analysis

**Abstract** We study the two analytical methods, the classical method of successive approximations (Picard method), Adomian decomposition method (ADM) see (Abbaoui and Cherruault, 1994; Adomian et al., 1992; Adomian, 1995) [1–3] and the (numerical method) predictor corrector method (PECE) for an initial value problem of arbitrary (fractional) orders differential equation (FDE). The existence and uniqueness of the solution will be proved and the convergence will be discussed for each method. Some examples will be studied.

**2000 Mathematical Subject Classification:** 26A33; 26A18; 39B12

Copyright 2015, Egyptian Mathematical Society. Production and hosting by Elsevier B.V. All rights reserved

## 1. Introduction

Let  $\alpha \in [0, 1)$ . In this paper, we study the existence and uniqueness of the solution of the initial value problem


$$\frac{dx}{dt} + D^\alpha x(t) = f(t, x(t)), \quad 0 < \alpha < 1, \quad (1)$$

$$x(0) = \tilde{x}. \quad (2)$$

\* Corresponding author.

E-mail addresses: [3adel@live.nl](mailto:3adel@live.nl) (A.S. Mohamed), [rony\\_695@yahoo.com](mailto:rony_695@yahoo.com) (R.A. Mahmoud).

Peer review under responsibility of Egyptian Mathematical Society.



We apply the three methods Adomian, Picard and predictor–corrector to obtain numerical solution of the problem (1) and (2).

Now, the definition of the fractional-order integral and differential operators are given by the following.

**Definition 1.** Let  $\beta$  be a positive real number, the fractional-order integral of order  $\beta$  of the function  $f$  is defined on the interval  $[0, T]$  by

$$I_0^\beta f(t) = \int_0^t \frac{(t-s)^{\beta-1}}{\Gamma(\beta)} f(s) ds$$

and the fractional-order derivative of the function  $f \in C^1[0, T]$  of order  $\alpha \in (0, 1]$  is defined by

$$D^\alpha f(t) = I^{1-\alpha} \frac{df}{dt}.$$

**2. Uniqueness theorem**

Now, the initial value problem (1) and (2) will be investigated under the following assumptions:

- (i)  $f : J = [0, T] \times D \rightarrow R$  is continuous where  $D$  is a closed subset of  $R$ ;
- (ii)  $f$  satisfies the Lipschitz condition with Lipschitz constant  $L$   
i.e

$$|f(t, x) - f(t, y)| \leq L|x - y|, \quad \forall (t, x), (t, y) \in I \times D.$$

Let  $C = C(J)$  be the space of all real-valued functions which are continuous on  $J$ .

**Definition 2.** By a solution of the problem (1) and (2) we mean a function  $x \in C[0, T]$ . This function satisfies the problem (1) and (2).

Let  $x(t)$  be a solution of the initial value problem (1) and (2). Integrating (1) we obtain

$$x(t) - \tilde{x} + I^{1-\alpha}(x(t) - \tilde{x}) = If(t, x),$$

then we have

$$x(t) = \tilde{x} \left( 1 + \frac{t^{1-\alpha}}{\Gamma(2-\alpha)} \right) - \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} x(s) ds + \int_0^t f(s, x(s)) ds. \tag{3}$$

Now let  $x \in C(J)$  be a solution of the integral Eq. (3), then

$$\begin{aligned} \frac{dx}{dt} &= 0 + \tilde{x} \frac{(1-\alpha)t^{-\alpha}}{\Gamma(2-\alpha)} - \frac{d}{dt} \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} x(s) ds \\ &\quad + \frac{d}{dt} \int_0^t f(s, x(s)) ds = \tilde{x} \frac{t^{-\alpha}}{\Gamma(1-\alpha)} \\ &\quad - \tilde{x} \frac{t^{-\alpha}}{\Gamma(1-\alpha)} - I^{1-\alpha} \frac{dx}{dt} + f(t, x) \end{aligned}$$

and

$$\frac{dx}{dt} + D^\alpha x(t) = f(t, x(t))$$

also

$$\begin{aligned} \tilde{x} &= \tilde{x} \left( 1 + \frac{t^{1-\alpha}}{\Gamma(2-\alpha)} \right)_{t=0} - \left( \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} x(s) ds \right)_{t=0} \\ &\quad + \left( \int_0^t f(s, x(s)) ds \right)_{t=0} \end{aligned}$$

then the problem (1) and (2) and the integral Eq. (3) are equivalent.

Comparison between analytical methods is studied in many papers, for examples [4-7].

Define the operator  $F$  as

$$\begin{aligned} (Fx)(t) &= \tilde{x} \left( 1 + \frac{t^{1-\alpha}}{\Gamma(2-\alpha)} \right) - \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} x(s) ds \\ &\quad + \int_0^t f(s, x(s)) ds, \quad \alpha > 0, \quad \forall x \in C. \end{aligned}$$

**Theorem 1.** Let the assumptions (i)-(ii) be satisfied if  $LT + \frac{T^{1-\alpha}}{\Gamma(2-\alpha)} < 1$ , then the initial value problem (1) has a unique solution  $x \in C$ .

**Proof.** Firstly we prove that  $F : C \rightarrow C$  is continuous.

let  $x \in C(J)$ ,  $t_1, t_2 \in J$  such that  $|t_2 - t_1| < \delta$

$$\begin{aligned} Fx(t_2) - Fx(t_1) &= \tilde{x} \frac{t_2^{1-\alpha}}{\Gamma(2-\alpha)} - \int_0^{t_2} \frac{(t_2-s)^{-\alpha}}{\Gamma(1-\alpha)} x(s) ds \\ &\quad + \int_0^{t_2} f(s, x(s)) ds - \tilde{x} \frac{t_1^{1-\alpha}}{\Gamma(2-\alpha)} \\ &\quad + \int_0^{t_1} \frac{(t_1-s)^{-\alpha}}{\Gamma(1-\alpha)} x(s) ds - \int_0^{t_1} f(s, x(s)) ds \end{aligned}$$

$$\begin{aligned} |Fx(t_2) - Fx(t_1)| &\leq |\tilde{x}| \left| \frac{t_2^{1-\alpha} - t_1^{1-\alpha}}{\Gamma(2-\alpha)} \right| - \int_0^{t_2} \frac{(t_2-s)^{-\alpha}}{\Gamma(1-\alpha)} |x(s)| ds \\ &\quad + \int_0^{t_1} \frac{(t_1-s)^{-\alpha}}{\Gamma(1-\alpha)} |x(s)| ds + \int_{t_1}^{t_2} |f(s, x(s))| ds \end{aligned}$$

$$\begin{aligned} \|Fx(t_2) - Fx(t_1)\| &= \max_{t \in J} |Fx(t_2) - Fx(t_1)| \\ &\leq \|\tilde{x}\| \left| \frac{t_2^{1-\alpha} - t_1^{1-\alpha}}{\Gamma(2-\alpha)} \right| - \|x\| \int_0^{t_2} \frac{(t_2-s)^{-\alpha}}{\Gamma(1-\alpha)} ds \\ &\quad + \|x\| \int_0^{t_1} \frac{(t_1-s)^{-\alpha}}{\Gamma(1-\alpha)} ds + k \int_{t_1}^{t_2} ds \\ &\leq \|\tilde{x}\| \left| \frac{t_2^{1-\alpha} - t_1^{1-\alpha}}{\Gamma(2-\alpha)} \right| \\ &\quad + \frac{\|x\|}{\Gamma(2-\alpha)} |t_2^{1-\alpha} - t_1^{1-\alpha}| + k|t_2 - t_1| \\ &\leq \frac{(\|\tilde{x}\| + \|x\|)}{\Gamma(2-\alpha)} |t_2^{1-\alpha} - t_1^{1-\alpha}| + k|t_2 - t_1| \leq \epsilon \end{aligned}$$

where

$$|f(t, x(t))| \leq k,$$

This proves that  $F : C[0, T] \rightarrow C[0, T]$ .

Now we prove that  $F$  is contraction, for this we have

$$\begin{aligned} Fx - Fy &= - \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} x(s) ds + \int_0^t f(s, x(s)) ds \\ &\quad + \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} y(s) ds - \int_0^t f(s, y(s)) ds \end{aligned}$$

$$\begin{aligned} |Fx - Fy| &\leq \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} |x(s) - y(s)| ds \\ &\quad + \int_0^t |f(s, x(s)) - f(s, y(s))| ds \\ &\leq \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} |x(s) - y(s)| ds + L \int_0^t |x(s) - y(s)| ds \\ &\leq \|x - y\| \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} ds + LT \|x - y\| \end{aligned}$$

$$\|Fx - Fy\| \leq \|x - y\| \left( LT + \frac{T^{1-\alpha}}{\Gamma(2-\alpha)} \right).$$

Since  $LT + \frac{T^{1-\alpha}}{\Gamma(2-\alpha)} < 1$ , then F is a contraction and F has a unique fixed point, thus there exists a unique solution  $x \in C(J)$  of the initial value problem (1) and (2).  $\square$

### 3. Method of successive approximations (Picard method)

Applying Picard method to integral Eq. (3), then the solution is constructed by the sequence

$$x_n(t) = \tilde{x} \left( 1 + \frac{t^{1-\alpha}}{\Gamma(2-\alpha)} \right) - \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} x_{n-1}(s) ds + \int_0^t f(s, x_{n-1}(s)) ds, \quad n = 1, 2, \dots \tag{4}$$

$$x_0 = \tilde{x} \left( 1 + \frac{t^{1-\alpha}}{\Gamma(2-\alpha)} \right)$$

All the functions  $x_n(t)$  are continuous functions and  $x_n$  can be written as a sum of successive differences

$$x_n = x_0 + \sum_{j=1}^n (x_j - x_{j-1})$$

This means that convergence of the sequence  $x_n$  is equivalent to convergence the infinite series  $\sum (x_j - x_{j-1})$  and the solution will be

$$x(t) = \lim_{n \rightarrow \infty} x_n(t)$$

i.e. if the finite series  $\sum (x_j - x_{j-1})$  converges, then the sequence  $x_n(t)$  will be converge to  $x(t)$ . To prove the uniform convergence of  $\{x_n(t)\}$ , we shall consider the associated series

$$\sum_{j=1}^{\infty} (x_j - x_{j-1})$$

from (4) for  $n = 1$ , we get

$$|x_1(t) - x_0| = \left| - \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} x_0(s) ds + \int_0^t f(s, x_0(s)) ds \right| \leq x_0 \frac{t^{1-\alpha}}{\Gamma(2-\alpha)} + MT$$

and

$$|x_1(t) - x_0| \leq x_0 \frac{T^{1-\alpha}}{\Gamma(2-\alpha)} + MT \tag{5}$$

from (4)

$$\begin{aligned} |x_2 - x_1| &\leq \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} |x_1(s) - x_0(s)| ds + \int_0^t |f(s, x_1(s)) - f(s, x_0(s))| ds \\ &\leq \left( \frac{x_0 T^{1-\alpha}}{\Gamma(2-\alpha)} + MT \right) \left( \frac{t^{1-\alpha}}{\Gamma(2-\alpha)} \right) + L \left( \frac{x_0 T^{1-\alpha}}{\Gamma(2-\alpha)} + MT \right) t \\ &\leq \left( \frac{x_0 T^{1-\alpha}}{\Gamma(2-\alpha)} + MT \right) \left( \frac{T^{1-\alpha}}{\Gamma(2-\alpha)} + LT \right) \end{aligned}$$

and

$$\begin{aligned} |x_3 - x_2| &\leq \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} |x_2(s) - x_1(s)| ds + \int_0^t |f(s, x_2(s)) - f(s, x_1(s))| ds, \\ &\leq \left( \frac{x_0 T^{1-\alpha}}{\Gamma(2-\alpha)} + MT \right) \left( \frac{T^{1-\alpha}}{\Gamma(2-\alpha)} + LT \right)^2 \end{aligned}$$

Repeating this technique, we obtain the general estimate for the terms of the series:

$$|x_n - x_{n-1}| \leq \left( \frac{x_0 T^{1-\alpha}}{\Gamma(2-\alpha)} + MT \right) \left( \frac{T^{1-\alpha}}{\Gamma(2-\alpha)} + LT \right)^{n-1}$$

Since  $\frac{T^{1-\alpha}}{\Gamma(2-\alpha)} + LT < 1$ , then the uniform convergence of

$$\sum_{n=1}^{\infty} (x_n - x_{n-1})$$

is proved and so the sequence  $\{x_n(t)\}$  is uniformly convergent.

### 4. Adomian decomposition method (ADM)

In this section, we shall study Adomian decomposition method (ADM) for the integral Eq. (3). The solution algorithm of the integral Eq. (3) using ADM is

$$x_0(t) = \tilde{x} \left( 1 + \frac{t^{1-\alpha}}{\Gamma(2-\alpha)} \right) \tag{6}$$

$$x_n(t) = - \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} x_{n-1}(s) ds + \int_0^t A_{n-1}(s) ds \tag{7}$$

where  $A_n$  Adomian polynomial of nonlinear term  $f(s, x)$ , which have the form

$$A_n = \frac{1}{n!} \frac{d^n}{d\lambda^n} \left( f \left( t, \sum_{i=0}^{\infty} \lambda^i x_i \right) \right)_{\lambda=0} \tag{8}$$

and the solution will be

$$x(t) = \sum_{i=0}^{\infty} x_i(t) \tag{9}$$

#### 4.1. Convergence analysis

**Theorem 2.** Let the solution of integral Eq. (3) be exist. If  $|x_1(t)| < k$ ,  $k$  is a positive constant, then the series solution (9) converges.

**Proof.** Define the sequence  $\{S_p\}$  such that  $S_p = \sum_{i=0}^p x_i(t)$  is the sequence of partial sums for the series  $\sum_{i=0}^{\infty} x_i(t)$  and we have

$$f(t, x) = \sum_{i=0}^{\infty} A_i,$$

let  $S_p$  and  $S_q$  be two partial sums with  $p > q$ , we are going to prove that  $\{S_p\}$  is the Cauchy sequence in the Banach space E.

$$\begin{aligned}
 S_p - S_q &= - \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} S_{p-1}(s) ds + \int_0^t \sum_{i=0}^p A_{i-1}(s) ds \\
 &\quad + \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} S_{q-1}(s) ds - \int_0^t \sum_{i=0}^q A_{i-1}(s) ds \\
 &= - \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} (S_{p-1}(s) - S_{q-1}(s)) ds \\
 &\quad + \int_0^t \left( \sum_{i=0}^q A_{i-1}(s) + \sum_{i=q+1}^p A_{i-1}(s) - \sum_{i=0}^q A_{i-1}(s) \right) ds
 \end{aligned}$$

$$\begin{aligned}
 \|S_p - S_q\| &\leq \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} |S_{p-1}(s) - S_{q-1}(s)| ds \\
 &\quad + \sup_{t \in I} \int_0^t \left| \sum_{i=q+1}^p A_{i-1}(s) \right| ds \\
 &\leq \|S_{p-1}(t) - S_{q-1}(t)\| \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} ds \\
 &\quad + \sup_{t \in I} \int_0^t \left| \sum_{i=q}^{p-1} A_i(s) \right| ds \\
 &\leq \|S_{p-1}(t) - S_{q-1}(t)\| \int_0^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} ds \\
 &\quad + \sup_{t \in I} |f(S, S_{p-1}) - f(S, S_{q-1})| \int_0^t ds \\
 &\leq \frac{T^{1-\alpha}}{\Gamma(2-\alpha)} \|S_{p-1} - S_{q-1}\| + LT \|S_{p-1} - S_{q-1}\| \\
 &\leq \left( LT + \frac{T^{1-\alpha}}{\Gamma(2-\alpha)} \right) \|S_{p-1} - S_{q-1}\| \\
 &\leq h \|S_{p-1} - S_{q-1}\|
 \end{aligned}$$

let  $p = q + 1$  then

$$\begin{aligned}
 \|S_{q+1} - S_q\| &\leq h \|S_q - S_{q-1}\| \leq h^2 \|S_{q-1} - S_{q-2}\| \\
 &\leq h^3 \|S_{q-2} - S_{q-3}\| \leq \dots \leq h^q \|S_1 - S_0\|
 \end{aligned}$$

from triangle inequality, we have

$$\begin{aligned}
 \|S_p - S_q\| &\leq \|S_{q+1} - S_q\| + \|S_{q+2} - S_{q+1}\| \\
 &\leq \dots \leq \|S_p - S_{p-1}\| \leq (h^q + h^{q+1} + \dots + h^{p-1}) \|S_1 - S_0\| \\
 &\leq h^q (1 + h + \dots + h^{p-q-1}) \|S_1 - S_0\| \\
 &\leq h^q \left( \frac{1 - h^{p-q}}{1 - h} \right) \|x_1\|
 \end{aligned}$$

where  $0 < h < 1$  and  $p > q$  imply that  $(1 - h^{p-q}) \leq 1$ . Consequently,

$$\|S_p - S_q\| \leq \frac{h^q}{1-h} \|x_1\| \leq \frac{h^q}{1-h} \sup_{t \in I} |x_1(t)|$$

but  $|x_1(t)| < k$  and as  $q \rightarrow \infty$ , then  $\|S_p - S_q\| \rightarrow 0$  and hence,  $\{S_p\}$  is a Cauchy sequence in this Banach space  $E$  and the series  $\sum_{i=0}^{\infty} x_i(t)$  converges.  $\square$

#### 4.2. Error analysis

**Theorem 3.** *The maximum absolute truncation error of the series solution (9) of the integral Eq. (3) is estimated to be*

$$\sup_{t \in I} \left| x(t) - \sum_{i=0}^q x_i(t) \right| \leq \frac{h^q}{1-h} \sup_{t \in I} |x_1(t)|$$

**Proof.** From theorem (2), we have

$$\|S_p - S_q\| \leq \frac{h^q}{1-h} \sup_{t \in I} |x_1(t)|$$

but  $S_p = \sum_{i=0}^p x_i(t)$  as  $p \rightarrow \infty$ , then  $S_p \rightarrow x(t)$  so,

$$\|x(t) - S_q\| \leq \frac{h^q}{1-h} \sup_{t \in I} |x_1(t)|$$

So, the maximum absolute truncation error in the interval  $I$  is

$$\sup_{t \in I} \left| x(t) - \sum_{i=0}^q x_i(t) \right| \leq \frac{h^q}{1-h} \sup_{t \in I} |x_1(t)|$$

and this completes the proof.  $\square$

### 5. Predictor–corrector method (PECE)

An Adams-type predictor–corrector method has been introduced in [8–12].

In this section, we use an Adams-type predictor–corrector method for the integral Eq. (3).

The product trapezoidal quadrature formula is used with  $t_j$  ( $j = 0, 1, \dots, k + 1$ ), taken with respect to the weight function  $(t_{k+1} - u)^{-\alpha}$ . In other words, one applies the approximation

$$\begin{aligned}
 \int_{t_0}^{t_{k+1}} (t_{k+1} - u)^{-\alpha} g(u) du &\approx \int_{t_0}^{t_{k+1}} (t_{k+1} - u)^{-\alpha} g_{k+1}(u) du \\
 &= \sum_{j=0}^{k+1} a_{j,k+1} g(t_j),
 \end{aligned}$$

where

$$a_{j,k+1} = \begin{cases} \frac{h^{1-\alpha}}{(1-\alpha)(2-\alpha)} [k^{2-\alpha} - (k+\alpha-1)(k+1)^{1-\alpha}] & \text{if } j = 0, \\ \frac{h^{1-\alpha}}{(1-\alpha)(2-\alpha)} & \text{if } j = k + 1 \end{cases}$$

and  $h$  is a step size, and for  $1 \leq j \leq k$

$$a_{j,k+1} = \frac{h^{1-\alpha}}{(1-\alpha)(2-\alpha)} [(k-j+2)^{2-\alpha} - 2(k-j+1)^{2-\alpha} + (k-j)^{2-\alpha}]$$

for  $\alpha = 0$  we have

$$a_{j,k+1} = \begin{cases} \frac{h}{2} & \text{if } j = 0, j = k + 1 \\ \frac{h}{2} [(k-j+2)^2 - 2(k-j+1)^2 + (k-j)^2] & \text{if } 1 \leq j \leq k \end{cases}$$

This yields the corrector formula, i.e. the fractional variant of the one-step Adams–Moulton method

$$\begin{aligned}
 x_{k+1} &= x_0(t_{k+1}) \left( 1 + \frac{t_{k+1}^{1-\alpha}}{\Gamma(2-\alpha)} \right) \\
 &\quad - \frac{1}{\Gamma(1-\alpha)} \left[ \sum_{j=0}^k a_{j,k+1} x(t_j) + a_{k+1,k+1} x_{k+1}^p \right] \\
 &\quad + \sum_{j=0}^k a'_{j,k+1} f(t_j, x(t_j)) + a'_{k+1,k+1} f(t_{k+1}, x_{k+1}^p)
 \end{aligned}$$

The remaining problem is the determination of the predictor formula that is need to calculate the value  $x_{k+1}^p$ .

The idea used to generalize the one-step Adam–Bashforth method is the same as the one described above for the Adams–Moulton technique: the integral on the right-hand side of Eq. (3) is replaced by the product rectangle rule, i.e,

$$\int_{t_0}^{t_{k+1}} (t_{k+1} - u)^{-\alpha} g(u) du \approx \sum_{j=0}^k b_{j,k+1} g(t_j),$$

where

$$b_{j,k+1} = \frac{h^{1-\alpha}}{1-\alpha} [(k-j+1)^{1-\alpha} - (k-j)^{1-\alpha}]$$

for  $\alpha = 0$  we have

$$b_{j,k+1} = h.$$

Thus, the predictor  $x_{k+1}^p$  is determined by the fractional Adams–Bashforth method:

$$x_{k+1}^p = x_0(t_{k+1}) \left( 1 + \frac{t_{k+1}^{1-\alpha}}{\Gamma(2-\alpha)} \right) - \frac{1}{\Gamma(1-\alpha)} \left[ \sum_{j=0}^k b_{j,k+1} x(t_j) \right] + \sum_{j=0}^k b_{j,k+1} f(t_j, x(t_j))$$

### 6. Numerical examples

In this section, we shall study some numerical examples applying Picard, ADM and Predictor–corrector methods and comparing the results.

**Example 1.** Consider the following FDE

$$\frac{dx}{dt} + D^{\frac{1}{2}} x(t) = 1 + \frac{2}{\sqrt{\pi}} t^{\frac{1}{2}} - t^2 + x^2, \quad x(0) = 0 \tag{10}$$

which has the exact solution  $x(t) = t$ .

Applying Picard method to Eq. (10), we get

$$x_n(t) = \left( t + \frac{4}{3\sqrt{\pi}} t^{\frac{3}{2}} - \frac{t^3}{3} \right) - \int_0^t \frac{(t-s)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})} x_{n-1}(s) ds + \int_0^t x_{n-1}^2(s) ds, \quad n = 1, 2, \dots$$

$$x_0(t) = \left( t + \frac{4}{3\sqrt{\pi}} t^{\frac{3}{2}} - \frac{t^3}{3} \right)$$

and the solution will be

$$x(t) = x_n(t).$$

Applying ADM to Eq. (10), we get

$$x_0(t) = \left( t + \frac{4}{3\sqrt{\pi}} t^{\frac{3}{2}} - \frac{t^3}{3} \right)$$

$$x_i(t) = - \int_0^t \frac{(t-s)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})} x_{i-1}(s) ds + \int_0^t A_{i-1}(s) ds, \quad i \geq 1.$$

where  $A_i$  are Adomian polynomials of the nonlinear term  $x^2$ , and the solution will be

$$x(t) = \sum_{i=0}^q x_i(t)$$

Applying PECE to Eq. (10), we have

$$x_0 = 0, \quad f(t, x) = 1 + \frac{2}{\sqrt{\pi}} t^{\frac{1}{2}} - t^2 + x^2$$

Table 1 shows a comparison between the absolute error of Picard (when  $n = 3$ ), ADM solutions (when  $q = 3$ ) and PECE solutions.

**Example 2.** Consider the following FDE

$$\frac{dx}{dt} + D^{\frac{2}{3}} x(t) = 2t - t^4 + \frac{2}{\Gamma(\frac{2}{3})} t^{\frac{4}{3}} + x^2, \quad x(0) = 0 \tag{11}$$

which has the exact solution  $x(t) = t^2$ .

Applying Picard method to Eq. (11), we get

$$x_n(t) = \left( t^2 + \frac{2}{\Gamma(\frac{10}{3})} t^{\frac{7}{3}} - \frac{t^5}{5} \right) - \int_0^t \frac{(t-s)^{-\frac{2}{3}}}{\Gamma(\frac{1}{3})} x_{n-1}(s) ds + \int_0^t x_{n-1}^2(s) ds, \quad n = 1, 2, \dots$$

$$x_0(t) = \left( t^2 + \frac{2}{\Gamma(\frac{10}{3})} t^{\frac{7}{3}} - \frac{t^5}{5} \right)$$

and the solution will be

$$x(t) = x_n(t).$$

Applying ADM to Eq. (11), we get

$$x_0(t) = \left( t^2 + \frac{2}{\Gamma(\frac{10}{3})} t^{\frac{7}{3}} - \frac{t^5}{5} \right)$$

**Table 1** Absolute error.

t	$ x_{Exact} - x_{ADM} $	$ x_{Exact} - x_{Picard} $	$ x_{Exact} - x_{PECE} $
0.1	0.000145565	0.000154652	0.00594708
0.2	0.000866898	0.00107201	0.00575016
0.3	0.00177981	0.00296762	0.00554446
0.4	0.00160474	0.00544699	0.00545929
0.5	0.00114624	0.00769084	0.00548365
0.6	0.00724476	0.0087612	0.00561132
0.7	0.0159463	0.00794191	0.00584349
0.8	0.0246302	0.00503403	0.0061882
0.9	0.0290786	0.000536499	0.00666045
1.0	0.0244311	0.00434315	0.00728311
1.1	0.00666801	0.00788871	0.00808866
1.2	0.0256868	0.00832175	0.00912196
1.3	0.070065	0.00454006	0.0104444
1.4	0.119007	0.00278959	0.0121396
1.5	0.160355	0.0097316	0.0143225
1.6	0.178695	0.00720761	0.171509
1.7	0.158033	0.02101	0.020844
1.8	0.085346	0.0998739	0.0257082
1.9	0.0455844	0.263316	0.0321751
2.0	0.230932	0.552159	0.0408581

**Table 2** Absolute error.

t	$ x_{Exact} - x_{ADM} $	$ x_{Exact} - x_{Picard} $	$ x_{Exact} - x_{PECE} $
0.1	0.0000996616	0.0000999158	0.0038934
0.2	0.000971827	0.000989338	0.00673041
0.3	0.00346046	0.00366554	0.00922231
0.4	0.00772792	0.00887093	0.0114889
0.5	0.0123999	0.0165692	0.0135783
0.6	0.014066	0.0254858	0.0155372
0.7	0.00780759	0.032944	0.0174238
0.8	0.0106784	0.0352666	0.0193149
0.9	0.041401	0.0289394	0.0213133
1.0	0.0760295	0.0124994	0.0235607
1.1	0.0960106	0.0112672	0.0262577
1.2	0.0761537	0.0337261	0.0296985
1.3	0.00391433	0.0409254	0.0343312
1.4	0.141526	0.0188498	0.0408647
1.5	0.293056	0.0321881	0.0504657
1.6	0.369605	0.0682921	0.0651329
1.7	0.261289	0.0428224	0.0884361
1.8	0.103847	0.562204	0.127046
1.9	0.683794	1.86137	0.194076
2.0	1.28107	4.28081	0.316867

$$x_i(t) = - \int_0^t \frac{(t-s)^{-\frac{2}{3}}}{\Gamma(\frac{1}{3})} x_{i-1}(s) ds + \int_0^t A_{i-1}(s) ds, \quad i \geq 1.$$

where  $A_i$  are Adomian polynomials of the nonlinear term  $x^2$ , and the solution will be

Applying PECE to Eq. (11), we have

$$x_0 = 0, \quad f(t, x) = 2t - t^4 + \frac{2}{\Gamma(\frac{7}{3})} t^{\frac{4}{3}} + x^2.$$

Table 2 shows a comparison between the absolute error of Picard (when  $n = 3$ ), ADM solutions (when  $q = 3$ ) and PECE solutions.

## 7. Conclusion

In this paper, we present two analytical methods and numerical method to solve fractional differential equation and the comparison between them in the above two examples. We see from these examples that picard and ADM methods give more accurate solution than PECE in small intervals but PECE gives more accurate solution in large interval.

## Acknowledgment

Thank you for the referees for their efforts.

## References

- [1] K. Abbaoui, Y. Cherruault, Convergence of Adomian's method applied to differential equations, *Comput. Math. Appl.* 28 (1994) 103–109.
- [2] G. Adomian, R. Rach, R. Mayer, Modified decomposition, *J. Math. Comput.* 23 (1992) 17–23.
- [3] G. Adomian, *Solving Frontier Problems of Physics: The Decomposition Method*, Kluwer, Dordrecht, 1995.
- [4] A.M.A. El-Sayed, H.H.G. Hashem, E.A.A. Ziada, Picard and Adomian methods for quadratic integral equation, *Comput. Appl. Math.* 29 (3) (2010) 2576–2580.
- [5] A.M.A. El-Sayed, H.H.G. Hashem, E.A.A. Ziada, Picard and Adomian decomposition methods for quadratic integral equation, *Comput. Appl. Math.* 33 (2014) 95–109, doi:10.1007/s40314-013-0045-3.
- [6] N. Bellomo, D. Sarafyan, On Adomian's decomposition method and some comparisons with Picard's iterative scheme, *J. Math. Anal. Appl.* 123 (1987) 389–400.
- [7] R. Rach, On the Adomian (decomposition) method and comparisons with Picard's method, *J. Math. Anal. Appl.* 128 (2) (1987) 480–483.
- [8] K. Diethelm, An algorithm for the numerical solution for differential equations of fractional order, *Electron. Transact. Numer. Anal.* 5 (1997) 1–6.
- [9] K. Diethelm, A. Freed, On the solution of nonlinear fractional order differential equations used in the modelling of viscoplasticity, in: F. Keil, W. Mackens, H. Vo, J. Werther (Eds.), *Scientific Computing in Chemical Engineering II—Computational Fluid Dynamics, Reaction Engineering, and Molecular Properties*, Springer, Heidelberg, 1999, pp. 217–224.
- [10] K. Diethelm, A. Freed, The FracPECE subroutine for the numerical solution of differential equations of fractional order, in: S. Heinzl, T. Plessner (Eds.), *Forschung und wissenschaftliches Rechnen 1998, Gesellschaft für Wissenschaftliche Datenverarbeitung, Göttingen*, 1999, pp. 57–71.
- [11] K. Diethelm, N.J. Ford, A.D. Freed, A predictor–corrector approach for the numerical solution of fractional differential equations, *Nonlinear Dyn.* 29 (2002) 3–22.
- [12] A.M.A. El-Sayed, A.E.M. El-Mesiry, H.A.A. El-Saka, On the fractional-order logistic equation, *Appl. Math. Lett.* 20 (2007) 817–823.