



## REVIEW ARTICLE

# Education in children's sleep hygiene: which approaches are effective? A systematic review<sup>☆,☆☆</sup>



Camila S.E. Halal<sup>a,b</sup>, Magda L. Nunes<sup>b,\*</sup>

<sup>a</sup> Grupo Hospitalar Conceição, Porto Alegre, RS, Brazil

<sup>b</sup> Faculdade de Medicina, Pontifícia Universidade Católica do Rio Grande do Sul (PUC-RS), Porto Alegre, RS, Brazil

Received 7 May 2014; accepted 14 May 2014

Available online 25 June 2014

### KEYWORDS

Sleep hygiene;  
Sleep education;  
School-aged;  
Child

### Abstract

**Aim:** To analyze the interventions aimed at the practice of sleep hygiene, as well as their applicability and effectiveness in the clinical scenario, so that they may be used by pediatricians and family physicians for parental advice.

**Source of data:** A search of the PubMed database was performed using the following descriptors: *sleep hygiene OR sleep education AND children or school-aged*. In the LILACS and SciELO databases, the descriptors in Portuguese were: *higiene E sono, educação E sono, educação E sono E crianças, e higiene E sono E infância*, with no limitations of the publication period.

**Summary of the findings:** In total, ten articles were reviewed, in which the main objectives were to analyze the effectiveness of behavioral approaches and sleep hygiene techniques on children's sleep quality and parents' quality of life. The techniques used were one or more of the following: positive routines; controlled comforting and gradual extinction or sleep remodeling; as well as written diaries to monitor children's sleep patterns. All of the approaches yielded positive results.

**Conclusions:** Although behavioral approaches to pediatric sleep hygiene are easy to apply and adhere to, there have been very few studies evaluating the effectiveness of the available techniques. This review demonstrated that these methods are effective in providing sleep hygiene for children, thus reflecting on parents' improved quality of life. It is of utmost importance that pediatricians and family physicians are aware of such methods in order to adequately advise patients and their families.

© 2014 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda.

Este é um artigo Open Access sob a licença de [CC BY-NC-ND](http://creativecommons.org/licenses/by-nc-nd/4.0/)

<sup>☆</sup> Please cite this article as: Halal CS, Nunes ML. Education in children's sleep hygiene: which approaches are effective? A systematic review. J Pediatr (Rio J). 2014;90:449–56.

<sup>☆☆</sup> Study conducted at Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, RS, Brazil.

\* Corresponding author.

E-mail: [nunes@puhrs.br](mailto:nunes@puhrs.br), [camilahalal.neuro@hotmail.com](mailto:camilahalal.neuro@hotmail.com) (M.L. Nunes).

**PALAVRAS-CHAVE**

Higiene do sono;  
Educação do sono;  
Infância;  
Criança

## Educação em higiene do sono na infância: quais abordagens são efetivas? Uma revisão sistemática da literatura

**Resumo**

**Objetivo:** Avaliar as intervenções visando práticas de higiene do sono em crianças, sua aplicabilidade e efetividade na prática clínica, para que as mesmas possam ser utilizadas na orientação dos pais pelos pediatras e médicos de família.

**Fonte dos dados:** Foi realizada busca na base de dados da Pubmed utilizando os descritores *sleep hygiene OR sleep education AND child or school-aged*, e nas bases Lilacs e Scielo, com as seguintes palavras-chave: *higiene E sono, educação E sono, educação E sono E crianças, e higiene E sono E infância*, não tendo sido limitado o período de busca.

**Síntese dos dados:** Foram revisados 10 artigos cujos objetivos eram analisar efetividade de abordagens comportamentais e de técnicas de higiene do sono sobre a qualidade do sono das crianças e na qualidade de vida dos pais. Foram utilizadas uma ou mais das seguintes técnicas: rotinas positivas, checagem mínima com extinção sistemática e extinção gradativa ou remodelamento do sono, bem como diários do padrão de sono. Todas as abordagens apresentaram resultados positivos.

**Conclusões:** Apesar de a abordagem comportamental no manejo do sono na faixa etária pediátrica ser de simples execução e adesão, existem poucos estudos na literatura que avaliaram sua efetividade. Os estudos revisados evidenciaram que estas medidas são efetivas na higiene e refletem em melhoria na qualidade de vida dos pais. É de fundamental importância os pediatras e médicos de família conhecerem estas abordagens, para que possam oferecer orientações adequadas a seus pacientes.

© 2014 Sociedade Brasileira de Pediatria. Publicado por Elsevier Editora Ltda.

Este é um artigo Open Access sob a licença de [CC BY-NC-ND](#)

**Introduction**

The prevalence of sleep disorders is high in childhood and may affect up to 30% of school-age children.<sup>1,2</sup> These disorders are important due to the effects they can have not only on the child, but also on their families and society.<sup>3</sup> Thus, a child with chronic sleep disorders may have learning and memory consolidation impairment at school, irritability and mood modulation alterations, difficulty sustaining attention, and behavioral alterations such as aggression, hyperactivity, or impulsivity.<sup>4-7</sup> Furthermore, the chronic sleep deficit lowers the threshold for accidental injury and promotes metabolic changes that, in the long term, can cause other conditions, such as overweight and its consequences.<sup>8,9</sup>

Pediatricians and family physicians play a key role in promoting quality of sleep in children.<sup>10,11</sup> For this purpose, they need to have knowledge of methods of sleep quality promotion, of physiology aspects and age-dependent sleep modifications, and of the importance of good-quality sleep in childhood.<sup>12</sup> A recent study demonstrated that, although 96% of American pediatricians consider it to be their role to advise parents about sleep hygiene methods, only 18% reported having received formal training on the subject.<sup>10</sup>

Moreover, in the United States, the Sleep in America Poll, conducted in 2004, including approximately 1,500 families of children up to 10 years old, demonstrated that only 13% of pediatricians asked parents about possible sleep disorders.<sup>13</sup> A survey conducted in pediatrics residency programs in the United States, Canada, Japan, India, Vietnam, South Korea, Singapore, and Indonesia observed that the mean time spent

in sleep education at the institutions in those countries was two hours during the training period, and that a quarter of the reported programs offered no instruction on the subject.<sup>14</sup>

Sleep disorders are divided into eight different categories, which include insomnia, sleep disordered breathing, hypersomnia of central origin, circadian rhythm disorders, parasomnias sleep-related movement disorders, unresolved issues and isolated symptoms (which include snoring, somniloquy, and benign myoclonus), and other sleep disorders.<sup>15</sup> The latter category includes sleep disorders considered to be physiological or environmental.<sup>16</sup> The environmental disorders, often of behavioral origin, can be prevented if properly managed through sleep hygiene measures.<sup>17</sup>

**Objective**

The objective of this article was to perform a systematic review of interventions aiming at sleep hygiene, and their applicability and effectiveness in pediatric clinical practice, so that they can be used in parental guidance.

**Methods**

Between the months of March and April of 2014, a search was conducted in the PubMed database using the following keywords *sleep hygiene OR sleep education AND children or school-aged*. Review articles were excluded, as well as those including participants aged 10 years or more, or populations

with comorbidities (hospitalization during the study, neurological or respiratory diseases, behavioral or psychiatric disorders).

Studies of children with diagnosis of parasomnias were also excluded from this analysis, as well as those in which the sleep hygiene method was not properly specified in the methods section. Only studies published in Portuguese, English, and Spanish were reviewed. The search totaled 6,621 articles, of which four were excluded as duplicates. Of the remaining 6,617, after reading the titles, 57 were considered possibly relevant and had their abstracts reviewed. Of these, 50 were excluded for not meeting the inclusion criteria, and seven were selected to be read as full texts. Of these, four were considered relevant for this study.<sup>18-21</sup>

The same search, but with descriptors in Portuguese, was conducted in the LILACS and SciELO databases. The keywords used were *higiene E sono, educação E sono, educação E sono E crianças, e higiene E sono E infância*. In the LILACS database, two references were found, neither of which met the scope of this review. In the SciELO database, the descriptors *higiene E sono* resulted in 20 articles, none of which met the criteria for inclusion in this study. In this same database, the descriptors *educação E sono E crianças* resulted in five articles, none of which met the scope of this review.

The analysis of the studies and their references also offered the possibility of access to other publications, and thus a total of six new references were sought and included in this review.<sup>22-27</sup> Fig. 1 shows the process of search, selection, and exclusion of articles present in the current literature.

### Definitions

The diagnosis of sleep disorders requires the presence of specific criteria, which must be present for a specific period of time and have negative consequences on the child and/or parents.<sup>1</sup> Thus, mild or moderate symptoms are not considered disorders, although they can cause some degree of impairment.<sup>28</sup>

The mean latency to sleep onset is usually about 19 minutes in children up to 2 years of age, and 17 minutes from the age of 3 until the beginning of adolescence.<sup>29</sup> In children, insomnia, defined as difficulty initiating or maintaining sleep and comprising a number of sub-classifications, usually falls under the diagnosis of behavioral insomnia.<sup>30</sup> That, in turn, is divided between 1) sleep-onset association type and 2) limit-setting type, or mixed form.

The most common subtype, ‘‘limit-setting type’’, comprises the attempt to postpone going to sleep or refusing to do so, characterized by crying, refusing to stay in bed, or requests for food, drink, or the reading of stories. In this context, it is customary to identify parents with inconsistent routines that tend to give in to their children’s requests.<sup>31</sup> In the ‘‘sleep-onset association’’ subtype, certain behaviors need to be repeated at every waking episode for the child to resume sleep. Thus, a child that associates sleep onset with the presence of one or both parents, food, or being rocked to sleep, when awakening during the night - even with normal awakenings expected for their age group - needs to have the ritual repeated to resume sleep.<sup>32</sup>




Figure 1 Article selection and exclusion process.

Parents' perception of their child having poor quality sleep is directly related to the number of night wakings and to how demanding the child is to initiate sleep at bedtime and retake sleep after night wakings.<sup>29,33</sup> A recent analysis of data from Brazilian birth cohort found, at 12 months, a prevalence of nocturnal awakenings of 64.4% in the two weeks preceding the study, with 56.5% of the children waking up every night, and most of them at least two times each night.<sup>34</sup>

The term "sleep hygiene" includes changes in sleep environment, as well as engaging the child and their parents on routine and practices that encourage sleep of good quality and sufficient duration. It also includes the practice of soothing activities during wakefulness aiming to propitiate sleep onset.<sup>11</sup> The most usual practices are having consistent bedtime and wake-up time for both the nocturnal and daytime sleep (among children at the age group where naps are considered physiological), establishing the appropriate place to initiate sleep, and avoiding environmental and behavioral associations with sleep onset (being rocked to sleep, parents laying on the child's bed until sleep onset, nursing to sleep, watching TV in bed, or drinking beverages rich in caffeine close to bedtime).<sup>17</sup>

Children who need behavioral associations to fall asleep, upon awakening during the night, will need these resources again in order to resume sleep.<sup>1</sup> The passive presence of a parent, however, appears to be positive at some age groups, as well as the use of the child's own resources, such as using a pacifier or thumb-sucking and sleeping with a transitional object.<sup>35</sup>

### Methods of sleep hygiene promotion

The most studied methods are discussed below. These strategies appear to work better in children aged 2 years and older, when a reward system can be used.<sup>2</sup> However, some studies have attempted to advise pregnant women or parents of infants in order to promote problem prevention.<sup>1,33</sup>

**Extinction:** Parents should put the child in bed at a pre-specified time and ignore the child until a certain time on the following morning, while monitoring for the possibility of injury. The method is based on eliminating the acts that reinforce certain behaviors (such as crying on awakening), aiming at their extinction over time.<sup>36</sup> The greatest difficulty in implementing this strategy is the parents' lack of consistency and the parental anxiety that is generated. As a result, some defend the strategy of extinction in the presence of parents, such that parents remain in the room but do not respond to the child's behavior.<sup>2,36</sup>

**Gradual extinction:** In spite of comprising different techniques, the gradual extinction method usually consists of ignoring the demands of the child for specific time periods; these periods are usually determined by the child's age and temperament and the parents' discretion in relation to how long they tolerate their child's crying. Parents should calm the child for short periods, which usually range from 15 seconds to one minute. The technique aims to promote the child's capacity to self-soothe and return to sleep, without undesirable associations or parental interference.<sup>1,2,36</sup>

When assessing 79 children with a mean age of 10.2 months (3-24 months) whose parents were instructed to

implement the gradual extinction technique during nocturnal sleep, Skuladottir et al. observed that the duration of nocturnal sleep increased from 10.27 hours to 10.57 hours ( $p < 0.001$ ) after the intervention, as well as reducing the frequency of nocturnal awakenings (from 4.57 to 1.57 per night,  $p < 0.001$ ).<sup>18</sup>

Eckerberg conducted a study to assess whether recommendations provided only in written form to the parents of children treated at a clinic for sleep disorders would work as well as clinical follow-up, which had been previously advocated.<sup>19</sup> Guidance to parents of children included in the study followed the gradual extinction method, the same provided by the physician during routine consultations. A total of 39 children between 4 and 30 months of age participated in the study, divided into an intervention group (written information sent by email, without contact with the clinician) and a control group (information given by the clinician). After the intervention, children from both groups fell asleep faster ( $p < 0.001$ ) and earlier ( $p < 0.01$ ), which amounted to 30 minutes earlier after a one-month intervention.

In both groups, there was also a significant reduction in nocturnal awakenings (from 4.6 to 4.2 awakenings in the control and from 3.3 to 2.8 in the intervention group,  $p < 0.001$ ) in the two weeks following the intervention. The probability of resume sleep on their own also increased after the intervention (2.1-fold in the control and 2.0-fold in the intervention group,  $p < 0.001$ ). After three months, this decrease continued in both groups, and there was an increase in the duration of nocturnal sleep (by 59 minutes in the control and by 72 minutes in the intervention group) and a decrease in time of wakefulness during the night (from 82 to 18 minutes,  $p < 0.001$ ), with no differences between the groups.

In an Australian case-control study carried out by Hiscock & Wake, 146 children between 7 and 9 months of age were recruited from an outpatient setting.<sup>22</sup> The intervention group received specialized guidance on the physiology of sleep and the application of the gradual extinction method, whereas the control group received a newsletter about normal sleep patterns at the age range of 6-12 months. Two months later, children in the intervention group had resolved more sleep problems than those in the control group ( $p = 0.005$ ), and the remaining problems were less intense in the intervention group. Maternal depressive symptoms decreased in both groups after two months, but decreased more significantly in the intervention group ( $p = 0.02$ ), the group whose mothers also reported that their own sleep was of better quality ( $p = 0.02$ ) at the end of the follow-up.

Aiming to compare the effectiveness of the gradual extinction and extinction methods, and of these in relation to no sleep hygiene method, Reid MJ et al. analyzed 43 children aged 16 to 48 months (14 extinction, 13 gradual extinction, and 16 controls) prior to intervention, and at 21 days and two months after intervention.<sup>23</sup> They observed that families allocated to the extinction group had more difficulty adhering to the method during the second week in relation to the gradual extinction group (interrupting the intervention on average 3.4 times each week, compared to 1.1 times in the other group,  $p = 0.02$ ). During the remaining time, adherence remained high and similar in both groups ( $p < 0.01$ ).

The intervention groups also had better assessments regarding quality at both the moment of sleep onset and sleep maintenance, when compared to the control group. In the subscale regarding quality of sleep in the CBCL (Child Behavior Checklist), both intervention groups also scored better in relation to the control group, and similarly to each other. Two months later, a new evaluation showed that the benefits remained in the groups that underwent interventions.

**Minimal checking with systematic extinction:** Similar to the extinction method, but the child can be checked every 5 to 10 minutes, and the parent can comfort the child quickly when necessary, arranging the covers, and making sure that there are no complications.<sup>2</sup>

Adachi et al. analyzed 99 children taken for childcare consultation at 4 months of age, randomly dividing them between intervention and control groups.<sup>20</sup> The intervention consisted of information about positive routines for initiating sleep, appropriate and inappropriate behaviors to resume sleep at night, and the method of minimal checking with systematic extinction. At the end of the study, the intervention group had a greater decrease in the rates of behaviors listed as "inadequate". The characteristic of "giving food or a diaper change immediately" decreased from 66.7% to 36.4% ( $p=0.001$ ), and the characteristic described as "holding and comforting immediately" decreased from 22.7% to 10.6% ( $p=0.021$ ). In the control group, the number of nocturnal awakenings increased significantly from 53% to 66.7% ( $p=0.022$ ), as did the number of awakenings with crying (from 8.1% to 19.4%,  $p=0.065$ ).

An intervention conducted by Hall et al. included 39 families of infants aged 6 to 12 months, whose parents sought help through a telephone answering service for parents of infants with difficulty sleeping.<sup>21</sup> The aim of the study was to analyze the improvement in the parents' quality of life, and at the end of the intervention, a significant improvement in the parents' quality of sleep was observed, as well as of symptoms of depressed mood and fatigue. The co-sleeping rates also decreased significantly (from 70% practicing them before the intervention to 74% not practicing them after the intervention,  $p<0.001$ ), with no change in the practice of breastfeeding.

**Positive routines:** This method consists of the development of routines preceding bedtime, comprising peaceful and pleasurable activities.<sup>37</sup> Another strategy that can be used is delaying the time to go to bed to ensure that, when lying down, the child will fall asleep quickly, until this habit of falling asleep quickly is consolidated. After that, start anticipating bedtime by 15 to 30 minutes on successive nights until the time considered appropriate is achieved. The child should not sleep during the day, except at the age groups in which daytime sleep is physiological.<sup>2,36</sup>

Positive routines are often used in combination with other methods of sleep hygiene. Adachi et al. included behavioral recommendations in their intervention group. At the end of the intervention, they found that the practice of positive routines characterized by "playing and stimulating during the day" ( $p=0.003$ ), "establishing a place to sleep" ( $p=0.008$ ), and "establishing a timetable to sleep and wake up" ( $p=0.007$ ) had increased significantly.<sup>20</sup>

Mindell et al. performed a study including children in two groups (7-18 months and 18-36 months), in which a routine

preceding sleep was implemented, consisting of a bath followed by massage and quiet activities, with a time period between the bath and turning off the lights of 30 minutes.<sup>24</sup> The routine significantly reduced problem behaviors in both groups, with decreased sleep latency and the number and duration of nocturnal awakenings ( $p<0.001$ ). The mothers in the group of children aged up to 18 months showed reduction in symptoms of stress, depression, anger, fatigue, lack of stamina, and confusion ( $p<0.001$ ) and, in the group older than 18 months, there was an improvement in the symptoms of stress, anger, fatigue, and confusion ( $p<0.001$ ).

This study had a long-term follow up, in which 65% of participants in the group aged up to 18 months were randomized into three groups: one received instructions exclusively via the internet, another received instructions via the internet in addition to those described in the previous study, and a third group was used as control.<sup>25</sup> After one year, the improvements observed in the two groups that received the interventions regarding sleep latency, difficulty falling asleep, number and duration of nocturnal awakenings, period of continuous sleep, and maternal confidence in relation to her child's sleep management persisted ( $p<0.001$ ). After three weeks of intervention, the quality of maternal sleep improved significantly ( $p<0.001$ ); after one year, however, it decreased back to levels close to those at the start of the intervention.

Aiming to compare the effectiveness between positive routines and gradual extinction in reducing temper tantrums with bouts of anger at bedtime, Adams and Rickert followed, for six weeks, two groups of children targeted for each intervention compared to a control group, in a total of 36 children (12 per group) aged between 18 and 48 months.<sup>26</sup> The children in the groups submitted to any of the interventions had significantly fewer bouts of anger, which were of shorter duration than the controls ( $p<0.05$  and  $p<0.001$ , respectively). There was no significant difference in response between the intervention groups, although children in the group submitted to positive routines showed faster favorable responses. Parents in the group allocated to implement positive routines also scored better on the Marital Adjustment Scale at the end of the intervention, which was validated for that population, and investigates the perception that the couple has of their relationship.<sup>38</sup>

**Programmed awakening:** It consists in waking up the child at night, between 15 and 30 minutes before the usual time of spontaneous awakening, and after that, comforting her to return to sleep. The number of programmed awakenings should vary with the usual number of spontaneous awakenings. Over time, it tends to extinguish spontaneous awakenings, and the process of reducing scheduled awakenings begins, resulting in increased sleep consolidation.<sup>1,2,36</sup>

Rickert & Johnson compared the methods of programmed awakening and systematic extinction with a control group in 33 children with a mean age of 20 months (6-54 months), randomly allocating them into three groups of 11 children (programmed awakening, systematic extinction, and a control group).<sup>27</sup> The intervention lasted eight weeks, and parents were re-contacted three and six weeks later. Children who experienced the interventions showed, at the end of the experiment, fewer nocturnal awakenings ( $p<0.05$ ),

**Table 1** Studies reviewed, authors, age range of study, number of participants, objectives, type of intervention, and outcomes.

Author/year of publication	Age range	n	Main objective	Sleep hygiene method	Results
Adachi et al., <sup>20</sup> 2009	4 months	Intervention: 99 Control: 95	- Decrease of inappropriate behavior, nighttime awakenings, and arousals with crying	- Positive routines - Minimal checking with systematic extinction	- Decrease of inappropriate behaviors - Prevention of increased nocturnal awakenings with age
Hall et al., <sup>21</sup> 2006	6 to 12 months	39	- Assessment of parents' quality of life after improvement of the child's sleep quality	- Minimal checking with systematic extinction	- Improvement of sleep quality and depressed mood symptoms of parents - Decrease in co-sleeping
Rickert & Johnson, <sup>27</sup> 1988	6-54 months	33 (11 in each intervention group, 11 in control group)	- Evaluation of the reduction in the number of nocturnal awakenings	- Programmed awakening - Systematic extinction - Control	- Lower number of nighttime awakenings in both groups that underwent interventions
Mindell et al., <sup>24</sup> 2009	7 to 18 months 18 to 36 months	206 (7-18 m) 199 (18-36 m)	- Changes in maternal mood after improved sleep quality of the child	- Positive routines	- Decreased sleep latency in children - Decrease in the duration of nocturnal awakenings - Decrease in depressive symptoms in mothers
Mindell et al., <sup>25</sup> 2011	18 to 48 months	171	- Change in sleep quality of child and mother  - Change in maternal self-confidence	- Positive routines	- Decrease in sleep latency, difficulty falling asleep, number and duration of nocturnal awakenings - Maternal self-confidence improvement - Temporary improvement of the quality of maternal sleep
Skuladottir et al., <sup>18</sup> 2005	3 to 24 months	79	- Changes in the nocturnal sleep pattern with improved daytime sleep pattern	- Positive routines for daytime sleep - Gradual extinction for nighttime sleep - Remodeling for daytime sleep	- Increase in nocturnal sleep - Decrease in nocturnal awakenings
Adams & Rickert, <sup>26</sup> 1989	18 and 48 months	36 (12 in each intervention group; 12 in control group)	- Effect on the number of temper tantrums	- Positive routines - Gradual extinction	- Lower number of temper tantrums and shorter episodes in both intervention groups - Best score in the parents' group positive routines in the Marital Adjustment Scale
Eckerberg, <sup>19</sup> 2002	4 to 30 months	39	- Effectiveness of written information over verbal instructions by the physician	- Gradual extinction	- Decrease in sleep latency in both groups - Decrease of awakenings in both groups - Increased chance to go back to sleep on their own in both groups
Hiscock & Wake, <sup>22</sup> 2002	7 to 9 months	Intervention: 75 Control: 71 (total 146)	- Effectiveness of the advice given by the physician in relation to written advice on sleep quality of the child and maternal depressive symptoms	- Gradual extinction	- Fewer sleep problems in the intervention group - Decrease in maternal depressive symptoms in the intervention group
Reid et al., <sup>23</sup> 1999	16 to 48 months	43 (14 in extinction group, 13 in gradual extinction group, and 16 controls)	- Comparison of the effectiveness of methods of sleep hygiene	- Extinction - Gradual extinction	- More difficulty in compliance in the extinction group during the second week - Improved sleep quality in both intervention groups compared to controls

although this decrease occurred faster in the group submitted to the extinction method. This difference remained statistically significant during the reassessments.

**Sleep remodeling:** Consists of not allowing naps to occur at times that can disrupt nocturnal sleep onset, which comprises four hours before bedtime in children at an age range that allows two naps per day, and six hours before bedtime in children who usually have one nap a day.<sup>18</sup>

The study developed by Skuladottir et al. used this technique for daytime naps.<sup>18</sup> As demonstrated above, they observed positive results regarding the duration of nocturnal sleep.

Table 1 summarizes the studies included in the present review by author, age group and sample size, objectives, type of intervention, and main results.

## Discussion

The number of studies available in the literature on interventions targeting sleep hygiene in children without comorbid conditions is scarce.<sup>18-27</sup> It is noteworthy that no Brazilian studies were found in this search. A recent cross-sectional study performed in the United States, with children aged 5 to 6 years from a low-income community consisting predominantly of Latino families, observed a four-fold higher prevalence of sleep alterations than normally expected at this age, suggesting that unfavorable socioeconomic conditions may contribute to poor sleep quality.<sup>39</sup>

This finding indicates the potential importance of this type of study and subsequent interventions in the Brazilian population, considering that the sleep quality improvement among children of different age ranges and socioeconomic levels could also contribute to the improvement of other indices of quality of life.

Some studies observed that groups of children receiving no intervention also obtained improvements in sleep quality indices at the reassessments. One possible explanation is the existence of an association between neural maturation and physiological circadian mechanisms, which in themselves act as a sleep regulator, improving sleep quality over time.<sup>1</sup> However, it is noteworthy that children who received interventions consistently showed more significant improvements regarding indices of sleep quality. This fact suggests the importance of the external environment on the sleep maturation process.

Information provided only in writing (brochures, newsletters) can be equally effective.<sup>19</sup> This is possibly due to the fact these can be consulted as often as parents deem necessary, and as doubts arise in the implementation of the techniques.

The interventions performed in the reviewed studies are simple and effective, and represent secondary educational measures for parents that do not imply additional cost to them or to the health system, because they basically consist of recommendations. It is possible that these interventions would actually imply cost savings to the system, as well to advised parents, whose children sleep better, and thus would probably have a lower chance of seeking specialized care, in addition to having better performance in their professional activities. The favorable results of all the interventions that objectively sought to analyze mood and quality of life of parents corroborate this hypothesis.

The reviewed studies addressed a broad age range, varying from 3 months to 4 years, mostly comprising children in their first year of life. The importance of this information lies in the fact that interventions in children younger than 2 years also appear to be effective, allowing early changes and preventing children's exposure to long periods of poor sleep.

The role of the health professional who works with children is to know the physiology of sleep and its physiological maturation process. The inclusion of questions about sleep quality and possible sleep impairment factors in the anamnesis, in addition to offering sleep hygiene guidelines regarding the prevention or treatment of pathological behaviors, should be part of every pediatric visit.

## Conflicts of interest

The authors declare no conflicts of interest.

## References

- Mindell JA, Kuhn B, Lewin DS, Meltzer LJ, Sadeh A. Behavioral treatment of bedtime problems and night wakings in infants and young children. *Sleep*. 2006;29:1263-76.
- Nunes ML, Cavalcante V. Avaliação clínica e manejo da insônia em pacientes pediátricos. *J Pediatr (Rio J)*. 2005;81:277-86.
- Karraker KH, Young M. Night waking in 6-month-old infants and maternal depressive symptoms. *J Appl Dev Psychol*. 2007;28:493-8.
- Hale L, Berger LM, LeBourgeois MK, Brooks-Gunn J. A longitudinal study of preschoolers' language-based bedtime routines, sleep duration, and wellbeing. *J Fam Psychol*. 2011;25:423-33.
- Blunden SL, Chapman J, Rigney GA. Are sleep education programs successful? The case for improved and consistent research efforts. *Sleep Med Rev*. 2012;16:355-70.
- Quach J, Gold L, Arnup S, Sia K-L, Wake M, Hiscock H. Sleep well-be well study: improving school transition by improving child sleep: a translational randomised trial. *BMJ Open*. 2013;3:e004009.
- Owens J. Classification and epidemiology of childhood sleep disorders. *Prim Care Clin Office Pract*. 2008;35:533-46.
- Cappuccio FP, Taggart FM, Kandala NB, Currie A, Peile E, Stranges S, et al. Meta-analyses of short sleep duration and obesity in children and adults. *Sleep*. 2008;31:619-26.
- Blunden SL, Chervin RD. Sleep problems are associated with poor outcomes in remedial teaching programmes: A preliminary study. *J Paed Child Health*. 2008;44:237-42.
- Faruqui F, Khubchandani J, Price JH, Bolyard D, Reddy R. Sleep disorders in children: a national assessment of primary care pediatrician practices and perceptions. *Pediatrics*. 2011;128:539-46.
- Gruber R, Cassoff J, Knäuper B. Sleep health education in pediatric community settings: rationale and practical suggestions for incorporating healthy sleep education into pediatric practice. *Pediatr Clin North Am*. 2011;58:735-54.
- Mindell JA, Moline ML, Zendell SM, Brown LW, Fry JM. Pediatricians and sleep disorders: training and practice. *Pediatrics*. 1994;94:194-200.
- National Sleep Foundation. Sleep in America Poll. 2004. [cited on 20 Mar 2014]. Available from: [www.sleepfoundation.org](http://www.sleepfoundation.org)
- Mindell JA, Bartle A, Ahn Y, Ramamurthy MB, Huong HTD, Kohyama J, et al. Sleep education in pediatric residency programs: a cross-cultural look. *BMC Res Notes*. 2013;6:130.
- Oliveiro B, Novelli L. Sleep disorders in children. *Clin Evid*. 2010;09:2304.

16. American Academy of Sleep Medicine. International classification of sleep disorders: diagnostic and coding manual. 2<sup>nd</sup> ed. Westchester: American Academy of Sleep Medicine; 2005.
17. Mindell JA, Meltzer LJ, Carskadon MA, Chervin RD. Developmental aspects of sleep hygiene: Findings from the 2004 National Sleep Foundation Sleep in America Poll. *Sleep Med*. 2009;10:771–9.
18. Skuladottir A, Thome M, Ramel A. Improving day and night sleep problems in infants by changing day time sleep rhythm: a single group before and after study. *Int J Nurs Studies*. 2005;42:843–50.
19. Eckerberg B. Treatment of sleep problems in families with small children: is written information enough? *Acta Paediatr*. 2002;91:952–9.
20. Adachi Y, Sato C, Nishino N, Ohryoji F, Hayama J, Yamagami T. A brief parental education for shaping sleep habits in 4-month-old infants. *Clin Med Res*. 2009;7:85–92.
21. Hall WA, Clauson M, Carty EM, Janssen PA, Saunders RA. Effects on parents of an intervention to resolve infant behavioral sleep problems. *Ped Nurs*. 2006;32:243–50.
22. Hiscock H, Wake M. Randomised controlled trial of behavioural infant sleep intervention to improve infant sleep and maternal mood. *BMJ*. 2002;324:1062–7.
23. Reid MJ, Walter AL, O’Leary SG. Treatment of young children’s bedtime refusal and nighttime wakings: a comparison of “standard” and graduated ignoring procedures. *J Abnorm Child Psychol*. 1999;27:5–16.
24. Mindell JA, Telofski LS, Wiegand B, Kurtz ES. A nightly bedtime routine: impact on sleep in young children and maternal mood. *Sleep*. 2009;32:599–606.
25. Mindell JA, Du Mond CE, Sadeh A, Telofski LS, Kulkarni N, Gunn E. Long-term efficacy of an internet-based intervention for infant and toddler sleep disturbances: one year follow-up. *J Clin Sleep Med*. 2011;7:507–11.
26. Adams LA, Rickert VI. Reducing bedtime tantrums: comparison between positive routines and graduated extinction. *Pediatrics*. 1989;84:756–61.
27. Rickert VI, Johnson CM. Reducing nocturnal awakening and crying episodes in infants and young children: a comparison between scheduled awakenings and systematic ignoring. *Pediatrics*. 1988;81:203–12.
28. Thorpy MJ. Classification of sleep disorders. *Neurotherapeutics*. 2012;9:687–701.
29. Galland BC, Taylor BJ, Elder DE, Herbison P. Normal sleep patterns in infants and children: A systematic review of observational studies. *Sleep Med Rev*. 2012;16:213–22.
30. Morgenthaler TI, Owens J, Alessi C, Boehlecke B, Brown TM, Coleman J, et al. Practice parameters for behavioral treatment of bedtime problems and night wakings in infants and young children. *Sleep*. 2006;29:1277–81.
31. Ward TM, Rankin S, Lee KA. Caring for children with sleep problems. *J Pediatr Nurs*. 2007;22:283–95.
32. Anuntaseree W, Mo-suwan L, Vasiknanonte P, Kuasirikul S, Ma-alee A, Choprapawan C. Night waking in Thai infants at 3 months of age: association between parental practices and infant sleep. *Sleep Med*. 2008;9:564–71.
33. Cook F, Bayer J, Le HN, Mensah F, Cann W, Hiscock H. Baby business: a randomised controlled trial of a universal parenting program that aims to prevent early infant sleep and cry problems and associated parental depression. *BMC Pediatr*. 2012;12:13.
34. Santos IS, Mota DM, Matijasevich A. Epidemiology of co-sleeping and nighttime waking at 12 months in a birth cohort. *J Pediatr (Rio J)*. 2008;84:114–22.
35. Moore M, Meltzer LJ, Mindell JA. Bedtime problems and night wakings in children. *Prim Care Clin Office Pract*. 2008;35:569–81.
36. Kuhn BR, Elliott AJ. Treatment efficacy in behavioral pediatric sleep medicine. *J Psychosom Res*. 2003;54:587–97.
37. Galland BC, Mitchell EA. Helping children sleep. *Arch Dis Child*. 2010;95:850–3.
38. Spanier GB. The measurement of marital quality. *Jour Sex Marit Ther*. 1979;5:288–300.
39. Sheares BG, Kattan M, Leu CS, Lamm CI, Dorsey KB, Evans D. Sleep problems in urban, minority, early-school-aged children more prevalent than previously recognized. *Clin Pediatr (Phila)*. 2013;52:302–9.