



Available online at www.sciencedirect.com

SciVerse ScienceDirect

Genomics Proteomics Bioinformatics 10 (2012) 364–367

---



---

**GENOMICS  
PROTEOMICS &  
BIOINFORMATICS**


---



---

www.elsevier.com/locate/gpb

Application Note

# DNA Barcode *ITS* Effectively Distinguishes the Medicinal Plant *Boerhavia diffusa* from Its Adulterants

Dhivya Selvaraj<sup>1</sup>, Dhivya Shanmughanandhan<sup>1</sup>, Rajeev Kumar Sarma<sup>1</sup>, Jijo C. Joseph<sup>1</sup>,  
Ramachandran V. Srinivasan<sup>2</sup>, Sathishkumar Ramalingam<sup>1,\*</sup>

<sup>1</sup> Plant Genetic Engineering Laboratory, Department of Biotechnology, Bharathiar University, Coimbatore 641046, India

<sup>2</sup> Department of Botany, Bharathiar University, Coimbatore 641046, India

Received 2 September 2011; revised 9 March 2012; accepted 21 March 2012

Available online 29 November 2012

---

## Abstract

*Boerhavia diffusa* (*B. diffusa*), also known as Punarnava, is an indigenous plant in India and an important component in traditional Indian medicine. The accurate identification and collection of this medicinal herb is vital to enhance the drug's efficacy and biosafety. In this study, a DNA barcoding technique has been applied to identify and distinguish *B. diffusa* from its closely-related species. The phylogenetic analysis was carried out for the four species of *Boerhavia* using barcode candidates including nuclear ribosomal DNA regions *ITS*, *ITS1*, *ITS2* and the chloroplast plastid gene *psbA-trnH*. Sequence alignment revealed 26% polymorphic sites in *ITS*, 30% in *ITS1*, 16% in *ITS2* and 6% in *psbA-trnH*, respectively. Additionally, a phylogenetic tree was constructed for 15 species using *ITS* sequences which clearly distinguished *B. diffusa* from the other species. The *ITS1* demonstrates a higher transition/transversion ratio, percentage of variation and pairwise distance which differentiate *B. diffusa* from other species of *Boerhavia*. Our study revealed that *ITS* and *ITS1* could be used as potential candidate regions for identifying *B. diffusa* and for authenticating its herbal products.

**Keywords:** Adulterant; *Boerhavia diffusa*; *ITS*; DNA barcoding; Punarnava

---

## Introduction

*Boerhavia* is one of the highly polymorphic genus in Nyctaginaceae family [1]. About 40 species are distributed in tropical, subtropical and temperate regions. Among these, 6 species are reported in India and *Boerhavia diffusa* (*B. diffusa*) is indigenous [2]. *B. diffusa* is described as Punarnava by an Indian system of medicine, Ayurveda [3]. Roots and whole plants of *B. diffusa* are used in the Ayurvedic and Unani systems of medicine in Arabian countries [4] and many tribal communities in India still use it for the treatment of jaundice and various other liver disorders. It has anti-inflammatory, diuretic, fibrinolytic, anti-convulsant properties [5–8] and also used as carminatives [9–10]. The two pharmaceutically important alkaloids, Punarna-

vine-1 and Punarnavine-2, belonging to the group of quinolizidine were separated from *B. diffusa* [11–12].

*B. diffusa* is known to be extensively adulterated with other species like *Boerhavia erecta*, *Boerhavia repanda*, *Boerhavia coccinea* and *Boerhavia verticillata*. *B. diffusa* have taxonomical conflicts with *B. coccinea*, *Boerhavia repens*, *Boerhavia tetrandra* and *Boerhavia albiflora*, making it difficult to distinguish from one another [13–14]. The species *B. verticillata* display similar morphological features and phytochemical properties with *B. diffusa*, but they differ by their habits [15]. Determination of plant specimens by DNA barcodes will be an effective, reliable and simple pharmacognostic tool to resolve the confusion in morphological identification. Due to different rates of evolution, nuclear ribosomal internal transcribed spacer (*ITS*) regions have become the routine marker in evolutionary studies at different taxonomic levels [16,17]. There is a report using the chloroplast intergenic spacer *psbA-trnH* for identifying

\* Corresponding author.

E-mail: rsathish@buc.edu.in (Ramalingam S).


the *Dendrobium* species of Chinese pharmacopoeia and *psbA-trnH* is recommended as an ideal DNA barcode candidate [18]. Recently the sequence variations are used to develop specific markers for the identification and authentication of drugs and herbal formulations [19].

The objective of the present study is to evaluate an ideal barcode candidate for distinguishing and authenticating the species *B. diffusa* from its common adulterants.

## Results

Genomic DNA was isolated from the species of *B. diffusa*, *B. repanda*, *B. erecta* and *B. verticillata* and used for PCR amplification of the *ITS* and *psbA-trnH*. The obtained sequences were submitted to GenBank. The size and accession number for the gene *ITS* and *psbA-trnH* is shown in Table S1. Additionally, *ITS* sequences from 11 species of *Boerhavia* were taken from the GenBank (Table S2) and used for sequence alignments.

Multiple sequence alignment and pairwise alignment analysis were performed for nuclear *ITS* and chloroplast *psbA-trnH* (Figure S1). The *ITS* region consists of *ITS1*, 5.8S *rDNA* and *ITS2*. The ribosomal sites of 5.8S *rRNA* and 28S *rRNA* are highly conserved. The regions *ITS1* and *ITS2* were compared by multiple sequence alignment, where *ITS1* showed more variation than *ITS2*. Phylogenetic analysis using *ITS1* and *ITS2* indicated *B. diffusa* and *B. erecta* in the same clade while *B. verticillata* and *B. diffusa* was shown in the same clade when using *psbA-trnH* region for phylogenetic analysis (Figure 1). The tree also constructed using *ITS* region clearly distin-



**Figure 1** Phylogenetic trees of the four *Boerhavia* species constructed using *ITS1*, *ITS2* and *psbA-trnH*

Phylogenetic trees were constructed by Minimum Evolution method using *ITS1*, *ITS2* and *psbA-trnH*, respectively, for the four species of *Boerhavia*, including *B. diffusa*, *B. erecta*, *B. repanda* and *B. verticillata*.


guished the morphologically similar species *B. diffusa* from the 14 other species of *Boerhavia* as shown in the Figure 2.

We further analyzed the nucleotide variations of *ITS* and *psbA-trnH* between different species. Percentage of variation shown in Figure 3 indicated that *ITS* demonstrated higher inter-specific divergence. The Wilcoxon rank test indicated significant variation between the species for *ITS1* when compared to *ITS2* and *psbA-trnH*. BLAST 1 and distance method also indicated that *ITS1* showed higher identification percentage at species level (Table 1).

## Discussion


Recent molecular methods like DNA barcoding have been extensively used for species identification, diversity, forensic medicine and ecological studies [20–21]. It also plays an important role in the identification of traditional medicinal herbs. *ITS2* has been effectively used in differentiating morphologically similar species like *Swartzia grandifolia* and *Swartzia longicarpa* and also in solving the controversial species *Caranga rosea* and *Caranga sinica* of the family Fabaceae [22]. Medicinal plant species like the family Polygonaceae [23] and the genus *Dendrobium* [24] have been identified using *ITS2* region. In addition, *ITS1* was used to demonstrate that species of *Amomum villosum* belongs to the family Zingiberaceae [25].

In our study, multiple sequence alignment of *ITS1* and *ITS2* from four *Boerhavia* species showed that *B. diffusa* had a unique basepair variation, which can distinguish it from the other three species, despite the fact that they share many morphological similarities. On the other hand, although *psbA-trnH* distinguishes some species of Polygonaceae [18], less sequence variation in *psbA-trnH* was revealed among the four species of *Boerhavia*. This result is consistent with a previous report that *psbA-trnH* does not show any variation for closely-related Cycad species [26]. Hence, *ITS1* may be a better barcode region for distin-



**Figure 2** Phylogenetic tree of the 15 *Boerhavia* species constructed using *ITS*

Phylogenetic tree was constructed by Minimum Evolution method for the 15 species of *Boerhavia* using *ITS* region.



**Figure 3** Sequence variation of *ITS* and *psbA-trnH* between four *Boerhavia* species

Percentage of nucleotide variations between different *Boerhavia* species, including *B. diffusa*, *B. erecta*, *B. repanda* and *B. verticillata*, was indicated for *ITS* (dark gray) and *psbA-trnH* (light gray), respectively.

**Table 1** Validation of *ITS1*, *ITS2* and *psbA-trnH* from four *Boerhavia* species

Gene name	Correct identification (%)		
	BLAST 1		Distance
	At genus level	At species level	At species level
<i>ITS1</i>	98	94	93.2
<i>ITS2</i>	99	92	91.1
<i>psbA-trnH</i>	94	91	76.86

guishing the species of *Boerhavia*, although *ITS2* has been widely used to distinguish the plant species [27]. This study clearly indicates that DNA barcoding using candidate like *ITS1* is a reliable method for differentiating *B. diffusa* from the other three species, which can also be applied to rapid identification of medicinal plants and their adulterants or substitutes.

## Materials and methods

### Sample collection

Four species of *Boerhavia* (*B. diffusa*, *B. erecta*, *B. repanda*, *B. verticillata*) were collected from the regions of Western Ghats (one of the hotspots), Coimbatore, India. The species were collected and identified by the taxonomist.

### DNA extraction, PCR amplification and DNA sequencing

Fresh leaves from each plant species were used for isolating total genomic DNA by CTAB method [28]. The *ITS* and *psbA-trnH* gene amplifications were performed using Taq DNA polymerase with the primers indicated below. The forward and reverse primers for *ITS* and *psbA-trnH* used were *ITS* F 5'-GGAAGGAGAAGTCGTAACAAGG-3', *ITS* R 5'-TCCTCCGCTTATTGATATGC-3' [28] and

*psbA-trnH* F 5'-GTTATGCATGAACGTAAGCTC-3', *psbA-trnH* R 5'-CGCGCATGGTGGATTCAAATCC-3', respectively. The forward primer of *ITS* region from [29] was modified at the position of 6 (T → G) and 8 (A → G). The PCR program was as follows, an initial denaturation at 94 °C 5 min, followed by 35 cycles of 94 °C 1 min, 57 °C 30 s, 72 °C 1 min and final extension at 72 °C 10 min. PCR products were resolved by gel electrophoresis, purified and subjected to sequencing. The obtained *ITS* and *psbA-trnH* sequences were deposited in the GenBank of NCBI database (GenBank Accessions: HQ386701, HQ386689, HQ386691, HQ386695, HQ386696, HQ407399, HQ386690 and JF423303) (Table S1). Sequences of *ITS* genes from 11 additional species of *Boerhavia* were obtained from GenBank (Table S2).

### Sequence alignment and phylogenetic analysis

The DNA sequences were compared and aligned using the programs ClustalW [30] and MULTALIGN (<http://www.multalin.toulouse.inra.fr/multalin/>). Further, the DNA sequences were subjected to BLAST (<http://www.ncbi.nlm.nih.gov/blast/blast.cgi>) for better identification of sequence at species level. Phylogenetic trees were constructed with the Minimum Evolution method using MEGA 4.0. The intra-specific variation between the species was calculated using MEGA 4.0 [31] and StatsDirect was used to calculate the Wilcoxon signed rank [32].

### Authors' contributions

SR supervised the research. RVS collected and identified the specimens. DS, DS, RKS and JCJ carried out the experimental study. Dhivya Selvaraj prepared the manuscript and SR revised it. All authors read and approved the final manuscript.

### Competing interests

The authors have declared that no competing interests exist.

### Acknowledgements

Dhivya Selvaraj thanks the University Grant Commission-Research Fellowship for Meritorious Students (UGC-RFMS). Sathishkumar Ramalingam thanks the financial support from UGC (Grant No. 34-272/2008(SR)).

### Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.gpb.2012.03.002>.

## References

- [1] Thakur RS, Puri HS, Hussain A. Major medicinal plants of India. Lucknow: Central Institute of Medicinal and Aromatic Plants; 1989, p. 585.
- [2] Chaudhary G, Dantu PK. Morphological, phytochemical and pharmacological, studies on *Boerhavia diffusa* L. J Med Plants Res 2011;5:2125–30.
- [3] Meena AK, Niranjana US, Yadav AK, Ajit K, Singh B, Kiran, et al. A quality assessment of *Boerhavia diffusa* Linn. Commonly known as “Punarnava” plant. Int J Pharmacog Phytochem Res 2010;2:25–8.
- [4] Nalamolu RK, Boini KM, Nammi S. Effect of chronic administration of *Boerhavia diffusa* Linn. Leaf extract on experimental diabetes in rats. Trop J Pharm Res 2007;3:305–9.
- [5] Rawat AKS, Mehrotra S, Tripathi SC, Shome U. Hepatoprotective activity of *Boerhavia diffusa* L. roots—a popular Indian ethnomedicine. J Ethnopharmacol 1997;56:61–6.
- [6] Olaleye MT, Akinmoladun AC, Ogunboye AA, Akindahunsi AA. Antioxidant activity and hepatoprotective property of leaf extracts of *Boerhavia diffusa* Linn against acetaminophen-induced liver damage in rats. Food Chem Toxicol 2010;48:2200–5.
- [7] Mudgal V. Studies on medicinal properties of *Convolvulus pluricaulis* and *Boerhavia diffusa*. Planta Med 1975;28:62–8.
- [8] Satyavati GV, Raina MK, Sharma M. Medicinal plants of India. New Delhi: Indian Council of Medical Research; 1976, p. 10–14.
- [9] Duhan A, Chauhan BM, Punia D. Nutritional value of some non-conventional plant foods of India. Plant Food Hum Nutr 1992;42:193–200.
- [10] Ujowundu CO, Igwe CU, Enemor VHA, Nwaogu LA, Okafor OE. Nutritive and anti-nutritive properties of *Boerhavia diffusa* and *Commelia nudiflora* leaves. Pak J Nutr 2008;7:90–2.
- [11] Nandi RP, Chatterjee SK. Occurrence of Punarnavines in *Boerhavia repens* Linn. Indian J Exp Biol 1974;12:509–11.
- [12] Awasthi LP, Verma HN. *Boerhavia diffusa* – a wild herb with potent biological and antimicrobial properties. Asian Agric Hist 2006;10:55–68.
- [13] Porcher RD. *Boerhavia diffusa* L. (*B. coccinea* Mill) (Nyctaginaceae) in the Carolinas. Castanea 1978;43:172–4.
- [14] Douglas NA, Manos PS. Molecular phylogeny of Nyctaginaceae: taxonomy, biogeography and characters associated with a radiation of xerophytic genera in North America. Am J Bot 2007;94:856–72.
- [15] Bajpai A, Ojha JK. Comparative studies of *Boerhavia diffusa* L. and *Boerhavia verticillata* Poir (Nyctaginaceae). Anc Sci Life 2000;19:105–9.
- [16] Balasubramani SP, Murugan R, Ravikumar K, Venkatasubramanian P. Development of ITS sequence based molecular marker to distinguish, *Tribulus terrestris* L. (Zygophyllaceae) from its adulterant. Fitoterapia 2010;81:503–8.
- [17] Bertini L, Amicucci A, Agostini D, Polidori E, Potenza L, Guidi C, et al. A new pair of primers designed for amplification of the ITS region in Tuber species. FEMS Microbiol Lett 1999;173:239–45.
- [18] Song J, Yao H, Li Y, Li X, Lin Y, Liu C, et al. Authentication of the family Polygonaceae in Chinese pharmacopoeia by DNA barcoding techniques. J Ethnopharmacol 2009;124:434–9.
- [19] Howard C, Bremner PD, Fowler MR, Isodo B, Scott NW, Slater A. Molecular identification of *Hypericum perforatum* by PCR amplification of the ITS and 5.8S rDNA region. Planta Med 2009;75:864–9.
- [20] Pereira F, Carneiro J, Amorim A. Identification of species with DNA-based technology: current progress and challenges. Recent Pat DNA Gene Seq 2008;2:187–99.
- [21] Ferri G, Corradini B, Alu M. Capillary electrophoresis of multigene barcoding chloroplast markers for species identification of botanical trace evidence. Methods Mol Biol 2012;830:253–63.
- [22] Gao T, Yao H, Song J, Liu C, Zhu Y, Ma X, et al. Identification of medicinal plants in the family Fabaceae using a potential DNA barcode ITS2. J Ethnopharmacol 2010;130:116–21.
- [23] Youngbae S, Kim S, Park CW. A phylogenetic study of Polygonum sect. Tovar (Polygonaceae) based on ITS sequences of nuclear ribosomal DNA. J Plant Biol 1977;40:47–52.
- [24] Yao H, Song JY, Ma XY, Liu C, Li Y, Xu HX, et al. Identification of Dendrobium species by a candidate DNA barcode sequence: the chloroplast psbA-trnH intergenic region. Planta Med 2009;75:667–9.
- [25] Qiao C, Han Q, Zhao Z, Wang Z, Xu L, Xu HX. Sequence analysis based on ITS1 region of nuclear ribosomal DNA of *Amomum villosum* and ten species of *Alpinia*. J Food Drug Anal 2009;17:142–5.
- [26] Sass C, Little DP, Stevenson DW, Specht CD. DNA barcoding in the Cycadales: testing the potential of proposed barcoding markers for species identification of cycads. PLoS One 2007;2:e1154.
- [27] Al-Qurainy F, Khan S, Tarroum M, Al-Hemaid FM, Ali MA. Molecular authentication of the medicinal herb *Ruta graveolens* (Rutaceae) and an adulterant using nuclear and chloroplast DNA markers. Genet Mol Res 2011;10:2806–16.
- [28] Khanuja SP, Shasany AK, Darokar MP, Kumar S. Rapid isolation of DNA from dry and fresh samples of plants producing large amounts of secondary metabolites and essential oil. Plant Mol Biol Rep 1999;17:1–7.
- [29] White TJ, Bruns TD, Lee S, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors. PCR protocols: a guide to methods and applications. New York: Academic Press; 1990. p. 315–22.
- [30] Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic Acids Res 1994;22:4673–80.
- [31] Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. Mol Biol Evol 2007;24:1596–9.
- [32] Buchan IE. The development of a statistical computer software resource for medical research. Liverpool, UK: University of Liverpool; 2000.