



Available at

[www.ElsevierMathematics.com](http://www.ElsevierMathematics.com)

POWERED BY SCIENCE @ DIRECT®

Discrete Applied Mathematics 134 (2004) 25–50

DISCRETE  
APPLIED  
MATHEMATICS[www.elsevier.com/locate/dam](http://www.elsevier.com/locate/dam)

# The maximum deviation just-in-time scheduling problem

N. Brauner<sup>a,\*</sup>, Y. Crama<sup>b</sup><sup>a</sup>Laboratoire Leibniz-IMAG, 46 avenue Félix Viallet, Grenoble cedex F-38031, France<sup>b</sup>Ecole d'Administration des Affaires, University of Liège, Bld. du Rectorat 7 (B31), Liège 4000, Belgium

Received 8 June 2001; received in revised form 11 October 2002; accepted 22 October 2002

---

## Abstract

This note revisits the maximum deviation just-in-time (MDJIT) scheduling problem previously investigated by Steiner and Yeomans. Its main result is a set of algebraic necessary and sufficient conditions for the existence of a MDJIT schedule with a given objective function value. These conditions are used to provide a finer analysis of the complexity of the MDJIT problem. The note also investigates various special cases of the MDJIT problem and suggests several questions for further investigation.

© 2003 Elsevier B.V. All rights reserved.

*Keywords:* Scheduling; Sequencing; Just-in-time; High multiplicity; Convex graphs; Balanced words

---

## 1. Introduction

This paper revisits an optimization model originally motivated by scheduling issues arising in just-in-time (JIT) manufacturing environments. This model, to be called the *maximum deviation just-in-time problem (MDJIT problem)*, has been analyzed in [25]. The authors gave a very interesting, quite complete analysis of the problem, including structural and algorithmic results. We believe, however, that several intriguing complexity issues are still open with respect to this problem (see [14] for additional open questions). The main purpose of the present note is to clarify some of these issues. On our way, we propose alternative proofs for some of the results stated in [25], we establish several new results and we propose a number of conjectures.

---

\* Corresponding author.

*E-mail addresses:* [nadia.brauner@imag.fr](mailto:nadia.brauner@imag.fr) (N. Brauner), [y.crama@ulg.ac.be](mailto:y.crama@ulg.ac.be) (Y. Crama).

Section 2 contains a more precise description of JIT scheduling problems and of related computational complexity issues. In Section 3, we focus on the maximum deviation JIT problem. We recast it as a matching problem in a bipartite graph and we derive necessary and sufficient conditions for the existence of a schedule with a given objective function value. In Section 4, we use the previous result to establish that the MDJIT problem is in Co-NP and to prove that the problem can be solved in polynomial time when the number of part types is fixed. Section 5 establishes lower and upper bounds on the optimal value of the MDJIT problem. Section 6 proposes some results and conjectures concerning the structure of instances with small deviation. Finally, Section 7 provides a complete solution of the MDJIT problem for the special case where there are only two distinct part types.

## 2. Just-in-time scheduling problems

### 2.1. Position of the problem

An instance of a generic just-in-time scheduling problem consists of a number  $n$  of different part types and of the demand  $d_i \in \mathbb{N}$  for part type  $i$  ( $i = 1, 2, \dots, n$ ). All part types are produced on the same equipment (typically, a mixed-model assembly line) and the production of each part requires one unit of time. We denote by  $D = \sum_{i=1}^n d_i$  the total demand and by  $r_i = d_i/D$  the *ideal production rate* for parts of type  $i$ . The term “ideal” refers here to the fact that, at each instant, we would like the line to have assembled part type  $i$  in proportion  $r_i$ . Such a schedule would be uniformly “leveled”. Obviously, perfectly leveled schedules are never attainable, but the aim of JIT control systems is to keep the actual production of each part as close as possible to its “ideal rate”. Monden [23] states that this is a main objective of Toyota’s JIT systems.

We can formulate this generic question as an optimization problem of the form [22,17,15]:

$$\text{minimize } \begin{cases} \max_{k,i} & \leftarrow \text{Maximum deviation} \\ \sum_k \sum_i & \leftarrow \text{Total deviation} \end{cases} F^i(x_{ik} - kr_i)$$

subject to

$$\sum_{i=1}^n x_{i,k} = k, \quad k = 1, \dots, D, \tag{1}$$

$$x_{i,D} = d_i, \quad i = 1, \dots, n, \tag{2}$$

$$0 \leq x_{i,k} - x_{i,k-1}, \quad i = 1, \dots, n; \quad k = 2, \dots, D, \tag{3}$$

$$x_{i,k} \in \mathbb{N}, \quad i = 1, \dots, n; \quad k = 1, \dots, D. \tag{4}$$

In this formulation,  $x_{i,k} = q$  if  $q$  parts of type  $i$  are produced in the first  $k$  periods. So, Eq. (1) means that  $k$  parts have to be produced during the first  $k$  periods, and Eq. (2) translates the demand constraints. Inequality (3) indicates that the number of parts of type  $i$  should not decrease with time. Furthermore, for each  $i$ , the function  $F^i$  is a nonnegative convex function such that  $F^i(x) = 0$  if and only if  $x = 0$ . Its interpretation is that  $F^i(x_{ik} - kr_i)$  penalizes the deviation between the actual production  $x_{ik}$  and the target  $kr_i$ .

The above formulation emphasizes the distinctive number-theoretic flavor of JIT scheduling problems: given  $n$  rational numbers  $r_1, \dots, r_n$  with common denominator  $D$ , the problem is to find  $nD$  integers  $x_{ik}$  which ‘optimally’ approximate the sequence  $(kr_i)$  under the ‘cardinality’ and ‘monotonicity’ restrictions (1)–(4). Diophantine approximation problems of a similar nature are investigated for instance in [11].

## 2.2. Literature review

We now briefly review some of the main results concerning the above JIT models. For more comprehensive surveys, we refer to [15,14].

The total deviation model has been proposed in [22], together with some heuristics for its resolution. Kubiak and Sethi [17,18] reformulated this model as an assignment problem. Their approach leads to an algorithm whose complexity is polynomial in  $n$  and  $D$ . Inman and Bulfin [13] also considered the total deviation objective with a slightly different penalty function. They described a heuristic which runs in time  $O(nD)$ . The heuristic is based on a reduction to the one-machine scheduling problem with penalties for earliness and tardiness.

The maximum deviation problem has been investigated in [25], with  $F^i(x) = |x|$ . The authors reduced the JIT problem to a one-machine scheduling problem with release dates and due dates, which they solved by an exact pseudo-polynomial algorithm with complexity  $O(D \log D)$ . This algorithm will be reviewed in Section 3.

Some of the connections between the above models are examined in [14]. For the sake of completeness, let us also mention that multilevel extensions of the basic model have been investigated in the literature. Kubiak [15] and Kubiak et al. [19] establish that several of these extensions (including total and maximum deviation objectives) are NP-hard.

Several authors have also noted the connections between JIT scheduling problems and apportionment problems, i.e. problems dealing with the allocation of seats of a legislature among the states or provinces of a nation, in close proportion to their respective populations; see e.g. [4,2]. From this connection, and from known results concerning apportionment problems, it is not too difficult to deduce that simple-minded procedures of a greedy nature do not provide an optimal solution of JIT scheduling problems (see the Appendix).

## 2.3. The maximum deviation JIT problem: complexity issues

Our main goal in this paper is to initiate further investigations into the complexity of JIT scheduling problems. We shall concentrate on (what seems to be) one of the

simplest variants of the problem, namely the maximum deviation (MDJIT) problem with  $F^i(x) = |x|$  previously investigated in [25].

The input of the generic JIT scheduling problem is essentially the list of integers  $d_1, d_2, \dots, d_n$ , so that its input size is  $O(\sum_{1 \leq i \leq n} \log d_i) = O(n \log D)$ . Hence, an algorithm which is polynomial in  $n$  and  $D$  is only *pseudo-polynomial*, but not polynomial in the input size. This limitation has already been noted by Kubiak [15], who also mentioned in his survey that the question of the exact complexity of JIT scheduling problems remains open. Similar issues actually arise for a larger class of so-called *high multiplicity* optimization problems—see for instance [24,12,6].

The recognition version of this problem can be stated as follows:

MDJIT:

*Input:*

- $n \in \mathbb{N}$ : number of part types;
- $d_i \in \mathbb{N}$ : demand for part type  $i$ ,  $i = 1, 2, \dots, n$ ;
- $B \in \mathbb{Q}$ : a bound.

*Question:* Does there exist an  $n \times D$  matrix  $X = (x_{i,k})$  such that:

$$\max_{1 \leq i \leq n, 1 \leq k \leq D} \left| x_{ik} - k \frac{d_i}{D} \right| \leq B, \quad (5)$$

$$\sum_{i=1}^n x_{i,k} = k, \quad k = 1, \dots, D, \quad (6)$$

$$x_{i,D} = d_i, \quad i = 1, \dots, n, \quad (7)$$

$$0 \leq x_{i,k} - x_{i,k-1}, \quad i = 1, \dots, n; \quad k = 2, \dots, D, \quad (8)$$

$$x_{i,k} \in \mathbb{N}, \quad i = 1, \dots, n; \quad k = 1, \dots, D. \quad (9)$$

It is actually interesting to observe that even this simple formulation of the problem is pseudo-polynomial in size, since it involves  $nD$  variables and  $O(nD)$  constraints. So, obtaining truly polynomial algorithms for JIT scheduling problems is far from trivial (if possible at all) and requires deep insight into the structural properties of the problems. In particular, it is not even obvious whether MDJIT is in NP (or in Co-NP), i.e. whether there exists a polynomial-size certificate for every “Yes” (or “No”) instance of MDJIT. We shall come back to this issue in Section 4.

#### 2.4. Notations

We shall use the following notations:

- $[x]$  is the rounding of  $x$  to the closest integer; when the fractional part of  $x$  is equal to  $1/2$ , we round downward unless otherwise specified; so,  $x - \frac{1}{2} \leq [x] < x + \frac{1}{2}$ ;
- $\lfloor x \rfloor$  is the largest integer smaller than  $x$ :  $x - 1 < \lfloor x \rfloor \leq x$ ;
- $\lceil x \rceil$  is the smallest integer larger than  $x$ :  $x \leq \lceil x \rceil < x + 1$ ;
- $[x_1 .. x_2]$  is the set of all integers between  $x_1$  and  $x_2$ .

### 3. Maximum deviation just-in-time problem

In this section, we establish several structural properties of the MDJIT problem. We consider the recognition version (5)–(9) of the problem for a fixed value of  $B$ . These results constitute an alternative approach to Steiner and Yeomans [25]. They lead to different insights and, in some cases, complete the arguments provided by these authors.

For  $i = 1, \dots, n$  and  $j = 1, \dots, d_i$ , we use the shorthand  $(i, j)$  to denote the  $j$ th part of type  $i$ . A *schedule* for the MDJIT problem is a bijection  $\{(i, j) | i = 1, \dots, n, j = 1, \dots, d_i\} \rightarrow [1 .. D]$ . We say that a schedule is *feasible* if inequality (5) holds for the implied variables  $X = (x_{i,k})$ .

The following statement is essentially due to Steiner and Yeomans [25]:

**Proposition 1.** *Consider an instance  $(n, d_1, d_2, \dots, d_n, B)$  of the MDJIT problem. A schedule for MDJIT is feasible if and only if, for all  $i = 1, \dots, n$  and  $j = 1, \dots, d_i$ , this schedule assigns part  $(i, j)$  to the interval  $[E(i, j) .. L(i, j)]$  where*

$$E(i, j) = \left\lceil \frac{j - B}{r_i} \right\rceil,$$

$$L(i, j) = \left\lfloor \frac{j - 1 + B}{r_i} + 1 \right\rfloor.$$

**Proof.** (a) Consider any feasible schedule and suppose that part  $(i, j)$  is produced at time  $k < (j - B)/r_i$  in this schedule. Then,  $x_{i,k} = j$  and

$$|x_{i,k} - kr_i| \geq x_{i,k} - kr_i > j - j + B = B,$$

contradicting feasibility. Therefore, part  $(i, j)$  cannot start before time  $(j - B)/r_i$ , and hence before time  $E(i, j) = \lceil (j - B)/r_i \rceil$ .

Similarly, assume that part  $(i, j)$  starts at time  $k > (j - 1 + B)/r_i + 1$ . Then, at time  $k - 1$ , there holds  $x_{i,k-1} = j - 1$  and

$$|x_{i,k-1} - (k - 1)r_i| \geq (k - 1)r_i - x_{i,k-1} > j - 1 + B - j + 1 = B,$$

a contradiction. This shows that part  $(i, j)$  cannot start after time  $L(i, j) = \lfloor (j - 1 + B)/r_i + 1 \rfloor$ . Hence, the necessary condition holds.

(b) Let us show that, if each part  $(i, j)$  is assigned to some time period in

$$[E(i, j) .. L(i, j)] \cap [1 .. D],$$

and all parts are assigned to different time periods, then the resulting schedule is feasible.

Consider a fixed part type  $i$  and a time period  $k$ . Let  $j \in \{1, 2, \dots, d_i\}$  be the number of parts of type  $i$  which have been produced up to (and including) time  $k$ , i.e.  $x_{i,k} = j$ . We must show that  $|j - kr_i| \leq B$ .

On one hand,  $k \geq E(i, j)$ . Therefore,

$$j - kr_i \leq j - E(i, j)r_i \leq j - \left( \frac{j - B}{r_i} \right) r_i = B. \tag{10}$$

If  $j = d_i$ , then  $j - kr_i = d_i - kr_i \geq 0$ , so (10) implies that  $|j - kr_i| \leq B$  as required.




Fig. 1. Convex bipartite graph for  $d_1 = d_2 = 3$ ,  $d_3 = 1$  and  $B = \frac{5}{7}$ .

On the other hand, if  $j < d_i$ , then  $k < L(i, j + 1)$  since the  $(j + 1)$ st part of type  $i$  is produced after time  $k$ . Thus,

$$kr_i - j \leq (L(i, j + 1) - 1)r_i - j \leq \left\lfloor \frac{j + B}{r_i} \right\rfloor r_i - j \leq j + B - j = B. \quad (11)$$

So,  $|j - kr_i| \leq B$  follows from (10) and (11).  $\square$

Steiner and Yeomans [25] rely on Proposition 1 to solve the MDJIT problem. First, they set up a bipartite graph  $G = (\mathcal{V}_1 \cup \mathcal{V}_2, E)$ , where  $\mathcal{V}_1 = [1..D]$ ,  $\mathcal{V}_2$  is the set of all parts  $(i, j)$ , for  $i = 1, \dots, n$ ,  $j = 1, \dots, d_i$ , and  $(k, (i, j))$  is an edge in  $E$  if and only if  $k \in [E(i, j)..L(i, j)]$ . For any subset  $X$  of vertices, denote by  $N(X)$  the neighborhood of  $X$ , i.e. the set of all vertices adjacent to at least one vertex in  $X$ . Observe that the neighborhood of every vertex  $(i, j) \in \mathcal{V}_2$  is an interval (namely, the interval  $[E(i, j)..L(i, j)]$ ). Therefore, in agreement with Glover's terminology [10], we say that the graph  $G$  is  $\mathcal{V}_1$ -convex.

Let us illustrate this concept on an example. Consider the instance  $n=3$ ,  $d_1 = d_2 = 3$ ,  $d_3 = 1$  and  $B = \frac{5}{7}$ . The convex bipartite graph  $G$  associated with this instance is represented in Fig. 1. Observe that this instance is feasible, as it admits the feasible production sequence  $(1, 2, 1, 3, 2, 1, 2)$ . Furthermore, this feasible sequence corresponds in a natural way to the perfect matching of  $G$  indicated by the thick edges in Fig. 1. This is no mere coincidence. Indeed, the following proposition is a simple corollary of Proposition 1 (it is implicit in [25]).

**Proposition 2.** *The MDJIT problem has a feasible solution if and only if the graph  $G$  has a perfect matching.*

**Proof.** Any solution of the MDJIT problem defines a perfect matching in  $G$ . More precisely, any such solution corresponds to an *order preserving* matching, i.e. a perfect matching such that, when  $j_1 < j_2$ , part  $(i, j_1)$  is matched to an earlier instant than part  $(i, j_2)$ .

Conversely, if  $G$  has a perfect matching  $M$ , then it necessarily has an order preserving one, which corresponds therefore to a feasible solution of MDJIT. Indeed, if  $(i, j_1)$  is matched to  $k_1$  and  $(i, j_2)$  is matched to  $k_2$  in  $M$ , where  $k_1 > k_2$ , then matching  $(i, j_1)$  to  $k_2$  and  $(i, j_2)$  to  $k_1$  is also feasible (this is due to the convexity of  $G$  and to the fact that both  $E(i, j)$  and  $L(i, j)$  are nondecreasing in  $j$ ).  $\square$

In view of the fact that  $G$  is convex, Steiner and Yeomans [25] suggest to use Glover’s *Earliest Due Date* algorithm [10] to check the existence of a perfect matching in  $G$ . The algorithm runs through the time instants  $k=1,2,\dots,D$ , in order, and assigns to  $k$  the part  $(i,j)$  with smallest value of  $L(i,j)$  among all the available parts such that  $(k,(i,j))\in E$  (see also [21,9,8] for successive improvements in the efficiency of Glover’s algorithm).

We now proceed to develop necessary and sufficient conditions for the existence of a perfect matching in  $G$ . These conditions build on Hall’s theorem, which implies that a bipartite graph  $H=(V_1\cup V_2,E)$  with  $|V_1|=|V_2|$  has a perfect matching if and only if

$$|N(X)|\geq |X| \quad \text{for all } X\subseteq V_1 \tag{12}$$

(see e.g. [5]).

We first show that, for convex graphs, Hall’s conditions can be restricted to bear on special subsets of vertices. Let us define the following sets. First,

$$\mathcal{I}_1 = \{I: I \text{ is an interval of } \mathcal{V}_1\}.$$

Then, for each  $I\in\mathcal{I}_1$ , let  $U(I)$  be the largest subset of  $\mathcal{V}_2$  whose neighborhood is completely contained in  $I$ , i.e.

$$U(I) = \{v\in\mathcal{V}_2: N(v)\subseteq I\}$$

(in terms of the original MDJIT problem,  $U(I)$  is the set of all parts which must be processed during the time interval  $I$ ). Finally, let

$$\mathcal{I}_2 = \{U(I): I\in\mathcal{I}_1\}.$$

**Proposition 3.** *A  $\mathcal{V}_1$ -convex bipartite graph  $G=(\mathcal{V}_1\cup\mathcal{V}_2,E)$  has a perfect matching if and only if*

$$\forall X\in\mathcal{I}_1\cup\mathcal{I}_2, \quad |N(X)|\geq |X|. \tag{13}$$

**Proof.** The necessity of condition (13) is a consequence of Hall’s theorem.

To establish sufficiency, suppose now that condition (13) is verified and that there exists  $X_1\subseteq\mathcal{V}_1$  such that  $|N(X_1)|<|X_1|$ . The set  $X_1$  is a union of disjoint intervals:

$$X_1 = I_1\cup I_2\cup\dots\cup I_p,$$

where  $I_i=[x_i..x'_i]$  and  $x'_i<x_{i+1}-1$  ( $i=1,\dots,p-1$ ). We assume that  $X_1$  is chosen so that  $p$  is minimal.

*Case 1:* Assume that  $N(I_i)\cap N(I_j)=\emptyset$  for all  $i,j\in\{1,\dots,p\}$  with  $i\neq j$ .

This implies that  $|N(X_1)|=\sum_{i=1}^p|N(I_i)|$ . But since  $I_i$  is an interval, condition (13) implies that  $|N(I_i)|\geq|I_i|$  for  $i=1,\dots,p$ . Therefore,

$$|N(X_1)|=\sum|N(I_i)|\geq\sum|I_i|=|X_1|,$$

which is in contradiction with the hypothesis on  $X_1$ .




Fig. 2. Convex bipartite graph for  $d_1 = d_2 = 3$ ,  $d_3 = 1$  and  $B = \frac{4}{7}$ .

*Case 2:* There exist  $i$  and  $j$  ( $i < j$ ) such that  $N(I_i) \cap N(I_j) \neq \emptyset$ .

Let  $u \in N(I_i) \cap N(I_j)$ . Since  $G$  is  $\mathcal{V}_1$ -convex,  $u \in N(I_i) \cap N(I_{i+1})$ . Therefore, we can suppose that  $j = i + 1$ . Let  $I$  be the interval nested between  $I_i$  and  $I_{i+1}$ , i.e.  $I = [x'_i + 1 .. x_{i+1} - 1]$ . Note that  $X_1 \cap I = \emptyset$ .

Let  $U = U(I)$  be the set of vertices of  $\mathcal{V}_2$  whose neighborhood is included in  $I$ . Since  $U \in \mathcal{S}_2$ , condition (13) implies that  $|U| \leq |N(U)|$ . Moreover,  $|N(U)| \leq |I|$  (by definition of  $U$ ), so that  $|U| \leq |I|$ . By definition of  $U$ , one has  $U \cap N(X_1) = \emptyset$  and by convexity of the graph one has  $N(I) \subseteq U \cup N(X_1)$ . Therefore,

$$|N(I) \cup N(X_1)| \leq |U \cup N(X_1)| = |U| + |N(X_1)| < |I| + |X_1| = |I \cup X_1|.$$

This implies that  $|N(X_1 \cup I)| < |X_1 \cup I|$ . Let  $Y = X_1 \cup I$ . The set  $Y$  is a union of  $p - 1$  disjoint intervals of  $\mathcal{V}_1$  and satisfies  $|N(Y)| < |Y|$ . This is in contradiction with the minimality of  $p$ .

Therefore, condition (13) implies Hall's condition (12), and thus (13) implies the existence of a perfect matching in  $G$ .  $\square$

One may be tempted to conjecture that the conditions on  $\mathcal{S}_2$  are superfluous in Proposition 3 and that Hall's conditions on intervals of  $\mathcal{V}_1$  would be sufficient, by themselves, to ensure the existence of a perfect matching in a  $\mathcal{V}_1$ -convex bipartite graph. But this conjecture actually fails even in the special case where the convex graph is associated with an instance of the MDJIT problem, as illustrated by the graph in Fig. 2. Indeed, this convex graph (which corresponds to the instance  $d_1 = d_2 = 3$ ,  $d_3 = 1$ ,  $B = \frac{4}{7}$ ) satisfies the conditions

$$\forall X \in \mathcal{S}_1, \quad |N(X)| \geq |X|,$$

but does not have a perfect matching.

On the other hand, conditions (13) can be strengthened by restricting them to those sets of  $\mathcal{S}_2$  whose neighborhood is an interval. Define

$$\mathcal{S}'_2 = \{U \in \mathcal{S}_2: N(U) \in \mathcal{S}_1\}.$$

Then, we can prove:

**Proposition 4.** *Conditions (14) and (15) are equivalent:*

$$\forall X \in \mathcal{S}_2, \quad |N(X)| \geq |X|, \tag{14}$$

$$\forall X \in \mathcal{S}'_2, \quad |N(X)| \geq |X|. \tag{15}$$



**Proof.** Since,  $\mathcal{S}'_2 \subseteq \mathcal{S}_2$ , (14) trivially implies (15).

Conversely, suppose that (15) holds and let  $U \in \mathcal{S}_2$ . If  $N(U)$  is an interval, then  $U \in \mathcal{S}'_2$  and  $|N(U)| \geq |U|$ . So, suppose that  $N(U)$  is a union of disjoint intervals:

$$N(U) = [x_1 \dots x'_1] \cup \dots \cup [x_p \dots x'_p]$$

with  $x'_i < x_{i+1} - 1$  ( $i = 1, \dots, p - 1$ ). Let  $Y_i = \{v \in \mathcal{V}_2: N(v) \subseteq [x_i \dots x'_i]\}$ . By definition,  $\bigcup_{i=1}^p Y_i \subseteq U$ . Suppose that there exists  $k \in U$  such that  $k \notin \bigcup_{i=1}^p Y_i$ . This would mean that  $k$  has neighbors in at least two distinct intervals  $[x_i \dots x'_i]$  and  $[x_j \dots x'_j]$ , with  $i < j$ . Since the graph is  $\mathcal{V}_1$ -convex, this would imply that all vertices between  $x'_i$  and  $x_j$  are in  $N(U)$ , a contradiction. Therefore  $\bigcup_{i=1}^p Y_i = U$ .

Since every element of  $[x_i \dots x'_i]$  is the neighbor of some element of  $U$ , there follows immediately that  $N(Y_i) = [x_i \dots x'_i]$  and hence  $Y_i \in \mathcal{S}'_2$ , for  $i = 1, \dots, p$ . Therefore,

$$|N(U)| = \sum_{i=1}^p |N(Y_i)| \geq \sum_{i=1}^p |Y_i| = |U|$$

and we conclude that (15) implies (14).  $\square$

**Remark 1.** From the above results, one easily concludes that, in the case of convex graphs, Hall’s conditions need only to be applied to those sets  $X$  such that  $X$  is either an interval in  $\mathcal{V}_1$ , or the neighborhood of an interval in  $\mathcal{V}_1$ . Although we shall not use it in this form, this compact statement may be of independent interest.

We are now in a position to apply Hall’s conditions to the convex bipartite graph associated with an instance of the MDJIT problem.

**Theorem 1.** *The MDJIT problem has a feasible solution if and only if, for all  $x_1, x_2$  in  $[1 \dots D]$  with  $x_1 \leq x_2$ , the following inequalities are both valid:*

$$\sum_{i=1}^n \max(0, \lfloor x_2 r_i + B \rfloor - \lceil (x_1 - 1)r_i - B \rceil) \geq x_2 - x_1 + 1, \tag{16}$$

$$\sum_{i=1}^n \max(0, \lceil x_2 r_i - B \rceil - \lfloor (x_1 - 1)r_i + B \rfloor) \leq x_2 - x_1 + 1. \tag{17}$$

**Proof.** We know that the MDJIT problem has a feasible solution if and only if  $G$  has a perfect matching. This is equivalent to

$$\forall X \in \mathcal{S}_1 \cup \mathcal{S}'_2, \quad |N(X)| \geq |X|. \tag{18}$$

We want to express these conditions algebraically.

(a) Let  $X = [x_1 \dots x_2]$  be an interval of  $\mathcal{V}_1$ . Part  $(i, j)$  is in the neighborhood of  $X$  if and only if

$$\begin{aligned} E(i, j) \leq x_2 \text{ and } L(i, j) \geq x_1 \\ \Leftrightarrow \frac{j - B}{r_i} \leq x_2 \text{ and } \frac{j - 1 + B}{r_i} + 1 \geq x_1 \\ \Leftrightarrow (x_1 - 1)r_i + 1 - B \leq j \leq x_2 r_i + B. \end{aligned}$$

For a given part type  $i \in \{1, \dots, n\}$ , the number of copies  $j$  which verify the previous inequality is

$$\max(0, \lfloor x_2 r_i + B \rfloor - \lceil (x_1 - 1)r_i + 1 - B \rceil + 1).$$

Therefore, for  $X = [x_1 \dots x_2] \in \mathcal{S}_1$ ,  $|N(X)| \geq |X|$  if and only if

$$\sum_{i=1}^n \max(0, \lfloor x_2 r_i + B \rfloor - \lceil (x_1 - 1)r_i - B \rceil) \geq x_2 - x_1 + 1.$$

(b) Given  $x_1, x_2$  in  $\mathcal{V}_1$ , with  $x_1 \leq x_2$ , and given  $(i, j) \in \mathcal{V}_2$ , the following equivalences hold:

$$\begin{aligned} [E(i, j) \dots L(i, j)] &\subseteq [x_1 \dots x_2] \\ \Leftrightarrow E(i, j) &\geq x_1 \text{ and } L(i, j) \leq x_2 \\ \Leftrightarrow E(i, j) &> x_1 - 1 \text{ and } L(i, j) < x_2 + 1 \\ \Leftrightarrow \frac{j - B}{r_i} &> x_1 - 1 \text{ and } \frac{j - 1 + B}{r_i} + 1 < x_2 + 1 \\ \Leftrightarrow (x_1 - 1)r_i + B &< j < x_2 r_i + 1 - B \\ \Leftrightarrow \lfloor (x_1 - 1)r_i + B + 1 \rfloor &\leq j \leq \lceil x_2 r_i - B \rceil. \end{aligned}$$

For a given part type  $i$ , the number of  $j$  which verify the previous equivalences is

$$\max(0, \lceil x_2 r_i - B \rceil - \lfloor (x_1 - 1)r_i + B \rfloor).$$

Thus, the cardinality of  $U([x_1 \dots x_2]) = \{(i, j) \in \mathcal{V}_2 : [E(i, j) \dots L(i, j)] \subseteq [x_1 \dots x_2]\}$  can be computed as

$$|U([x_1 \dots x_2])| = \sum_{i=1}^n \max(0, \lceil x_2 r_i - B \rceil - \lfloor (x_1 - 1)r_i + B \rfloor). \quad (19)$$

(c) Assume now that inequalities (17) hold for all values of  $x_1 \leq x_2$  and consider  $X \in \mathcal{S}'_2$ . By definition,  $N(X)$  is an interval of  $\mathcal{V}_1$ , say  $N(X) = [x_1 \dots x_2]$ . So,  $X \subseteq U([x_1 \dots x_2])$ . In view of (19) and (17), we conclude that

$$|X| \leq |U([x_1 \dots x_2])| \leq x_2 - x_1 + 1 = |N(X)|$$

as required by (18).

(d) Conversely, assume next that  $|X| \leq |N(X)|$  for all  $X \in \mathcal{S}'_2$ . Then, we know from Proposition 4 that  $|X| \leq |N(X)|$  for all  $X \in \mathcal{S}_2$ . Consider now  $x_1, x_2$  in  $\mathcal{V}_1$  with  $x_1 \leq x_2$ . Let  $I = [x_1 \dots x_2]$  and let  $U = U(I)$ . By definition of  $U$ ,  $N(U) \subseteq I$ , and by definition of  $\mathcal{S}_2$ ,  $U \in \mathcal{S}_2$ . So,

$$\begin{aligned} |U| &= \sum_{i=1}^n \max(0, \lceil x_2 r_i - B \rceil - \lfloor (x_1 - 1)r_i + B \rfloor) \quad (\text{by (19)}) \\ &\leq |N(U)| \quad (\text{because } U \in \mathcal{S}_2) \end{aligned}$$

$$\begin{aligned} &\leq |I| \quad (\text{because } N(U) \subseteq I) \\ &= x_2 - x_1 + 1, \end{aligned}$$

and (17) follows.  $\square$

In Section 5, we will see that the MDJIT problem always has a solution with  $B < 1$  (see Theorem 5). If we slightly anticipate on this result, we can reformulate Theorem 1 in a simpler form by relying on the next observation.

**Proposition 5.** *When  $B < 1$ , the following statements (a)–(d) are equivalent:*

(a) *for all  $x_1, x_2$  in  $[1 .. D]$  with  $x_1 \leq x_2$ ,*

$$\sum_{i=1}^n (\lfloor x_2 r_i + B \rfloor - \lceil (x_1 - 1) r_i - B \rceil) \geq x_2 - x_1 + 1, \tag{20}$$

(b) *for all  $x_1, x_2$  in  $[1 .. D]$  with  $x_1 \leq x_2$ ,*

$$\sum_{i=1}^n \max(0, \lfloor x_2 r_i + B \rfloor - \lceil (x_1 - 1) r_i - B \rceil) \geq x_2 - x_1 + 1, \tag{21}$$

(c) *for all  $x$  in  $[1 .. D]$ ,*

$$\sum_{i=1}^n \lfloor x r_i + B \rfloor \geq x, \tag{22}$$

(d) *for all  $x$  in  $[1 .. D]$ ,*

$$\sum_{i=1}^n \lfloor x r_i + B \rfloor \geq x, \tag{23}$$

$$\sum_{i=1}^n \lceil x r_i - B \rceil \leq x. \tag{24}$$

**Proof.** We are going to show that the following chain of implications is valid:

$$(a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d) \Rightarrow (a).$$

The first implication (a)  $\Rightarrow$  (b) is trivial. To derive the next implication, just set  $x_1 = 1$  in (21) and observe that  $\lfloor x_2 r_i + B \rfloor \geq 0$  and  $\lceil -B \rceil = 0$  (since  $B < 1$ ).

The third implication is obtained by verifying that (23) holds for all  $x \in [1 .. D]$  if and only if (24) holds for all  $x \in [1 .. D]$  (just write out each inequality at the point  $x' = D - x$ , the case  $x = D$  being trivial).

Finally, to derive implication (d)  $\Rightarrow$  (a), just write (23) at  $x = x_2$ , write (24) at  $x = x_1 - 1$  and subtract the resulting inequalities.  $\square$

As a corollary, we obtain the simpler form of Theorem 1:

**Theorem 2.** *When  $B < 1$ , the MDJIT problem has a feasible solution if and only if, for all  $x_1, x_2$  in  $[1 .. D]$  with  $x_1 \leq x_2$ , the following inequalities are both valid:*

$$\sum_{i=1}^n \lfloor x_1 r_i + B \rfloor \geq x_1, \quad (25)$$

$$\sum_{i=1}^n \max(0, \lfloor x_2 r_i - B \rfloor - \lfloor (x_1 - 1) r_i + B \rfloor) \leq x_2 - x_1 + 1. \quad (26)$$

**Proof.** This is an immediate consequence of Theorem 1 and Proposition 5.  $\square$

In view of the equivalence of statements (c) and (d) in Proposition 5 and of the easy observation that (24) arises by setting  $x_1 = 1$  and by dropping the max-operator in (26), one may be led to conjecture that inequalities (26) actually are redundant in Theorem 2 and that the following compact statement holds true:

**Conjecture 1.** *The MDJIT problem has a feasible solution if and only if the following inequality is valid for all  $x \in [1 .. D]$ :*

$$\sum_{i=1}^n \lfloor x r_i + B \rfloor \geq x. \quad (27)$$

However, this conjecture is wrong, as evidenced by the instance  $I = (d_1 = 3, d_2 = 3, d_3 = 1, B = \frac{4}{7})$ . Indeed, conditions (27) hold for this instance (for all  $x \in [1 .. 7]$ ), but the instance is infeasible since the graph in Fig. 2 has no perfect matching or, alternatively, since inequality (26) fails when  $x_1 = x_2 = 4$ .

It may be interesting to note that Conjecture 1 is closely related to the so-called “Deficiency cases” described in [25]. The authors assert that the EDD “algorithm can stop at time  $k < D - 1$  for one of two reasons”: if less than  $k$  parts are available to schedule in  $[1 .. k]$ , or more than  $k$  parts must be scheduled in  $[1 .. k]$ . We observe that, for the instance  $I = (d_1 = 3, d_2 = 3, d_3 = 1, B = \frac{4}{7})$ , neither of the deficiency cases holds, even though  $I$  is infeasible.

#### 4. Complexity results

As discussed in Section 2.3, the complexity of the MDJIT problem is not exactly known. Namely, it would be possible for its recognition version to be solvable in polynomial time despite the fact that formulation (5)–(9) is of pseudo-polynomial size (see [6] for a finer discussion of this issue). Such a result, if true, would necessarily imply that the MDJIT problem is both in NP and in Co-NP. Here, we prove that at least one of these conditions holds:

**Theorem 3.** *The MDJIT problem is in Co-NP.*

**Proof.** If an instance of the MDJIT problem is not feasible, then, by Theorem 2, there exist values of  $x_1$  and  $x_2$  such that one of inequalities (25) or (26) is not valid. Given  $x_1$  and  $x_2$ , this can be checked in time  $O(n \log D)$ .  $\square$

We can also use Theorem 2 to reduce the MDJIT problem to an integer linear program (ILP) whose size is polynomial in  $n$  and  $\log D$ . Indeed, consider an instance  $(d_1, \dots, d_n, B)$  of MDJIT. By Theorem 2, the instance has no feasible solution if and only if either there exists  $x \in [1 .. D]$  such that

$$\sum_{i=1}^n \lfloor xr_i + B \rfloor < x \tag{28}$$

or there exist  $x_1, x_2 \in [1 .. D]$  with  $x_1 \leq x_2$  such that

$$\sum_{i=1}^n \max(0, \lfloor x_2 r_i - B \rfloor - \lfloor (x_1 - 1) r_i + B \rfloor) > x_2 - x_1 + 1. \tag{29}$$

For  $i = 1, \dots, n$ , introduce the decision variable  $X_i$  to represent the quantity  $\lfloor xr_i + B \rfloor$ . Then, inequality (28) holds exactly when the following system of inequalities is feasible:

$$\begin{aligned} \sum_{i=1}^n X_i &< x \\ X_i &> xr_i + B - 1 \quad (i = 1, \dots, n), \\ x &\in [1 .. D], \\ X_i &\in \mathbb{N} \quad (i = 1, \dots, n). \end{aligned}$$

Observe that, from expression (5) (or from the definition of the “deviation” concept), it is clear that we can restrict the values of  $B$  to integer multiples of  $1/D$ . Thus, let  $B = \beta/D$ , where  $\beta$  is an integer. After some manipulations, the above system can be rewritten as

$$\sum_{i=1}^n X_i \leq x - 1, \tag{30}$$

$$DX_i \geq xd_i + \beta - D + 1 \quad (i = 1, \dots, n), \tag{31}$$

$$x \leq D, \tag{32}$$

$$x, X_i \in \mathbb{N} \quad (i = 1, \dots, n). \tag{33}$$

In particular, the smallest value of  $B$  such that inequality (25) holds for all values of  $x$  can be computed by solving the ILP

$$\max \left( \frac{\beta}{D} + \frac{1}{D} \right) \quad \text{subject to (30) – (33) and } \beta \in \mathbb{N}.$$

Inequality (29) can be handled in a similar way, except that the presence of the max-operators creates some additional difficulties. So, for  $i = 1, \dots, n$ , introduce the decision variables  $x_1, x_2, X_i^1, X_i^2$  and  $X_i$ , where  $X_i^1$  stands for  $\lfloor (x_1 - 1)r_i + B \rfloor$ ,  $X_i^2$  stands for  $\lceil x_2 r_i - B \rceil$  and  $X_i$  stands for  $\max(0, X_i^2 - X_i^1)$ . For each  $i$ , we also introduce a binary variable  $\delta_i$  which takes value 1 when  $X_i = X_i^2 - X_i^1$  and value 0 when  $X_i = 0$ .

Inequality (29) holds exactly when the following system of inequalities is feasible:

$$\begin{aligned} \sum_{i=1}^n X_i &> x_2 - x_1 + 1, \\ X_i^1 &> (x_1 - 1)r_i + B - 1 \quad (i = 1, \dots, n), \\ X_i^2 &< x_2 r_i - B + 1 \quad (i = 1, \dots, n), \\ X_i &\geq X_i^2 - X_i^1 \quad (i = 1, \dots, n), \\ X_i &\geq 0 \quad (i = 1, \dots, n), \\ X_i &\leq X_i^2 - X_i^1 + D(1 - \delta_i) \quad (i = 1, \dots, n), \\ X_i &\leq \delta_i D \quad (i = 1, \dots, n), \\ x_1 &\leq x_2, \\ x_1, x_2 &\in [1 .. D], \\ X_i^1, X_i^2, X_i &\in \mathbb{N}, \quad \delta_i \in \{0, 1\} \quad (i = 1, \dots, n). \end{aligned}$$

After some manipulations, we obtain again that the smallest value of  $B$  such that inequality (26) holds for all values of  $(x_1, x_2)$  can be computed by solving an appropriate ILP problem. So, we conclude that the optimization version of the MDJIT problem (i.e., the problem of finding the smallest value of  $B$  such that (5)–(9) is feasible) can be reduced to a pair of integer linear programs involving  $O(n)$  variables. In particular:

**Theorem 4.** *For fixed  $n$ , the minimum value of  $B$  such that the MDJIT problem is feasible can be computed in time polynomial in  $O(\log D)$ .*

**Proof.** This is a straightforward consequence of the above discussion and of Lenstra's results [20] on integer programming problems with a fixed number of variables.  $\square$

In Section 7, we will show that the MDJIT problem is actually very easy to solve when  $n = 2$ . But for fixed  $n > 2$ , we are not aware of a more direct proof of the fact that MDJIT is polynomially solvable.

These observations also leave open the more challenging question: is it possible to solve the (recognition version of the) MDJIT problem in time polynomial in  $n$  and  $\log D$  or is the problem co-NP-complete?

**5. Bounds on the maximum deviation**

In this section, we provide an upper bound and several lower bounds on the optimal value of the MDJIT problem.

*5.1. Upper bound*

Steiner and Yeomans [25] proved the interesting fact that the optimal value of the MDJIT problem is always less than or equal to 1. The same bound can also be deduced from the description of the quota apportionment method (see [3]; in fact, the quota method can be interpreted as a version of the EDD algorithm, applied with the bound  $B = 1$ ). Here, we state a slightly stronger version of this result, whose proof is a rather easy consequence of Theorem 2.

**Theorem 5.** *The optimal value  $B^*$  of the MDJIT problem satisfies*

$$B^* \leq 1 - \frac{1}{D}.$$

**Proof.** Let  $B = 1 - 1/D$ . Since  $B < 1$ , we are in a position to apply Theorem 2, i.e., we only need to prove that inequalities (25) and (26) hold for this value of  $B$ .

We first prove that, for any integer  $x \geq 0$ , one has

$$\lfloor xr_i + B \rfloor \geq xr_i. \tag{34}$$

Indeed, if  $xr_i$  is an integer, then  $\lfloor xr_i + B \rfloor = xr_i$ . If  $xr_i$  is not an integer, then  $\{xr_i\} \geq 1/D$  (where  $\{\alpha\}$  denotes the fractional part of  $\alpha$ ). This implies

$$\{xr_i\} + B \geq \frac{1}{D} + 1 - \frac{1}{D} = 1$$

and hence

$$\lfloor xr_i + B \rfloor = \lfloor xr_i \rfloor + 1 > xr_i,$$

which establishes (34). Inequality (25) follows now immediately by summing (34) over  $i = 1, \dots, n$ .

For inequality (26), fix  $x_1, x_2 \in [1 .. D]$ , with  $x_1 \leq x_2$ , and consider the set  $J \subseteq \{1, 2, \dots, n\}$  defined by

$$i \in J \Leftrightarrow \lceil x_2 r_i - B \rceil - \lfloor (x_1 - 1)r_i + B \rfloor \geq 0.$$

Note that substituting  $D - x_2$  for  $x$  in inequality (34) leads to  $\lceil x_2 r_i - B \rceil \leq x_2 r_i$  for  $i = 1, \dots, n$ . Then, we derive successively

$$\sum_{i=1}^n \max(0, \lceil x_2 r_i - B \rceil - \lfloor (x_1 - 1)r_i + B \rfloor) = \sum_{i \in J} (\lceil x_2 r_i - B \rceil - \lfloor (x_1 - 1)r_i + B \rfloor)$$

$$\begin{aligned} &\leq \sum_{i \in J} (x_2 r_i - (x_1 - 1) r_i) \\ &\leq x_2 - x_1 + 1 \end{aligned}$$

and hence inequality (26) holds.  $\square$

Note that the bound  $1 - 1/D$  is attained when  $d_i = 1$  for all  $i = 1, \dots, n$ .

Theorem 5 has interesting side-implications regarding the periodicity of optimal sequences (this issue has also been considered, for instance, in [22,16], for total deviation objective functions). Assume that  $kr_i$  is an integer for some  $k \in [1..D]$  and  $i \in \{1, \dots, n\}$ . Then, in every optimal solution of MDJIT, the number of parts of type  $i$  produced up to time  $k$  must be equal to  $x_{i,k} = kr_i$ , since otherwise the deviation at time  $k$  would be at least  $1 > B^*$ . Now, if  $\gcd(d_1, \dots, d_n, D) = \alpha$  and  $D = \alpha\Delta$ , then  $\Delta r_i = d_i/\alpha$  is integral for *all* values of  $i$ . Thus, necessarily,  $d_i/\alpha$  parts of type  $i$  must have been produced up to time  $\Delta$ , for *all*  $i \in \{1, \dots, n\}$ . A similar argument also applies at times  $2\Delta, 3\Delta \dots$  and this implies that the optimal schedule must consist of  $\alpha$  subsequences, where each subsequence involves exactly  $d_i/\alpha$  parts of type  $i$  for  $i = 1, \dots, n$ . Pushing the reasoning a little bit further actually leads to the conclusion that, when  $\alpha > 1$ , there exists an optimal production sequence of the form  $(S, S, \dots, S)$ , where  $S$  is an optimal production sequence for the reduced instance  $(d_1/\alpha, \dots, d_n/\alpha)$  (we omit the details). This result has been obtained earlier in [26].

## 5.2. Lower bounds

We now turn to lower bounds on the optimal value  $B^*$ . We start with an easy result.

**Proposition 6.** *Assume that  $d_1 \leq d_2 \leq \dots \leq d_n$ . Then, the optimal value  $B^*$  of the MDJIT problem satisfies*

$$r_i \leq B^* \quad \text{for } i = 1, 2, \dots, n-1, \quad (35)$$

$$1 - r_i \leq 2B^* \quad \text{for } i = 1, 2, \dots, n-1, \quad (36)$$

$$r_n \leq 2B^*, \quad (37)$$

$$1 - r_n \leq B^*, \quad (38)$$

$$\frac{n-1}{2n-1} \leq B^*. \quad (39)$$

**Proof.** (a) In any feasible schedule, some part type  $i$  must be produced in the first time period  $k = 1$ . From inequality (5), we deduce that  $|1 - r_i| = 1 - r_i \leq B^*$  and that  $|0 - r_j| = r_j \leq B^*$  for all  $j \neq i$ . Conditions (35) and (38) follow.



(b) If MDJIT has a feasible schedule, then necessarily  $E(i, j) \leq L(i, j)$  for all parts  $(i, j)$ . Hence:

$$\begin{aligned} 0 &\leq L(i, j) - E(i, j) \\ &\leq \frac{j - 1 + B^*}{r_i} + 1 - \frac{j - B^*}{r_i} \\ &= \frac{2B^* + r_i - 1}{r_i} \end{aligned}$$

and condition (36) follows. Moreover, condition (37) also holds, since

$$2B^* \geq 1 - r_1 = \sum_{i=2}^n r_i \geq r_n.$$

(c) For the last bound, just add the inequalities (36) ( $i = 1, 2, \dots, n - 1$ ) to inequality (38).  $\square$

Interestingly, Kovalyov et al. [14] mention that, for small examples, the optimal value of MDJIT very often coincides with the lower bound (38). In their computational experiments, however, they provide examples showing that the bound is not always attained. As a matter of fact, combining bounds (37) and (38) immediately implies that (38) cannot be tight as soon as  $r_n > \frac{2}{3}$ . In particular, the instance  $d_1 = 1, d_2 = 3$ , whose optimal value is  $B^* = \frac{1}{2}$ , suffices to show that none of the lower bounds in Proposition 6 needs to be attained (see also Theorem 9).

**Theorem 6.** Let  $\Delta_i = D/\text{gcd}(d_i, D)$  ( $i = 1, \dots, n$ ). The optimal value  $B^*$  of the MDJIT problem satisfies

$$B^* \geq \frac{1}{\Delta_i} \left\lfloor \frac{\Delta_i}{2} \right\rfloor \quad \text{for } i = 1, \dots, n.$$

**Proof.** Consider some part type  $i$ . Its ideal production rate is  $r_i = d_i/D = \delta_i/\Delta_i$  where  $\text{gcd}(\Delta_i, \delta_i) = 1$ . Note that any feasible solution  $(x_{ik})$  of the MDJIT problem satisfies  $|x_{ik} - kr_i| \geq |[kr_i] - kr_i|$  for all  $k \in [1..D]$ . We shall prove the theorem by showing that there exists a value of  $k$  such that  $|[kr_i] - kr_i| = 1/\Delta_i \lfloor \Delta_i/2 \rfloor$ .

If  $\Delta_i$  is even, then, for  $k = \Delta_i/2$ ,

$$|kr_i - [kr_i]| = \left| \frac{\Delta_i}{2} \frac{\delta_i}{\Delta_i} - \left\lfloor \frac{\Delta_i}{2} \frac{\delta_i}{\Delta_i} \right\rfloor \right| = \left| \frac{\delta_i}{2} - \left\lfloor \frac{\delta_i}{2} \right\rfloor \right|.$$

Since  $\text{gcd}(\delta_i, \Delta_i) = 1$  and  $\Delta_i$  is even,  $\delta_i$  is odd and hence  $|\delta_i/2 - \lfloor \delta_i/2 \rfloor| = 0.5$ . Therefore, for  $\Delta_i$  even, the optimal value of the objective function is at least  $1/\Delta_i \lfloor \Delta_i/2 \rfloor = 0.5$ .

Suppose now that  $\Delta_i$  is odd. We prove that there exists a value of  $k$  such that  $|[kr_i] - kr_i| = (\Delta_i - 1/2\Delta_i)$ . Since  $\delta_i$  and  $\Delta_i$  are relatively prime, Bezout's identity implies that there exist two integers  $u$  and  $v$  such that  $u\Delta_i + v\delta_i = 1$ . Multiply this

inequality by  $(\Delta_i - 1)/2\Delta_i$  and set  $k = \lfloor v \rfloor (\Delta_i - 1)/2$  thus obtaining

$$k \frac{\delta_i}{\Delta_i} = \lfloor u \rfloor \frac{\Delta_i - 1}{2} \pm \frac{\Delta_i - 1}{2\Delta_i}.$$

Since  $(\Delta_i - 1)/2\Delta_i < 0.5$ , one has  $\lfloor k(\delta_i/\Delta_i) \rfloor = \lfloor u \rfloor (\Delta_i - 1)/2$  and  $|\lfloor k(\delta_i/\Delta_i) \rfloor - k(\delta_i/\Delta_i)| = (\Delta_i - 1)/2\Delta_i$ .  $\square$

We will see in Section 7 that the bound presented in Theorem 6 is attained for  $n=2$ .

## 6. Small deviations

Observe that the bound in Theorem 6 is usually close to  $1/2$ , and that  $B^* \geq 1/2$  as soon as there exists some  $i$  such that  $\Delta_i$  is even. Note also that the bound (39) goes to  $1/2$  when  $n$  goes to  $+\infty$ . These observations suggest to look more closely at those instances for which the maximum deviation does not exceed the value  $1/2$ .

In the remainder of this section, we state a few conjectures regarding the structure of the instances with  $B^* \leq 1/2$ , and we identify all instances with optimal value  $B^* < 1/2$  for  $n \leq 6$ . In order to formulate these statements in the simplest possible form, we restrict our attention to *standard* instances of the MDJIT problem, i.e. instances  $(d_1, \dots, d_n)$  such that  $d_1 \leq d_2 \leq \dots \leq d_n$  and  $\gcd(d_1, \dots, d_n, D) = 1$  (remember the comments following Theorem 5).

### 6.1. Instances with $B^* < 1/2$

When  $B^* < 1/2$ , the condition  $|x_{ik} - kr_i| \leq B^*$  forces  $x_{ik}$  to be equal to  $\lfloor kr_i \rfloor$ , so that the problem becomes highly constrained. As a matter of fact, we conjecture that, for  $n > 2$ , only a handful of very special instances have this property, namely the instances of the form  $(1, 2, 4, \dots, 2^{n-1})$  (the case  $n=2$  will be dealt with in the next section). We will prove this conjecture for  $n \leq 6$  using connections with so-called balanced words [28].

**Conjecture 2.** For  $n > 2$ , a standard instance  $(d_1, \dots, d_n)$  of the MDJIT problem has optimal value  $B^* < \frac{1}{2}$  if and only if  $d_i = 2^{i-1}$  for  $i = 1, 2, \dots, n$ , and  $B^* = (2^{n-1} - 1)/(2^n - 1)$ .

We first establish the sufficiency of the conditions.

**Proposition 7.** The instance  $(d_1, \dots, d_n)$ , with  $d_i = 2^{i-1}$  for  $i = 1, 2, \dots, n$ , has optimal value  $B^* = (2^{n-1} - 1)/(2^n - 1)$ .

**Proof.** Consider the instance  $d_i = 2^{i-1}$  for  $i = 1, \dots, n$ , and let  $B^* = (D - 1)/2D = (2^{n-1} - 1)/(2^n - 1) < \frac{1}{2}$ . We want to show that this value  $B^*$  is feasible. In view of Theorem 6, this will imply that  $B^*$  is optimal.

The ideal rates are defined by  $r_i = 2^{i-1}/(2^n - 1)$ . Let  $s_{ij} = 2^{n-i}(2j - 1)$  for  $i \in [1 .. n]$  and  $j \in [1 .. 2^{i-1}]$ . We first prove that

$$s_{ij} \geq \frac{j - B^*}{r_i} \quad \text{and} \quad s_{ij} \leq \frac{j - 1 + B^*}{r_i} + 1.$$

Indeed,

$$\begin{aligned} s_{ij} - \frac{j - B^*}{r_i} &= \frac{2^{n-1}(2j - 1) - j(2^n - 1) + 2^{n-1} - 1}{2^{i-1}} \\ &= \frac{j - 1}{2^{i-1}} \geq 0 \end{aligned}$$

and

$$\begin{aligned} \frac{j - 1 + B^*}{r_i} + 1 - s_{ij} &= \frac{(j - 1)(2^n - 1) + 2^{n-1} - 1 + 2^{i-1} - 2^{n-1}(2j - 1)}{2^{i-1}} \\ &= \frac{2^{i-1} - j}{2^{i-1}} \geq 0. \end{aligned}$$

Since  $s_{ij}$  is integer, one has  $E(i, j) = \lceil (j - B^*)/r_i \rceil \leq s_{ij} \leq \lfloor (j - 1 + B^*)/r_i + 1 \rfloor = L(i, j)$ . Moreover,  $s_{ij} \neq s_{i', j'}$  for  $i \neq i'$  or  $j \neq j'$  and  $1 \leq s_{ij} \leq D = 2^n - 1$ . Therefore, by Proposition 1, there exists a feasible schedule: just produce part  $(i, j)$  at time  $s_{ij}$  for  $i \in [1 .. n]$  and  $j \in [1 .. 2^{i-1}]$ . This completes the proof. Note, incidentally, that the quantity  $s_{ij}$  corresponds exactly to the ideal position for copy  $j$  of product  $i$ , as used for instance in [17].  $\square$

In order to prove the unicity of the instance in Conjecture 2 for  $n \leq 6$ , let us introduce some terminology from [28]. A *balanced word* on  $\{1, 2, \dots, n\}$  is an infinite sequence  $\sigma = (s_1, s_2, \dots)$  such that

- (1)  $s_j \in \{1, 2, \dots, n\}$  for all  $j \in \mathbb{N}_0$ , and
- (2) if  $\sigma_1$  and  $\sigma_2$  are two subsequences consisting of  $t$  consecutive elements of  $\sigma$  ( $t \in \mathbb{N}$ ), then the number of occurrences of  $i$  in  $\sigma_1$  and  $\sigma_2$  differs by at most 1, for all  $i = 1, 2, \dots, n$ .

If  $\sigma$  is a balanced word on  $\{1, 2, \dots, n\}$ , then the *density* of  $i$  in  $\sigma$  is

$$\theta_i = \lim_{t \rightarrow \infty} \frac{|\{j \in [1 .. t]: s_j = i\}|}{t},$$

for  $i = 1, 2, \dots, n$  (it can be shown that the limit always exists).

**Proposition 8.** *Let  $S$  be an optimal sequence with  $B^* < 1/2$  for an instance  $(d_1, \dots, d_n)$  of MDJIT. Then, the infinite sequence obtained by repeating  $S$  is a balanced word with distinct densities  $(d_1/D, \dots, d_n/D)$ .*

**Proof.** Consider an optimal sequence  $S$  for the instance  $(d_1, d_2, \dots, d_n)$ , with  $B^* < 1/2$ , and consider the infinite sequence  $\sigma$  obtained by repeating  $S$ , i.e.  $\sigma = (S, S, S, \dots)$ . We want to show that  $\sigma$ , viewed as a sequence on  $\{1, 2, \dots, n\}$ , is a balanced word. Let

$t \in \mathbb{N}$ , let  $\sigma_1, \sigma_2$  be two subsequences consisting of  $t$  consecutive elements of  $\sigma$ , and let us establish condition (2) in the definition of balanced words.

Assume that  $\sigma_j$  ranges from time  $t_j + 1$  to time  $t_j + t$ , for  $j = 1, 2$ . Fix  $i \in \{1, 2, \dots, n\}$  and denote by  $|I|$  the number of occurrences of  $i$  in any time interval  $I$ . Then, for  $j = 1, 2$ ,

$$\begin{aligned} |[t_j + 1, t_j + t]| &= |[1, t_j + t]| - |[1, t_j]| \\ &< \left( (t_j + t)r_i + \frac{1}{2} \right) - \left( t_j r_i - \frac{1}{2} \right) \\ &= tr_i + 1 \end{aligned}$$

and similarly

$$\begin{aligned} |[t_j + 1, t_j + t]| &> \left( (t_j + t)r_i - \frac{1}{2} \right) - \left( t_j r_i + \frac{1}{2} \right) \\ &= tr_i - 1. \end{aligned}$$

Thus,  $|[t_1 + 1, t_1 + t]|$  and  $|[t_2 + 1, t_2 + t]|$  are two integers lying strictly between  $tr_i - 1$  and  $tr_i + 1$ . There follows that  $|[t_1 + 1, t_1 + t]|$  and  $|[t_2 + 1, t_2 + t]|$  differ at most by 1, and hence  $\sigma$  is balanced.

For  $i = 1, 2, \dots, n$ , the density of  $i$  in  $\sigma$  is equal to  $r_i = d_i/D$ . If  $r_i = r_j$  for  $i \neq j$ , then  $x_{ik} = [kr_i] = [kr_j] = x_{jk}$  for all  $k$ , which is clearly impossible. Hence, the densities in  $\sigma$  are pairwise distinct.  $\square$

Some attention has been devoted in number theory to the properties of balanced words and related concepts, such as Beatty covers and Sturmian words. Without going into the details, let us just mention here that Beatty covers and Sturmian words are special types of balanced words. Fraenkel conjectured that, when  $n \geq 3$ , every Beatty cover on  $n$  letters with distinct densities has densities  $2^{i-1}/(2^n - 1)$ , for  $i = 1, \dots, n$ .

Tijdeman [28] presents a thorough discussion of the state-of-the-art concerning this conjecture. Although the conjecture is still open, it is known to hold when  $n \leq 6$ , even in the more general case of balanced words (see [27] for  $n = 3$ , [1] for  $n = 4$ , [28] for  $n = 5$  and [28,29] for  $n = 6$ ). As a corollary, we obtain:

**Theorem 7.** *Conjecture 2 is valid for  $n \leq 6$ .*

**Proof.** Fraenkel's conjecture holds for balanced words when  $n \leq 6$ . Applying its conclusions to the infinite sequence constructed as in Proposition 8, we directly obtain Conjecture 2.  $\square$

More generally, Conjecture 2 would follow (for all values of  $n$ ) from the validity of Fraenkel's conjecture for balanced words. Although the latter conjecture is still open, Brauner et al. [7] have very recently been able to establish the validity of Conjecture 2 for all values of  $n$  by direct arguments.

6.2. Instances with  $B^* \leq 1/2$

If we only impose  $B^* \leq 1/2$ , then the structure of feasible instances becomes more complex, as  $x_{ik}$  may now be equal either to  $\lceil kr_i \rceil = kr_i - 1/2$  or to  $\lfloor kr_i \rfloor + 1 = kr_i + 1/2$  when  $kr_i$  is half-integral. The following statement describes a few classes of instances with optimal value  $B^* \leq 1/2$ . Although we are not able to establish that these classes cover all possible cases, we conjecture that they indeed do so.

**Conjecture 3.** *A standard instance  $(d_1, \dots, d_n)$  of the MDJIT problem has optimal value  $B^* \leq 1/2$  if and only if it satisfies one of the following conditions:*

[1] *the demands are successive powers of two:*

$$d_i = 2^{i-1} \quad \text{for all } i \in [1..n],$$

[2]  *$d_1$  and  $d_2$  are arbitrary and for  $i \geq 3$ ,  $d_i$  is the sum of all demands with smaller index:*

$$d_i = \sum_{j < i} d_j = 2^{i-3}(d_1 + d_2) \quad \text{for all } i \in [3..n],$$

[3]  *$d_1 = 1, d_2 = 2, d_3 = 9$  and for  $i \geq 4$ ,  $d_i$  is the sum of all demands with smaller index:*

$$d_1 = 1; \quad d_2 = 2; \quad d_3 = 9; \quad d_i = \sum_{j < i} d_j = 2^{i-4} \times 12 \quad \text{for all } i \in [4..n],$$

[4]  *$d_1 = 2, d_2 = 3, d_3 = 7$  and for  $i \geq 4$ ,  $d_i$  is the sum of all demands with smaller index:*

$$d_1 = 2; \quad d_2 = 3; \quad d_3 = 7; \quad d_i = \sum_{j < i} d_j = 2^{i-4} \times 12 \quad \text{for all } i \in [4..n],$$

[5] *there exists an index  $l \in [3..n]$  such that*

$$d_i = 2^{i-1} \quad \text{for all } i \in [1..l],$$

$$d_i = \sum_{j < i} d_j = 2^{i-l-1}(2^l - 1) \quad \text{for all } i \in [l+1..n],$$

[6] *there exists an index  $l \in [3..n]$  such that*

$$d_i = 2^{i-1} \quad \text{for all } i \in [1..l-1],$$

$$d_l = 2^{l-1} + 1,$$

$$d_i = \sum_{j < i} d_j = 2^{i-1} \quad \text{for all } i \in [l+1..n].$$

Observe that, except for the instances satisfying condition [1], which arise directly from Conjecture 2, all other instances are constructed following the same general pattern: they consist of an initial segment  $(d_1, \dots, d_l)$  where  $l$  can take different values,

and each of the remaining demands  $d_i$  ( $i > l$ ) is simply the sum of the previous ones ( $d_1, \dots, d_{i-1}$ ). Based on this observation, the sufficiency of each condition in Conjecture 3 can be verified using case-by-case analysis on the initial segments, together with the general result below (we skip the details).

**Proposition 9.** *Let  $I = (n, d_1, \dots, d_n, B)$  be a feasible instance of the MDJIT problem with  $B \geq 1/2$ . Then, the following instance, involving an additional part type, is also feasible:*

$$I' = (n + 1, d_1, \dots, d_n, d_{n+1} = \sum_{i=1}^n d_i, B).$$

**Proof.** Let  $(x_{ik})$  be a solution of the MDJIT for the instance  $I$ . For each  $k = 0, 1, \dots, D - 1$ , one can determine the index of the part type which is produced at time  $k + 1$ , namely the unique index  $i_k$  such that

$$x_{i_k, k+1} = x_{i_k, k} + 1 \quad \text{and} \quad x_{i, k+1} = x_{i, k} \quad \text{for all} \quad i \neq i_k$$

(for simplicity of notations, we assume here that  $x_{i0} = 0$ ).

For the extended instance, define a solution  $x'_{ij}$  ( $i = 1, 2, \dots, n + 1, j = 1, 2, \dots, 2D$ ) as follows:

$$\begin{aligned} \text{for } k = 1 \dots D \text{ and } i = 1 \dots n : \quad & x'_{i, 2k} = x_{ik}, \\ & x'_{n+1, 2k} = k, \\ \text{for } k = 0 \dots D - 1 \text{ and } i \neq i_k : \quad & x'_{i, 2k+1} = x_{ik}, \\ \text{for } k = 0 \dots D - 1 \text{ and } i = i_k : \quad & x'_{i, 2k+1} = x_{ik} \quad \text{if } |x_{ik} - (k + 1/2)r_i| \leq B, \\ & = x_{ik} + 1 \quad \text{otherwise} \\ \text{for } k = 0 \dots D - 1 \text{ and } i = i_k : \quad & x'_{n+1, 2k+1} = k + 1 \quad \text{if } |x_{ik} - (k + 1/2)r_i| \leq B, \\ & = k \quad \text{otherwise.} \end{aligned}$$

It remains to prove that  $x'_{ik}$  is a solution of the MDJIT problem for the instance  $I'$ . Since  $x_{i, k+1} \geq x_{ik}$  for any  $i$  and  $k$ , one may verify that  $x'_{i, k+1} \geq x'_{ik}$  for any  $i$  and  $k$ . Moreover,  $x'_{i, 2D} = x_{i, D} = d_i$  and  $x'_{n+1, 2D} = D = d_{n+1}$  imply equality (2). We will now prove that equality (1) applies for  $x'_{i, k}$ :

$$\begin{aligned} \sum_{i=1}^{n+1} x'_{i, 2k} &= \sum_{i=1}^n x_{i, k} + k \\ &= 2k \quad \text{for } k = 1, 2, \dots, D, \end{aligned}$$

$$\sum_{i=1}^{n+1} x'_{i, 2k+1} = \sum_{i=1, i \neq i_k}^n x_{i, k} + x'_{i_k, 2k+1} + x'_{n+1, 2k+1}$$

$$\begin{aligned}
 &= \sum_{i=1, i \neq i_k}^n x_{i,k} + x_{i_k,k} + k + 1 \\
 &= 2k + 1 \quad \text{for } k = 0, 1, \dots, D - 1.
 \end{aligned}$$

Note that for the instance  $I'$ , the ideal production rate  $r'_i$  of part  $i$  is equal to  $r_i/2$ , where  $r_i$  is the rate for part  $i$  in the instance  $I$  ( $i = 1, 2, \dots, n$ ), and  $r'_{n+1} = 1/2$ . Using these remarks, one can verify that  $|x'_{ik} - kr'_i| \leq B$  for  $i = 1, 2, \dots, n + 1$  and  $k = 1, 2, \dots, 2D$ . This concludes the proof.  $\square$

### 7. Two-part type problem

In this last section, we consider the MDJIT problem for two part types ( $n=2$ ) with ideal production rates  $r_1$  and  $r_2 = 1 - r_1$ . We assume without loss of generality that  $0 < r_1 \leq 0.5$ . The optimization version of the MDJIT problem can be written as

*Input:*  $d_1, d_2 \in \mathbb{N}$ .

As usual, denote the total demand by  $D = d_1 + d_2$  and the ideal production rate for part type 1 by  $r_1 = d_1/D$ . Then the ideal production rate for part type 2 satisfies  $r_2 = 1 - r_1$ .

*Question:* Find a  $2 \times D$  matrix  $X = (x_{i,k})$  which minimizes the maximum deviation  $\max_{1 \leq k \leq D} (|x_{1k} - kr_1|, |x_{2k} - kr_2|)$  subject to

$$x_{1,k} + x_{2,k} = k, \quad k = 1, \dots, D, \tag{40}$$

$$x_{i,D} = d_i, \quad i = 1, 2, \tag{41}$$

$$0 \leq x_{i,k} - x_{i,k-1}, \quad i = 1, 2, \quad k = 2, \dots, D, \tag{42}$$

$$x_{i,k} \in \mathbb{N} \quad i = 1, 2, \quad k = 1, \dots, D. \tag{43}$$

**Theorem 8.** *The matrix  $X$  defined by  $x_{1,k} = [kr_1]$  and  $x_{2,k} = k - [kr_1]$  ( $k = 1, \dots, D$ ) is an optimal solution of the 2-part type MDJIT problem.*

**Proof.** Note that  $x_{2,k}$  might be different from  $[kr_2]$  whenever the fractional part of  $x_{1,k}$  is 0.5. One can observe that for  $k = 1, \dots, D$ ,

$$x_{1,k} + x_{2,k} = [kr_1] + k - [kr_1] = k,$$

$$x_{i,D} = \left\lceil D \frac{d_i}{D} \right\rceil = d_i,$$

$$x_{1,k} - x_{1,k-1} = [kr_1] - [(k-1)r_1] \geq 0$$

and

$$x_{2,k} - x_{2,k-1} = k - [kr_1] - k + 1 + [(k-1)r_1]$$

$$\begin{aligned}
&= 1 + [(k-1)r_1] - [kr_1] \\
&\geq 0.
\end{aligned}$$

Therefore,  $X$  satisfies constraints (40)–(43). We now prove that  $X$  is an optimal solution for the 2-part type problem. Clearly,

$$|x_{1,k} - kr_1| = |[kr_1] - kr_1| \leq 1/2$$

and

$$|x_{2,k} - kr_2| = |k - [kr_1] - k + kr_1| = |-[kr_1] + kr_1| \leq 1/2.$$

Let  $X' = (x'_{1,k}; x'_{2,k})$  be another feasible solution of the 2-part MDJIT problem. We want to show that  $X'$  has maximum deviation larger than or equal to  $1/2$ , which implies that  $X'$  is not better than  $X$ . Assume that  $X'$  differs from  $X = (x_{1,k}; x_{2,k})$  at period  $l$ . Because of constraint (40), it must be the case that  $x'_{1,l}$  is not equal to  $x_{1l} = [lr_1]$ . Thus, by definition of the  $[\cdot]$  operator,  $x'_{1,l}$  is at distance at least  $1/2$  from  $lr_1$ , i.e.  $|x'_{1,l} - lr_1| \geq 1/2$ , as needed.  $\square$

**Remark 2.** Note that the matrix defined in Theorem 8 actually minimizes the deviation  $x_{ik} - kr_i$  for all  $k$  and  $i$ . Therefore, it is optimal for the maximum deviation problem and for the total deviation problem with any penalty function  $F^i$  such as introduced in Section 2.1.

Theorem 8 solves the two-part type MDJIT problem in polynomial time, in the following sense: at every instant  $k$ , the theorem allows to determine efficiently which part type should be produced at time  $k$ . Furthermore, the optimal value of the two-part type problem can be computed very easily, as shown by our next result.

**Theorem 9.** Let  $\Delta = D/\text{gcd}(d_1, D) = D/\text{gcd}(d_2, D)$ . The optimal value  $B^*$  of the objective function of the 2-part type MDJIT problem is

$$B^* = \frac{1}{\Delta} \left\lfloor \frac{\Delta}{2} \right\rfloor.$$

**Proof.** By Theorem 6, we know that  $B^* \geq 1/\Delta \lfloor \Delta/2 \rfloor$ . There remains to prove that  $B^* \leq 1/\Delta \lfloor \Delta/2 \rfloor$ , i.e., that  $|[kr_i] - kr_i| \leq 1/\Delta \lfloor \Delta/2 \rfloor$  for  $i = 1, 2$  and for every  $k$ .

If  $\Delta$  is even, then  $|kr_i - [kr_i]| \leq 0.5 = 1/\Delta \lfloor \Delta/2 \rfloor$ . Suppose now that  $\Delta$  is odd and let us prove that  $|[kr_i] - kr_i| \leq (\Delta - 1)/2\Delta$ .

After simplification, the ideal rate  $r_i = d_i/D$  can be rewritten as  $\delta_i/\Delta$ . Therefore, the deviation  $|[kr_i] - kr_i|$  is an integral multiple of  $1/\Delta$ , say  $|[kr_i] - kr_i| = c/\Delta$  with  $c \in \mathbb{N}$ . Suppose that  $c/\Delta = 0.5$ . Then,  $\Delta = 2c$ , in contradiction with the hypothesis that  $\Delta$  is odd.

Thus,  $c/\Delta < 0.5$ , which implies that  $2c < \Delta$  and hence  $2c \leq \Delta - 1$ . Therefore  $c/\Delta \leq (\Delta - 1)/2\Delta$  as required.  $\square$



## 8. Conclusions

In this paper, we have revisited one of the most basic scheduling models of JIT production systems. In spite of its apparent simplicity, this model is not completely understood, yet. In particular, its computational complexity is not exactly known. We have shown that the model is in co-NP and that it is polynomially solvable when the number of part-types is fixed, but its general version may still turn out to be either co-NP-complete or polynomially solvable. We believe that the algebraic characterization of feasible instances presented in Theorem 2 may provide a useful tool for the analysis of such issues. Finally, obtaining a full description of instances with small max-deviation also presents an interesting challenge for future research.

## Acknowledgements

The authors are grateful to Alexandre Grigoriev, Joris van de Klundert and András Sebő for useful discussions regarding the topic of this note, to Gerhard Woeginger for pointing out the relevance of balanced sequences, and to the anonymous referees for their constructive comments. This work was carried out while the first author was visiting the University of Liège in the framework of a postdoctoral project supported by the European Network DONET (contract number ERB TMRX-CT98-0202). The second author acknowledges the partial financial support of ONR (grant N00014-92-J-1375), NSF (grant DMS-98-06389), DIMACS, and research grants from the Natural Sciences and Engineering Research Council of Canada (NSERC).

## Appendix.

Greedy algorithm. One may want to suggest the following greedy algorithm for the MDJIT problem: at each instant  $k$ , produce the part for which  $x_{i,k-1}$  is the farthest from  $kr_i$ . But this algorithm is not optimal, as shown by the following simple example. Assume that 6 parts of type  $A$ , 6 parts of type  $B$  and 1 part of type  $C$  are to be produced. Then, the greedy algorithm would schedule the part of type  $C$  at time 5 since  $\frac{5}{13} - 0 > 5 * \frac{6}{13} - 2$ , thus leading to a maximum deviation of  $\frac{10}{13}$  (at time 6). But the optimal solution  $ABABABCABABAB$ , where part  $C$  is produced at time 7, has a maximum deviation of  $\frac{9}{13}$ . (The famous Alabama paradox similarly establishes the shortcomings of greedy approaches in the apportionment context.) Note also that the greedy algorithm is pseudo-polynomial, but not polynomial.

## References

- [1] E. Altman, B. Gaujal, A. Hordijk, Balanced sequences and optimal routing, J. ACM 47 (2000) 752–775.
- [2] M. Balinski, N. Shahidi, A simple approach to the product rate variation problem via axiomatics, Oper. Res. Lett. 22 (1998) 129–135.
- [3] M. Balinski, H.P. Young, The quota method of apportionment, Amer. Math. Monthly 82 (1975) 701–730.

- [4] J. Bautista, R. Companys, A. Corominas, A note on the relation between the product rate variation (PRV) problem and the apportionment problem, *J. Oper. Res. Soc.* 47 (1996) 1410–1414.
- [5] J.A. Bondy, U.S.R. Murty, *Graph Theory with Applications*, North-Holland, New York, 1976.
- [6] N. Brauner, Y. Crama, J. van de Klundert, A. Grigoriev, A note on the complexity of high-multiplicity optimization problems, Working Paper GEMME, University of Liège, Liège, 2001.
- [7] N. Brauner, V. Jost, W. Kubiak, On symmetric Fraenkel’s and small deviations conjecture, Working paper, Les cahiers du laboratoire Leibniz, No 54, Grenoble, France, 2002.
- [8] H. Gabow, R.E. Tarjan, A linear-time algorithm for a special case of disjoint set union, *J. Comput. System Sci.* 30 (1985) 209–221.
- [9] G. Gallo, An  $O(n \log n)$  algorithm for the convex bipartite matching problem, *Oper. Res. Lett.* 3 (1984) 31–34.
- [10] F. Glover, Maximum matching in a convex bipartite graph, *Naval Res. Logistics Quart.* 4 (1967) 313–316.
- [11] M. Grötschel, L. Lovász, A. Schrijver, *Geometric Algorithms and Combinatorial Optimization*, Springer, Berlin, 1993.
- [12] D.S. Hochbaum, R. Shamir, Strongly polynomial algorithms for the high multiplicity scheduling problem, *Oper. Res.* 39 (1991) 648–653.
- [13] R.R. Inman, R.L. Bulfin, Sequencing JIT mixed-model assembly lines, *Manage. Sci.* 37 (1991) 901–904.
- [14] M.Y. Kovalyov, W. Kubiak, J.S. Yeomans, A computational study of balanced JIT optimization algorithms, *INFOR* 39 (2001) 299–316.
- [15] W. Kubiak, Minimizing variation of production rates in just-in-time systems: a survey, *European J. Oper. Res.* 66 (1993) 259–271.
- [16] W. Kubiak, Cyclic just-in-time sequences are optimal, Preprint, Memorial University of Newfoundland, St. John’s, Canada, 2000.
- [17] W. Kubiak, S.P. Sethi, A note on level schedules for mixed-model assembly lines in just-in-time production systems, *Manage. Sci.* 37 (1991) 121–122.
- [18] W. Kubiak, S.P. Sethi, Optimal just-in-time schedules for flexible transfer lines, *Internat. J. Flexible Manuf. Systems* 6 (1994) 137–154.
- [19] W. Kubiak, G. Steiner, J.S. Yeomans, Optimal level schedules for mixed-model, multi level just-in-time assembly systems, *Ann. Oper. Res.* 69 (1997) 241–259.
- [20] H.W. Lenstra Jr., Integer programming with a fixed number of variables, *Math. Oper. Res.* 8 (1983) 538–548.
- [21] W. Lipski Jr., F.P. Preparata, Efficient algorithms for finding maximum matchings in convex bipartite graphs and related problems, *Acta Inform.* 15 (1981) 329–346.
- [22] J. Miltenburg, Level schedules for mixed-model assembly lines in just-in-time production systems, *Manage. Sci.* 35 (1989) 192–207.
- [23] Y. Monden, *Toyota Production System*, Institute of Industrial Engineers, Norcross, GA, 1983.
- [24] H.N. Psaraftis, A dynamic programming approach for sequencing groups of identical jobs, *Oper. Res.* 28 (1980) 347–359.
- [25] G. Steiner, J.S. Yeomans, Level schedules for mixed-model, Just-in-time processes, *Manage. Sci.* 39 (1993) 728–735.
- [26] G. Steiner, J.S. Yeomans, Optimal level schedules in mixed-model, multi level JIT assembly systems with pegging, *European J. Oper. Res.* 95 (1996) 38–52.
- [27] R. Tijdeman, On complementary triples of Sturmian bisequences, *Indagationes Math.* 7 (1996) 419–424.
- [28] R. Tijdeman, Exact covers of balanced sequences and Fraenkel’s conjecture, in: F. Halter-Koch, R.F. Tichy (Eds.), *Algebraic Number Theory and Diophantine Analysis*, Walter de Gruyter, Berlin, 2000, pp. 467–483.
- [29] R. Tijdeman, Fraenkel’s conjecture for six sequences, *Discrete Math.* 222 (2000) 223–234.