

7. Steele TW, Lanser J, Sangster N. Isolation of *Legionella longbeachae* serogroup I from potting mixes. *Appl Environ Microbiol* 1990; 56: 49–53.
8. O'Connor BA, Carman J, Eckert K et al. Does using potting mix make you sick? Results from a *Legionella longbeachae* case-control study in South Australia. *Epidemiol Infect* 2007; 135: 34–39.
9. Helbig JH, Benson RF, Pelaz C et al. Identification and serotyping of atypical *Legionella pneumophila* strains isolated from human and environmental sources. *J Appl Microbiol* 2007; 102: 100–105.
10. Velonakis EN, Kiouisi IM, Koutis C et al. First isolation of *Legionella* species, including *L. pneumophila* serogroup I, in Greek potting soils: possible importance for public health. *Clin Microbiol Infect* 2010; 16: 763–766.
11. Yu VL, Plouffe JF, Pastoris MC et al. Distribution of *Legionella* species and serogroups isolated by culture in patients with sporadic community-acquired legionellosis: an international collaborative survey. *J Infect Dis* 2002; 186: 127–128.
12. Wallis L, Robinson P. Soil as a source of *Legionella pneumophila* Sg I. *Aust N Z J Public Health* 2005; 29: 518–520.

Clonal dissemination of *Klebsiella pneumoniae* ST258 harbouring KPC-2 in Argentina

S. A. Gomez^{1,*†}, F. G. Pasteran^{1†}, D. Faccone¹, N. Tijet², M. Rapoport¹, C. Lucero¹, O. Lastovetska^{2,3,4}, E. Albornoz¹, M. Galas¹, KPC Group[‡], R. G. Melano^{2,3,4}, A. Corso¹ and A. Petroni¹

1) Servicio Antimicrobianos, Instituto Nacional de Enfermedades Infecciosas (INEI)-ANLIS 'Dr Carlos G. Malbrán', Ciudad Autónoma de Buenos Aires, Argentina, 2) Ontario Agency for Health Protection and Promotion, Public Health Laboratories, 3) Department of Laboratory Medicine & Pathobiology, University of Toronto and 4) Department of Microbiology, Mount Sinai Hospital, Toronto, ON, Canada

Abstract

The present work describes the abrupt emergence of *Klebsiella pneumoniae* carbapenemase (KPC) and characterizes the first 79 KPC-producing enterobacteria from Argentina (isolated from 2006 to 2010). The emergence of *bla*_{KPC-2} was characterized by two patterns of dispersion: the first was the sporadic occurrence in diverse enterobacteria from distant geographical regions, harbouring plasmids of different incompatibility groups and *bla*_{KPC-2} in an unusual genetic environment flanked by *ISKpn8-Δbla*_{TEM-1} and *ISKpn6*-like. *bla*_{KPC-2} was associated with IncL/M transferable plasmids; the second was the abrupt clonal spread of *K. pneumoniae* ST258 harbouring *bla*_{KPC-2} in Tn4401a.

Keywords: *bla*_{KPC-2}, Enterobacteriaceae, ST11, ST258, Tn4401a

Original Submission: 14 December 2010; **Revised Submission:** 3 June 2011; **Accepted:** 6 June 2011

Editor: P. Tassios

Article published online: 11 June 2011

Clin Microbiol Infect 2011; 17: 1520–1524

10.1111/j.1469-0691.2011.03600.x

Corresponding author: S. A. Gomez, Servicio Antimicrobianos, Instituto Nacional de Enfermedades Infecciosas (INEI)-ANLIS 'Dr Carlos G. Malbrán', Av. Velez Sarsfield 563, 1282AFF, Ciudad Autónoma de Buenos Aires, Argentina
E-mail: sgomez@anlis.gov.ar

*Research Career, Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET).

†Sonia A. Gomez and Fernando G. Pasteran contributed equally to this work.

‡Authors are listed in the Acknowledgements.

The emergence and rapid dissemination of *Klebsiella pneumoniae* carbapenemase (KPC)-producing Gram-negative bacilli has become an important therapeutic and infection control problem in the healthcare setting [1]. The worldwide spread of KPC-producing *K. pneumoniae* strains has revealed the successful dissemination of a major clone defined as sequence type (ST)258 [2–4]. *bla*_{KPC} has been commonly associated with Tn4401, which is possibly responsible for its acquisition [5,6]. A different *bla*_{KPC} environment was described in plasmids from Chinese enterobacteria, where *bla*_{KPC-2} was associated with a Tn801-like transposon [7]. Moreover, ST11 was demonstrated to be the dominant clone of KPC-producing *K. pneumoniae* in China, where ST258 was not detected [8].

We previously reported the emergence of *bla*_{KPC} in Enterobacteriaceae from Argentina (2006), and designed an algorithm to detect class A carbapenemases and metallo-β-lactamases at the level of the clinical microbiology laboratory [9,10]. Since 2008, as the National Reference Laboratory in Antimicrobial Resistance (NRLAR), we have used the algorithm to implement active surveillance of such carbapenemases among 432 laboratories across Argentina that participate in the National Quality Control Programme in Bacteriology (Argentinean Ministry of Health). All enterobacteria were screened through the algorithm: isolates that showed (i) imipenem inhibition zones ≤21 mm, (ii) resistance or intermediate resistance to expanded-spectrum cephalo-




FIG. 1. Species distribution of the 77 *bla*_{KPC-2}-producing isolates in the different time periods. The first two KPC-producing isolates collected in 2006 (*Citrobacter freundii* and *Klebsiella pneumoniae* NON-ST258) were excluded. For each time period, the number of hospitals where *bla*_{KPC-2}-producing isolates were recovered, and the numbers of *Kpn*-ST258, *Kpn*-NON-ST258 and other enterobacteria, are shown. *Kpn*, *K. pneumoniae*.

sporins and (iii) positive synergy between the disks of carbapenems and 3-aminophenyl-boronic acid were suspected of KPC production and sent to the NRLAR. Possible interference by AmpC was overcome by using the oxacillin-based modified Hodge test [11].

In this study, we included the first two KPC producers detected in Argentina in 2006 [9], plus the 77 consecutive isolates (one per patient) that were confirmed at the NRLAR by PCR as KPC producers (subsequently characterized as *bla*_{KPC-2} by sequencing). These 77 isolates were referred between January 2008 and July 2010 from 30 hospitals from five distant geographical regions (no. isolates/no. hospitals): Buenos Aires City (54/19), Buenos Aires Province (16/8), Mendoza (4/1), Neuquén (2/1), and Formosa (1/1).

From January 2008 to July 2009, isolates of *Citrobacter freundii* (*n* = 2), *K. pneumoniae* (*n* = 5), *Serratia marcescens* (*n* = 1), *Enterobacter cloacae* (*n* = 2) and *Escherichia coli* (*n* = 1) were recovered from six hospitals (four regions). However, from August 2009, the number of *bla*_{KPC-2} isolates and the number of hospitals affected increased dramatically, mostly owing to *K. pneumoniae* recovered from Buenos Aires City and Province (Fig. 1).

Pulsed-field gel electrophoresis (PFGE) was performed as previously described [12], and the results were analysed according to the van Belkum guidelines [13]. Five types were observed among the 69 *K. pneumoniae* isolates: K1 (63 isolates); K2 (three isolates); and K3–K5 (one isolate each). No genotypic relationship was found among the three *C. freundii* or the five *E. cloacae* isolates.

Multilocus sequence typing (MLST) [14] was performed according to the MLST Database (<http://www.pasteur.fr/recherche/genopole/PF8/mlst/>). Six representative strains of

K. pneumoniae PFGE type K1 (different hospitals, isolated in 2008–2010) were identified as ST258. The minor PFGE types belonged to three STs: K2 and K5 to ST11, a single-locus variant of ST258; K3 to ST476; and K4 to the new ST526 described here.

The genetic surroundings of *bla*_{KPC-2} were studied by PCR mapping and sequencing (see primers used in the Supporting Information). Two different genetic environments were found (Fig. 2). All of the 63 *K. pneumoniae* clone K1/ST258 isolates harboured *bla*_{KPC-2} in the Tn4401a isoform [5]. To analyse the remaining 16 clinical isolates that did not harbour Tn4401, we constructed *Eco*RI-based DNA libraries [15] from *C. freundii* M9169 [9] and *E. cloacae* M11180. The *bla*_{KPC-2} environment found was very similar to the so-called Variant I (the accession number was not provided), a variant of the region found in plasmid pKP048 from a Chinese *K. pneumoniae* isolate [7]. Therefore, we named the Argentinian *bla*_{KPC-2} environment Variant Ia (Fig. 2a). However, we found two key differences: a composite transposon (partial sequence) from plasmid pFBAOT6 of *Aeromonas punctata* [16], which disrupts *tnpA* of the Tn801-like transposon, and an additional 671-bp fragment between ISKpn8 and *bla*_{KPC-2}, which contains a truncated *bla*_{TEM-1}.

By PCR mapping (Fig. 2b), Variant Ia was detected in 14 of 16 isolates, whereas the remaining two showed the new Variant Ib (Fig. 2a). As a whole, Variants Ia and Ib were found in five enterobacterial species, including *K. pneumoniae* ST11, ST476, and ST526. Interestingly, ST11 was also described in two KPC-producing PFGE clones from Brazil, but these harboured a Tn4401 [17], in contrast to the ST11 Argentinian and Chinese [8] isolates. As essentially the same *bla*_{KPC-2} environment (Variants Ia and Ib) was found in different species,




FIG. 2. Genetic environment of *bla*_{KPC-2} in enterobacteria from Argentina. Genes (open reading frames) and their corresponding transcriptional orientations are indicated by horizontal broad arrows. Filled and empty triangles represent inverted repeats of insertion sequences (ISs) or Tn3-group transposons, respectively. (a) The Tn4401a sequence was from plasmid pNYC (GenBank accession number EU176011) and the region of the plasmid pKP048 depicted corresponds to coordinates 15 736 to 25 596 (complement) in GenBank FJ628167. Variant 1a represents the sequence obtained from *Citrobacter freundii* M9169 and *Enterobacter cloacae* M11180 [9] (10 047 bp, GenBank JN048639 and JN048640, respectively). Δbla_{TEM-1} indicates a deletion of 10 bp upstream of the *bla_{TEM-1}* start plus the first 291 nucleotides of the encoding gene for the Tn3 sequence. Only the 58-bp deleted fragment in Variant 1b is shown, indicating the Tn801-like inverted repeat right (underlined text) and its A/T-rich pentameric target site duplication (TSD). The grey-shaded areas show identical regions among the compared structures. Thick lines with single or double arrowheads indicate the different ISs or transposons found, and thin lines with double arrowheads indicate the identity with plasmid pFBAOT6 from *Aeromonas punctata* (GenBank CR376602). (b) PCR mapping used for characterizing Tn4401a-harbours (upper) or Variant 1a/b-harbours (lower). The amplicons obtained in the different PCR assays and the lengths (bp) of the different amplicons are represented by thick lines. The letters and numbers at the left of each line indicate the different PCR used (primers are listed in the Supporting Information). Amplicons 7 and 8 were sequenced in all isolates to discriminate between Variants 1a and 1b.

we speculated that a unique plasmid could be responsible for such interspecies dissemination. However, a plasmid content analysis [18] of the 16 isolates harbouring Variants 1a and 1b

showed a range of plasmid sizes (7–80 kb), with several isolates harbouring more than one plasmid (data not shown). In addition, a Southern blot analysis with a *bla*_{KPC} probe showed

that *bla*_{KPC-2} was associated with plasmids of different sizes (data not shown). Therefore, PCR-based replicon typing [19] was performed on the 16 clinical isolates. We detected a single Inc group in 11 isolates: IncHI2 (*n* = 3), IncL/M (*n* = 3), and IncA/C (*n* = 5). Two isolates produced amplicons for three Inc groups (*Escherichia coli*, IncFIA–IncFIB–IncFrepB; *K. pneumoniae*, K5/ST11, IncL/M–IncA/C–IncFIIS) and three isolates gave negative results with all of the Inc primers tested (see Supporting Information).

Biparental conjugation [15] of six representative ST258 isolates (from different hospitals) did not yield transconjugants. Interestingly, three of 11 selected isolates (different hospitals and species) harbouring Variants 1a and 1b yielded transconjugants: two *C. freundii* (Variants 1a and 1b) and one *S. marcescens* (Variant 1a). β -Lactam MICs [20] of the transconjugants were lower than those of the parental strains (MIC range, mg/L): cefotaxime, 1–16 vs. 8–64; ceftazidime, 2–16 vs. 4–64; imipenem, 0.5–2 vs. 2–16; meropenem, 0.125–0.5 vs. 2–16; and ertapenem 0.25–1 vs. 2–8 mg/L. Replicon typing analysis allowed us to associate *bla*_{KPC-2} with IncL/M transferable plasmids in two transconjugants, whereas the remaining one was negative for all of the Inc primers tested, as was its parental strain.

The present work shows the results of a 3-year study (2008–2010) conducted in Argentina by the NRLAR to detect, characterize and warn of KPC emergence. The dissemination of *bla*_{KPC-2} occurred in two phases (Fig. 1): (i) from January 2008 to July 2009, with the sporadic occurrence of *bla*_{KPC-2} in diverse enterobacteria obtained from distant locations; and (ii) since August 2009, with a six-fold increase in the number of *bla*_{KPC-2} isolates, 94% of which were ST258. Therefore, the abrupt dissemination of KPC in Argentina could be explained by the surge of a unique clone of the international ST258. Moreover, the fact that Tn4401a was only found among these isolates reinforces the notion of clonal expansion in our hospital settings. Conversely, phase I of the local KPC emergence was driven by a diversity of enterobacterial species harbouring plasmids of several incompatibility groups. However, these isolates harboured *bla*_{KPC-2} in a unique and unusual genetic environment (Variants 1a and 1b) that was transferable by conjugation. These facts strongly suggest that not only plasmid transferability but also *bla*_{KPC-2} mobilization among different plasmids (i.e. transposition) could have played a major role in phase I of the Argentinean KPC emergence, leading to dissemination of Variants 1a and 1b in diverse clones and species.

To conclude, the actual KPC epidemiology in Argentina is complex and diverse. Cooperative efforts and strict infection control policies are still required in all institutions afflicted by dissemination of KPC.

The nucleotide sequences obtained here have been assigned GenBank accession numbers JN048639 and JN048640 (Variant 1a), and JN048641 and JN048642 (Variant 1b).

Acknowledgements

This study was supported in part by a grant from ANPCYT (PICT 2008-0286) Buenos Aires, Argentina, to S. A. Gomez, by the regular Federal Budget of the Ministry of Health of Argentina. S. A. Gomez is a member of the Carrera del Investigador Científico, CONICET, Argentina. We specially thank A. Carattoli for the kind provision of the positive controls for replicon typing experiments.

KPC Group participants include the following: Clínica AMEB PBA, A. Perez Ghiglia; Clínica y Maternidad Suizo Argentina, A. Rodríguez; FLENI, N. Orellana; Hospital General de Agudos 'Dr T. Alvarez', L. Rodríguez; Hospital Zonal General de Agudos 'Prof. Dr Ramón Carrillo', Ciudadela, N. Dainese; Hospital Central de San Isidro, M. E. Ochiuzzi; Hospital de Vicente López, V. Berdiñas; Hospital Español de Mendoza, R. A. Bucciarelli; Hospital General de Agudos 'Dr J. A. Fernández', L. Errecalde; Hospital General de Agudos 'Dr C. Argerich', N. Gomez; Hospital General de Agudos 'P. Piñero', F. Amalfa; Hospital Italiano, G. Grecco; Hospital de la Madre y el Niño de Formosa, M. S. Vivaldo; Hospital Militar Central, S. Perret; Hospital Nacional 'Prof. A. Posadas', A. Di Bella; Hospital General de Agudos 'J. A. Penna', M. C. Perrone; Hospital Provincial de Neuquén 'Dr Castro Rendón', M. R. Nuñez; Hospital General de Agudos 'J. M. Ramos Mejía', A. P. Meo; Hospital General de Agudos 'Donación F. Santojanni', C. Alfonso; Hospital de Gastroenterología 'B. Udaondo', C. Brotto; Hospital Universitario Fundación Favaloro, P. Andres; Clínica Basilea, P. Andres; Hospital Universitario Austral, V. Vilches; Laboratorios Clínicos Menéndez, G. Cervantes; Sanatorio de la Trinidad San Isidro, M. Orduña; Sanatorio Méndez, A. Vinante; Sanatorio de la Trinidad Palermo, D. Stepanik; Sanatorio de la Trinidad Mitre, A. Di Martino; Sanatorio de la Providencia, E. Berardinelli.

Transparency Declaration

The authors have no conflicts of interest to declare.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Figure S1. Pulse field gel electrophoresis patterns of *bla*_{KPC-2} isolates.

Table S1. Primers used in PCR mapping of *bla*_{KPC-2} genetic environments.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

References

- Hirsch EB, Tam VH. Detection and treatment options for *Klebsiella pneumoniae* carbapenemases (KPCs): an emerging cause of multidrug-resistant infection. *J Antimicrob Chemother* 2010; 65: 1119–1125.
- Kitchel B, Rasheed JK, Patel JB *et al.* Molecular epidemiology of KPC-producing *Klebsiella pneumoniae* isolates in the United States: clonal expansion of multilocus sequence type 258. *Antimicrob Agents Chemother* 2009; 53: 3365–3370.
- Leavitt A, Carmeli Y, Chmelnitsky I, Goren MG, Ofek I, Navon-Venezia S. Molecular epidemiology, sequence types, and plasmid analyses of KPC-producing *Klebsiella pneumoniae* strains in Israel. *Antimicrob Agents Chemother* 2010; 54: 3002–3006.
- Lopez JA, Correa A, Navon-Venezia S *et al.* Intercontinental spread from Israel to Colombia of a KPC-3-producing *Klebsiella pneumoniae* strain. *Clin Microbiol Infect* 2011; 17: 52–56.
- Naas T, Cuzon G, Villegas MV, Lartigue MF, Quinn JP, Nordmann P. Genetic structures at the origin of acquisition of the β -lactamase *bla*_{KPC} gene. *Antimicrob Agents Chemother* 2008; 52: 1257–1263.
- Nordmann P, Cuzon G, Naas T. The real threat of *Klebsiella pneumoniae* carbapenemase-producing bacteria. *Lancet Infect Dis* 2009; 9: 228–236.
- Shen P, Wei Z, Jiang Y *et al.* Novel genetic environment of the carbapenem-hydrolyzing β -lactamase KPC-2 among Enterobacteriaceae in China. *Antimicrob Agents Chemother* 2009; 53: 4333–4338.
- Qi Y, Wei Z, Ji S, Du X, Shen P, Yu Y. ST11, the dominant clone of KPC-producing *Klebsiella pneumoniae* in China. *J Antimicrob Chemother* 2011; 66: 307–312.
- Pasteran FG, Otaegui L, Guerriero L *et al.* *Klebsiella pneumoniae* carbapenemase-2, Buenos Aires, Argentina. *Emerg Infect Dis* 2008; 14: 1178–1180.
- Pasteran F, Mendez T, Guerriero L, Rapoport M, Corso A. Sensitive screening tests for suspected class A carbapenemase production in species of Enterobacteriaceae. *J Clin Microbiol* 2009; 47: 1631–1639.
- Pasteran F, Mendez T, Rapoport M, Guerriero L, Corso A. Controlling false-positive results obtained with the Hodge and Masuda assays for detection of class A carbapenemase in species of Enterobacteriaceae by incorporating boronic acid. *J Clin Microbiol* 2010; 48: 1323–1332.
- Melano R, Corso A, Petroni A *et al.* Multiple antibiotic-resistance mechanisms including a novel combination of extended-spectrum β -lactamases in a *Klebsiella pneumoniae* clinical strain isolated in Argentina. *J Antimicrob Chemother* 2003; 52: 36–42.
- van Belkum A, Tassios PT, Dijkshoorn L *et al.* Guidelines for the validation and application of typing methods for use in bacterial epidemiology. *Clin Microbiol Infect* 2007; 13 (suppl 3): 1–46.
- Diancourt L, Passet V, Verhoef J, Grimont PA, Brisse S. Multilocus sequence typing of *Klebsiella pneumoniae* nosocomial isolates. *J Clin Microbiol* 2005; 43: 4178–4182.
- Tijet N, Andres P, Chung C *et al.* rmtD2, a new allele of a 16S rRNA methylase gene, has been present in Enterobacteriaceae isolates from Argentina for more than a decade. *Antimicrob Agents Chemother* 2011; 55: 904–909.
- Rhodes G, Parkhill J, Bird C *et al.* Complete nucleotide sequence of the conjugative tetracycline resistance plasmid pFBAOT6, a member of a group of IncU plasmids with global ubiquity. *Appl Environ Microbiol* 2004; 70: 7497–7510.
- Cuzon G, Naas T, Truong H *et al.* Worldwide diversity of *Klebsiella pneumoniae* that produce β -lactamase *bla*_{KPC-2} gene. *Emerg Infect Dis* 2010; 16: 1349–1356.
- Nakamura M, Sato S, Ohya T, Suzuki S, Ikeda S. Plasmid profile analysis in epidemiological studies of animal *Salmonella typhimurium* infection in Japan. *J Clin Microbiol* 1986; 23: 360–365.
- Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 2005; 63: 219–228.
- Clinical and Laboratory Standards Institute. *Performance standards for antimicrobial susceptibility testing; twentieth informational supplement*. CLSI M100-S20-U Vol. 30 No. 15. Wayne, PA: CLSI, 2010.

Genetic diversity of carbapenem-hydrolysing β -lactamases in *Acinetobacter baumannii* from Romanian hospitals

R. A. Bonnin¹, L. Poirel¹, M. Licker² and P. Nordmann¹

- 1) Service de Bactériologie-Virologie, INSERM U914 'Emerging Resistance to Antibiotics', Hôpital de Bicêtre, Assistance Publique/Hôpitaux de Paris, Faculté de Médecine et Université Paris-Sud, K. Bicêtre, France and
- 2) Department of Microbiology, University of Medicine and Pharmacy Victor Babes, Timisoara, Romania

Abstract

Thirteen carbapenem-resistant *Acinetobacter baumannii* isolates, collected in Romania during 2009–2010, were investigated to identify the mechanism(s) responsible for carbapenem resistance. Genotyping was performed by pulsed-field gel electrophoresis, multiplex PCR sequence typing and multilocus sequence typing. Eleven non-clonally related isolates harboured the *bla*_{OXA-23} gene on their chromosome within a Tn2008 transposon structure. The two remaining isolates harboured a *bla*_{OXA-58} gene that was either plasmid or chromosome borne. Two isolates co-expressed OXA-23 together with the extended-spectrum β -lactamase PER-1. This study constitutes the first report of OXA-58 and OXA-23-producing *A. baumannii* isolates in Romania.

Keywords: *Acinetobacter baumannii*, carbapenemase, extended-spectrum β -lactamase, OXA-23, OXA-58