

Contents lists available at [ScienceDirect](http://ScienceDirect)

## Physics Letters B

[www.elsevier.com/locate/physletb](http://www.elsevier.com/locate/physletb)

## Supermodels for early LHC

Christian W. Bauer<sup>a,b</sup>, Zoltan Ligeti<sup>a,b,\*</sup>, Martin Schmaltz<sup>a,b,c</sup>, Jesse Thaler<sup>a,b</sup>, Devin G.E. Walker<sup>a,b,d</sup><sup>a</sup> Theoretical Physics Group, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, United States<sup>b</sup> Berkeley Center for Theoretical Physics, University of California, Berkeley, CA 94720, United States<sup>c</sup> Physics Department, Boston University, Boston, MA 02215, United States<sup>d</sup> Center for the Fundamental Laws of Nature, Jefferson Physical Laboratory, Harvard University, Cambridge, MA 02138, United States

## ARTICLE INFO

## Article history:

Received 24 March 2010

Received in revised form 13 May 2010

Accepted 13 May 2010

Available online 20 May 2010

Editor: B. Grinstein

## Keywords:

Supermodels

Early LHC

## ABSTRACT

We investigate which new physics signatures could be discovered in the first year of the LHC, beyond the expected sensitivity of the Tevatron data by the end of 2010. We construct “supermodels”, for which the LHC sensitivity even with only  $10 \text{ pb}^{-1}$  useful luminosity is greater than that of the Tevatron with  $10 \text{ fb}^{-1}$ . The simplest supermodels involve  $s$ -channel resonances in the quark–antiquark and especially in the quark–quark channels. We concentrate on easily visible final states with small standard model backgrounds, and find that there are simple searches, besides those for  $Z'$  states, which could discover new physics in early LHC data. Many of these are well-suited to test searches for “more conventional” models, often discussed for multi- $\text{fb}^{-1}$  data sets.

© 2010 Elsevier B.V. Open access under [CC BY license](http://creativecommons.org/licenses/by/3.0/).

## 1. Introduction

In this Letter, we explore the new physics discovery potential of the first LHC run. The latest schedule calls for collisions at 7 TeV throughout much of 2010 and 2011, with the hope of delivering about  $100 \text{ pb}^{-1}$  integrated luminosity in 2010 and  $1 \text{ fb}^{-1}$  by the end of 2011 [1,2]. However, the useful luminosity for physics analyses may be less than the delivered luminosity projected, especially during 2010. We therefore find it interesting to study the sensitivity of the first run as a function of LHC luminosity and energy.

In particular, it has been stated that the first tens of  $\text{pb}^{-1}$  of good data to be analyzed by ATLAS and CMS would essentially correspond to an “engineering run” with only the capability to “rediscover” the standard model. Generally, the expectation is that order  $100 \text{ pb}^{-1}$  of good data will be necessary for the LHC to have sensitivity to plausible new physics scenarios. Here we take a fresh look at the new physics capabilities of a  $10 \text{ pb}^{-1}$  low-luminosity data set, and allow ourselves to contemplate new physics which is not motivated by model building goals such as unification, weak scale dark matter, or solving the hierarchy problem.

We find that, setting such model building prejudices aside, there is a set of interesting new physics scenarios that could give rise to a clean, observable signal in early LHC data, while not being detected with  $10 \text{ fb}^{-1}$  of Tevatron data (roughly the pro-

jected integrated luminosity at the end of 2010). These models are consistent with previous experiments such as LEP II, precision electroweak constraints, and flavor physics. Moreover, these scenarios have similar signatures to “well-motivated” new physics models that require higher luminosity for discovery.

To set the stage, recall that the production cross sections for new hypothetical particles can be quite large. For example, QCD pair production of 500 GeV colored particles have cross sections in the pb range, such that tens of such particles could be produced in early LHC. Of course, in order for the new particles to be observable, they must have sufficiently large branching fractions to final states with distinctive signatures and controllable standard model backgrounds. Also, the new particles should not be ruled out by current or future Tevatron searches, implying that the cross section times integrated luminosity at the LHC should be larger than the corresponding quantity at the Tevatron.


Thus, the four criteria for a new physics scenario to be discovered in early LHC with low luminosity are:

1. Large enough LHC cross section to produce at least 10 signal events<sup>1</sup> with  $10 \text{ pb}^{-1}$  of data;
2. Small enough Tevatron cross section to evade the projected 2010 Tevatron sensitivity with  $10 \text{ fb}^{-1}$ ;
3. Large enough branching fraction to an “easy” final state with essentially no backgrounds;
4. Consistency with other existing bounds.

\* Corresponding author at: Theoretical Physics Group, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, United States.

E-mail address: [zligeti@lbl.gov](mailto:zligeti@lbl.gov) (Z. Ligeti).

<sup>1</sup> While fewer events might be sufficient for discovery, we shall demand 10 events to allow for  $\mathcal{O}(1)$  uncertainties in our analysis.



**Fig. 1.** LHC parton luminosities as defined in Eq. (1), as functions of the partonic invariant mass. The solid (dashed) curves are for the 7 TeV (10 TeV) LHC. The up quark has been chosen as a representative quark, and each curve includes the contribution from the  $CP$  conjugate initial partons.

We call a new physics scenario satisfying these conditions a *supermodel*.


The classic example that comes to mind as a candidate supermodel is a TeV-scale  $Z'$  boson [3]. Assuming the  $Z'$  mass exceeds the Tevatron reach, but is light enough and has large enough couplings so that it can be produced copiously at the LHC, then it can be discovered through its decay to electron and muon pairs. Such leptonic final states are “easy” to reconstruct with a peak in the invariant mass distribution, which reduces the already low standard model backgrounds.

As we will see, however, a typical leptonically decaying  $Z'$  is actually not a supermodel. First, since the  $Z'$  is produced via the quark–antiquark initial state, the Tevatron is quite competitive with the LHC. Second, the leptonic branching fraction is severely bounded by LEP II data, which restricts the couplings of the  $Z'$  to leptons. It is therefore nontrivial to find supermodels that are as discoverable as a standard  $Z'$  but consistent with known bounds on new physics.

The remainder of the Letter is organized as follows. In Section 2, we identify new particle production channels with sufficiently large LHC cross sections and for which the LHC has an advantage over the Tevatron. Assuming perturbative couplings, we find that  $s$ -channel production of quark–quark ( $qq$ ) or quark–antiquark ( $q\bar{q}$ ) resonances are the best starting points for early LHC supermodels. In Section 3, we construct explicit models where these resonances can decay to interesting and easily reconstructable final states. While a standard  $Z'$  does not work, generalized  $Z'$  scenarios can be supermodels, as are scenarios involving diquarks. We conclude in Section 4.

## 2. Production modes

In this section, we discuss which production modes have the potential to be supermodels, deferring detailed model building to Section 3. Since the expected integrated luminosity at the Tevatron ( $\sim 10 \text{ fb}^{-1}$ ) is orders of magnitude larger than our  $10 \text{ pb}^{-1}$  benchmark luminosity for early LHC analysis, and since  $p\bar{p}$  parton luminosities are not so different from  $pp$  parton luminosities, one must consider sufficiently heavy new particles to evade the Tevatron reach. We will find that the most promising perturbative



**Fig. 2.** Ratios of the parton luminosities for 7 TeV (solid) and 10 TeV (dashed) LHC compared to the 1.96 TeV Tevatron, as functions of the partonic invariant mass. When this ratio is above the  $10^3$  horizontal dashed line, the LHC with  $10 \text{ pb}^{-1}$  will have greater sensitivity than the Tevatron with  $10 \text{ fb}^{-1}$ .

scenarios accessible with  $10 \text{ pb}^{-1}$  of LHC data are  $qq$  and  $q\bar{q}$  resonances.

To begin, we plot in Fig. 1 the LHC parton luminosities, defined as


$$\mathcal{F}_{ij}(\hat{s}, s) = \int_{\hat{s}/s}^1 dx_i \frac{\hat{s}}{x_i s} f_i(x_i) f_j[\hat{s}/(x_i s)], \quad (1)$$

and in Fig. 2 the ratios of each parton luminosity at the LHC and the Tevatron. In Eq. (1),  $\sqrt{s}$  is the center of mass energy of the collider,  $\sqrt{\hat{s}}$  is the invariant mass of the two interacting partons, and  $f_i(x_i)$  are the parton distribution functions evaluated at a momentum fraction  $x_i$  and scale  $\sqrt{\hat{s}}$ . We use the CTEQ-5L parton distribution functions [4] implemented in Mathematica. (For similar plots using CTEQ-6L1 [5], see Ref. [6].) We checked that MSTW 2008 [7] gives fully compatible results at the level of accuracy we require.

It is often stated that the LHC is essentially a gluon collider, so one might think that processes initiated by gluons would be the best starting points for constructing supermodels. However, Fig. 1 shows that the  $gg$  parton luminosity only dominates for small invariant mass, where the initial LHC data set cannot compete with the Tevatron. As seen in Fig. 2, only at large invariant masses do the LHC parton luminosities become sufficiently enhanced compared to the Tevatron. (The enhancement of the  $q\bar{q}$  channel is the smallest, so it is harder for the LHC to compete in cases where the initial  $q\bar{q}$  state contributes.) To build supermodels, we must explore the possible LHC cross sections in the region with large enough enhancements compared to the Tevatron. We will emphasize this point in the next subsection by showing why QCD pair production is not a supermodel, and then go on to consider supermodels constructed from  $s$ -channel resonances.

### 2.1. QCD pair production?

A simple process initiated by gluons is QCD pair production of new colored particles. For not too heavy states, it can have a cross section above a pb, yielding  $\mathcal{O}(10)$  events with  $10 \text{ pb}^{-1}$  of LHC data. However, it is easy to show that such processes are generi-



**Fig. 3.** LHC reach for pair production of a single flavor of heavy quark as a function of energy and luminosity. Each contour corresponds to the production of 10 events at the LHC for the indicated quark mass. The red region corresponds to quark masses which the Tevatron would be able to rule out with  $10 \text{ fb}^{-1}$ , because it would produce 10 or more events. The intersection of the straight dashed lines touches the contour corresponding to the maximum quark mass ( $\sim 400 \text{ GeV}$ ) probed by the 7 TeV LHC with  $10 \text{ pb}^{-1}$  of data. One sees that the early LHC is generically not sensitive to QCD pair production of quarks with masses beyond the Tevatron reach. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this Letter.)

cally not supermodels. For concreteness, we study the production of a color-triplet quark  $Q$ . We assume that it always decays to a highly visible final state, and that reconstruction efficiencies are perfect. One can then use the standard QCD diagrams to calculate the largest value of  $m_Q$  for which the Tevatron would observe  $10 Q\bar{Q}$  pair production events with  $10 \text{ fb}^{-1}$  of data. In this idealized example, the hypothetical Tevatron bound is  $m_Q \gtrsim 500 \text{ GeV}$ . The same exercise can be repeated for the LHC as a function of the center of mass energy and integrated luminosity, and the result is shown in Fig. 3.

To reach the Tevatron sensitivity for QCD pair production at a 7 TeV LHC, the required luminosity is about  $50 \text{ pb}^{-1}$ . While this is likely within the reach of an early LHC run, the LHC will not easily surpass Tevatron bounds in this channel, and it is unlikely that a  $5\sigma$  LHC discovery is possible without the Tevatron already having seen some events. (The same holds for colored scalar pair production [8–10].) This conclusion is only bolstered when realistic branching fractions to visible final states and signal efficiencies are taken into account.

The primary reason why QCD pair production is not a supermodel is that the same final state can also be produced from the  $q\bar{q}$  initial state, where the LHC has less of an advantage over the Tevatron. The situation can be improved if there is a large multiplicity of near-degenerate new colored states or if the new states are color octets (like gluinos in supersymmetry). Then the total cross sections are larger by a multiplicity factor and the LHC reach can surpass that of the Tevatron (where the cross section is more strongly suppressed at higher masses). As an example, leptoquark pair production [11] yields the easily reconstructable final state of two leptons and two jets, so this could be a supermodel with a sufficiently large multiplicity of such leptoquarks.

In any case, because QCD pair production is quite well-studied in specific new physics scenarios and because the early LHC advantage over the Tevatron can only be marginal, we will not consider it to be a supermodel in this Letter. In Section 3.4, though, we show that pair production through an intermediate resonance can give rise to supermodels.

## 2.2. Resonance production

While pair production of new colored particles is not a supermodel, production of an  $s$ -channel resonance has the potential to be a supermodel, as long as the resonance has renormalizable couplings to the partonic initial states. Recall that parametrically the production cross section for a single resonance is enhanced over pair production by a phase space factor of  $16\pi^2$ . Moreover, unlike QCD pair production where  $SU(3)$  gauge invariance relates the  $gg$  and  $q\bar{q}$  scattering amplitudes, single resonance production can be dominated by one partonic initial state.

In the narrow width approximation, we can parametrize generic single resonance production by

$$\sigma(p_i p_j \rightarrow X) = [g_{\text{eff}}^2]_{ij} \delta(\hat{s} - m_X^2), \quad (2)$$

where  $p_{i,j}$  denote the two partons which participate in the hard scattering,  $m_X$  is the mass of the resonance, and  $[g_{\text{eff}}^2]_{ij}$  encodes all information about the production of resonance  $X$  from the two partons, including couplings, polarization, and color factors. Using the parton luminosities defined in Eq. (1), the hadronic cross section is

$$\sigma(pp \rightarrow X) = \frac{1}{m_X^2} \sum_{ij} [g_{\text{eff}}^2]_{ij} \mathcal{F}_{ij}(m_X^2, s). \quad (3)$$

For the resonances considered in this Letter, one production channel dominates, allowing us to drop the  $ij$  label from  $g_{\text{eff}}^2$ .

For reasonably narrow resonances with dimension four couplings,  $g_{\text{eff}}^2$  can be order 1, which is the case for the  $q\bar{q}$  and  $qq$  initial states. However, for the  $gg$  or  $gg$  initial states  $SU(3)$  gauge invariance forbids renormalizable couplings to a single resonance. For example, for the  $gg$  initial state, the lowest dimension operator allowed is a dimension five coupling to a scalar or pseudoscalar:


$$\frac{g_s^2}{16\pi^2 \Lambda} X \text{Tr}(G_{\mu\nu} G^{\mu\nu}). \quad (4)$$

The coefficient of the operator has been estimated assuming perturbative physics at  $\Lambda \sim \text{TeV}$ , with the  $1/(16\pi^2)$  factor coming from a loop. This gives a rather suppressed effective coupling  $[g_{\text{eff}}^2]_{gg} \sim [1/(16\pi^2)(m_X/\Lambda)]^2 \sim [1/(16\pi^2)]^2$ , where we have assumed  $m_X$  is around a TeV. If there is strong dynamics involving  $X$  at the TeV scale, then the coefficient in Eq. (4) can be enhanced up to its naive dimensional analysis value  $g_s^2/(4\pi\Lambda)$ . However, such strong dynamics near the TeV scale is constrained by precision measurements, and we will adopt the perturbative estimate  $g_{\text{eff}}^2 \sim [1/(16\pi^2)]^2$  for both  $gg$  and  $qq$  resonances.

In Fig. 4, we show our estimate of the generic early LHC reach in  $m_X$ , as a function of the energy and luminosity, for the four resonance channels using

$$g_{\text{eff}}^2 = \begin{cases} 1, & qq \text{ or } q\bar{q} \text{ resonances,} \\ [1/(16\pi^2)]^2, & gg \text{ or } gg \text{ resonances.} \end{cases} \quad (5)$$

As in Fig. 3, we assume 100% branching fraction of  $X$  to highly visible final states and assume perfect detector efficiency, though we will relax these assumptions below. Note that for a charged resonance (that is produced from  $qq$  and  $qg$  initial states), the search strategy for the charge conjugate resonance arising from  $\bar{q}\bar{q}$  and  $\bar{q}g$




**Fig. 4.** LHC reach for single resonance production as a function of energy and luminosity. As in Fig. 3, the contours show the production of 10 events for a given resonance mass, the red regions show the Tevatron sensitivity with  $10 \text{ fb}^{-1}$ , and the intersection of the dashed lines shows the maximum resonance mass which can be probed by the 7 TeV LHC with  $10 \text{ pb}^{-1}$  data. The expected couplings for perturbative new physics in Eq. (5) are included. One sees that the early LHC can exceed the Tevatron sensitivity for  $q\bar{q}$  and especially for  $q\bar{q}$  resonances. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this Letter.)

initial states is equivalent, and any analysis will include both. The plots in Fig. 4 use the  $u$  quark parton distribution function for simplicity (rather than that of the  $d$  or both) and include the  $uu + \bar{u}\bar{u}$  and  $ug + \bar{u}\bar{g}$  initial states instead of  $uu$  and  $ug$  only.

We see that while gluons are by far the most abundant partons at small  $x$ , scattering initiated by gluons does not lead to very large resonance cross sections at the LHC, if perturbative couplings are assumed. In the  $u\bar{u}$  and especially in the  $uu$  channels, shown in the two upper plots of Fig. 4, the first LHC run even with modest energy and luminosity will supersede the Tevatron. Thus,  $q\bar{q}$  and  $q\bar{q}$  resonances are the most suitable starting points for constructing supermodels, examples of which will appear in Section 3.

### 2.3. Production of $q\bar{q}$ and $q\bar{q}$ resonances

The plots in Fig. 4 give a rough idea of the LHC discovery potential for  $s$ -channel resonances. They are valid for a particular value of the effective coupling,  $g_{\text{eff}}^2$ , and assume that the  $X$  resonance is observed with 100% efficiency. For the two most promising scenarios of  $q\bar{q}$  and  $q\bar{q}$  resonances, we are interested in the dependence of the reach on  $g_{\text{eff}}^2$  and on branching fractions/efficiencies. Here, we introduce a new kind of plot which is convenient for reading off cross sections at the LHC and comparing them to the Tevatron for variable couplings and detection efficiencies. In Fig. 5, we plot in the LHC energy vs. resonance mass plane the contours of con-



**Fig. 5.** The LHC reach for  $uu$  and  $u\bar{u}$  resonances in the LHC energy vs. resonance mass plane. The solid lines are contours of constant LHC production cross sections for  $g_{\text{eff}}^2 = 1$ , and the dashed green lines are contours of constant LHC to Tevatron cross section ratios. The blue shaded regions show where the discovery reach of a  $10 \text{ pb}^{-1}$  LHC run is beyond that of the  $10 \text{ fb}^{-1}$  Tevatron. The green regions show where the LHC sensitivity is greater than that of the Tevatron, but the Tevatron can also see at least 10 events.

stant production cross section and contours of constant ratio of LHC vs. Tevatron cross section.

The solid curves (with positive slopes) in Fig. 5 show contours of constant LHC cross sections for  $g_{\text{eff}}^2 = 1$ . From these, one can read off how many events are produced for a given LHC luminosity as a function of the resonance mass and the LHC energy. For example, assuming 100% visible decay rate and detection efficiency, the region to the right of the curve labeled “ $10^0 \text{ pb}$ ” will yield at least 10 events with  $10 \text{ pb}^{-1}$  of LHC data. For a concrete model, with a different value of  $g_{\text{eff}}^2$ , branching ratio  $\mathcal{B}$  to easily reconstructable final state(s), and detector/reconstruction efficiency  $\text{Eff}_{\text{LHC}}$ , the LHC with a given energy and  $\mathcal{L}_{\text{LHC}}$  luminosity can see  $N$  or more events to the right of the solid curve labeled

$$\frac{N}{\mathcal{L}_{\text{LHC}} \times g_{\text{eff}}^2 \times \mathcal{B} \times \text{Eff}_{\text{LHC}}} \quad (6)$$

For example, if  $g_{\text{eff}}^2 = 0.1$  and  $\mathcal{B} \times \text{Eff}_{\text{LHC}} = 10\%$ , then 10 or more events can be observed in the region to the right of the curve labeled “ $10^2 \text{ pb}$ ” (“ $10^1 \text{ pb}$ ”) with  $10 \text{ pb}^{-1}$  ( $100 \text{ pb}^{-1}$ ) of LHC data.

The dashed curves (with negative slopes) in Fig. 5 show contours of constant ratio of LHC vs. Tevatron cross sections. From these, one can read off the relative advantage of the LHC compared to the Tevatron for a given model. For many resonances, the same initial state dominates the production of a resonance at the Tevatron and the LHC, so that the factors of  $g_{\text{eff}}^2$  cancel in the ratio of the two cross section. If, in addition, the same final states are searched for at the Tevatron and at the LHC and the detection efficiencies are similar, then these cancel in the ratio as well. Therefore, the LHC has better sensitivity than the Tevatron in the region above the dashed curve labelled by

$$\mathcal{L}_{\text{TEV}}/\mathcal{L}_{\text{LHC}} \quad (7)$$

For example, the LHC with  $10 \text{ pb}^{-1}$  ( $100 \text{ pb}^{-1}$ ) will produce more events than the Tevatron with  $10 \text{ fb}^{-1}$  in the region above the dashed curve labeled “ $10^3$ ” (“ $10^2$ ”). Any significant differences in

detection efficiencies between the two colliders can easily be included by multiplying Eq. (7) by  $\text{Eff}_{\text{TEV}}/\text{Eff}_{\text{LHC}}$ .


Thus, it is a “wedge” bounded by a solid and a dashed curve which defines the region in which the LHC has better sensitivity than the Tevatron and yields at least a certain number of events. For example, the wedge to the right of the intersection of the “ $10^0 \text{ pb}$ ” and the “ $10^3$ ” curves gives the region for which at least 10 events are produced with  $10 \text{ pb}^{-1}$  of LHC data and the number of events at the LHC is greater than that at the Tevatron. These regions are shaded in Fig. 5. (To include model specific effects, replace the “ $10^0 \text{ pb}$ ” solid curve by the  $10^0 \text{ pb}/(g_{\text{eff}}^2 \mathcal{B} \text{Eff}_{\text{LHC}})$  one.)

At the intersection of a solid and a dashed curve, the ratio of their labels gives the Tevatron cross section, and can be used to estimate the Tevatron discovery reach. The intersection of any “ $10^{n+a} \text{ pb}$ ” solid curve with a “ $10^a$ ” dashed curve corresponds to the same fixed Tevatron cross section of  $10^n \text{ pb}$  for arbitrary  $a$ . Since the Tevatron cross section does not depend on the LHC energy, these intersections lie on a horizontal line. The corresponding value of the resonance mass is the one for which the Tevatron with  $10 \text{ fb}^{-1}$  data produces  $10^{4+n}$  events. For example, for masses below the straight line across the intersection of the “ $10^0 \text{ pb}$ ” and the “ $10^3$ ” curves (i.e.  $n = -3$ ), the Tevatron will also produce at least 10 events with  $10 \text{ fb}^{-1}$  data. While everywhere in the full wedges shaded in Fig. 5 the LHC sensitivity is better than that of the Tevatron, below this straight line the Tevatron sensitivity is good enough that it can make a discovery prior to the LHC.<sup>2</sup> Therefore, it is only the part of the “wedge” above this straight line, shaded blue, which is the true LHC discovery region.

Using these plots, one can also estimate the minimum value of  $m_X$  and  $g_{\text{eff}}^2 \mathcal{B} \text{Eff}_{\text{LHC}}$  for a scenario to be a supermodel. Take a  $q\bar{q}$  resonance as an example. A 7 TeV and  $10 \text{ pb}^{-1}$  early LHC

<sup>2</sup> For a sequential  $Z'$ , taking  $g_{\text{eff}}^2 \mathcal{B} \text{Eff}_{\text{LHC}} \sim 0.01$  for the decay to a lepton pair, this predicts a  $10 \text{ fb}^{-1}$  Tevatron bound near 1.2 TeV; crude, but reasonable [12].





**Fig. 6.** Three typical event topologies for weakly coupled resonance decay. Double lines denote new particles, while single lines are detectable final states, either standard model particles or new (quasi-)stable states. The ellipses indicate that the secondary resonances may decay to two or more particles.

run supersedes the Tevatron sensitivity for a  $m_X \gtrsim 1.4$  TeV (the value of the “ $10^3$ ” dashed curve at 7 TeV). We can then read off that  $g_{\text{eff}}^2 \mathcal{B} \text{Eff}_{\text{LHC}} \gtrsim 0.1$  is required to observe at least 10 events. For larger initial LHC luminosity, this minimum of course gets smaller.

### 3. Example supermodels

Considering production cross sections alone,  $q\bar{q}$  and  $qq$  resonances emerged as the two best starting points for constructing early LHC supermodels. In this section, we consider concrete supermodel examples to see what kind of final states can be obtained from the decay of these resonances. Since we are interested in final states that involve the cleanest signatures and least background contamination, we concentrate on decay chains yielding (at least) two charged leptons (or two other stable charged particles) in the final state.

Because there is a plethora of possible decay patterns for a  $qq$  or  $q\bar{q}$  resonance, we find it convenient to classify the decay modes of new resonances in terms of three basic decay topologies, which often appear in perturbative new physics scenarios. These are depicted in Fig. 6. Double lines denote new resonances and solid lines are detectable final states, either standard model particles or new (quasi-)stable states.

- Topology A: The resonance  $X$  decays directly to two detectable final state particles. Note that a three-body decay for the resonance will generically not compete with the decay back to QCD partons, since the branching fraction to a three-body final state is suppressed by a phase space factor compared to the two-body decay channel back to the initial state.
- Topology B: The resonance  $X$  decays to one detectable final state and one new secondary resonance  $Y$ . Subsequently  $Y$  can decay to two or more standard model particles. Note that  $Y$  always has a decay mode back through the production diagram with a virtual  $X$ .
- Topology C: The resonance  $X$  decays to two new secondary resonances  $Y_1$  and  $Y_2$ , possibly of different masses. These secondary resonances can each decay to two or more detectable states. This topology can be used for example to “resurrect” pair production of colored particles as a supermodel.

The classic case of topology A is a leptonically decaying  $Z'$ , and this is often described as the simplest new physics channel to discover at a hadron collider. However, we will argue below that a

standard  $Z'$  is not a supermodel if one takes into account the indirect constraints coming from LEP II. After dismissing the standard  $Z'$  case, we will give a (non-exhaustive) list of supermodels exhibiting the three above topologies.

While some of the example supermodels are simple variants on well-motivated models, the case of a  $qq$  resonance (or diquark) is less well-known (though we will show that such a particle can arise in the minimal supersymmetric standard model; see also Ref. [13]). Despite the unfamiliar quantum numbers of the diquark, the final states achievable in diquark decays can be shared by more familiar resonances, albeit with smaller non-supermodel cross sections. In this way, searches for supermodels in early LHC data can anticipate searches that require higher luminosity.

For reference, the standard model quantum numbers are:

Field	Spin	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$
$q$	1/2	$\mathbf{3}$	$\mathbf{2}$	1/6
$u^c$	1/2	$\bar{\mathbf{3}}$	–	–2/3
$d^c$	1/2	$\bar{\mathbf{3}}$	–	1/3
$l$	1/2	–	$\mathbf{2}$	–1/2
$e^c$	1/2	–	–	1
$h$	0	–	$\mathbf{2}$	1/2

#### 3.1. The case against a standard $Z'$

For a  $q\bar{q}$  resonances to be supermodel, it must have a large branching fraction to visible final states. In particular, since a  $q\bar{q}$  resonance can have zero electric charge, it is natural for such a resonance to decay via topology A to pairs of oppositely charged leptons, in particular  $e^+e^-$  and  $\mu^+\mu^-$ . However, the same resonance also induces a low energy effective four-lepton vertex, and such operators are severely constrained by LEP II. As recently emphasized in Ref. [14], once the LEP II bound is imposed, the branching fraction of the  $q\bar{q}$  resonance to  $l^+l^-$  has to be too small to realize a supermodel.

As a concrete example, consider a new  $U(1)$  gauge boson  $Z'$ :

Field	Spin	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$
$Z'$	1	–	–	–

with couplings to standard model fermions  $\psi_i$  of the form<sup>3</sup>

$$\mathcal{O}_{Z'} = g_i Z'_\mu \bar{\psi}_i \bar{\sigma}^\mu \psi_i, \quad (10)$$

where  $g_i$  are the corresponding coupling constants. For simplicity, imagine a common coupling  $g_Q$  to quarks and  $g_L$  to leptons.<sup>4</sup>

In the narrow width approximation, the production cross section for this  $Z'$  resonance is

$$\sigma(q\bar{q} \rightarrow Z') = \frac{\pi}{3} g_Q^2 \delta(\hat{s} - m_{Z'}^2), \quad (11)$$

and the branching ratio of this  $Z'$  to charged leptons is<sup>5</sup>

$$\mathcal{B}(Z' \rightarrow l^+l^-) = \frac{2}{3} \frac{g_L^2}{2g_L^2 + 6g_Q^2}. \quad (12)$$

The coupling to  $Z'$  to leptons is strongly constrained by LEP II limits on four-fermion operators [15],

<sup>3</sup> Here and below, we use 2-component fermion notation.

<sup>4</sup> While such a choice of  $Z'$  charges is typically anomalous (unless  $g_Q = -3g_L$ ), such anomalies can always be canceled with new fermions (spectators) whose masses result from the same  $U(1)$  symmetry breaking which gives the  $Z'$  its mass.

<sup>5</sup> We are assuming Dirac neutrino masses for definiteness, and we do not consider final state  $\tau$ 's to be easily reconstructable, hence the overall factor of 2/3.

$$\frac{g_L^2}{m_{Z'}^2} \leq \frac{4\pi}{(10 \text{ TeV})^2}. \quad (13)$$

Putting this together, we find

$$g_{\text{eff}}^2 \times \mathcal{B} = \frac{2\pi}{9} \frac{g_Q^2 g_L^2}{2g_L^2 + 6g_Q^2} \lesssim \left( \frac{m_{Z'}}{8 \text{ TeV}} \right)^2. \quad (14)$$

Using Fig. 5 and Eq. (6), one finds that there is no value of  $m_{Z'}$  for which  $10 Z' \rightarrow \ell^+ \ell^-$  events could be seen with  $10 \text{ pb}^{-1}$  of data, even with a center of mass energy of 14 TeV.

There are ways to evade this conclusion. Since the LEP II bound only applies for the electron coupling, one could imagine coupling the  $Z'$  only to muons. However, such flavor non-universal couplings typically require significant fine-tuning to avoid constraints from flavor changing neutral currents. Alternatively, one might consider the production of a  $W'$ -like resonance, i.e., a resonance with electric charge 1, which decays to a charged lepton and a neutrino. However, typically such  $W'$  models have an accompanying  $Z'$ -like state, which faces the strong LEP II bounds.

### 3.2. Decays to quasi-stable particles

While the decay of a  $Z'$  to standard model charged leptons does not give a viable supermodel example of topology A, one could imagine a  $q\bar{q}$  resonance that instead decayed with a large branching fraction to new quasi-stable charged particles. Since ATLAS and CMS trigger on penetrating charged particles as if they were muons [16], such scenarios are as visible in the early LHC data as a  $Z'$  decaying to muons.

One simple choice is to take the above  $Z'$  boson with  $g_L = 0$ , and introduce  $N_E$  vector-like “leptons”,  $E$  and  $E^c$ , with coupling  $g_E$  to the  $Z'$  and non-zero electric charge.

Field	Spin	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$
$E$	1/2	-	-	-1
$E^c$	1/2	-	-	1

(15)

The  $E$  and  $E^c$  fields can have an approximate  $\mathbf{Z}_2$  symmetry to make them long-lived, but they must eventually decay to avoid cosmological bounds on absolutely stable charged relics. For this reason, the hypercharge for  $E$  and  $E^c$  has been chosen such that the quasi-stable lepton can decay via mixing with the standard model  $e^c$ .

The branching fraction of the  $Z'$  to the stable charged states is

$$\mathcal{B}(Z' \rightarrow E^+ E^-) = \frac{N_E g_E^2}{N_E g_E^2 + 18g_Q^2}. \quad (16)$$

For large enough values of  $N_E g_E^2$ , the branching fraction can be order 1. Because the stable “leptons” are being produced from a resonance, they will typically have a velocity of

$$\beta \simeq \sqrt{1 - \frac{4m_E^2}{m_{Z'}^2}}, \quad (17)$$

so depending on the relevant mass ratio  $m_E/m_{Z'}$ , a standard “slow muon” cut of  $\beta < 0.9$  [17] will not be particularly effective at capturing the signal. In such cases, these stable charged particles should be treated like ordinary muons (without imposing any kind of  $dE/dx$  quality cut tailored to muons) to reconstruct a  $Z'$  resonance.

Alternatively, one could consider a  $Z'$  that decayed to quasi-stable colored particles that then form  $R$ -hadron-like bound states with QCD partons. Such  $R$ -hadron final states could potentially be visible in early LHC data, though charge flipping interactions [16] complicate both triggering and momentum reconstruction.

### 3.3. Fun with diquarks

From Fig. 5, one sees that  $qq$  resonances can yield an impressive early LHC reach. Such resonances are known as diquarks, and they have spin zero or one, carry baryon number  $2/3$ , and electric charge  $4/3$ ,  $1/3$  or  $-2/3$ . They may transform as a  $\mathbf{6}$  or  $\bar{\mathbf{3}}$  of color. Their couplings are necessarily non-trivial in flavor space because the initial quarks carry flavor. Flavor changing neutral currents impose constraints on couplings of new states with masses of order TeV and large couplings to first generation quarks. To be safe, we consider diquarks with the same flavor quantum numbers as the quarks which produce them, allowing the couplings of the diquark to quarks to be flavor invariant. (Other authors [18] considered diquarks whose couplings to the quarks do not preserve the full flavor symmetries of the SM and can give rise to dangerous flavor violation.)

To be concrete, we consider a spin-zero and color-six diquark, with couplings to the  $SU(2)$  singlet up-type quarks only and symmetric in flavor indices.

Field	Spin	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$
$D$	0	$\mathbf{6}$	-	$4/3$

(18)

The production operator can be written as

$$\mathcal{O}_D = \frac{\kappa_D}{2} D u^c u^c, \quad (19)$$

where  $u^c$  are the up-type singlet quarks and  $D$  is the diquark. The normalization of the coupling constant  $\kappa_D$  depends on the normalization of the color matrices  $R^a$  with which we expand  $D = D^a R^a$ . We use orthonormal matrices such that  $\text{Tr}(R^a R^b) = \delta^{ab}$ . Then the partonic cross section is<sup>6</sup>

$$\sigma(uu \rightarrow D) = \frac{\pi}{6} \kappa_D^2 \delta(\hat{s} - m_D^2). \quad (20)$$

If Eq. (19) were the only available coupling for the diquark, then any produced diquark would simply decay back to the initial state with a partial width given by  $\Gamma = \kappa_D^2 m_D / (16\pi)$ . To be a supermodel, the diquark has to have a large branching fraction to a visible final state. By color conservation, diquark decays must yield at least two jets in the final state, so the most  $Z'$ -like decay possible for a diquark (in the sense that it yields two oppositely charged leptons in the final state) is

$$D \rightarrow 2j + \ell^+ \ell^-. \quad (21)$$

Such a final state can appear via topology B or C, though we will consider the case of topology B since it requires fewer new degrees of freedom.

For example, we can introduce a vector-like fermion  $L$  and  $L^c$ , with the quantum numbers:

Field	Spin	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$
$L$	1/2	$\mathbf{6}$	-	$7/3$
$L^c$	1/2	$\bar{\mathbf{6}}$	-	$-7/3$

(22)

Given its quantum numbers,  $L/L^c$  would be called a “leptodiquark”. The diquark can then decay via the operator

$$\bar{\kappa}_D D L^c e^c \quad (23)$$

with a decay width of  $\Gamma = \bar{\kappa}_D^2 m_D / (16\pi)$ . Thus, as long as  $\bar{\kappa}_D \gtrsim \kappa_D$ , the diquark preferentially decays to the leptodiquark and a lepton. The  $L^c$  will finally decay via the operators in Eqs. (19) and (23) as

<sup>6</sup> Recently radiative corrections were calculated, giving a  $K$ -factor of 1.3 [19].

$$L^c \rightarrow \bar{e}^c u^c u^c \quad (24)$$

through an off-shell diquark, leading to the full decay chain:

$$uu \rightarrow D \begin{array}{l} \searrow \ell^- L \\ \searrow \ell^+ 2j. \end{array} \quad (25)$$

The final state therefore has two jets plus an opposite sign lepton pair, arranged in topology B.

While this diquark–leptodiquark system may strike the reader as baroque, the identical decay topology appears in the case of a  $W'_R$  gauge boson [20,21], where the diquark plays the role of the  $W'_R$  and the leptodiquark plays the role of a right-handed neutrino. However, discovering a left–right symmetric model through this channel typically requires  $1 \text{ fb}^{-1}$  [20,21] of LHC data, whereas the diquark–leptodiquark example motivates a search for the  $2j + \ell^+ \ell^-$  final state in early LHC data.

### 3.4. Resurrecting pair production

In Section 2.1, we argued that QCD pair production of new colored resonances was not a supermodel. However, topology C allows for pair production of new resonances via a supermodel resonance.

For example, using either a  $q\bar{q}$  or a  $qq$  resonance, one can produce vector-like up-type quarks  $U$  and  $U^c$  with quantum numbers:

Field	Spin	$SU(3)_c$	$SU(2)_L$	$U(1)_Y$
$U$	1/2	$\mathbf{3}$	–	2/3
$U^c$	1/2	$\bar{\mathbf{3}}$	–	–2/3

They can be produced via the  $Z'$  through

$$g_U Z'_\mu (\bar{U} \bar{\sigma}^\mu U - \bar{U}^c \bar{\sigma}^\mu U^c), \quad (27)$$

or via the diquark through

$$\frac{\tilde{\kappa}_D}{2} D U^c U^c. \quad (28)$$

If these new colored particles were exactly stable, they would form  $R$ -hadron-like bound states as mentioned above, leading to topology A. However, the heavy  $U/U^c$  quarks may decay via small CKM-like mixings with the standard model quarks, leading to

$$U \rightarrow Z + u/c/t, \quad U \rightarrow W + d/s/b. \quad (29)$$

Such decays are not ideal for making a supermodel, since the  $W$  ( $Z$ ) boson only has 22% (7%) branching fraction to electrons and muons. However, if the  $U/U^c$  quarks only couple to other standard model fermions through higher-dimension operators like

$$\frac{1}{\Lambda^2} (\bar{u}^c \sigma^\mu U^c) (\bar{e}^c \sigma_\mu e^c), \quad \frac{1}{\Lambda^3} (\bar{d}^c \sigma^\mu U^c) (\bar{\ell} h^\dagger \sigma_\mu e^c), \quad (30)$$

then each  $U/U^c$  decay can lead to leptons via

$$U \rightarrow e^+ e^- + u/c/t, \quad U \rightarrow e^+ \nu + d/s/b. \quad (31)$$

Operators like Eq. (30) can always be generated through sufficiently creative model building, and the scale  $\Lambda$  can be made sufficiently large to evade flavor constraints while still being small enough to give prompt decay.

Another option to force leptons to appear in the final state is to have a resonance decay to pairs of colored particles that also

carry lepton number, such as leptoquarks or even the leptodiquarks of Eq. (22).<sup>7</sup> Alternatively, through a  $q\bar{q}$  resonance, one can pair produce color singlet objects with lepton number, such as “sleptons” [22], as long as the  $q\bar{q}$  resonance does not couple to standard model leptons directly.

Finally, a neutral  $q\bar{q}$  resonance can dominantly decay to two secondary resonances that carry no standard model charges. These secondary resonances have a near infinity of possible decay modes, opening up a huge range of final state possibilities. Such scenarios will be supermodels as long as the secondary resonances have an  $\mathcal{O}(1)$  branching fraction to highly visible final states.

## 4. Conclusions

In this Letter, we investigated some new physics scenarios which could be probed by a low energy and low luminosity initial LHC data set, and which will not have been ruled out by the Tevatron and other measurements. We call such scenarios *supermodels*; they are not necessarily motivated by usual model building goals such as solving the hierarchy problem, but are constructed to demonstrate that high production cross sections and clean experimental signatures are possible for early LHC.

Assuming perturbative couplings, we found that  $s$ -channel production of  $qq$  or  $q\bar{q}$  resonances are the most promising supermodel scenarios. To supersede the Tevatron sensitivities in searches for pair-produced particles or single resonances produced from an initial state involving gluon(s), the LHC typically needs higher integrated luminosity than considered in this Letter. Not unexpectedly, given a  $pp$  collider, we found that resonances that couple to  $qq$  initial states have by far the largest LHC cross sections, and in this channel there is a large space of supermodels (see Fig. 5).

We explored various decay topologies of the produced resonances that lead to easily identifiable final states containing a pair of charged leptons or other (quasi-)stable charged particles. While the possibilities for such decays are only limited by theorists imagination and model building ingenuity, we presented some simple examples focusing on decay topologies that also arise in more standard extensions of the standard model. While the supermodels exhibited here might not be as attractive as the name suggests, many of the same final state signatures are useful search channels for “well motivated” models as well. Therefore, searching for supermodels with the early LHC data may benefit finding prettier models when larger data samples become available.

Finally, the space of interesting early LHC searches would be extended if (i) nonperturbative couplings are considered; (ii) pair production is enhanced by high particle multiplicities; (iii) one compares to the currently published Tevatron bounds (some of which utilize less than  $100 \text{ pb}^{-1}$  of data) instead of the 2010 Tevatron sensitivity with  $10 \text{ fb}^{-1}$  of data; (iv) the early LHC data used for analysis approaches or goes beyond  $100 \text{ pb}^{-1}$ .

## Acknowledgements

We thank Mina Arvanitaki, Beate Heinemann, and Matthew Schwartz for helpful conversations. This work was supported in part by the Director, Office of Science, Office of High Energy Physics of the U.S. Department of Energy under contract DE-AC02-05CH11231. M.S. is supported by DE-FG02-01ER-40676. J.T. acknowledges support from the Miller Institute for Basic Research

<sup>7</sup> To produce such states from the diquark resonance, one needs two different fields of opposite lepton number but same baryon number, i.e. a leptoquark and an antilepto-quark.



in Science. D.W. was supported by a University of California Presidential Fellowship. M.S. and J.T. thank the Aspen Center for Physics for their hospitality while this work was in preparation.

## References

- [1] S. Myers, LHC machine status report, 101st LHCC Meeting, Geneva, Switzerland, 5–6 May 2010, <http://indico.cern.ch/conferenceDisplay.py?confId=92525>.
- [2] LHC Performance Workshop, Chamonix, France, 25–29 January 2010, <http://indico.cern.ch/conferenceOtherViews.py?view=standard&confId=67839>.
- [3] See the review  $Z'$ -boson searches, C. Amsler, et al., Particle Data Group, Phys. Lett. B 667 (2008) 1.
- [4] H.L. Lai, et al., CTEQ Collaboration, Eur. Phys. J. C 12 (2000) 375, hep-ph/9903282, and <http://www.phys.psu.edu/~cteq/>.
- [5] J. Pumplin, D.R. Stump, J. Huston, H.L. Lai, P.M. Nadolsky, W.K. Tung, JHEP 0207 (2002) 012, hep-ph/0201195.
- [6] C. Quigg, arXiv:0908.3660.
- [7] A.D. Martin, W.J. Stirling, R.S. Thorne, G. Watt, Eur. Phys. J. C 63 (2009) 189, arXiv:0901.0002, and <http://projects.hepforge.org/mstwpdf/>.
- [8] A.V. Manohar, M.B. Wise, Phys. Rev. D 74 (2006) 035009, hep-ph/0606172.
- [9] B.A. Dobrescu, K. Kong, R. Mahbubani, Phys. Lett. B 670 (2008) 119, arXiv:0709.2378.
- [10] C.P. Burgess, M. Trott, S. Zuberi, arXiv:0907.2696.
- [11] See the review “Leptoquarks” in C. Amsler, et al., Particle Data Group, Phys. Lett. B 667 (2008) 1.
- [12] T. Aaltonen, et al., CDF Collaboration, Phys. Rev. Lett. 102 (2009) 091805, arXiv:0811.0053.
- [13] V.D. Angelopoulos, J.R. Ellis, H. Kowalski, D.V. Nanopoulos, N.D. Tracas, F. Zwirner, Nucl. Phys. B 292 (1987) 59.
- [14] E. Salvioni, G. Villadoro, F. Zwirner, JHEP 0911 (2009) 068, arXiv:0909.1320; E. Salvioni, A. Strumia, G. Villadoro, F. Zwirner, arXiv:0911.1450.
- [15] J. Alcaraz, et al., LEP Collaborations: ALEPH, DELPHI, L3, OPAL, LEP Electroweak Working Group, hep-ex/0612034.
- [16] M. Fairbairn, A.C. Kraan, D.A. Milstead, T. Sjostrand, P. Skands, T. Sloan, Phys. Rep. 438 (2007) 1, hep-ph/0611040.
- [17] CMS Collaboration, CMS PAS EXO-08-003, <http://cms-physics.web.cern.ch/cms-physics/public/EXO-08-003-pas.pdf>, 2008.
- [18] Z. Chacko, R.N. Mohapatra, Phys. Rev. D 59 (1999) 055004, hep-ph/9802388; O. Cakir, M. Sahin, Phys. Rev. D 72 (2005) 115011, hep-ph/0508205; K.S. Babu, R.N. Mohapatra, S. Nasri, Phys. Rev. Lett. 97 (2006) 131301, hep-ph/0606144; K.S. Babu, P.S. Bhupal Dev, R.N. Mohapatra, Phys. Rev. D 79 (2009) 015017, arXiv:0811.3411; C.R. Chen, W. Klemm, V. Rantala, K. Wang, Phys. Rev. D 79 (2009) 054002, arXiv:0811.2105.
- [19] R.N. Mohapatra, N. Okada, H.B. Yu, Phys. Rev. D 77 (2008) 011701, arXiv:0709.1486; T. Han, I. Lewis, T. McElmurry, JHEP 1001 (2010) 123, arXiv:0909.2666.
- [20] G. Aad, et al., ATLAS Collaboration, arXiv:0901.0512.
- [21] G.L. Bayatian, et al., CMS Collaboration, J. Phys. G 34 (2007) 995.
- [22] M. Baumgart, T. Hartman, C. Kilic, L.T. Wang, JHEP 0711 (2007) 084, hep-ph/0608172.