

EXPERIMENTAL STUDY

Effect of a combination of calorie-restriction therapy and Lingguizhugan decoction on levels of fasting blood lipid and inflammatory cytokines in a high-fat diet induced hyperlipidemia rat model

Wang Yuanyuan, Jin Minghua, Zhang Lina, Li Suhua, Zhai Jiayu, Shen Yongzhi, Chen Chunyu, Qin Jian

Wang Yuanyuan, Jin Minghua, Zhang Lina, Li Suhua, Zhai Jiayu, Shen Yongzhi, Chen Chunyu, Qin Jian, Department of Traditional Chinese Medicine, First Affiliated Hospital, Sun Yat-Sen University, Guangzhou 510080, China

Correspondence to: Professor Qin Jian, Department of Traditional Chinese Medicine, First Affiliated Hospital, Sun Yat-Sen University, Guangzhou 510080, China. himy-box@yeah.net

Telephone: +86-18902233983; +86-13760831583

Accepted: May 9, 2014

Abstract

OBJECTIVE: To combine calorie-restriction therapy (CRT) with Traditional Chinese Medicine (TCM) using Lingguizhugan Decoction and note the effects on expression of inflammatory cytokines [tumor necrosis factor (TNF)- α , high-sensitivity C-reactive protein (hs-CRP), interleukin (IL)-6] in high-fat-diet induced hyperlipidemia in rats.

METHODS: Forty rats were divided randomly and equally into four groups: control group A (regular feeding for 5 weeks); model group B (high-fat diet for 5 weeks), calorie-limited group C (high-fat diet for 5 weeks) and TCM calorie-limit group D (high-fat diet for 5 weeks). After modeling, groups C and D were fasted for 3 days: group C with simple fasting, and group D with TCM fasting. The motion as well as changes in color, body weight, food intake, plasma lipids [low-density lipoprotein-cholesterol (LDL-C), high-density lipoprotein-cholesterol, total cholesterol (TC) and triglyceride (TG) along with TNF- α , hs-CRP and IL-6] were measured before and after intervention.

RESULTS: Modeled rats were established after five weeks. After 3 days of fasting, compared with group B, blood lipid levels (TG, TC, LDL-C) of groups C and D decreased dramatically. Those of group D decreased more obviously than those of group C. However, the values of TNF- α , hs-CRP and IL-6 showed no obvious difference between each of the three treatment groups.

CONCLUSION: CRT can reduce fasting blood lipid levels in rats, but not by reducing the levels of inflammatory cytokines.

© 2015 JTCM. All rights reserved.

Key words: Hyperlipidemias; Tumor necrosis factor-alpha; Interleukin-6; Lingguizhugan decoction; Calorie-restriction therapy

INTRODUCTION

Hyperlipidemia is an independent risk for atherosclerosis and is the pathologic basis of cardiovascular and cerebrovascular diseases. In recent years, there has been an improvement in people's living standards. Consumption of high-calorie, high-fat diets and less exercise have led to a higher incidence of obesity, hypertension, diabetes mellitus (DM), hyperlipidemia, coronary heart disease and cerebrovascular diseases.^{1,2} Strategies for controlling diets and improving lifestyle have become a necessary means for the treatment of obesity and related metabolic disorders.

Calorie-restriction therapy (CRT) has been demonstrated to decrease body weight, improve insulin resistance, and regulate immune disorders. Studies have also shown that CRT is effective against metabolic diseases.¹

³ Through long-term clinical experiments,⁴ our research

team found that pure CRT had lipid-lowering effects in hyperlipidemic patients but, during treatment, many patients felt hungry and weak, and that their tongues were significantly thicker after treatment. We added Lingguizhugan Decoction (LGZGD) to CRT to create Traditional Chinese Medicine (TCM)-CRT. This new "hybrid" therapy not only lowered lipid profiles in blood, it also reduced the prevalence of metabolic-related factors and their side effects. We hypothesized that the mechanism of action of these effects may involve vascular inflammatory cytokines, and wished to test our theory in animals, hence the study we describe here.

MATERIALS AND METHODS

Ethical approval of the study protocol

The study protocol was approved by the Ethics Committee of the First Affiliated Hospital (Sun Yat-Sen University, Guangzhou, China).

Animals and diets

Male Wistar rats [specific pathogen-free; 8 weeks; (245 ± 8) g] were purchased from the Experimental Animal Center of Guangdong Province [certificate SCXK (Guangdong) 2009-0011].

All rats lived in their own cage. They were placed in a room with a 12-h light-dark cycle and had free access to food and water. The basic diet and high-fat diet were purchased from Guangdong Province Animal Experiments (Guangdong, China). The high-fat diet contained 55% more fat than a normal diet (15% lard, 21% sucrose, 9% egg-yolk powder, 0.5% bile salts).

Drugs and reagents

LGZGD comprising Fuling (*Poria*), Guizhi (*Ramulus Cinnamomi*), Baizhu (*Rhizoma Atractylodis Macrocephalae*), and stir-frying with liquid adjuvant Gancao (*Radix Glycyrrhizae*) in the proportion 4:3:2:2 was obtained from 999 Medicine Group (Shenzhen, China).

Assay kits for tumor necrosis factor (TNF)-and interleukin (IL)-6 were obtained from Ray Biotech (Atlanta, GA, USA). An assay kit for high-sensitivity C-reactive protein was purchased from Millipore (Bedford, MA, USA). A Microplate Reader (Sunrise™; Tecan, Männedorf, Switzerland), an ultrapure water system for trace organic analyses (Yee Young Enterprises, Nanjing, China) and a Thermostatic Incubator (Jiangnan Instrument Factory, Ningbo, China) were also purchased.

Doses of LGZGD

Doses of LGZGD were based on those stated in the Chinese Pharmacopoeia.⁵ We determined the daily doses for adults according to a mean adult body weight of 60 kg as: Fuling (*Poria*) 12 g, Guizhi (*Ramulus Cinnamomi*) 9 g, Baizhu (*Rhizoma Atractylodis Macrocephalae*) 6 g, and stir-frying with liquid adjuvant Gancao (*Radix Glycyrrhizae*) 6 g.

Grouping and feeding of the rats

First, rats were allowed to adapt to a normal diet for one week. Rats were then divided into four groups (A-D) of ten in a 1:1:1:1 ratio randomly using Matlab software. Group-A rats were fed a normal diet and the other three groups were fed a high-fat diet for five weeks. We evaluated the modeling effects at the fifth weekend.

After modeling, groups A and B were administered the original diet for three days. Group C underwent simple restriction of calorie intake for three days (i.e., no food, free access to water, and gavage using physiologic (0.0%) saline (9 mg/100 g) every day). Group D underwent calorie restriction and LGZGD administration for three days [no food, free access to water, and gavage with LGZGD (9 mg/100 g) every day].

Measurements

General condition of rats before and after treatment: motion as well as changes in color, weight, and food intake of rats was recorded. We measured the blood lipid profiles of all rats at the fifth weekend as well as before and after restriction of calorie intake. After restriction of calorie intake and an overnight fast, 3 mL of blood was obtained from the tail vein. Blood was centrifuged at 1000 × g for 10 min at room temperature. Serum was separated from plasma and sent (as 1-mL samples) to Affiliated Hospital Laboratory for determination of levels of triglycerides (TG), total cholesterol (TC), high-density lipoprotein-cholesterol (HDL-C), low-density lipoprotein-cholesterol (LDL-C) and hs-CRP. The remaining 2-mL samples were sent to Sun Yat-Sen Center for measurement of levels of inflammatory factors.

TNF-α and IL-6

According to a series of standard concentrations and the corresponding values of optical density at 450 nm (OD₄₅₀) we created a TNF-α standard curve using Origin 7.5 software (OriginLab, Northampton, MA, USA) to obtain the equation:

$$Y = 2.1637 + (0.025742 - 2.16372)/(1 + (X/68212)^{0.82886}),$$

where Y is OD₄₅₀ value and X is the TNF-α standard concentration.

We followed an identical procedure to an equation for IL-6:

$$Y = 5.0013 + (-0.012815 - 5.0013)/(1 + (X/36451)^{0.93208}),$$

where Y is OD₄₅₀ value and X is the IL-6 standard concentration.

Using the standard curves, we could obtain the values of the samples in pg/mL at an absorbance of 450 nm.

Statistical analyses

SPSS v18.0 (IBM, Armonk, NY, USA) was used for statistical analyses. Measurement data for mean ± standard deviation ($\bar{x} \pm s$), and the Student's *t*-test was used to compare between the two groups. Multiple sets of single-factor analysis of variance were used to compare multiple groups. $P \leq 0.05$ was considered significant.

RESULTS

General conditions

No deaths were observed in groups A and C. However, one rat died in group B and one rat died in group D. The rat from group B started to fast without an obvious reason in the fourth week of the experiment, started to look peaky gradually, and then died. Another rat in group D died for no obvious reason in the fifth week before calorie intake-limiting treatment. We took blood samples from this rat for analyses. Thirty-eight rats were assessed in total.

In the first four weeks, there was no difference in food intake, activity or color between rats in groups A-D. However, after the fourth week, the activities and food intake, as well as the colors of rats from groups B-D rats, became worse than those of group A. After three days of treatment, rats from group C showed low activity and a more yellowish color than rats of group B, but rats of group D showed much better activity and color than those of group C.

Body weight

At week-5, the body weight of the control group demonstrated a linear upward trend. However, in groups B-D, body weight began to decline in week-4 ($P < 0.01$) and rats looked anorexic. However, between the two calorie-restriction groups, a significant difference in body weight was not observed ($P > 0.05$). After calorie restriction, rats in group C also exhibited anorexia, but group-D rats became stronger than group-C rats (Table 1).

Changes in blood lipid levels

At the fifth weekend, serum levels of TG and cholesterol

of the other three groups were significantly higher than those of group A ($P < 0.01$), but there were no significant differences in levels of HDL-C or LDL-C ($P > 0.05$). After three days of calorie restriction, levels of TG, cholesterol, HDL-C and LDL-C of groups C and D were significantly lower than those of group B ($P < 0.01$). Also, levels of TG, cholesterol, HDL-C and LDL-C of group D were reduced further than those of group C ($P < 0.01$) (Table 2).

Comparison of levels of TNF- α , hs-CRP and IL-6

Before CRT, compared with the control group, levels of TNF-, hs-CRP and IL-6 of groups B-D were increased significantly ($P < 0.05$). However, after CRT, compared with the model group, levels of TNF- and hs-CRP of groups C and D showed no significant differences, and even the IL-6 levels of these two groups showed a decreasing trend but without significant differences between them, or between them and the model group ($P > 0.05$) (Table 3).

DISCUSSION

The present study showed that CRT and TCM-CRT can reduce serum lipid levels (TG, cholesterol, LDL-C, HDL-C) significantly and also the body weight of rats. Group-C rats showed fatigue, drowsiness and lethargy, but group-D rats treated with LGZGD showed none of these features, and even exhibited better activity and fur gloss than group-C rats.

CRT can improve hyperlipidemia significantly. However, TCM-CRT not only elicited better effects, but also reduced the side effects of CRT, so what was the therapeutic mechanism? Taking into account the close correlation of hyperlipidemia and inflammatory factors, we

Table 2 Comparisons of lipid levels between the four groups (mmol/L)

Group	n	TC	TG	HDL-C	LDL-C
A	10	1.622±0.250	1.062±0.461	0.773±0.090	0.252±0.052
B	9	2.082±0.380 ^a	1.660±0.553 ^a	0.854±0.073 ^a	0.500±0.152 ^a
C	10	1.181±0.301 ^b	0.943±0.212 ^c	0.543±0.103 ^c	1.165±0.064 ^b
D	9	1.383±0.275 ^b	0.861±0.184 ^b	0.584±0.101 ^b	0.193±0.047 ^b

Notes: A: control group (standard food for 5 weeks); B: model group (high-fat diet for 5 weeks); C: intervention group (high-fat diet for 5 weeks); D: intervention group (high-fat diet for 5 weeks). TC: total cholesterol; TG: triglyceride; LDL-C: low-density lipoprotein-cholesterol; HDL-C: high-density lipoprotein-cholesterol. Compared with group A, ^a $P < 0.01$; compared with group B, ^c $P < 0.05$; compared with group B, ^b $P < 0.01$.

Table 1 Comparison of the body weight of rats between the four groups (g)

Group	n	First Day	First weekend	Third weekend	Fifth weekend	After fasting
A	10	196±10	225±16	254±25	275±34 ^a	269±48 ^b
B	9	202±12	230±18	255±27	292±56 ^a	301±55 ^b
C	10	205±9	222±12	242±31	305±57 ^a	289±64 ^b
D	9	197±11	228±15	251±38	309±65 ^a	281±70 ^b

Notes: A: control group (standard food for 5 weeks); B: model group (high-fat diet for 5 weeks); C: intervention group (high-fat diet for 5 weeks); D: intervention group (high-fat diet for 5 weeks). Compared with the first day of this group, ^a $P < 0.05$; compared with the control group, ^b $P < 0.01$.

Table 3 Comparison of the levels of TNF- α , hs-CRP and IL-6 between the four groups (pg/mL)

Group	n	TNF- α	hs-CRP	IL-6
A	10	932.071 \pm 110.942	0.132 \pm 0.024	551.691 \pm 72.762
B	9	852.362 \pm 121.763 ^a	0.173 \pm 0.044 ^a	579.151 \pm 177.363 ^a
C	10	836.801 \pm 35.412 ^b	0.183 \pm 0.054 ^b	518.002 \pm 52.001 ^b
D	9	891.041 \pm 108.473 ^c	0.171 \pm 0.031 ^c	512.930 \pm 88.912 ^c

Notes: A: control group (standard food for 5 weeks); B: model group (high-fat diet for 5 weeks); C: intervention group (high-fat diet for 5 weeks); D: intervention group (high-fat diet for 5 weeks). TNF- α : tumor necrosis factor- α ; hs-CRP: high sensitivity C-reactive protein; IL-6: interleukin-6. Compared with group A, ^a $P < 0.05$; compared with group B, ^b $P > 0.05$; compared with group C, ^c $P > 0.05$.

studied the mechanism of action of this treatment. However, after treatment, levels of TNF- α , hs-CRP, and IL-6 in groups C and D showed no significant improvement than before treatment ($P \geq 0.05$). Hence, there was no significant correlation between TCM-CRT and anti-inflammatory effects.

Previously, we undertook another study focusing on CRT in insulin-resistant rats.² We discovered that blood glucose and fasting insulin levels were improved after TCM-CRT, and that peroxisome proliferator-activated receptor (PPAR)- γ activity was also inhibited. TCM-CRT might inhibit PPAR- γ activity to improve insulin resistance. PPAR- γ is a key factor of adipocyte differentiation, and adipocyte differentiation may have played a key part in our study. Excessive activation of PPAR- γ expression in adipose tissue and fat cells might be involved in obesity.⁶⁻⁸ Hence, inhibition of PPAR- γ activity in TCM-CRT may be based on inhibition of the differentiation of fat cells.⁹⁻¹¹

We have found that, in clinical trials, CRT can significantly improve hyperlipidemia, early DM, insulin resistance and other related metabolic disorders, but its mechanism of action has not have been explored until now. Comprehensive experiments in two animal studies have shown that the mechanism underlying this effect may not reduce the expression of inflammatory factors in the body,^{2,6} but might inhibit the differentiation of fat cells. After this treatment, patients changed to eating a healthy diet. We noted the blood levels of glucose and lipids did not rebound. This phenomenon may also have been due to reactivation of the differentiation of fat cells.¹¹⁻¹³ Calorie restriction may complete the entire process of apoptosis and neogenesis in the body. The more "comfortable" feeling in the body after treatment may be associated with this correlation.¹³⁻¹⁵ Despite our interesting observations, more clinical and animal studies are needed to support these findings.

REFERENCES

- 1 **Ke B**, Qin J. fasting therapy and its research progress. *Shenzhen Chuan Tong Zhong Yi* 2009; 19(1): 50.
- 2 **Wang YY**, Jin MH, Ke B, Qin J. Effects of Lingguizhugan decoction combined calorie restriction on the insulin resistance and PPAR- γ of model rats. *Zhong Guo Zhong Xi Yi Jie He* 2013; 3(33): 355-357.
- 3 **Qin J**. Effects of the modified Lingguizhugan decoction combined with short-term very low calorie diets on glyce-mic control in newly diagnosed type 2 diabetics. *J Tradit Chin Med* 2011; 3(1): 51.
- 4 **Jin M**, Wang Y, Ou B, Meng J, Ke B, Qin J. Effects of Danqidihuang granules on glucolipid metabolism in insu-lin resistantrats. *J Tradit Chin Med* 2012; 32(4): 627-630.
- 5 **Liu T**, Yang SY. Fat diet induced the phlegm animal mod-el and evaluation. *Changchun Zhong Yi Yao* 2009; 25(3): 333-335.
- 6 **Chen DS**, Qin J. TCM fasting therapy to rebuild spleen governs the transportation Theoretical Analysis and prac-tice. *Zhong Yi Lin Chuang Yan Jiu* 2010; 4(2): 10-12.
- 7 **Ke B**, Qin J, Zhang JJ. Short-term fasting physiological and biochemical parameters in rats. *Shi Zhen Guo Yi Guo Yao* 2011; 11 (22): 2778-2780.
- 8 **Miles PD**, Barak Y, He W, Evans RM, Olefsky JM. Improved insulin-sensitivity in mice heterozygous for PPAR- γ deficiency. *J Clin Invest* 2000; 105(3): 287-292.
- 9 **Sakamoto A**, Hongo M, Furuta K, Saito K, Nagai R, Ishizaka N. Pioglitazone ameliorates systolic and diastolic cardiac dysfunction in rat model of angiotensin II-induced hypertension. *Int J Cardiol* 2013; 9(31): 409-415.
- 10 **Tian AP**, Guo SS, Chen YT, Shen ZF. Experimental re-search of high heat feed induced insulin resistance animal models. *Zhong Guo Yao Xue* 2006; 41(11): 827-831.
- 11 **Vazquez LA**, Pazos F, Amado JA, et al. Effects of changes in body weight and insulin resistance on inflammation and endothelial function in morbid obesity after bariatric surgery. *J Clin Endocrinol Metab* 2005; 90(1): 316-322.
- 12 **Liu W**, Li J, Shen B, et al. Research progresses of insu-lin resistance animal models. *Zhong Yi Yao Xin Xi* 2008; 15 (6): 9-11.
- 13 **Reimer MK**, Ahren B. Altered β -cell distribution of pdx-1 and GLUT-2 after a short-term challenge with a high-fat diet in C57BL/6Jmice. *Diabetes* 2002; 51: S138-S143.
- 14 **Korach-André M**, Gao J, Gounarides JS, Deacon R, Islam A, Laurent D. Relation-ship between visceral adiposity and intramyocellular lipid content in two models of insu-lin resistance. *Am J Physiol Endocrinol Metab* 2005; 288 (1): E106-E116.
- 15 **Liu L**, Zhang N. Clinical research progress for insulin resistance of hypertension in Traditional Chinese Medicine. *Zhong Yi Yao Xin Xi* 2009; 26(6): 123-125.