

and essentially stops transcription initiation. Inhibited transcription can be recovered upon gyrase binding and reaction on the DNA. Furthermore, using single-cell mRNA counting fluorescence in situ hybridization (FISH) assay, we find the extent of transcriptional bursting depends on the intracellular gyrase concentration. These findings prove that transcriptional bursting of highly expressed genes in bacteria is primarily caused by reversible switching between different chromosomal supercoiling levels.

2451-Pos Board B143

Watching the RNA Polymerase Transcription by Time-Dependent Soak-Trigger-Freeze X-Ray Crystallography

Katsuhiko Murakami.

Penn State University, University Park, PA, USA.

The challenge for structural biology is to understand atomic-level macromolecular motions during enzymatic reaction. X-ray crystallography can reveal high resolution structures; however, one perceived limitation is that it reveals only static views. Here we use time-dependent soak-trigger-freeze X-ray crystallography, namely, soaking nucleotide and divalent metal into the bacteriophage RNA polymerase (RNAP)-promoter DNA complex crystals to trigger the nucleotidyl transfer reaction and freezing crystals at different time points, to capture real-time intermediates in the pathway of transcription. In each crystal structure, different intensities and shapes of electron density maps corresponding to the nucleotide and metal were revealed at the RNAP active site which allow watching the nucleotide and metal bindings and the phosphodiester bond formation in a time perspective. Our study provides the temporal order of substrate assembly and metal co-factor binding at the active site of enzyme which completes our understanding of the two-metal-ion mechanism and fidelity mechanism in single-subunit RNAPs. The nucleotide-binding metal (MeB) is coordinated at the active site prior to the catalytic metal (MeA). MeA coordination is only temporal, established just before and dissociated immediately after phosphodiester bond formation. We captured these elusive intermediates exploiting the slow enzymatic reaction in crystallo. These results demonstrate that the simple time-dependent soak-trigger-freeze X-ray crystallography offers a direct means for monitoring enzymatic reactions. [Ref: Basu, R.S., and K.S. Murakami (2013). *J Biol Chem.*, 288, 3305-3311]

2452-Pos Board B144

Structural and Dynamic Regulation of TFIID-Mediated Transcription Initiation Complex Assembly by the Tumor Suppressor p53 Protein

Anna Piasecka¹, Lihua Song¹, Michael Cianfrocco², Vincent Wong¹, Shenglong Wang¹, Joseph Hargitai¹, William Rice³, Eva Nogales⁴, Robert A. Coleman¹, Wei-Li Liu¹.

¹Albert Einstein College of Medicine, Bronx, NY, USA, ²Harvard Medical School, Boston, MA, USA, ³New York Structural Biology Center, New York, NY, USA, ⁴Howard Hughes Medical Institute, University of California, Berkeley, Berkeley, CA, USA.

p53 plays a central role in tumor suppression. To quickly respond to diverse stress stimuli, p53 binds specific elements in various target promoters to induce vast gene networks for maintaining cellular integrity. p53 stimulates transcription in part by aiding promoter recruitment of the transcription machinery. TFIID, a key component within the transcription machinery, is responsible for binding specific core promoter DNA sequences and recruiting other basal factors including RNA Polymerase II to initiate transcription. However, the exact mechanism underlying how p53 facilitates TFIID-mediated transcription is unclear.

Each p53 target gene has a unique arrangement of p53-responsive and core promoter elements. How these various arrangements on different gene promoters regulate the structural architecture of TFIID and the positioning of p53-specific elements remains unknown. Moreover, structural information of p53 bound to its various target promoters and other factors remains elusive. Therefore, we aim to decipher the molecular mechanism underlying p53's ability to stimulate transcription by revealing the biochemical, structural and dynamic basis of TFIID bound to p53 and promoter DNA.

To this end, we established unique protein purification strategies to generate high-purity native TFIID complex bound to p53/TFIIA/native promoter DNA. We next determined the 3D structures of TFIID/p53/TFIIA co-complexes on two distinct p53 target gene promoters via single particle cryo-electron microscopy. Strikingly, we discovered a common mode of TFIID binding to different types of promoters. Our biochemical studies showed that p53 significantly promotes TFIID's interaction with DNA. To further mechanistically dissect TFIID's enhanced promoter recognition/binding directed by p53, we examined the dynamic interaction between p53, TFIID and promoter DNA via single molecule TIRF microscopy. Taken together, our structural and functional studies elucidate how p53 facilitates TFIID-mediated transcription initiation complex assembly on different p53 target gene promoters.

2453-Pos Board B145

Single Molecule Probing of p53's Ability to Dynamically Regulate Chromatin Structure

Vincent Wong¹, Yu-Jen Chen¹, Charles Kenworthy¹, Lihua Song¹, Gina Dailey², Wei-Li Liu¹, Robert A. Coleman¹.

¹Albert Einstein College of Medicine, Bronx, NY, USA, ²Howard Hughes Medical Institute, University of California-Berkeley, Berkeley, CA, USA.

The tumor suppressor p53 protein is a transcriptional activator that binds to its response elements (REs) on target promoters and activates expression of a large number of genes involved in tumor suppression. Previous ChIP-seq studies indicate that p53 binds to its REs embedded in regions densely populated with nucleosomes. However, it is unknown if nucleosomes help or hinder binding of p53 to its response elements to regulate transcription.

To decipher the interaction of p53 with nucleosomes, we have utilized a combination of Next Generation Proteomics, bioinformatics, bulk biochemical and real-time single molecule FRET assays. Our proteomic assays show that p53 interacts with peptides that have strong homology to histones H2A, H2B, and H4. Biochemical assays indicate that p53 can stably interact with histone peptides in the absence of DNA. Genomic maps of p53 REs and nucleosome positions further reveal that p53 REs cluster specifically within 2 regions of the nucleosome. Furthermore these clustered p53 REs are adjacent to histone regions identified in our proteomic studies, suggesting localization of two physiologically relevant binding platforms for p53 on the nucleosome.

We have also established a single molecule FRET assay to characterize dynamic structural changes in nucleosomes with and without p53 REs. Interestingly, our studies reveal that p53 can bind to our biochemically defined nucleosomal surface and dynamically alter the structure of nucleosomal DNA independently of the presence of a p53 RE. Intriguingly, p53 has a higher affinity for its RE when incorporated into nucleosomes assembled with native DNA compared to REs on naked DNA. Thus, our studies indicate that histone/p53 contacts may enhance p53 directed transcription by creating a stable platform for p53 promoter recruitment.

2454-Pos Board B146

Substantial Nucleotide Selection Prior to Full Insertion of the Nucleotide in T7 RNA Polymerase Elongation

Baogen Duan, Shaogui Wu, Yu Jin.

Beijing Computational Science Research center, No. 3 He-Qing Road, Hai-Dian District, Beijing, China.

Nucleotide selection is essential for fidelity control in gene replication and transcription. Previous work on T7 RNA polymerase suggests that a small post-translocation free energy stabilizes Tyr639 in the active site to facilitate the nucleotide selection. Currently, we implement atomistic molecular dynamics simulations and demonstrate that Tyr639 is indeed stabilized by $\sim 2k_B T$ favorable association with the end base pair of the DNA-RNA hybrid to serve for nucleotide 'gating' from pre-insertion to insertion. Upon the nucleotide pre-insertion, a relative binding free energy above thermal fluctuation level arises against the miscoded nucleotide, primarily due to electrostatic screening from charged residues that assist the nucleotide binding. Interestingly, the pre-insertion of a right nucleotide marginally destabilizes Tyr639, while a wrong nucleotide pre-insertion substantially stabilizes Tyr639 to hinder further nucleotide insertion. The activation barrier of the miscoded nucleotide insertion under an O-helix rotation rises significantly above that of the right nucleotide. The selection against deoxyribonucleotide can be even strong and arises essentially due to steric detection from Tyr639. Our studies suggest that substantial nucleotide selection in T7 RNAP happens upon the nucleotide pre-insertion and during the insertion, prior to full insertion of the nucleotide for base pairing and chemical addition.

2455-Pos Board B147

Complete Dissection of Transcription Elongation Reveals Slow Translocation of RNA Polymerase II in a Linear Ratchet Mechanism

Manchuta Dangkulwanich¹, Toyotaka Ishibashi¹, Shixin Liu¹,

Maria L. Kireeva², Lucyna Lubkowska², Mikhail Kahlev³, Carlos J. Bustamante¹.

¹UC Berkeley, Berkeley, CA, USA, ²National Cancer Institute, Frederick, MD, USA, ³National Cancer Institute, Frederick, CA, USA.

During transcription elongation, RNA polymerase has been assumed to attain equilibrium between pre- and post-translocated states rapidly relative to the subsequent catalysis. Under this assumption, recent single-molecule studies proposed a branched Brownian ratchet mechanism that necessitates a putative secondary nucleotide binding site on the enzyme. By challenging individual yeast Pol II with a nucleosomal barrier, we separately measured the forward and reverse translocation rates. Surprisingly, we found that the forward translocation rate is comparable to the catalysis rate. This finding reveals a linear,