



# The Influence of Target Angular Velocity on Visual Latency Difference Determined using the Rotating Pulfrich Effect

R. W. D. NICKALLS\*

Received 16 May 1995; in revised form 6 September 1995; in final form 22 January 1996

Visual latency difference was determined directly in normal volunteers, using the rotating Pulfrich technique described by Nickalls [*Vision Research*, 26, 367–372 (1986)]. Subjects fixated a black vertical rod rotating clockwise on a horizontal turntable turning with constant angular velocity (16.6, 33.3 or 44.7 revs/min) with a neutral density filter (OD 0.7 or 1.5) in front of the right eye. For all subjects the latency difference associated with the 1.5 OD filter was significantly greater ( $P < 0.001$ ) with the rod rotating at 16.6 rev/min than at 33.3 revs/min. The existence of an inverse relationship between latency difference and angular velocity is hypothesized. Copyright © 1996 Published by Elsevier Science Ltd.

Pulfrich Latency Illusion Stereopsis Tracking

## INTRODUCTION

The Pulfrich effect is a remarkable visual illusion, seen when a moving object is viewed binocularly with a neutral density filter in front of one eye (Pulfrich, 1922). For example, if a swinging pendulum is viewed in this way from a direction at right angles to its motion, then the pendulum bob appears to describe an elliptical orbit.

Although the Pulfrich effect has been extensively analysed for a pendulum and simple harmonic motion (Lit, 1949; Weale, 1954; Trincker, 1953; Levick *et al.*, 1972), a number of other manifestations of the Pulfrich phenomenon have also been investigated. These include the 'rotating' Pulfrich effect (Nickalls, 1986a, b); apparent bending of unevenly illuminated rods (Barlow & McNaughton, 1980); a paradoxical decrease in apparent size when the target appears to come towards the observer (Weale, 1954; Spiegler, 1983); and an apparent hyperbolic path when the target moves with constant velocity in a plane which intersects the pupillary plane of the observer (Spiegler, 1986). Pulfrich effects have also been described in association with concentric rotation (Prestrude & Baker, 1968); bouncing balls (Wilson, 1965); motion of the observer (Enright, 1970), as well as in a variety of medical conditions which affect

the optic nerve (Larkin *et al.*, 1994), retina (Hofeldt *et al.*, 1985) or retinal illumination (Sokol, 1976).

The magnitude of the Pulfrich effect has been found to be a function of the plane of motion (Spiegler, 1986); the degree of binocular intensity difference (Lythgoe, 1938; Lit, 1949); viewing distance (Lit & Hyman, 1951), target size (Spiegler, 1983); target thickness (Lit, 1960c); and target velocity (Lit, 1960a, b, 1964; Spiegler, 1983).

## Mechanism

The mechanism underlying the Pulfrich phenomenon is not clear. The classical explanation proposed by Fertsch [see Pulfrich (1922)], is that the phenomenon is due to a unilateral increase in visual latency resulting from the decrease in retinal image intensity due to the filter (Williams & Lit, 1983; Carney *et al.*, 1989). Compelling evidence in support of a temporal delay model arises from the demonstration that a unilateral light-attenuating filter is able to delay a unilaterally time-advanced sequence of random-dot stereograms sufficiently to restore depth perception (Julesz & White, 1969; Ross & Hogben, 1975). A saccadic-suppression model has also been suggested (Harker, 1967) in order to explain the apparent asymmetrical path associated with a pendulum described by Trincker (1953).

However, there are difficulties associated with the classical temporal delay model since the Pulfrich effect can be seen even with intermittent (stroboscopic) target presentation, possibly owing to some form of interaction

\*Department of Anaesthesia, City Hospital, Nottingham, U.K. [Email dick.nickalls@nottingham.ac.uk].

(e.g. lateral inhibition) between successive inputs to the eyes (Lee, 1970). It is possible that the filter could introduce a spatial disparity by causing fusion of non-corresponding discrete positions, but this is unlikely in the case where the temporal interval between the motion samples is greater than the delay between the eyes (Morgan & Thompson, 1975). Furthermore, a Pulfrich-type effect can be produced without a delay by artificially increasing the target persistence in one eye (Morgan, 1975).

#### *Preliminary study*

In preliminary studies involving the "rotating" Pulfrich effect (Nickalls, 1986a) it was noticed, contrary to expectation, that the latency difference for a given illumination appeared to vary significantly with turntable speed. In view of this discrepancy, the present study was designed to investigate the influence of turntable speed on latency difference.

### METHODS

Latency difference was determined directly using the technique described by Nickalls (1986a), which makes use of a "rotating" Pulfrich effect. With this technique, an observer with a neutral density filter in front of the right eye, binocularly fixates a horizontally clockwise rotating target from within the plane of rotation. By varying the viewing distance, the observer identifies a null-position (known as "transition") at which the target appears not to rotate at all, but appears to move only from side-to-side. The latency difference can then be calculated from the viewing distance at transition [see Eq. (1)].

#### *Apparatus*

The rotating target used in the present study was a black vertical rod (1.5 mm diameter) mounted 11.9 cm from the centre of a horizontal clockwise-rotating turntable, and was clearly visible against a white background. Rotational cues from both the turntable and the ends of the rod were screened out by viewing through a 5 cm wide horizontal slit as described by Nickalls (1986a).

The turntable was mounted at eye-level, on a trolley which ran backwards and forwards on a straight 3.5 m track. The observer (fixed) was positioned at one end of the track, and was able to vary the position of the trolley, and hence the viewing distance, by turning a small hand-wheel. The subject's head was immobilized using chin and forehead rests in the usual way.

The angular velocity of the turntable (Garrard SP mark 2 record player) was determined using a diffuse-scan opto-switch, and displayed continuously in revolutions per minute (rev/min;  $\Omega$ ) to one decimal place. The overall mean rev/min (range) for each turntable speed for all observations described in this paper are as follows: 16.6 rev/min (16.3–16.9); 33.3 rev/min (33.0–33.9); 44.7 rev/min (44.3–45.1). The maximum variation in turntable speed during a set of 10 observations (see Procedure) was  $\pm 0.3$  rev/min.

#### *Illumination*

The illumination was the same for all observations. The illumination of both the front screen and the background screen was from above in order to maintain a uniform luminance throughout the full range of movement of the turntable apparatus.

The luminance of the front screen and background (both white) was measured using a narrow angle 40A Opto-meter (Model R, United Detector Technology Inc.) which incorporated a silicon PIN photodiode with a foot-lambert lens. All luminance readings were made from a distance of 20 cm in front of the front screen; background readings were made through the viewing slit. The mean (range) luminance of both the front screen and the background screen over the full range of viewing distance was 114 cd/m<sup>2</sup> (109–121;  $n = 8$ ) and 124 cd/m<sup>2</sup> (116–130;  $n = 8$ ), respectively.

#### *Filters*

Two different Wrattan neutral-density filters (Kodak) were used, having optical densities (OD) of 0.7 and 1.5. During each experiment one of these filters was placed in front of the right eye, using special goggles which prevented any extraneous non-filtered light from reaching the filtered eye.

#### *Separation of rotation centres of the eyes*

The semi-separation ( $a$ ) of the rotation centres of the two eyes [required for Eq. (1)], was determined by measuring the inter-pupillary distance (IPD) when the eyes were both parallel and at right angles to the line joining the two eyes. The IPD was measured using a corneal-reflection pupillometer (Essilor Ltd., Bristol, U.K.). The IPD values presented in the Tables are the mean (rounded to the nearest 1 mm) of five sequential measurements.

#### *Transition viewing distance*

The viewing distance at "transition" ( $d_T$ ) was measured from the centre of the turntable to the line joining the centres of rotation of the two eyes. This was done by first measuring the distance from the centre of the turntable to the front of the cornea. An additional 1.5 cm was then added to this value to account for the distance between the front of the cornea and the rotation centre of the eye (Fry & Hill, 1962). For all studies the viewing distance at transition was within the range 90–270 cm.

#### *Latency difference*

Each of the latency difference determinations presented in the Tables is the mean of 10 sequential measurements. The latency difference ( $\Delta t$  sec) was derived from three parameters namely:

- (i) the viewing distance ( $d_T$  cm) at which the "transition" null-point is perceived;
- (ii) the value of half the separation of the rotation centres of the eyes ( $a$  cm); and
- (iii) the angular velocity of the turntable ( $\Omega$  rev/min).




FIGURE 1. Variation of latency difference ( $\Delta t$ ) with turntable speed ( $\Omega$ ) and transition distance ( $d_T$ ) as described by Eq. (1) with  $a = 3.2$  cm (IPD = 6.4 cm).

The latency difference was calculated using Eq. (1), in which the angular velocity ( $\Omega$ ) is in rev/min (see Appendix A for derivation).

$$\Delta t = (1/3\Omega)\tan^{-1}(a/d_T) \quad (1)$$

Equation (1) is depicted graphically in Fig. 1. Owing to the non-linear nature of Eq. (1) it follows that for a given number of observations, the mean  $d_T$  does not correlate accurately with the associated mean  $\Delta t$ . Consequently, the mean  $\Delta t$  values for each subject given in the tables are derived from the individual  $\Delta t$  values.

In addition, the format of Eq. (1) is significant in that a relatively large error in the viewing distance at transition ( $d_T$ ) is associated with only a very small error in the calculated latency difference ( $\Delta t$ ). For example, if  $a = 3.2$  cm,  $\Omega = 33$  rev/min and  $d_T = 120$  cm, then an error of  $\pm 1$  cm in  $d_T$  is associated with an error of only  $\pm 0.13$  msec in  $\Delta t$ .

**Subjects**

A number of studies were performed on a total of 14 normal experienced volunteers whose ages ranged from

TABLE 1. A typical series of 10 sequential measurements of visual latency difference (subject SY; age 30 yr; IPD 64 mm)

Initial turntable position	OD = 1.5		
	$\Omega$ (rpm)	$d_T$ (cm)	$\Delta t$ (msec)
Near	33.6	117.5	15.48
Far	33.5	116.3	15.68
Near	33.5	114.8	15.89
Far	33.5	110.7	16.48
Near	33.4	125.6	14.57
Far	33.5	109.6	16.64
Near	33.5	126.4	14.43
Far	33.4	113.7	16.09
Near	33.3	118.3	15.51
Far	33.3	107.1	17.13
<i>n</i>	10	10	10
Mean	33.4	116.0	15.8
SEM	0.03	2.01	0.27

22 (PO)-54 (RC) years. All subjects had normal depth perception as determined using the Wirt Fly test.

**Procedure**

The latency difference was determined for a number of combinations of turntable speed (16.6, 33.3, 44.7 mean rev/min) and filter density (OD 0.7, 1.5).

The subjects were investigated in two sessions. During one session, the following combinations of filter optical density and turntable speed were used: (0.7 OD/33.3 rev/min; 0.7 OD/44.7 rev/min; 1.5 OD/33.3 rev/min). During the other session, the following combinations were used: (1.5 OD/16.6 rev/min; 1.5 OD/33.3 rev/min). Note that the only combinations of filter density and turntable speed which could be used, were those for which the viewing distance at transition was within the range of the physical track (3.5 m) that the turntable

TABLE 2. Comparison of latency difference determinations in 10 subjects ( $\Omega = 33.3$  rev/min) using two different neutral density filters

Subject	IPD (mm)	OD = 1.5 $\Omega = 33.3$ rev/min		OD = 0.7 $\Omega = 33.3$ rev/min	
		$d_T$ (cm) Mean $\pm$ SEM (range)	$\Delta t$ (msec) Mean $\pm$ SEM (range)	$d_T$ (cm) Mean $\pm$ SEM (range)	$\Delta t$ (msec) Mean $\pm$ SEM (range)
PO	60.9	142.8 $\pm$ 2.3 (125.5-160.7)*	12.3 $\pm$ 0.21 (10.8-14.0)*	228.1 $\pm$ 6.2 (197.4-260.6)	7.8 $\pm$ 0.21 (6.7-8.9)
RWDN	66.4	119.9 $\pm$ 1.0 (112.1-130.1)*	15.9 $\pm$ 0.14 (14.5-17.1)*	234.9 $\pm$ 3.5 (206.2-261.4)*	8.2 $\pm$ 0.12 (7.4-9.3)*
EAN	61.3	120.5 $\pm$ 2.5 (99.6-135.8)*	14.7 $\pm$ 0.31 (13.1-17.5)*	229.0 $\pm$ 3.3 (210.1-244.4)	7.7 $\pm$ 0.11 (7.2-8.4)
RC	68.0	111.4 $\pm$ 1.9 (96.1-128.3)*	17.6 $\pm$ 0.30 (15.1-20.3)*	214.1 $\pm$ 3.6 (195.3-239.1)	9.1 $\pm$ 0.15 (8.2-10.0)
SM	61.9	130.3 $\pm$ 2.0 (106.4-151.9)*	13.7 $\pm$ 0.22 (11.7-16.7)*	210.2 $\pm$ 10.1 (171.0-263.7)	8.6 $\pm$ 0.41 (6.7-10.3)
MC	61.2	129.2 $\pm$ 3.3 (109.0-162.4)*	13.8 $\pm$ 0.33 (10.9-16.2)*	219.0 $\pm$ 12.6 (168.1-261.8)	8.3 $\pm$ 0.50 (6.7-10.5)
IJ	57.9	115.0 $\pm$ 2.7 (104.7-134.6)	15.2 $\pm$ 0.34 (12.9-16.6)	205.9 $\pm$ 9.1 (168.0-245.5)	8.6 $\pm$ 0.37 (7.1-10.4)
AM	64.0	121.5 $\pm$ 4.5 (105.5-153.5)	15.6 $\pm$ 0.52 (12.2-17.7)	190.4 $\pm$ 11.4 (143.4-246.8)	10.1 $\pm$ 0.61 (7.5-13.0)
PN	63.5	100.2 $\pm$ 1.0 (97.1-107.6)	17.9 $\pm$ 0.16 (16.8-18.5)	198.9 $\pm$ 5.0 (169.8-219.6)	9.1 $\pm$ 0.23 (8.2-10.6)
MK	64.4	141.4 $\pm$ 5.8 (123.5-185.3)	13.2 $\pm$ 0.48 (9.9-14.9)	202.5 $\pm$ 8.2 (169.8-244.2)	9.2 $\pm$ 0.36 (7.5-10.9)
<i>n</i>		10	10	10	10
Mean		123.2	15.0	213.3	8.7
SEM		4.16	0.58	4.60	0.23

The mean within-subject difference in  $\Delta t$  is significant ( $P < 0.001$ ). Each determination is the mean of 10 sequential measurements. \*indicates pooled data from two separate determinations. These data are shown in Fig. 2.

TABLE 3. Repeat determinations of latency difference in nine subjects (OD = 1.5; Ω = 33.3 rev/min)

Subject	IPD (mm)	OD = 1.5, Ω = 33.3 rev/min		Time interval (days)	Difference (msec) Δt <sub>1</sub> - Δt <sub>2</sub>
		Δt <sub>1</sub> msec (first test) Mean ± SEM (range)	Δt <sub>2</sub> msec (second test) Mean ± SEM (range)		
PO	60.9	12.9 ± 0.25 (11.6-14.0)	11.7 ± 0.21 (10.8-12.6)	206	+1.2
RWDN	66.4	16.1 ± 0.16 (15.5-17.1)	15.8 ± 0.23 (14.5-16.8)	197	+0.3
EAN	61.3	13.8 ± 0.19 (13.1-14.8)	15.6 ± 0.41 (13.5-17.5)	203	-1.8
RC	68.0	16.7 ± 0.28 (15.1-17.9)	18.5 ± 0.34 (16.7-20.3)	119	-1.8
SM	61.9	13.7 ± 0.21 (12.7-14.9)	13.7 ± 0.40 (11.7-16.7)	269	0.0
MC	61.2	14.6 ± 0.34 (12.8-16.2)	13.1 ± 0.47 (10.9-15.4)	439	+1.5
MH	57.9	16.6 ± 0.41 (14.3-18.8)	15.6 ± 0.23 (14.9-17.2)	175	+1.0
SY	64.0	15.8 ± 0.27 (14.4-17.1)	16.7 ± 0.13 (16.3-17.4)	27	-0.9
SR	63.5	17.2 ± 0.47 (15.5-19.8)	16.3 ± 0.32 (14.8-18.1)	1	+0.9
			<i>n</i>	9	9
			Mean	182	0.04
			SEM	—	0.42

The mean within-subject difference in Δt is not significant (P > 0.1). Each determination is the mean of 10 sequential measurements.

moved on [see Nickalls (1986a) for details of the laboratory setup].

In nine subjects repeat latency difference determinations were made for the 1.5 OD/33.3 rev/min combination (see Table 3) following a mean interval of 182 days (range: 1 day-14 months) in order to check reproducibility. A repeat determination for the 0.7 OD/33.3 rev/min combination was made in one subject (RWDN).

The illumination was the same for all observations. Each subject was given 20 min to dark adapt to each filter [see Standing *et al.* (1968)].

For each combination of turntable speed and filter density, the latency difference was determined as the mean of a series of 10 sequential measurements. For each measurement of latency difference, the turntable was initially positioned at either the near or the far point of the track (i.e. either close to or far away from the subject).

The subject was then asked to fixate the horizontally rotating target, and at the same time to adjust the position of the turntable using the small hand wheel, until the null-point (transition) was identified. During each series of 10 measurements a bracketing technique was used, whereby the initial position of the turntable was alternately varied from the point nearest the observer on one measurement (near), to the point farthest from the observer for the next measurement (far). A typical series of 10 sequential measurements is shown in Table 1.

**RESULTS**

Each determination of latency difference is the mean of 10 sequential measurements (see Table 1). The range of the calculated SEM for Δt for all subjects was 0.11-1.80 msec. The data are shown in Tables 1-5 and in Figs

TABLE 4. Comparison of latency difference determinations in 10 subjects

Subject	IPD (mm)	OD = 0.7 Ω = 33.3 rev/min		OD = 0.7 Ω = 44.7 rev/min	
		d <sub>T</sub> (cm) Mean ± SEM (range)	Δt(msec) Mean ± SEM (range)	d <sub>T</sub> (cm) Mean ± SEM (range)	Δt(msec) Mean ± SEM (range)
PO	60.9	228.1 ± 6.2 (197.4-260.6)	7.8 ± 0.22 (6.7-8.9)	195.3 ± 5.7 (171.0-224.0)	6.7 ± 0.19 (5.8-7.6)
RWDN	66.4	234.9 ± 3.5 (206.2-261.4)*	8.2 ± 0.12 (7.4-9.3)*	171.1 ± 2.4 (157.4-181.7)	8.4 ± 0.12 (7.9-9.1)
EAN	61.3	229.0 ± 3.3 (210.1-244.4)	7.7 ± 0.11 (7.2-8.4)	178.3 ± 2.8 (165.6-191.3)	7.4 ± 0.12 (6.9-7.9)
RC	68.0	214.1 ± 3.6 (195.3-239.1)	9.1 ± 0.15 (8.2-10.0)	180.5 ± 3.3 (163.0-195.5)	8.1 ± 0.15 (7.4-8.9)
SM	61.9	210.2 ± 10.1 (171.0-263.7)	8.6 ± 0.41 (6.7-10.3)	179.4 ± 14.2 (143.5-229.1)†	7.5 ± 0.57 (5.7-9.2)†
MC	61.2	219.0 ± 12.6 (168.1-261.8)	8.3 ± 0.50 (6.7-10.5)	198.2 ± 11.3 (150.1-234.1)	6.8 ± 0.42 (5.6-8.7)
IJ	60.3	205.9 ± 9.1 (168.0-245.5)	8.6 ± 0.38 (7.1-10.4)	193.0 ± 7.3 (155.5-224.4)	6.8 ± 0.27 (5.8-8.3)
AM	65.1	190.4 ± 11.4 (143.4-246.8)	10.1 ± 0.61 (7.5-13.0)	165.6 ± 8.0 (132.7-199.5)	8.6 ± 0.41 (7.0-10.5)
PN	62.7	198.9 ± 5.0 (169.8-219.6)	9.1 ± 0.23 (8.2-10.6)	182.7 ± 3.0 (171.9-199.6)	7.3 ± 0.12 (6.9-7.8)
MK	64.4	202.5 ± 8.2 (169.8-244.2)	9.2 ± 0.36 (7.5-10.9)	190.1 ± 10.1 (144.4-252.5)	7.4 ± 0.38 (5.4-9.5)
	<i>n</i>	10	10	10	10
	Mean	213.3	8.7	183.4	7.5
	SEM	4.60	0.23	3.4	0.21
		Mean angular velocity at the eye = 7.9 deg/sec r.m.s.		Mean angular velocity at the eye 12.3 deg/sec r.m.s.	

The mean within-subject difference in Δt is significant (P < 0.001). Each determination is the mean of 10 measurements (†n = 6). \*Indicates pooled data from two separate determinations.

TABLE 5. Comparison of latency difference determinations in 11 subjects

Subject	IPD (mm)	OD = 1.5 $\Omega = 16.6$ rev/min		OD = 1.5 $\Omega = 33.3$ rev/min	
		$d_T$ (cm) Mean $\pm$ SEM (range)	$\Delta t$ (msec) Mean $\pm$ SEM (range)	$d_T$ (cm) Mean $\pm$ SEM (range)	$\Delta t$ (msec) Mean $\pm$ SEM (range)
PO	60.9	204.0 $\pm$ 7.1 (171.6–240.2)	17.3 $\pm$ 0.62 (14.5–20.3)	142.8 $\pm$ 2.3 (125.5–160.7)*	12.3 $\pm$ 0.21 (10.8–14.0)*
RWDN	66.4	179.4 $\pm$ 9.4 (143.2–221.2)	21.7 $\pm$ 1.10 (17.4–26.4)	119.9 $\pm$ 1.0 (112.1–130.1)*	15.9 $\pm$ 0.14 (14.5–17.1)*
EAN	61.3	156.4 $\pm$ 4.3 (135.9–182.8)	22.5 $\pm$ 0.62 (19.2–25.6)	120.5 $\pm$ 2.5 (99.6–135.8)*	14.7 $\pm$ 0.31 (13.1–17.5)*
RC	68.0	166.4 $\pm$ 4.8 (143.5–195.1)	23.6 $\pm$ 0.64 (20.2–27.1)	111.4 $\pm$ 1.9 (96.1–128.3)*	17.6 $\pm$ 0.30 (15.1–20.3)*
SM	61.9	188.8 $\pm$ 8.5 (152.3–238.2)	19.3 $\pm$ 0.85 (15.0–23.5)	130.3 $\pm$ 2.0 (106.4–151.9)*	13.7 $\pm$ 0.22 (11.7–16.7)*
MC	61.2	182.7 $\pm$ 10.8 (140.7–223.1)	20.1 $\pm$ 1.20 (15.8–25.3)	129.2 $\pm$ 3.3 (109.0–162.4)*	13.8 $\pm$ 0.33 (10.9–16.2)*
MH	57.9	142.2 $\pm$ 3.7 (120.3–156.1)	23.5 $\pm$ 0.68 (21.0–27.7)	103.6 $\pm$ 1.6 (88.7–116.0)*	16.1 $\pm$ 0.26 (14.3–18.8)*
SY	64.0	168.2 $\pm$ 3.0 (150.1–182.8)	21.9 $\pm$ 0.41 (20.0–24.7)	112.8 $\pm$ 1.3 (105.3–126.4)*	16.3 $\pm$ 0.18 (14.4–17.4)*
SR	63.5	146.8 $\pm$ 9.2 (113.2–199.5)	25.6 $\pm$ 1.52 (18.1–32.7)	109.4 $\pm$ 1.8 (92.4–123.4)*	16.8 $\pm$ 0.30 (14.8–19.8)*
MK	64.4	197.4 $\pm$ 16.7 (131.0–269.2)	20.0 $\pm$ 1.71 (13.8–27.9)	141.4 $\pm$ 5.8 (123.5–185.3)	13.2 $\pm$ 0.48 (9.9–14.9)
NGH	63.1	162.5 $\pm$ 12.7 (114.7–223.8)	23.6 $\pm$ 1.80 (16.3–31.3)	100.8 $\pm$ 1.4 (90.8–106.2)	18.0 $\pm$ 0.26 (17.1–20.0)
<i>n</i>		11	11	11	11
Mean		172.3	21.7	120.2	15.3
SEM		6.0	0.72	4.31	0.57
		Mean angular velocity at the eye = 4.9 deg/sec r.m.s.		Mean angular velocity at the eye = 14.1 deg/sec r.m.s.	

The mean within-subject difference in  $\Delta t$  is significant ( $P < 0.001$ ). Each determination is the mean of 10 measurements. \*Indicates pooled data from two separate determinations.

2–4. Paired and unpaired data were analysed using a two-tailed Student's *t*-test.

*Influence of filter density on latency difference ( $\Omega = 33.3$  rev/min)*

These data, which are presented in order to serve as a comparison with other latency difference studies in the literature, are presented in Table 2 and Fig. 2.

These results indicate that for each subject ( $\Omega = 33.3$  rev/min), the latency difference using the 1.5 OD filter was greater than that that using the 0.7 OD filter. The mean ( $\pm$  SEM) within-subject difference in  $\Delta t$  ( $6.3 \pm 0.52$  msec;  $n = 10$ ) is significant ( $P < 0.001$ ). In this group the mean ( $\pm$  SEM) latency difference using the 1.5 OD filter ( $15.0 \pm 0.58$  msec;  $n = 10$ ) is significantly different ( $P < 0.001$ ) from that using the 0.7 OD filter ( $8.7 \pm 0.23$  msec;  $n = 10$ ).

In nine subjects repeat latency difference determinations were made (see Table 3) following a mean interval

of 182 days (range: 1–439). There was no significant within-subject difference between the two determinations ( $P > 0.1$ ); the mean ( $\pm$  SEM) within-subject difference in  $\Delta t$  being  $0.04 \pm 0.42$  msec ( $n = 9$ ). All repeat determinations were made using the same turntable speed (33.3 rev/min), neutral-density filter (1.5 OD), and illumination.

*Influence of angular velocity on latency difference*

The latency difference data are presented in terms of turntable speed (Fig. 3) and in terms of mean (r.m.s.) angular velocity at the eye (Fig. 4).

The mean angular velocity at the eye (deg/sec) of the rotating rod was determined as the root mean square (r.m.s.) angular velocity, and shown in Tables 4 and 5. This was calculated from the turntable speed and the mean viewing distance at transition, using the formula for the instantaneous angular velocity described in Appendix B.




FIGURE 2. Influence of optical density on visual latency difference in 10 subjects ( $\Omega = 33.3$  rev/min). For data see Table 2.




FIGURE 3. Influence of turntable speed on visual latency difference in 11 subjects. For data see Tables 4 and 5.




FIGURE 4. Influence of angular velocity at the eye on visual latency difference in 11 subjects. For data see Tables 4 and 5.




FIGURE 5. Hypothetical relationship between turntable angular velocity and visual latency difference. The bars indicate the range of data points given in Tables 4 and 5 and shown in Fig. 3.

**0.7 OD filter.** These results (see Table 4 and Figs 3 and 4) indicate that with the exception of one subject (RWDN), both the latency difference and the viewing distance at transition associated with the slower turntable speed (33.3 rev/min), were greater than those associated with the faster turntable speed (44.7 rev/min).

The mean ( $\pm$  SEM) within-subject difference in  $\Delta t$  ( $1.2 \pm 0.21$  msec;  $n = 10$ ) is significant ( $P < 0.001$ ). In this group, the mean ( $\pm$  SEM) latency difference determination at 33.3 rev/min ( $8.7 \pm 0.23$  msec;  $n = 10$ ) is significantly different ( $P < 0.001$ ) from that determined at 44.7 rev/min ( $7.5 \pm 0.21$  msec;  $n = 10$ ).

**1.5 OD filter.** These results (see Table 5 and Figs 3 and 4) indicate that for each subject, both the latency difference and the viewing distance at transition associated with the slower turntable speed (16.6 rev/min), were greater than those associated with the faster turntable speed (33.3 rev/min).

The mean ( $\pm$  SEM) within-subject difference in  $\Delta t$  ( $6.4 \pm 0.35$  msec;  $n = 11$ ) is significant ( $P < 0.001$ ). In this group the mean ( $\pm$  SEM) latency difference determination at 33.3 rev/min ( $15.3 \pm 0.57$  msec;  $n = 11$ ) is significantly different ( $P < 0.001$ ) from that determined at 16.6 rev/min ( $21.7 \pm 0.72$  msec;  $n = 11$ ).

## DISCUSSION

### Variation of latency difference with optical density

This study indicates that the variation of latency difference with filter density (see Fig. 2) for the given illumination using the rotating Pulfrich effect, is in close agreement with both

- (i) the data of Prestrude and Baker (1968) using concentric rotating lines with similar filters and illumination; and
- (ii) the data of Standing *et al.* (1968) using similar filters and a vertical rod.

These results therefore further validate the use of the rotating Pulfrich technique (Nickalls, 1986a) for the measurement of visual latency differences.

In addition the present study also indicates that latency

difference determinations made using the rotating Pulfrich technique for the 1.5 OD/33.3 rev/min combination under identical circumstances, are reproducible over many months (see Table 3).

### Variation of latency difference with angular velocity

The major finding of this study is that for the interocular luminance differences used, visual latency difference was found to vary significantly with both turntable speed and mean (r.m.s.) angular velocity at the eye.

When using the 1.5 OD filter, there was a significant inverse relationship between latency difference and both turntable speed and mean (r.m.s.) angular velocity at the eye (see Figs 3 and 4). Similarly, the data obtained using the 0.7 OD filter was also in keeping with this inverse relationship, with the mean latency difference at 33.3 rev/min being significantly greater than that at 44.7 rev/min.

### Other relevant studies

The most relevant study in the literature appears to be that of Lit (1960a), who investigated the relationship between latency difference and angular velocity using a black vertical rod moving with constant linear horizontal velocity (range 1.5–31.8 deg/sec), at four separate binocular illuminance differences.

In this study Lit observed a significant non-linear inverse relationship between latency difference and target angular velocity at the eye for angular velocities less than about 20–25 deg/sec, which became progressively more pronounced as the binocular illuminance difference was increased. For any given difference in binocular illuminance, visual latency difference decreased progressively to a plateau as the angular velocity at the eye increased.

However, Lit gives no estimate of the precision of his observations. Furthermore, Lit determined the apparent displacement of the moving target using an adjustable pointer which was fixated by the subject, and in view of the difference between foveal and extra-foveal latency, this may have introduced some error [see Nickalls (1986a)].

Significantly, similar but rather more subjective evidence for an increase in visual latency as angular velocity decreases has also been described in association with a number of "sensation-time" experiments; for example those by Fröhlich (1923), Holz (1934) and others, which have been well summarized by Lit (1960a).

### Hypothesis

The present study shows conclusively that there is a significant inverse relationship between visual latency difference and turntable angular velocity [and hence with mean (r.m.s.) target angular velocity at the eye] within the parameter range studied. In view of these findings, and those of Lit (1960a), Fröhlich (1923) and Holz (1934), the author suggests the hypothesis that for a given inter-ocular illuminance difference there exists a continuous inverse relationship between visual latency difference and turntable angular velocity as shown in Fig. 5.

The mechanism by which velocity influences visual latency difference is not clear. However, it has recently been shown that motion produces equivalent spatial blur which is velocity dependent (Pääkkönen & Morgan, 1994), and it is possible, therefore, that there may be an association between smaller blur (slow velocity) and longer latency. Alternatively, this effect may be related to properties of the different motion sensor systems which process slow and fast velocities [see Hawken *et al.* (1994)].

### REFERENCES

- Barlow, H. B. & McNaughton, P. A. (1980). Illusory curvature caused by retinal delay. *Journal of Physiology, London*, 308, 11P–12P.
- Carney, T., Paradiso, M. A. & Freeman, R. D. (1989). A physiological correlate of the Pulfrich effect in cortical neurons of the cat. *Vision Research*, 29, 155–165.
- Enright, J. T. (1970). Distortions of apparent velocity: A new optical illusion. *Science*, 168, 464–467.
- Fröhlich, F. W. (1923). Über die Abhängigkeit der Empfindungszeit und des zeitlichen Verlaufes der Gesichtsempfindung von der Intensität Dauer und Geschwindigkeit der Belichtung. *Zeitung Sinnesphysiologie*, 55, 1–46.
- Fry, G. A. & Hill, W. W. (1962). The center of rotation of the eye. *American Journal of Optometry and Archives of the American Academy of Optometry*, 39, 581–595.
- Harker, G. S. (1967). A saccadic suppression explanation of the Pulfrich phenomenon. *Perception and Psychophysics*, 2, 423–426.
- Hawken, M. J., Gegenfurtner, K. R. & Tang, C. (1994). Contrast dependence of colour and luminance in human vision. *Nature*, 367, 268–270.
- Hofeldt, A. J., Leavitt, J. & Behrens, M. M. (1985). Pulfrich stereoisolation phenomenon in serous sensory retinal detachment of the macula. *American Journal of Ophthalmology*, 100, 576–580.
- Holz, J. (1934). Der Stereoeffekt Pulfrich's und die Empfindungszeit. *Zeitung Biologie*, 95, 502–516.
- Julesz, B. & White, B. (1969). Short term visual memory and the Pulfrich phenomenon. *Nature*, 222, 639–641.
- Larkin, E. B., Dutton, G. N. & Heron, G. (1994). Impaired perception of moving objects after minor injuries to the eye and midface: The Pulfrich phenomenon. *British Journal of Oral and Maxillofacial Surgery*, 32, 360–362.
- Lee, D. N. (1970). A stroboscopic stereophenomenon. *Vision Research*, 10, 587–593.
- Levick, W. R., Cleland, B. G. & Coombs, J. S. (1972). On the apparent orbit of the Pulfrich pendulum. *Vision Research*, 12, 1381–1388.
- Lit, A. (1949). The magnitude of the Pulfrich stereophenomenon as a function of binocular differences of intensity at various levels of illumination. *American Journal of Psychology*, 62, 159–181.
- Lit, A. (1960a). The magnitude of the Pulfrich stereophenomenon as a function of target velocity. *Journal of Experimental Psychology: Human Perception*, 59, 165–175.
- Lit, A. (1960b). Effect of target velocity in a frontal plane on binocular spatial localisation at photopic retinal illuminance levels. *Journal of the Optical Society of America*, 50, 970–973.
- Lit, A. (1960c). Magnitude of the Pulfrich stereophenomenon as a function of target thickness. *Journal of the Optical Society of America*, 50, 321–327.
- Lit, A. (1964). Equidistance settings at photopic retinal illuminance levels as a function of target velocity in a frontal plane. *Journal of the Optical Society of America*, 54, 83–88.
- Lit, A. & Hyman, A. (1951). The magnitude of the Pulfrich stereophenomenon as a function of distance of observation. *American Journal of Optometry*, 28, 564–580.
- Lythgoe, R. J. (1938). Some observations on the rotating pendulum. *Nature*, 141, 474.
- Morgan, M. J. (1975). Stereoillusion based on visual persistence. *Nature*, 256, 639–640.
- Morgan, M. J. & Thompson, P. (1975). Apparent motion and the Pulfrich effect. *Perception*, 4, 3–18.
- Nickalls, R. W. D. (1986a). The rotating Pulfrich effect, and a new method of determining visual latency differences. *Vision Research*, 26, 367–372.
- Nickalls, R. W. D. (1986b). A new line and conic theorem having an interesting visual correlate. *Mathematical Gazette*, 70, 27–29.
- Pääkkönen, A. K. & Morgan, M. J. (1994). Effects of motion on blur discrimination. *Journal of the Optical Society of America (A)*, 3, 992–1002.
- Pulfrich, C. (1922). Die Stereoskopie im Dienste der isochromen und heterochromen Photometrie. *Naturwissenschaften*, 10, 553–564.
- Prestrude, A. M. & Baker, H. D. (1968). New method of measuring visual-perceptual latency differences. *Perception and Psychophysics*, 4, 152–154.
- Ross, J. & Hogben, J. H. (1975). The Pulfrich effect and short-term memory in stereopsis. *Vision Research*, 15, 1289–1290.
- Sokol, S. (1976). The Pulfrich stereo-illusion as an index of optic nerve dysfunction. *Survey of Ophthalmology*, 20, 432–434.
- Spiegler, J. B. (1983). Distance, size, and velocity changes during the Pulfrich effect. *American Journal of Optometry and Physiological Optics*, 60, 902–907.
- Spiegler, J. B. (1986). Apparent path of a Pulfrich target as a function of the slope of its plane of motion: A theoretical note. *American Journal of Optometry and Physiological Optics*, 63, 209–216.
- Standing, L. G., Dodwell, P. C. & Lang, D. (1968). Dark adaptation and the Pulfrich effect. *Perception and Psychophysics*, 4, 118–120.
- Trincker, D. (1953). Hell–Dunkel–Anpassung und raumliches Sehen. *Pflügers Archives gesicht Physiology*, 257, 48–69.
- Weale, R. A. (1954). Theory of the Pulfrich effect. *Ophthalmologica*, 128, 380–388.
- Williams, J. M. & Lit, A. (1983). Luminance-dependent visual latency for the Hess effect, the Pulfrich effect, and simple reaction time. *Vision Research*, 23, 171–179.
- Wilson, G. S. (1965). An investigation of the Pulfrich effect. *British Journal of Physiological Optics*, 22, 208–237.

*Acknowledgements*—The author would like to thank Professor A. L. Crombie (Department of Ophthalmology, University of Newcastle-upon-Tyne, U.K.) for help and encouragement with this study; Mr P. Byrnes and Mr P. Oliver (Department of Medical Physics, General Hospital, Newcastle-upon-Tyne, U.K.) for making the apparatus; and Dr K. Mitchell (Department of Medical Physics, Royal Victoria Infirmary, Newcastle-upon-Tyne, U.K.) for assistance with luminance measurements.

## APPENDIX A

*Latency Difference*

The following relationship for the rotating Pulfrich effect, between the latency difference ( $\Delta t$  sec), the viewing distance ( $d_T$  cm) at which the 'transition' null-point is perceived, the value of half the separation of the rotation centres of the eyes ( $a$  cm), and the angular velocity of the turntable ( $\omega$  deg/sec), was derived by Nickalls (1986a).

$$\Delta t = (2/\omega)\tan^{-1}(a/d_T)$$

However, in the present study the angular velocity of the turntable was calibrated in revolutions per minute ( $\Omega$  rev/min). Since 1 rev/min = 6 deg/sec, then the above equation (where  $\omega$  is in deg/sec) can be modified to become Eq. (1) where  $\Omega$  is in rev/min, as follows.

$$\Delta t = (1/3\Omega)\tan^{-1}(a/d_T)$$

## APPENDIX B

*Target angular velocity at eye*

Let the eyes be in the same plane as the rotating rod, and let the centre of rotation O lie in the subject's sagittal plane. Let the radius of rotation be  $r$  about the centre of rotation O. If O is a distance  $d$  from the eye ( $d > r$ ), then it can be shown that the instantaneous angular velocity of the rod at the eye ( $d\psi/dt$ ) is given by:

$$\frac{d\psi}{dt} = \omega \left[ \frac{r(d \cos \theta - r)}{r^2 + d^2 - 2rd \cos \theta} \right]$$

where the angle  $\theta$  defines the instantaneous position of the rod about the centre of rotation O, and  $\omega$  is the speed of rotation of the rod. The rod is at nearest approach to the eye when  $\psi = \theta = 0$ .