

Desensitization and Resensitization Kinetics of Glutamate Receptor Channels from *Drosophila* Larval Muscle

M. Heckmann and J. Dudel

Physiologisches Institut der Technischen Universität, 80802 München, Germany

ABSTRACT Outside-out patches from wild-type *Drosophila* larval muscle were exposed to L-glutamate (glu) using a piezo-driven application system. Glu receptor channels opened and desensitized in response to rapid applications of 10 mM glu. Desensitization was fitted with an exponential function with a mean time constant of desensitization (τ_d) of 15 ms in response to 10 mM glu. The τ_d was concentration dependent and decreased to 6 ms (on average) with 0.7 mM glu and increased again to 12 ms (on average) in response to 0.5 mM glu. Desensitization in response to longer applications of glu was almost complete, but surprisingly, even a 1-ms pulse of 3 mM glu produced about 30% desensitization. In the presence of low glu concentrations, the response to a pulse was reduced and was about halved by preequilibration with 30 μ M glu. Recovery from desensitization was not concentration dependent and was fitted with an exponential function with a mean time constant of 150 ms. During recovery the channels rarely opened. Kinetic schemes were fitted to these results, and a circular reaction scheme was found to fit the data best. An important feature of the scheme is desensitization from a lower ligated closed state. This allows substantial desensitization of synaptic receptor channels in response to quantal release of transmitter, in part without opening of the channels. Desensitization reduces the probability of the channels opening in response to a subsequent release for a period of time determined by the rate of recovery from desensitization and might serve as a form of molecular short-term memory.

INTRODUCTION

Glutamate (glu) serves as an excitatory neurotransmitter in vertebrate and invertebrate nervous systems and activates a number of different receptors. Fast postsynaptic currents after glu release at a synapse are mediated by glu receptor channels (Usherwood, 1994; Edmonds et al., 1995). Glu receptor channels can be subdivided into two main subtypes, the AMPA/KA (α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid/kainate) and NMDA (*N*-methyl-D-aspartic acid) receptor channels (Seeburg, 1993; Wisden and Seeburg, 1993; Hollmann and Heinemann, 1994). Postsynaptic currents mediated by AMPA/KA receptor channels rise and decay faster than those mediated by NMDA receptor channels (Jonas and Spruston, 1994), and rapid desensitization within milliseconds in response to glu appears to be typical for AMPA/KA receptor channels (Franke et al., 1987; Dudel et al., 1988, 1990b; Colquhoun et al., 1992). The time course of desensitization and resensitization of AMPA/KA receptor channels might therefore influence the transfer of information at a synapse (Dudel et al., 1992; Trussell et al., 1993).

In the present study we investigated the desensitization and resensitization kinetics of glu receptor channels from *Drosophila* larval muscle. These channels are the functional equivalent of vertebrate muscle nicotinic acetylcholine re-

ceptors and are considered to be distantly related to vertebrate AMPA/KA receptor channels (Schuster et al., 1991). To gain insight into the kinetics of these channels, we applied glu rapidly in a number of protocols to outside-out patches of muscle membrane. We explored several reaction schemes in computer simulations and found a scheme that agrees with our experimental data. Based on our results, desensitization is likely to be of functional relevance at glutamatergic synapses with similar receptor channels.

MATERIALS AND METHODS

Preparation

Wild-type *Drosophila melanogaster* strain Oregon R was reared on standard *Drosophila* medium in bottles as described (Heckmann and Dudel, 1995). Third-instar larvae were dissected, pinned on a thin layer of sylgard (Jan and Jan, 1976), and moved to the stage of an upright microscope fitted with a 20 \times 0.4 NA objective (LD Achroplan, 20 mm working distance; Zeiss, Oberkochen, Germany). The preparation was bathed in Schneiders *Drosophila* medium (Life Technologies GmbH, Eggenstein, Germany) containing \sim 30 μ g/ml collagenase type 1A (Sigma-Aldrich Chemie GmbH, Deisenhofen, Germany) for approximately 15 min before patching and then superfused with an extracellular solution containing 135 mM NaCl, 5 mM KCl, 4 mM MgCl₂, 2 mM CaCl₂, and 5 mM HEPES, with the pH adjusted to 7.4 with 5 N NaOH.

Electrophysiology

The measurements were made at room temperature (18–22°C). Patch pipettes were pulled from thick-walled borosilicate glass tubes with filament (outer diameter 2.0 mm, internal diameter 1.16 mm; Clark, Pangbourne, England), with a DMZ-Universal Puller (Zeitz Instruments, Augsburg, Germany). The pipettes had resistances of \sim 5 M Ω when filled with intracellular solution containing 150 mM K-propionate, 5 mM Na-propionate, 10 mM MgCl₂, 1 mM CaCl₂, 10 mM EGTA, and 10 mM Tris-maleate buffer, with the pH adjusted to 7.4 with 5 N NaOH. Outside-out

Received for publication 31 October 1996 and in final form 7 February 1997.

Address reprint requests to Dr. Manfred Heckmann, Physiologisches Institut der Technischen Universität München, Biedersteiner Strasse 29, D-80802 München, Germany. Tel.: 0049-89-4140-3144; Fax: 0049-89-4140-3377; E-mail: heckmann@physiol.med.tu-muenchen.de.

© 1997 by the Biophysical Society

0006-3495/97/05/2160/10 \$2.00

patches were obtained as described previously (Hamill et al., 1981). They were taken from muscles 6 and 7 of the abdominal segments of the larvae (Atwood et al., 1993) and were voltage clamped to -60 mV.

Fast application of agonists

Glu was applied with a liquid filament switch (Franke et al., 1987). A P 810.30 piezo element (Physik Instrumente, Waldbronn, Germany) held in a brass cylinder was used with a glass tube (outer diameter 0.5 mm, wall thickness 0.175 mm; Hilgenberg, Malsfeld, Germany) connected to it. The tube was perfused with extracellular solution, to which glu was added at the desired concentration. The performance of the application system was tested by perfusing the glass tube with diluted extracellular solution and switching the ionic concentration at the tip of patch pipettes. The 10–90% rise and decay times of this change in liquid junction current with open pipettes were below 100 μ s.

Data collection and analysis

We used an Axopatch 200A amplifier (Axon Instruments, Foster City, CA) and filtered the data at 5 kHz bandwidth (-3 dB) with an 8-pole low-pass Bessel filter. The data were recorded with ISO₂ software (MFK Computer, Niederhausen, Germany) and a PC 486 (sampling rate of 50 kHz, 12-bit resolution) and stored on the hard disk of a computer. Averaging and fitting of exponential curves was done off line with ISO₂.

Computer simulations

Simulations of reaction schemes and sets of rate constants were performed with a program using a variable-coefficient ordinary differential equation solver, with fixed-leading coefficient implementation (Brown et al., 1989) for solving the sets of differential equations on a workstation.

RESULTS AND DISCUSSION

Several paths to desensitization

Rapid application of glu to outside-out patches from *Drosophila* larval muscle elicits openings of glu receptor channels. Responses to 100-ms applications of 10 mM glu to a patch held at -60 mV are shown in Fig. 1. Two single responses and the average current response to 100 consecutive applications at an interval of 1 Hz are shown. In this patch up to two channels opened simultaneously. The channel openings started with a very short delay after the beginning of the applications. For some milliseconds the channels flickered rapidly between open and closed conformations and then remained closed in the continued presence of glu. The average current reached a peak value of -8 pA. The 10–90% current rise time was 200 μ s, and the current decay was fitted well by a single exponential function with a time constant of desensitization (τ_d) of 10 ms.

We investigated the dependence of channel open probability and current rise time on glu concentration (Scheme 1). R represents a closed channel, A represents a glu molecule and AR, A₂R, A₃R, A₄R, A₅R, and A₅O represent a closed





FIGURE 1 Desensitization of glutamate receptor channels on an outside-out patch. Two single responses to 100-ms pulses of 10 mM glutamate and the average current response to 100 applications are shown. The decay of the average current was fitted by a single exponential function with a time constant of 10 ms. The patch was held at -60 mV, the pulse frequency was 1 Hz, and the data were filtered to a final cutoff frequency of 5 kHz. The time course of the application is indicated at the top.

and an open channel with one, two, three, four, and five glu molecules bound. The five binding steps were necessary to reproduce the current increase proportional to the power $n = 4.4$ of the glu concentration between 0.3 and 0.5 mM glu. The units of the rate constants are s^{-1} , and the binding rate constants are $M^{-1} s^{-1}$ (Heckmann et al., 1996b). We increased the channel opening rate constant, β , from 15,000 s^{-1} to 19,000 s^{-1} , because the latter value reproduces our data slightly better.

In response to 10 mM glu in 23 patches we observed values for the τ_d ranging from 7 to 29 ms, with a mean value of 14.6 ± 5.4 ms. The τ_d was thus quite variable between patches but was well fitted with one exponential, indicating that individual patches contained channels with similar kinetics. Variability of the τ_d from patch to patch has been reported for channels from a number of preparations (Dudel et al., 1990a,c; Franke et al., 1993; von Beckerath et al., 1995; Heckmann et al., 1996a) and might be explained by differences in subunit combinations (Seeburg, 1993; Lomeli et al., 1994) or reversible modifications like phosphorylation or dephosphorylation of the channels (Huganir and Greengard, 1990). A reversible mechanism seems more likely in our case, because we observed that the τ_d increased or decreased occasionally during experiments (data not shown).



SCHEME 1




FIGURE 3 The time constant of desensitization is concentration dependent. (A) Plot of the time constant of desensitization (τ_d) versus glutamate concentration on a logarithmic scale. Average currents from 50–300 single responses to 100-ms pulses of 0.2–10 mM glutamate concentration were evaluated. The results are from 30 patches. (B) Plot of the rate of desensitization, $1/\tau_d$, versus the peak open probability of the channel (p_o). The p_o with 10 mM Glu was taken to be 0.6; each symbol represents a different patch ($n = 6$).

mM glu, respectively. In response to 0.5 mM glu concentrations, the τ_d increased again to an average value of 11.6 ± 5.7 ms, $n = 9$. We compared the τ_d values by using a rank-sum test allowing multiple comparisons (Dunn,

1964; Rosner, 1990) and obtained a $p < 0.01$ for the responses to 10 mM not to be different from those to 1 mM glu, and a $p < 0.06$ for the responses to 1 mM not to be different from those to 0.5 mM glu. The τ_d values measured




FIGURE 4 Modeling the concentration dependence of desensitization. Average current responses of a patch to 60, 95, and 391 applications of 10, 0.7, and 0.2 mM glutamate, respectively, are shown in the left column, normalized with respect to the peak current. The time constants of desensitization were 24, 10, and 14 ms with 10, 0.7, and 0.2 mM glutamate, respectively. The pulse duration was 150 ms. Dotted lines mark the baseline current level; the other experimental conditions were as in Fig. 1. The results of computer simulations of this experiment with Schemes 2, 3, and 4 are shown in the other columns. The simulated current traces are scaled with respect to the experimental data obtained in response to 10 mM glutamate. The concentration dependence of the average peak current is fitted reasonably by all three reaction schemes, but only Scheme 4 reproduces the time course of desensitization over the whole concentration range. We used the following rate constants of desensitization and resensitization for the simulations: $d_{+1} = 100 \text{ s}^{-1}$, $d_{+2} = 500 \text{ s}^{-1}$, $d_{-1} = 1.2 \text{ s}^{-1}$, and $d_{-2} = 6 \text{ s}^{-1}$.




FIGURE 5 Channel openings during recovery from desensitization are rare. In response to 200-ms applications of 10 mM glutamate, up to 20 channels opened simultaneously in this patch. After 5 out of 100 pulses, we observed brief openings of channels after the end of the application. Two traces with channel openings during recovery from desensitization are shown. These “reopenings” are shown again at higher amplification and on an increased time scale in the insets. The filter cutoff frequency in this experiment was 2 kHz; the other experimental conditions were as in Fig. 1.

pulse of low glu concentration. The reason is that the channels should desensitize predominantly to A_5D and A_3D during the high and low concentration pulses, respectively. The rate of recovery from A_5D has to be lower than that from A_3D to reproduce the desensitization in response to a long pulse. For a reasonable fit of current traces like the ones shown in Figs. 1, 2, and 4, values of about 1 s^{-1} and 6 s^{-1} are necessary for d_{-1} and d_{-2} .


We investigated the time course of recovery from desensitization in response to 100-ms double pulses of 10 and 0.7 mM glu. Average current responses from one patch in response to pulses of 10 mM glu are shown in Fig. 6 A. With only 30 ms between the first and second pulses, the response to the second pulse was markedly reduced because of desensitization of the channels. After 1 s the response to the second pulse reached the amplitude of the first response. Results from four such experiments with pulses of 10 and 0.7 mM glu are shown as mean values \pm standard deviation in Fig. 6, B and C. Recovery was fitted with a single exponential function with a time constant of about 150 ms in response to pulses of the two concentrations. The data are

shown together with results from simulations of these experiments with Scheme 4 and Scheme 5, which will be introduced below. Recovery from desensitization after the 0.7 mM glu pulses is well reproduced by Scheme 4, but this mechanism does not reproduce the recovery after the 10 mM glu pulses.

A circular reaction scheme


To overcome the two deficits of Scheme 4, the excessive number of reopenings, and the too slow recovery from A_5D , an alternative route for recovery from A_5D must be provided. A simple solution is to introduce an additional desensitized state A_4D (Scheme 5) that connects the states A_3D and A_5D in a circular reaction scheme (Katz and Thesleff, 1957; Cachelin and Colquhoun, 1989; Franke et al., 1993; Heckmann et al., 1996a). All we know about the rate constants for the additional binding and dissociation steps is that they should reduce the number of reopenings and speed up the time course of recovery from A_5D . For a circular interconversion the principle of microscopic reversibility requires, in addition, that under equilibrium conditions, the transition frequencies (the product of the rate constants) be the same in both directions (Luger, 1995). Microscopic reversibility can be achieved with the rate constants determined for Scheme 4 by simply assuming that the binding and dissociation rate constants of the desensitized states are the same as those of the respective R states. With these rate constants Scheme 5 reproduced the time course of recovery from desensitization (Fig. 6, A and B) and allowed recovery from desensitization with fewer reopenings.

Some glu receptor channels are desensitized by transmitter concentrations much lower than those needed to open the channels (Dudel et al., 1990b; Colquhoun et al., 1992; Heckmann et al., 1996a). To test for this effect we added glu in low concentrations to the solution superfusing the patches between applications and recorded the responses to 10 mM glu pulses. The average response to 10 mM glu was reduced to half-amplitude by about 30 μM glu (data not shown). In the left column of Fig. 7 are shown the average current responses from one experiment recorded without and during superfusion of the patch with 0.1 mM glu, which reduced the response to about one-quarter the control value. A simulation of this experiment with Scheme 5 reproduced this result and is shown in the next column. Scheme 3



SCHEME 4

FIGURE 7 Low glutamate concentrations desensitize the channels. Average current responses of a patch to applications of 10 mM glutamate are shown in the left column. The lower trace was recorded while 100 μ M glutamate superfused the patch between the pulses, and the upper trace was recorded without such a pre-desensitization. The other experimental conditions were as in Fig. 1. The results of computer simulations of this experiment with Schemes 5 and 3 are shown in the other columns. The simulated current traces are scaled with respect to the experimental data obtained without pre-desensitization.



Why do the channels desensitize so much in response to the brief pulse? A closer look at Scheme 5 helps to understand this. The time course of the probabilities of five of the nine states of Scheme 5 in response to a 1-ms pulse of 3 mM glu are plotted in Fig. 9 A. Before the pulse, the channel is in the unligated closed state (R). The probability of being in this state decreases rapidly after the beginning of the pulse, and the probabilities of the triple ligated closed state (A_3R) and then the open state (A_5O) increase. The probabilities of the desensitized states A_3D , A_4D , and A_5D remain low during the pulse. After the pulse, as the probability of the state A_3R increases again, the probability of the triple ligated desensitized state (A_3D) rises. Most of the desensiti-

zation therefore occurs after the pulse during dissociation of glu and results from desensitization from a lower ligated state combined with sufficiently slow dissociation of glu from the channels. In response to a 1-ms pulse of a lower concentration (for example, 0.5 mM glu, as illustrated in Fig. 9 B), the open state reaches a maximum probability of only 2%; however, the probability of A_3D reaches 10%.

What is the advantage of this kinetic mechanism? The release of glu might be viewed as serving two roles. The first is to create a fast-rising excitatory postsynaptic current of sufficient size and duration to transmit excitation. The concentration of glu in the synaptic cleft probably reaches millimolar concentrations within less than one-tenth of a millisecond within about 100 nm of the point of release (Clements, 1996). According to our kinetic mechanism, receptor channels within this area will open with about 50% probability. Farther from the center of release, lower peak glu concentrations are reached in the synaptic cleft. Because five glu must bind to open the channels, whereas three suffice to desensitize them, the probability of the channels desensitizing falls more slowly than the probability of opening with increasing distance from the center of glu release. Therefore it might actually be that at a synapse more channels are desensitized than opened by glu, depending on the structural relation between the presynaptic release site and postsynaptic receptor channel localization.




FIGURE 8 Desensitization by brief pulses of glutamate. Two 1-ms pulses of 3 mM Glu were applied to a patch at an interval of 10 ms. The current trace is the average of 100 single responses; the other experimental conditions were as in Fig. 1. The result of a computer simulation of this experiment with Scheme 5 is shown scaled with respect to the peak current value of the response to the first pulse in the experimental average.

A testable hypothesis

We attempted to find a reaction scheme that allows us to reproduce the data with a minimum number of states and rate constants. Rate constants might have to be changed and states and pathways added as additional information becomes available. Scheme 5 is intended to serve as a working hypothesis, a summary of previous studies, and a guide for further investigations. The five glu binding sites probably




FIGURE 9 Desensitization of the channels by brief pulses. The results of computer simulations of the time course of the probabilities of the states R, A₃R, A₅O, A₅D, and A₃D in response to a 1-ms pulse of 3 mM glutamate (A) and of the states R, A₃R, A₅O, and A₃D in response to 0.5 mM glutamate (B) with Scheme 5 are shown. Note the extent of desensitization after the pulses (A₃D).

reside in different subunits of the channel (Sutcliffe et al., 1996). The subunits each have the same rate constants of binding and unbinding of glu, but only with three and five glu bound is the channel subject to appreciable desensitization. This may indicate that the channel is a heteropentamer with three nondesensitizing and two desensitizing subunits and might be supported by investigations of the molecular structure of the channel. However, even a homopentamer could change its ability to desensitize with the number of bound glu molecules.

We thank Dr. J. Eilers and Dr. M. Larkum for comments on the manuscript. This work was supported by a grant from the Deutsche Forschungsgemeinschaft (SFB 391).

REFERENCES

- Atwood, H. L., C. K. Govind, and C. F. Wu. 1993. Differential ultrastructure of synaptic terminals on ventral longitudinal abdominal muscles in *Drosophila* larvae. *J. Neurobiol.* 24:1008–1024.
- Brown, P. N., G. D. Byrne, and A. C. Hindmarsh. 1989. VODE: a variable coefficient ODE solver. *SIAM J. Sci. Statist. Comput.* 10:1038–1051.
- Cachelin, A. B., and D. Colquhoun. 1989. Desensitization of the acetylcholine receptor of frog end-plates measured in a Vaseline-gap voltage clamp. *J. Physiol. (Lond.)* 415:159–188.
- Clements, J. D. 1996. Transmitter timecourse in the synaptic cleft: its role in central synaptic function. *Trends Neurosci.* 19:163–171.
- Colquhoun, D., P. Jonas, and B. Sakmann. 1992. Action of brief pulses of glutamate on AMPA/kainate receptors in patches from different neurones of rat hippocampal slices. *J. Physiol. (Lond.)* 458:261–287.
- Dudel, J., C. Franke, and H. Hatt. 1990a. A family of glutamatergic, excitatory channel types at the crayfish neuromuscular junction. *J. Comp. Physiol. A.* 166:757–768.
- Dudel, J., C. Franke, and H. Hatt. 1990b. Rapid activation, desensitization, and resensitization of synaptic channels of crayfish muscle after glutamate pulses. *Biophys. J.* 57:533–545.
- Dudel, J., C. Franke, and H. Hatt. 1992. Rapid activation and desensitization of transmitter-liganded receptor channels by pulses of agonists. In *Ion Channels*, Vol. 3. T. Narahashi, editor. Plenum Press, New York. 207–260.
- Dudel, J., C. Franke, H. Hatt, R. L. Ramsey, and P. N. Usherwood. 1988. Rapid activation and desensitization by glutamate of excitatory, cation-selective channels in locust muscle. *Neurosci. Lett.* 88:33–38.
- Dudel, J., C. Franke, H. Hatt, R. L. Ramsey, and P. N. Usherwood. 1990c. Glutamatergic channels in locust muscle show a wide time range of desensitization and resensitization characteristics. *Neurosci. Lett.* 114:207–212.
- Dudel, J., C. Franke, and W. Luboldt. 1993. Reaction scheme for the glutamate-ergic, quisqualate type, completely desensitizing channels on crayfish muscle. *Neurosci. Lett.* 158:177–180.
- Dunn, O. J. 1964. Multiple comparisons using rank sums. *Technometrics.* 6:241–252.
- Edmonds, B., A. J. Gibb, and D. Colquhoun. 1995. Mechanisms of activation of glutamate receptors and the time course of excitatory synaptic currents. *Annu. Rev. Physiol.* 57:495–519.
- Franke, C., H. Hatt, and J. Dudel. 1987. Liquid filament switch for ultra-fast exchanges of solutions at excised patches of synaptic membrane of crayfish muscle. *Neurosci. Lett.* 77:199–204.
- Franke, C., H. Parnas, G. Hovav, and J. Dudel. 1993. A molecular scheme for the reaction between acetylcholine and nicotinic channels. *Biophys. J.* 64:339–356.
- Hamill, O. P., A. Marty, E. Neher, B. Sakmann, and F. J. Sigworth. 1981. Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflügers Arch.* 391:85–100.
- Heckmann, M., J. Büfler, C. Franke, and J. Dudel. 1996a. Kinetics of homomeric GluR6 glutamate receptor channels. *Biophys. J.* 71:1743–1750.
- Heckmann, M., and J. Dudel. 1995. Recordings of glutamate-gated ion channels in outside-out patches from *Drosophila* larval muscle. *Neurosci. Lett.* 196:53–56.
- Heckmann, M., F. Parzefall, and J. Dudel. 1996b. Activation kinetics of glutamate receptor channels from wild-type *Drosophila* muscle. *Pflügers Arch.* 432:1023–1029.
- Hestrin, S. 1992. Activation and desensitization of glutamate-activated channels mediating synaptic currents in the visual cortex. *Neuron.* 9:991–999.
- Hollmann, M., and S. Heinemann. 1994. Cloned glutamate receptors. *Annu. Rev. Neurosci.* 17:31–108.
- Huganir, R. L., and P. Greengard. 1990. Regulation of neurotransmitter receptor desensitization by protein phosphorylation. *Neuron.* 5:555–567.
- Jan, L. Y., and Y. N. Jan. 1976. Properties of the larval neuromuscular junction in *Drosophila melanogaster*. *J. Physiol. (Lond.)* 262:189–214.
- Jonas, P., G. Major, and B. Sakmann. 1993. Quantal components of unitary EPSCs at the mossy fibre synapse on CA3 pyramidal cells of rat hippocampus. *J. Physiol. (Lond.)* 472:615–663.
- Jonas, P., and N. Spruston. 1994. Mechanisms shaping glutamate-mediated excitatory postsynaptic currents in the CNS. *Curr. Opin. Neurobiol.* 4:366–372.
- Katz, B., and S. Thesleff. 1957. A study of the desensitization produced by acetylcholine at the motor end-plate. *J. Physiol. (Lond.)* 138:63–80.
- Läuger, P. 1995. Conformational transitions of ionic channels. In *Single-Channel Recording*. B. Sakmann and E. Neher, editors. Plenum Press, New York. 651–662.
- Lomeli, H., J. Mosbacher, T. Melcher, T. Hoyer, J. R. Geiger, T. Kuner, H. Monyer, M. Higuchi, A. Bach, and P. H. Seeburg. 1994. Control of

- kinetic properties of AMPA receptor channels by nuclear RNA editing. *Science*. 266:1709–1713.
- Raman, I. M., and L. O. Trussel. 1995. The mechanism of α -amino-3-hydroxy-5-methyl-4-isoxazolepropionate receptor desensitization after removal of glutamate. *Biophys. J.* 68:137–146.
- Rosner, B. 1990. *Fundamentals of Biostatistics*. PWS-Kent Publishing Company, Boston.
- Schuster, C. M., A. Ultsch, P. Schloss, J. A. Cox, B. Schmitt, and H. Betz. 1991. Molecular cloning of an invertebrate glutamate receptor subunit expressed in *Drosophila* muscle. *Science*. 254:112–114.
- Seeburg, P. H. 1993. The TIPS/TINS lecture: the molecular biology of mammalian glutamate receptor channels. *Trends Pharmacol. Sci.* 14: 297–303.
- Standley, C., T. M. Norris, R. L. Ramsey, and P. N. Usherwood. 1993. Gating kinetics of the quisqualate-sensitive glutamate receptor of locust muscle studied using agonist concentration jumps and computer simulations. *Biophys. J.* 65:1379–1386.
- Sutcliffe, M. J., Z. G. Wo, and R. E. Oswald. 1996. Three-dimensional models of non-NMDA glutamate receptors. *Biophys. J.* 70:1575–1589.
- Trussell, L. O., S. Zhang, and I. M. Raman. 1993. Desensitization of AMPA receptors upon multiquantal neurotransmitter release. *Neuron*. 10:1185–1196.
- Usherwood, P. N. R. 1994. Insect glutamate receptors. *Adv. Insect. Physiol.* 24:309–341.
- von Beckerath, N., H. Adelsberger, F. Parzefall, C. Franke, and J. Dudel. 1995. GABAergic inhibition of crayfish deep extensor abdominal muscle exhibits a steep dose-response relationship and a high degree of cooperativity. *Pflügers Arch.* 429:781–788.
- Wisden, W., and P. H. Seeburg. 1993. Mammalian ionotropic glutamate receptors. *Curr. Opin. Neurobiol.* 3:291–298.