

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Biochimica et Biophysica Acta 1742 (2004) 21–36

<http://www.elsevier.com/locate/bba>

Review

The mammalian TRPC cation channels

Guillermo Vazquez, Barbara J. Wedel, Omar Aziz, Mohamed Trebak, James W. Putney Jr.*

The Calcium Regulation Section, National Institute of Environmental Health Sciences, National Institutes of Health,
Department of Health and Human Services, 111 TW Alexander Dr., Research Triangle Park, NC 27709, United States

Received 3 August 2004; received in revised form 27 August 2004; accepted 28 August 2004

Available online 11 September 2004

Abstract

Transient Receptor Potential-Canonical (TRPC) channels are mammalian homologs of Transient Receptor Potential (TRP), a Ca^{2+} -permeable channel involved in the phospholipase C-regulated photoreceptor activation mechanism in *Drosophila*. The seven mammalian TRPCs constitute a family of channels which have been proposed to function as store-operated as well as second messenger-operated channels in a variety of cell types. TRPC channels, together with other more distantly related channel families, make up the larger TRP channel superfamily. This review summarizes recent findings on the structure, regulation and function of the apparently ubiquitous TRPC cation channels.

Published by Elsevier B.V.

Keywords: Ion channel; Calcium channel; Non-selective cation channel; TRP channel; TRPC channel; Capacitative calcium entry; Store-operated channel; Second messenger-operated channel; Inositol trisphosphate receptor; Diacylglycerol; Phospholipase C

1. Origin, classification and nomenclature of TRPCs

The recognition that the protein product derived from the transient receptor potential (*trp*) gene from *Drosophila melanogaster* encoded a PLC-activated Ca^{2+} permeable channel eventually led to identification of seven TRP homologs in mammals [1,2]. Today, the TRP superfamily of channel forming proteins is comprised of these seven homologs of *Drosophila* TRP together with more distantly related channel genes identified through other searches for channels with specific functions [3–8]. TRP superfamily proteins fall into one of the three major families on a phylogenetic basis: the TRPC or canonical TRP family, with seven members (TRPC1 through TRPC7), which are the most closely related to the original *Drosophila* TRP channel; the TRPV family, with six members (TRPV1–6) named after the first group

member, the vanilloid receptor; and the TRPM family, with eight members (TRPM1–8), named after the original member, melastatin (Fig. 1). Other more distantly related members of the superfamily have been identified as genes whose dysfunction forms the bases for certain inherited diseases. The TRP superfamily and the corresponding family and subfamily designations follow a recent consensus nomenclature [9] that will be used throughout this review.

TRP superfamily members are cation channels involved in a continually growing number of cellular functions [6], and overall share 20–60% homology. Based on structural and functional similarities, the TRPC family can be further subdivided into four different subfamilies: TRPC1, TRPC2, TRPC3, 6 and 7 and TRPC4 and 5 (Fig. 1). TRPC2 is a pseudogene in humans, and in old world monkeys and apes [10], but TRPC2 apparently forms fully regulated channels in other mammalian species (see below). TRPC3, 6 and 7 form a closely related subfamily, sharing a high degree of amino acid identity (70–80%) and functional, regulatory and pharmacological similarities. A similarly close structural, and apparently func-

* Corresponding author. Tel.: +1 919 541 1420; fax: +1 919 541 7879.

E-mail address: putney@niehs.nih.gov (J.W. Putney).




Fig. 1. Structural relatedness of the three major TRP channel families. A multiple alignment of the ion channel domains of TRPC, TRPM and TRPV family members was carried out with VectorNTI software (AlignX) which uses the Clustal W algorithm. Kimura's correction was applied [140]. The phylogenetic tree was built using the Neighbor Joining method of Saitou and Nei [141]. All sequences with the exception of TRPC2 (mouse, pseudogene in human) are human. Calculated distance values are displayed in parentheses.

tional, relationship also exists between TRPC4 and TRPC5.

2. Structure–function relationships of TRPC channels

All members of the TRPC family are believed to share a common topology, based on the original analysis by Vannier et al. [11]. The cytoplasmic N- and C-termini are separated by six predicted transmembrane domains (TM1–TM6), including a putative pore region between TM5 and TM6. However, there is uncertainty as to the exact assignment of the first two transmembrane segments [11,12]. The structure shown in Fig. 2 is based on the assignments given in Ref. [11]; alternative assignments are shown in the bottom half of Fig. 2, and the reader is referred to Ref. [12] for a discussion of this issue. The N-terminus is composed of three to four ankyrin repeats, a predicted coiled coil region and a putative caveolin binding region. The cytoplasmic C-terminus includes the TRP signature motif (EWKFAR), a highly conserved proline rich motif, the CIRB (*calmodulin/IP₃ receptor binding*) region and a predicted coiled-coil region. An extended C-terminus containing a PDZ binding motif is unique to TRPC4 and TRPC5. The following section summarizes insights gained from structure–function studies of different TRPC isoforms.

Ankyrin repeats fold into a helix-loop-helix-beta-hairpin/loop structure with several repeats stacked against one another to form a single domain with a protein-binding interface (recently reviewed in Ref. [13]). Unlike other

protein–protein interaction domains, ankyrin repeats do not recognize a conserved sequence or structure, but in contrast can accommodate a variety of binding partners mainly by variation of surface exposed residues. It is therefore hard to predict possible binding partners for the TRPC ankyrin repeats. Based on truncation studies, the ankyrin repeats appear to be required for correct targeting of TRPC3 to the plasma membrane [14]. While N-terminal deletion of TRPC3 up to the ankyrin repeats yields functional and plasma membrane targeted channels, further deletion of the ankyrin repeats results in accumulation of the protein in intracellular compartments. A similar trafficking defect has been described for a TRPC6 mutant lacking 131 N-terminal amino acid residues that include the first ankyrin repeat [14]. Not all ankyrin repeats seem to be crucial for functional expression of channels, as is the case for TRPC1, since a splice variant missing the third out of four ankyrin repeats forms channels that are activated in response to store depletion [15].

The coiled-coil (CC) is a ubiquitous protein motif that is commonly used to control oligomerisation. CC sequences consist of several heptad repeats—seven residue patterns—folding into an alpha-helix. Two (dimer), three (trimer) or more of these alpha-helices are wound around each other to form a supercoil. As mentioned above, there is one CC in each of the cytosolic N- and C-termini of TRPCs (CC-N and CC-C). It is therefore conceivable that CC domains contribute to homo- and heteromerization of TRPCs to form ion channel tetramers with distinct characteristics, or to link TRPC to other coiled domain containing proteins. A




Fig. 2. Structure of TRPC channels. Top: Structural features of TRPC channels. See text for description of structural features. ANK, ankyrin-like repeats; CC-N and CC-C are N-terminal and C-terminal coiled-coil domains, respectively. The region shaded in gray at the extreme C-terminus is specific for TRPC 4 and 5. Bottom: The arrangement of the first three hydrophobic segments of TRPCs is uncertain; three alternatives are shown (A, B and C). See Refs. [11,12]. The acronyms are: PDZ-B, PDZ binding domain; CC-N, N-terminal coiled coil region; CC-C, C-terminal coiled coil region; ANK1–4, ankyrin repeats 1–4; CIRB, calmodulin/IP₃ receptor binding region; PLC, phospholipase C; IP₃R, IP₃ receptor; CaM, calmodulin; LFW, amino acid motif conserved in the putative pore region of all TRPCs.

recent study showed that CC-N of TRPC1 was able to homodimerize based on a yeast two-hybrid screen while the C-terminus including CC-C was unable to dimerize [16]. The microtubule destabilizing phosphoprotein stathmin has recently been pulled out of a yeast two-hybrid screen using the N-terminus of TRPC5 as a bait, and CC-N was found to be responsible for the interaction, presumably via the predicted CC region of stathmin [17]. This interaction seemed to be specific for TRPC4 and TRPC5 since the N-termini of TRPC4 and 5 but not TRPC1 or TRPC7 were able to interact with the stathmin helical domain. TRPC5 appears to regulate axon length and growth cone morphology in hippocampal neurons where its activation and associated Ca²⁺ influx may prevent random axonal outgrowth. The C-terminal CC region can be deleted in TRPC3 without loss of plasma membrane targeting and channel activation when compared with the wild type [14]. However, in TRPC1, calmodulin (CaM) binding to the predicted CC-C domain has been reported and its deletion resulted in

diminished Ca²⁺-dependent inactivation of store-operated Ca²⁺ entry [18].

Caveolins are enriched in plasma membrane microdomains called caveolae (for a recent review see Ref. [19]). Co-immunoprecipitation or co-localization with caveolins has been shown for TRPC1 and TRPC3 [20–22]. Caveolin-1 binding usually occurs at aromatic-rich (θ) regions that contain a specific spacing following the consensus sequences θXθX₄θ, θX₄θX₂θ, or θXθX₄θX₂θ [23]. A similar motif is conserved in all members of the TRPC family located in the cytosolic N-terminus adjacent to the first transmembrane domain [20]. Deletion of this region prevents TRPC1 from targeting to the plasma membrane and the respective mutant dominant-negatively affects store-operated Ca²⁺ entry [20].

A systematic analysis of the predicted pore region of TRPC family members has not yet been carried out. However, several studies have examined the effect of point mutations within the predicted pore region and extrac-

ellular loops bordering the pore. Change of a conserved LFW motif in the predicted pore helix of TRPC5 and TRPC6 to AAA resulted in dominant negative mutants [24,25]. A means to discriminate endogenous receptor operated/store operated channels from overexpressed TRPCs has been the fact that the latter are insensitive to inhibition by lower micromolar concentrations of lanthanides. In the case of TRPC4 and TRPC5, higher micromolar concentrations of lanthanides potentiate currents, while they inhibit overexpressed TRPC6 [26]. Jung et al. [26] identified charged residues in the pore loops of TRPC5 that when mutated from E to Q (E543Q and E595Q/E598Q) caused a loss of lanthanide potentiation. Likewise, E576K or D581K mutants of TRPC1 exhibited decreased store-mediated Ca^{2+} but not Na^{+} current suggesting that these residues play an important role in calcium selectivity [27]. This is an interesting finding since conventional thought of the function of a nonselective cation channel would not predict that mutations would affect Ca^{2+} permeability selectively; in other words, the mutation was in reality a gain of function mutation converting a nonselective channel to one selective for monovalent cations. Interestingly, E576 is conserved between TRPC1, 4 and 5 and part of a CX_4CE motif unique to these isoforms. Since these cysteines likely face the extracellular space, they may form disulfide bridges thus potentially explaining preferred heteromer formation between TRPC1, 4 and 5 on one hand and the ‘cysteine-less’ TRPC3, 6 and 7 on the other [25].

A proline-rich motif (LPXPFXXXPSPK) downstream of the EWKFAR motif is conserved in all members of the TRPC family and is thought to be responsible for interaction with Homer (for TRPC1) and/or immunophilins [28,29]. Homer proteins dimerize via a CC motif while the EVH domain of each monomer is able to bind to a consensus PPXXF motif in their target molecules thus linking, for example, cell surface receptors and intracellular Ca^{2+} release channels. In analogy to this scheme Homer binding to TRPC1 and the IP_3 receptor would make it a linker involved in the proposed conformational coupling model of TRP channel activation (discussed in a subsequent section). In support of this hypothesis, mutations within the proline-rich motif disrupted Homer binding and resulted in channels with high constitutive activity and greatly reduced agonist regulation [28]. A direct interaction of TRPC channels with IP_3 receptors will be discussed separately (Section 4.2). It is, however, important to note that a TRPC1 mutant lacking the entire cytoplasmic C-terminus was apparently able to function as a store-operated channel [30].

Recently an interaction of TRPC with the immunophilins FKBP12 and FKBP52 has been demonstrated [29]. Immunophilins are peptidyl-prolyl *cis-trans* isomerases binding to a XP motif in their binding partners. Pull-down experiments showed that FKBP12 preferentially interacted with TRPC3, 6 and 7, while FKBP52 preferred TRPC1, 4 and 5. Mutation of the first proline in the proline rich motif of

TRPCs disrupted that interaction. Furthermore, FK506, which displaces immunophilins from their targets, was able to inhibit agonist-induced currents through TRPC6 channels [29].

Evidence for functional roles of the CIRB (CaM/ IP_3 receptor binding) region is discussed in Section 4.2.3. As mentioned above, TRPC4 and 5 share an extended C-terminus terminating in a PDZ binding motif. A part of this region is missing in a TRPC4 β splice variant, and this variant has been proposed to have autoinhibitory function [31]. This region has also been shown to interact with the C-terminus of IP_3 receptors and CaM [32,33]. Little is known about the function of this region in TRPC5 and the degree of homology between the C-termini of TRPC4 and TRPC5 is very low. The PDZ binding motif TRL of TRPC4 and TRPC5 seems to be responsible for the interaction with the adaptor protein, NHERF, possibly linking the channels to PLC β and the cytoskeleton [34]. Deletion of the PDZ binding motif of TRPC4 not only reduced the surface expression of TRPC4 but also changed it from a general membrane distribution to a predominant expression in cell outgrowths [35].

Recently, different patterns of N-glycosylation have been summoned to explain different levels of constitutive activity within members of the TRPC3, 6 and 7 family. TRPC3, which is supposedly monoglycosylated in the first extracellular loop, shows high constitutive activity, while TRPC6, dually glycosylated in the first and second extracellular loop is tightly regulated. By converting TRPC3 into the TRPC6-like dually glycosylated version and likewise converting TRPC6 into the monoglycosylated TRPC3-like version, Dietrich et al. [36] were able to alter constitutive activity as hypothesized. It is worth noting, however, that TRPC7 exhibits considerable basal activity despite the fact that it has a predicted N-glycosylation site in the first and second extracellular loop like TRPC6 [37].

The major structural features of TRPC channel subunits are summarized in Fig. 2.

3. Pharmacology and electrophysiological properties

3.1. TRPC1

TRPC1 was the first member of the TRPC family to be cloned [1,2], and the first shown by electrophysiological measurements to form a calcium-permeable cation channel [15]. In this initial study, expression of TRPC1 in Chinese hamster ovary cells (CHO) resulted in a linear nonselective cation current, activated by intracellular infusion (via the patch pipet) of either IP_3 or thapsigargin to deplete intracellular calcium stores. This current was shown to be sensitive to Gd^{3+} (20 μM). Noise analysis provided evidence that the current was passed through single channels with a conductance of 16 pS. The current–voltage relationship of these channels was shown to be predom-

inantly inward, with little outward current [38,39]. In this respect, the TRPC1 current was described as I_{crac} -like [40] (the properties of I_{crac} are discussed in a subsequent section dealing with regulation of TRPC channels, and the behavior of store-operated channels), although unlike I_{crac} , the current was not Ca^{2+} -selective.

3.2. TRPC2

There has been little direct work to date regarding the electrophysiological characteristics of TRPC2. Work on mice vomeronasal neural dendrites has shown that knocking out the TRPC2 gene resulted in the near loss of a diacylglycerol (DAG)-activated linear current, which was not activated by store depletion [41], suggesting that TRPC2 may be a component of a DAG-operated channel *in situ*. However, using fluorescence rather than electrophysiological methodology, Vannier et al. [42] reported that expressed TRPC2 appeared to behave as a store-operated channel.

3.3. TRPC3, 6 and 7

TRPC3, 6 and 7 appear to form nonselective cation channels that show both inward and outward rectification, at negative and positive voltages, respectively. The current–voltage relationships for these TRPCs demonstrate a far greater outward current (at positive voltages) than inward current (at negative voltages) with a reversal potential at around 0 mV.

Zhu et al. [43] first described TRPC3 as an agonist-activated channel when expressed in HEK293 cells and demonstrated that it was relatively insensitive to low concentrations of Gd^{3+} , but was blocked by concentrations in the 100 μM range. Hofmann et al. [44] later demonstrated that TRPC3 could also be activated directly by DAG analogues.

TRPC3 channels have also been shown to have two potential conductance states when expressed in HEK293 cells, one of 66pS and one of 17pS [45]. In this study it was shown that these channels were activated by calcium store depletion in intact cells, consistent with an earlier report by Preuß et al. [46], or by addition of recombinant IP_3 receptors in excised patches. In a subsequent study, however, activation of TRPC3 was shown to be dependent solely upon DAG with no requirement for either IP_3 or the IP_3 receptor [47].

Since their discovery, these channels have been shown to be sensitive to a variety of agents with relative sensitivities varying dramatically depending upon expression system. In HEK293 cells, TRPC3 was inhibited by high concentrations of: SKF96365, verapamil, La^{3+} , Gd^{3+} and Ni^{2+} [43] and was partially inhibited by moderate (30 μM) concentrations of 2-aminoethoxydiphenyl borane (2APB) [48]; in DT40 cells by low concentrations of Gd^{3+} [48]; and in CHO cells they are reported to be sensitive to low concentrations of La^{3+} , Gd^{3+} and

SKF96365 [49]. However, in the latter case, it appeared that the lanthanides acted by first entering the cell and blocking from the inside. This requirement for penetration, which does not normally occur with lanthanides, may explain the relative insensitivity of TRPC3 to lanthanides in other cellular systems.

TRPC6 has also been shown to be a DAG-activated channel [44] and is very similar to TRPC3 and 7 with regard to its current–voltage relationship. The single channel conductance was 35pS, with no second conductance state observed [44]. TRPC6 does, however, possess a unique characteristic to this TRPC subfamily. When expressed in HEK293 cells, TRPC6 currents are enhanced by the cation channel blocker flufenamate while TRPC3 and 7 currents are not [50].

TRPC7, the final member of this subfamily, demonstrates properties very similar to TRPC3 and 6 with regard to its voltage–current relationship, and activation by DAG [37]. TRPC7 has demonstrable sensitivity to SKF96365 and, in a similar fashion to TRPC3 and 6, is relatively insensitive to lanthanides.

The differences between the three channel types may lie in their ion selectivity, in that TRPC6 is reported to be somewhat Ca^{2+} -selective, while TRPC3 and TRPC7 do not appear to be. Reported values for ($P_{\text{Na}}/P_{\text{Ca}}$) are 1\1.5, 1\5 and 1\2 for TRPC3, 6 and 7, respectively (cf. [51]).

3.4. TRPC4 and 5

TRPC4 was originally cloned by Philipp et al. [52]. These investigators reported that, when expressed in mammalian cells, TRPC4 resulted in a novel, inwardly-rectifying Ca^{2+} -selective current that was activated by either IP_3 or thapsigargin-induced store depletion. This role of TRPC4 in store-operated entry has since been questioned with reports of TRPC4 being activated by G-protein coupled receptors [53]. Interestingly, TRPC4 currents have been shown to be potentiated by lanthanides [53], in contrast to their inhibitory effects on endogenous store-operated channels or other TRPCs. TRPC4 has a reported single channel conductance of 41 pS [53], with a current–voltage relationship similar to that of TRPC3, 6 and 7, having a greater current at positive voltages than at negative voltages.

TRPC5 was originally cloned by the same group who first cloned TRPC4 [54], and was shown to have properties expected of a store-operated channel. However, subsequent studies from other laboratories demonstrated that TRPC5 behaved as a receptor-operated cation channel [53,55]. The current–voltage relationship of this channel indicates that far more current is passed at negative voltages (inward current) than at positive. In a similar fashion to TRPC4, TRPC5 currents are also potentiated by lanthanides [53]. The single channel conductance of TRPC5 has been estimated at around 63 pS [53]. The reported ion selectivities of TRPC4 and 5 vary (see Ref. [6]).

4. Signaling mechanisms and regulation of TRPC channels

4.1. Store depletion

In most types of cells, membrane receptors coupled to phosphoinositide-specific phospholipase C (PLC) promote inositol 1,4,5-trisphosphate (IP_3)-mediated release of Ca^{2+} from endoplasmic reticulum (ER) and subsequent Ca^{2+} entry across the plasma membrane which occurs, predominantly, through a process known as capacitative calcium entry (CCE) or store-operated calcium entry [56]. Under physiological conditions, CCE is triggered when IP_3 discharges Ca^{2+} from ER stores; it is the subsequent reduction in ER Ca^{2+} content that signals to the plasma membrane for activation of store-operated channels (SOCs). When measured by electrophysiological techniques, in hematopoietic cells and perhaps some other cell types, CCE corresponds to a highly Ca^{2+} -selective current termed the calcium-release activated calcium current, or I_{crac} [40]. In other cell types, most notably smooth muscle, the currents underlying CCE appear to be less Ca^{2+} -selective [57,58], indicating that no one gene product is likely to be responsible for store-operated currents in all cell types. Neither the precise activation mechanism nor the molecular identity of the SOCs is known with certainty [59,60]. The fact that activation of *Drosophila* TRP channel was found to require activation of the PLC cascade (see Ref. [5] and references therein) originally suggested that the mammalian counterparts might be molecular candidates for SOCs. Consequently, a number of studies on the molecular nature of SOCs have focused primarily on mammalian TRP channels, particularly those from the TRPC family. As a result, most of the members of the TRPC family, either endogenous or ectopically expressed in cell lines, have been implicated in store-operated cation entry.

There is a considerable body of evidence supporting the idea that TRPC1 constitutes, or is a part of, SOCs (reviewed in Ref. [61]). Ectopic expression of TRPC1 in many cell lines has resulted in either nonselective [15] or Ca^{2+} -selective [62] cation channels activated by either agonist stimulation or pharmacological depletion of stores with thapsigargin. The most compelling evidence for TRPC1 as a subunit of SOCs is derived from studies using TRPC1 antisense constructs. This strategy proved to be efficient in reducing SOC entry in human submandibular gland cells [62], HEK293 cells [63], *Xenopus* oocytes [64], human lung epithelial A249 cells [65] and pulmonary artery smooth muscle cells [66]. Additionally, genetic disruption of the TRPC1 gene significantly reduced I_{crac} in chicken DT40 B lymphocytes [67]. Other studies provided evidence for TRPC1 as part of endogenous SOCs by using antibodies directed against extracellular loops of TRPC1. Extracellular application of these antibodies significantly reduced SOC entry in vascular smooth muscle cells [68] and human platelets [69].

TRPC2, which in humans appears to have degenerated to a pseudogene [2], has been shown to be functional in mouse, rat and other mammalian species. Ectopic expression of two splice variants of mouse TRPC2 (TRPC2a and TRPC2b) in COS-M6 cells increased the magnitude of SOC entry induced by muscarinic receptor stimulation [42], although the constitutive activity of the overexpressed proteins was not assayed. Importantly, an antibody targeting an extracellular site of mouse TRPC2b, near the pore region, blocked thapsigargin-evoked Ca^{2+} entry and the acrosome reaction in sperm [70], suggesting that TRPC2 might be a subunit of endogenous SOCs in sperm. Two additional splice variants of mouse TRPC2 (TRPC2 α and TRPC2 β) cloned by a different lab [71] were devoid of any measurable function upon heterologous expression, in principle due to a plasma membrane trafficking defect. Expression of a TRPC2 antisense construct in CHO cells, which mainly express TRPC1 and TRPC2, significantly reduced SOC entry induced by thapsigargin [72].

TRPC3 is the most extensively characterized of the TRPC family members, although its role in CCE has been controversial. While earlier observations suggested a role for TRPC3 in CCE when transiently expressed in HEK293 cells [46,73], subsequent studies showed that this likely reflected constitutive channel activity, rather than a regulated mode of Ca^{2+} entry [43]. TRPC3 constitutive activity has frequently caused misinterpretation of studies aimed to address the role of store depletion in channel regulation (see Ref. [48]). This is due to an apparent increase in thapsigargin-induced Ca^{2+} entry, which results from diminished ER Ca^{2+} buffering capacity, and the concomitant enhancement of constitutive channel activity. A considerable body of literature now supports the idea that TRPC3 commonly behaves as a receptor-activated channel that cannot be activated by store depletion (discussed in Refs. [44,74–77]). An interesting exception occurs with ectopic expression of TRPC3 in the avian B lymphocyte DT40 cell line. In these cells, TRPC3 is clearly activated by store depletion, but only when expressed at very low levels [48,78,79], as conditions that provide higher levels of channel protein expression give rise to the receptor-regulated mode of TRPC3 activation [79,80], indicating that the coupling mode of TRPC3 might be related to its expression level. Channel expression level was also shown to have a critical impact on the coupling mechanism for TRPV6 activation [81]. A thorough discussion of the potential relationship between channel expression level and channel regulation has recently been published [82].

Evidence for store-operated regulation of other members of the TRPC3, 6 and 7 subfamily is also controversial. Transient ectopic expression of mouse TRPC7 in HEK293 cells resulted in a constitutively active cation channel, whose activity was further stimulated by activation of the PLC-linked P2Y purinergic receptor, but not by store depletion alone [37]. In contrast, Riccio et al. [83] reported

that stable expression of human TRPC7 in HEK293 cells results in a cation channel activated by store depletion. The authors suggested that the different coupling modes of activation for mouse TRPC7 versus the human ortholog could be due to a difference in the amino acid at position 111 (proline in their human TRPC7 clone vs. leucine in mouse TRPC7). However, in the study by Riccio et al., TRPC7 constitutive activity was not assessed. Thus, based on prior experiences with other TRPCs, it must be said that for now there is no convincing evidence that TRPC7 behaves differently from its close relatives, TRPC3 and TRPC6.

Although conflicting results exist regarding the role of PLC-derived products, and more recently, phosphorylation (see below) in regulation of TRPC6, the literature is relatively consistent with regard to the role of store depletion on channel activity. Both mouse and human TRPC6, when ectopically expressed in cell lines, appear to behave as nonselective cation channels whose activation is independent of store depletion but linked to PLC-derived products [44,50,84] (for a review, see Ref. [74]).

TRPC4 and TRPC5 are structurally similar, and thus it is not surprising that, in a given study in a given laboratory, they appear to be regulated similarly. Ectopic expression of bovine TRPC4 in both HEK293 and CHO cells increased SOC entry in response to either IP₃ or thapsigargin [52,85]. In addition, bovine TRPC4 markedly augmented I_{crac} currents when expressed in RBL cells [85]. Rat TRPC4 increased thapsigargin-induced inward calcium-dependent chloride currents, an indirect reflection of SOC entry, when expressed in *Xenopus* oocytes [64,86]. A TRPC4 antisense construct markedly reduced an I_{crac} -like current in SBAC cells, an adrenal cortical cell line, suggesting TRPC4 is part of the endogenous CRAC channels in these cells [87]. Similarly, vascular endothelial cells derived from TRPC4 knock-out (TRPC4^{-/-}) mice showed impaired store operated calcium entry [88]. In a recent study of the properties of lung endothelial cells derived from the same TRPC4^{-/-} mice, Tiruppathi et al. [89] expanded the observations of Freichel et al. [88], and found that absence of TRPC4 was correlated with a loss of endothelial cell responses to thrombin, suggesting a key involvement of TRPC4 in microvascular permeability. More recently, TRPC4 antisense oligonucleotides were shown to partially inhibit store-operated calcium entry in mouse mesangial cells, suggesting that TRPC4 might also form part of endogenous SOCs in that cell type [90].

In contrast with these findings, mouse TRPC4 expressed in HEK293 cells, adrenal chromaffin cells and PC12 cells was found to form agonist-PLC regulated cation channels. The channels could not be activated by IP₃, DAG or store depletion [53,91]. In another study, human TRPC4 was not regulated by either store depletion or PLC activation, and showed only constitutively active when expressed in HEK293 cells [77]. Stable expression of a TRPC4 antisense construct in the same cell line did not alter store-operated

entry [92]. Fewer reports have dealt with regulation of TRPC5 compared to TRPC4. Almost simultaneously, two different labs reported the cloning of mouse TRPC5, its transient expression in HEK293 cells rendering a receptor-activated channel, independent of store depletion for one group [55] and store-operated for the other [54]. Subsequently, in one instance in which one laboratory compared both TRPC4 and TRPC5 in the same study, the same mode of regulation was observed—PLC-dependent activation not related to store depletion [53]. These authors expressed mouse TRPC5 in CHO and HEK293 cells, and found that the channel was activated by either G-protein-coupled receptor or receptor tyrosine kinase stimulation but not by IP₃, DAG or store depletion. Upon ectopic expression in cell lines, TRPC5 showed significant constitutive activity, similar to that seen for other TRPC channels. Unlike TRPC4, expression of mouse TRPC5 in *Xenopus* oocytes significantly increased muscarinic receptor-induced calcium-dependent chloride currents, but not those stimulated by thapsigargin, indicating, again, that TRPC5 was receptor-activated rather than store-operated [93].

4.2. Interaction between IP₃ receptors and TRPC channels

The inositol 1,4,5 trisphosphate receptor (IP₃R) is a calcium release channel found predominantly in the membrane of the endoplasmic reticulum (ER). IP₃R releases calcium from the ER in response to the second messenger IP₃ generated by the enzymatic activity of phospholipase C (PLC). One of the hypotheses for the mechanism of activation of capacitative calcium entry was originally proposed by Irvine [94] and termed “conformational coupling”. According to this model, further developed by Berridge [95], SOCs in the plasma membrane interact with IP₃ receptors in the endoplasmic reticulum resulting in physical coupling of Ca²⁺ release and Ca²⁺ entry. With the discovery of TRPC channels and their subsequent proposal as candidates for capacitative calcium entry channels, investigators assessed the possibilities of a direct interaction between TRPC channels and the IP₃ receptor (for review, see Ref. [96]). Although its physiological significance is still a matter of great debate, physical interactions between the IP₃R and virtually every member of the TRPC family have been reported. Tang et al. [97] described CaM and IP₃R binding sites in all TRPC proteins and suggested a reciprocal regulation of TRPC channels by IP₃R and CaM. The affinity of IP₃R peptide in displacing CaM was found to be lower in TRPC1,2,4,5 than in TRPC3,6,7 [97]. Specific findings relating to interactions between IP₃ receptors and the TRPC sub-families are discussed below.

4.2.1. TRPC1

Rosado and Sage [98] showed that in human platelets TRPC1 and type II IP₃R co-immunoprecipitate when internal calcium stores are depleted by thapsigargin and

that xestospongins, an IP₃R antagonist, blocks CCE. In a subsequent study, the same authors showed that xestospongins also blocks the thapsigargin-induced association of TRPC1 with type II IP₃R suggesting that TRPC1 is a component of CCE channels, and that the mechanism of activation involves interaction with IP₃ receptors [99]. Rosado and Sage showed that the association with TRPC1 and IP₃R is reversible; the TRPC1–IP₃R complex dissociates upon refilling of the stores. They treated platelets with jasplakinolide, a drug that increases peripheral actin accumulation, and found that this blocked CCE and prevented the association of TRPC1 and IP₃R (see also ref. [100]). In a subsequent study, again in platelets, Rosado et al. [69] showed that store depletion with an agonist (i.e., thrombin), as seen with thapsigargin, caused reversible coupling of the type II IP₃R with the Ca²⁺ entry channel, TRPC1. A more recent study from the same laboratory [101] reported a direct interaction between type II IP₃ receptor and TRPC1 within 1 s upon stimulation of human platelets with thrombin, consistent with a gating mechanism of the TRPC1 channels by IP₃R. Mehta et al. [102] reported a Rho-induced association of IP₃R with TRPC1 that was dependent on actin filament polymerization, and upon store depletion either with agonist or thapsigargin. These authors subsequently concluded that Rho activation signals interaction of IP₃R with TRPC1 at the plasma membrane of endothelial cells, and triggers Ca²⁺ entry following store depletion. Similarly, Yuan et al. [28] proposed that the adaptor protein Homer facilitates a physical association between TRPC1 and the IP₃R that is required for TRPC1 activation. Lockwich et al. [21] showed that endogenous TRPC1 in HSG cells is Triton X-100-insoluble, suggesting association with the cytoskeleton and lipid rafts; TRPC1 was found associated with caveolin, type III IP₃R and Gαq, suggesting the existence of a signalplex comprised of TRPC1 and several players of the PLC pathway.

4.2.2. TRPC2

Tang et al. [97] showed that TRPC2, like all members of the TRPC family, has the potential to interact with the IP₃R. The IP₃R-binding domain was also found to interact with CaM in a Ca²⁺-dependent manner with an affinity of 10 nM and CaM inhibits the TRPC–IP₃R interaction. The authors then concluded that both CaM and IP₃R play reciprocal roles in controlling the activation of all TRPC channels. Brann et al. [103] showed an overlapping of immunoreactivity for the type III IP₃R with that of TRPC2 in the epithelium of the vomeronasal organ of the rat. This observation was further confirmed by co-immunoprecipitation of type III IP₃R and TRPC2 from vomeronasal organ lysates.

4.2.3. TRPC3, 6 and 7

Although evidence for interaction with IP₃R has been reported for all members of this subfamily [97], most of the evidence is derived from studies with TRPC3 channels. The

interaction between TRPC3 and IP₃R was reviewed earlier by Trebak et al. [74]. Presuming that TRPC proteins might function as store-operated channels, Kiselyov et al. [45] used a HEK293 cell line stably expressing TRPC3 (T3–9, from Zhu et al. [43]) to demonstrate “conformational coupling” of TRPC3 and the IP₃R. As pointed out by Trebak et al. [74], a large body of evidence, including data from Zhu et al. [43] on T3–9 cells, showed that TRPC3 overexpressed in HEK293 behaves as a non-store-operated, receptor-activated channel. It is therefore unclear why, in the electrophysiological studies by Kiselyov et al. [45], TRPC3 exhibited characteristics of store-operated channels. In the studies of Kiselyov et al. [45], T3–9 cells stimulated by carbachol or thapsigargin showed increased single channel activity that was absent in wild-type HEK293 cells. Channel activity declined after patch excision, but was reactivated by IP₃. After extensive washing of the patches, IP₃ could no longer activate TRPC3 channels. In this situation, channels could be activated by a combination of IP₃ and IP₃R. The ability of IP₃ to activate TRPC3 channels in excised patches was blocked by the IP₃R antagonists, heparin and xestospongins C. The authors concluded that TRPC3 channels are store-operated and that the mechanism of activation involves conformational coupling to the IP₃R in its IP₃ bound state [45]. However, in a series of studies Trebak et al. [47,48] evaluated the involvement of TRPC3 in store-operated Ca²⁺ entry and the role of IP₃ in TRPC3 activation. When expressed in HEK293 cells, TRPC3 was shown to form channels that are receptor-operated but not activated by store depletion [48] in agreement with Zhu et al. [43]. In a subsequent study Trebak et al. [47] demonstrated that DAG is sufficient to activate TRPC3 and that neither IP₃ nor G proteins are required for TRPC3 activation in HEK293 cells, including the T3–9 cell line used by Kiselyov et al. [45]. Furthermore, the IP₃R antagonist, heparin, was unable to block TRPC3 activation in response to agonist. In an earlier study, Inoue et al. [50] showed that TRPC6 could not be activated by IP₃ and that DAG was sufficient for TRPC6 activation. A report by Albert and Large [104] describing a native TRPC6-like calcium channel in rabbit portal vein myocytes showed that those TRPC6-like channels could not be activated by IP₃; however, they report that IP₃ appears to synergize with DAG to produce greater TRPC6-like channel activation.

Kiselyov et al. [105] further identified a TRPC3 binding domain in the C-terminus of the IP₃R. Transiently expressed TRPC3 increased the basal rate of Ba²⁺ entry in HEK293 cells, and this was further increased by co-expression of the IP₃R binding domain or an IP₃R construct lacking the transmembrane domain, but not by the full-length IP₃R protein. This indicated that forms of IP₃R that are uncoupled from intracellular stores activate TRPC3, while the full-length receptor may require store depletion or some additional signal.

Boulay et al. [106] also investigated the interaction between human type 3 IP₃R and human TRPC3. IP₃R

proteins could be recovered in TRPC3 and TRPC6 immunoprecipitates. The authors identified the minimal mutual binding domains in TRPC3 and IP₃R: two sequences in IP₃R that interact with the C-terminus of TRPC3, designated F2q and F2g, corresponding to amino acids 669–698 and 751–821, respectively. F2q is modestly conserved among human IP₃Rs, while F2g is well conserved. For TRPC3, a small fragment designated C8, corresponding to amino acids 777–797 in human TRPC3, interacted with the N-terminus of IP₃R. This sequence is well conserved among TRPC3, TRPC6 and TRPC7. Transient expression of the C7 fragment (54 amino acids containing the C8 peptide) significantly inhibited entry due to either agonist or thapsigargin in wild-type HEK293 cells. Transient expression of an IP₃R fragment, which contains F2q, inhibited both agonist and thapsigargin-induced entry. These experiments are suggestive of an involvement of TRPC3 and IP₃R in native store-operated Ca²⁺ entry in wild-type HEK293.

Zhang et al. [107] identified a sequence in the C-terminus of TRPC3 corresponding to amino acids 761–795 (termed CIRB domain for CaM/IP₃ receptor binding domain, discussed above) that binds CaM, and overlaps with the C8 region identified as the IP₃R binding region by Boulay et al. [106]. CaM was found to compete with the IP₃R peptide, F2q, for binding to CIRB and activation of TRPC3 channels was also observed with a CaM binding peptide, and with calmidazolium, a CaM antagonist [107].

A pharmacological study by Ma et al. [76] investigated the involvement of IP₃R in store-operated Ca²⁺ entry and TRPC3-mediated cation entry by examining the actions of 2APB, a purported membrane permeant IP₃R antagonist [108]. These authors observed ATP-stimulated entry of Sr²⁺ in a HEK293 cell line stably transfected with TRPC3, but not in control cells; 2APB blocked this entry. Furthermore, 2APB blocked capacitative Ca²⁺ entry in response to agonist or thapsigargin in control HEK293 cells. Ma et al. [76] suggested that the action of 2APB is consistent with the “conformational coupling” hypothesis, assuming that 2APB is acting by blocking IP₃ receptors. However, it is now clear that 2APB blocks the endogenous store-operated Ca²⁺ pathway by acting directly on SOCs [109–115]. It is currently unclear whether the effect of 2APB on TRPC3 channels is also due to a direct action on the channels, unrelated to IP₃R. A study by Li et al. [116] focused on the regulation of endogenous TRPC3 channels. They found that the neurotrophin BDNF activates a Ca²⁺-dependent nonselective cation current in pontine neurons, reminiscent of that activated by TRPC3. The BDNF-induced current could be blocked by the PLC antagonist U73122 as well as by IP₃R antagonists, xestospongin C and heparin. While thapsigargin did not activate this current, including IP₃ in the patch pipette did. The data from Li et al. [116] are consistent with an endogenous TRPC3 that is activated by a PLC- and IP₃-

dependent coupling mechanism, not related to store depletion.

In summary, the exact role of IP₃R in TRPC3 or SOC activation is still unclear. In a DT40 cell line lacking all forms of IP₃R, TRPC3 is activated by receptor agonists to the same extent as in wild-type cells [80] and this cell line also exhibits normal Ca²⁺ entry following passive depletion of the stores by thapsigargin [117,118]. Furthermore, the physiological significance of the biochemical interactions is uncertain. The effects of the TRPC3 and IP₃R peptides are small [106]. As shown by Trebak et al. [47], TRPC3 channels expressed in HEK293 cells are activated independently of IP₃ and G proteins; in addition, heparin could not block TRPC3 activation via G protein-coupled agonists. Nonetheless, Tang et al. [97] showed that all TRPC family members (1 through 7) associate with IP₃R (see also Ref. [32]). Therefore, interaction with the IP₃ receptor might play a role in assembling signaling complexes that involve phospholipase C, underlying IP₃ receptors, and TRPC channels.

4.2.4. TRPC4 and 5

The activation mechanism of the two members of this subfamily was reviewed recently by Plant and Schaefer [119]. Using a yeast two-hybrid assay and glutathione-S-transferase pulldown experiments, Mery et al. [32] reported that the C-terminus of an alternatively spliced variant of TRPC4, the alpha isoform, but not the beta, associates in vitro with the C-terminal domain of IP₃R receptors, types 1, 2 and 3. Subsequently, Tang et al. [97] confirmed these findings and discovered an additional CaM/IP₃R binding region common to all members of the TRPC family, as discussed above. In the presence of Ca²⁺, the TRPC4–IP₃R interaction was inhibited by CaM. A synthetic peptide representing a TRPC-binding domain of IP₃R inhibited the binding of CaM to TRPC3, 6 and 7 more effectively than that to TRPC1, TRPC2 and TRPC4 and 5. In inside-out membrane patches, TRPC4 was activated strongly by calmidazolium, an antagonist of CaM. Kanki et al. [93] showed that mouse TRPC5 expressed in *Xenopus* oocytes was activated by agonist but not by depletion of stores with thapsigargin. The agonist activation of TRPC5 was inhibited by the PLC inhibitor, U73122 and by the IP₃R antagonist xestospongin C. The authors also showed that the IP₃R agonist, adenophostin A, activated TRPC5 in a dose-dependent manner. However, the effect of injection of IP₃ itself on TRPC5 activity was apparently not examined. TRPC5 activation by adenophostin A was blocked by a pre-injection of either xestospongin C or a peptide mimicking the IP₃ binding domain of *Xenopus* IP₃R [93]. These authors concluded that the IP₃R is essential for TRPC5 activation. This conclusion conflicts with the report that TRPC5 transfected into the DT40 cell line lacking all isoforms of the IP₃R showed normal TRPC5 activation [120].

4.3. Role of membrane lipids in TRPC activation

When expressed in different cell lines, only members of the TRPC3, 6 and 7 subfamily have been consistently shown to be activated by membrane permeant DAG analogs such as 1-oleoyl-2-acetyl-*sn*-glycerol (OAG). These channels can also be activated by DAG lipase and DAG kinase inhibitors, but not by monoacylglycerols. This provides a possible mechanism of activation of these channels by PLC-coupled receptors, independently of IP₃R or store depletion.

Hofmann et al. [44] reported that exogenously applied DAG analogues can activate TRPC6 and TRPC3. These authors demonstrated that both TRPC6 and TRPC3, expressed transiently in COS cells, are activated by DAG, most efficiently by the membrane permeant DAG analog, OAG. They also demonstrated that the effects of DAG on TRPC6 are not mediated by protein kinase C (PKC). Using whole-cell patch clamp recordings and single cell fluorimetry, Trebak et al. [47] showed that receptor-mediated activation of PLC activates TRPC3 via DAG production independently of IP₃. They also showed that phorbol esters inhibit OAG-mediated activation of TRPC3 suggesting a negative regulation of TRPC3 by PKC [47]. The activation of TRPC3-mediated cation entry by DAG has been confirmed by several groups using different expression systems [44,49,76–78,80].

Ma et al. showed that in TRPC3-transfected HEK293 cells, TRPC3 was activated after exogenous application of OAG or a DAG lipase inhibitor (RHC80267) to the cells, while control cells showed no response to either of these stimuli. In addition, OAG could activate TRPC3 after 2APB treatment whereas ATP failed to activate TRPC3 in the presence of 2APB. The authors concluded that DAG activates TRPC3 by a mechanism distinct from activation by IP₃R [76]. However, as discussed in the IP₃R section, it is likely that the effect of 2APB on TRPC3 channels is due to a direct action on the channels themselves, unrelated to IP₃R, as it is the case for native SOCs.

TRPC6 appeared to function as a receptor-regulated, but not a store-regulated cation channel. Hofmann et al. [44] showed that neither thapsigargin, ionomycin, PIP₂ nor IP₃ activated TRPC6. The PLC inhibitor, U73122, blocked agonist activation of TRPC6 leading the authors to conclude that TRPC6 is activated through PLC production of DAG, not by IP₃ or IP₃R, and not directly by G-proteins. Following TRPC6 transfection into superior cervical ganglion neurons by Delmas et al. [121], OAG, but not thapsigargin, activated a nonselective cation current, confirming that TRPC6 is not sensitive to store depletion. Inoue et al. [50] confirmed that TRPC6 is activated by DAG in a PKC-independent manner and is insensitive to activation by thapsigargin or IP₃.

Okada et al. [37] showed that OAG and the DAG lipase inhibitor RHC80267 activated TRPC7-mediated cation entry. The effect of OAG was PKC-independent but PMA blocked activation of TRPC7 by OAG suggesting a negative

regulation of TRPC7 by PKC as described for TRPC3 and TRPC6 [47,50]. Reciprocally, a PKC inhibitor, BIM I, caused a slight potentiation of OAG effects in TRPC7-mediated cation entry [37].

The mechanisms of activation of TRPC1, TRPC2 and TRPC4 and 5 are much less certain. Ectopically expressed TRPC4 and 5 were shown to be insensitive to DAG although their activation clearly depended upon phospholipase C [53]. Lintschinger et al. [39] reported that TRPC1 was only activated by DAG when associated with TRPC3; however, subsequent work by Hofmann et al. [25] cast doubt on the likelihood of co-assembly of TRPC3 and TRPC1 subunits. The evidence that TRPC1 may be a component of store-operated channels, or regulated by IP₃R, has been discussed above. The mechanism of activation of TRPC2 is still largely unknown. The sensitivity of TRPC2 to DAG when ectopically expressed has apparently not been investigated. Recently, Lucas et al. [41] identified a native calcium-permeable channel in vomeronasal neuron dendrites that was activated by DAG, independently of Ca²⁺, IP₃, arachidonic acid and PKC. When the TRPC2 gene was deleted in mice, a severe deficit in the native DAG-activated channel was observed suggesting the involvement of TRPC2 protein as a component of this DAG-activated channel. Based on antisense studies, TRPC4 has been implicated in arachidonic acid-mediated Ca²⁺ entry in mammalian cells [92]. This activation was PKC-independent.

4.4. Calcium and CaM

Calcium is known to influence TRPC activity, and there is evidence that CaM is involved in this regulation. Initial studies on TRPC3 channels concluded that TRPC3 could function as a calcium-activated channel [75]. However, stimulation by raising Ca²⁺ was only seen in whole cell mode, and not with single channels in excised patches. It is possible that this effect may reflect Ca²⁺ activation of PLC. Nonetheless, there is considerable evidence for Ca²⁺ and CaM regulation of TRPCs, more commonly in an inhibitory mode.

CaM, a soluble Ca²⁺ binding and Ca²⁺ regulatory protein, is known to be involved in the regulation of many cellular functions, including channel activity [122]. CaM has been shown to bind to *Drosophila* TRP [123,124] at two distinct sites, both of which are capable of regulating channel activity. Zhang et al. [107] demonstrated that TRPC3 bound calcium-CaM at a site that overlapped with the IP₃ binding domain. In this study it was shown that TRPC3 activity could be increased by protocols that displaced CaM from TRPC3 and also by calmidazolium. This CaM IP₃ binding domain (CIRB) was later shown to be present on other TRPC channels, with CaM binding in a calcium-dependent manner [97]. In the presence of calcium, this binding was inhibitory, although the apparent affinity of TRPC channels for binding differed for each TRPC channel

type. Interestingly, when the CIRB region was deleted from TRPC3 channels, a loss of function was observed that was due to incorrect trafficking of the channel [14].

Although most studies have implicated Ca^{2+} and CaM as inhibitory regulators of TRPC channels, in one study, TRPC6 appeared to require CaM function for positive regulation. Boulay [125] reported that CaM binding to TRPC6 was inhibited by the CaM inhibitors calmidazolium and trifluoperazine. These inhibitors also prevented agonist-activated entry through TRPC6 channels, but not the endogenous store-operated channels. However, in an earlier study, the closely related channel, TRPC3, was shown to be unaffected by the CaM inhibitor calmidazolium [126].

4.5. Phosphorylation of TRPC channels

Despite the existence of multiple potential phosphorylation sites in the primary sequence of all TRPCs, few studies have specifically addressed the role of kinases in TRPC channel activity. Phorbol 12-myristoyl 13-acetate (PMA)-induced activation of PKC has been shown to block OAG-induced activation of ectopically expressed mouse TRPC7 [37] and rat TRPC6 [127], but, curiously, reported to have no effect on human TRPC6 [128]. Another PKC activator, phorbol 12,13-dibutyrate, was reported to significantly reduce spontaneous inward currents mediated by mouse TRPC7 in HEK293 cells [37]. Recent work provided pharmacological evidence that PKC negatively modulates DAG-activation of the human isoform of TRPC3 [47]. The authors showed that in TRPC3-expressing HEK293 cells, activation of PKC with PMA completely abrogated the ability of OAG to activate TRPC3, whereas down-regulation of PMA-sensitive PKC isoforms prevented blockade by the phorbol ester. Similar findings were reported by Venkatachalam et al. [120], who showed that TRPC5 and TRPC4 were also inhibited by phorbol esters. In their studies, exogenous application of OAG, or pharmacological inhibition of DAG metabolism, prevented activation of human TRPC4 and TRPC5 when expressed in HEK293 cells or DT40 B lymphocytes.

Resolution of the temporal interrelation between DAG-induced activation of TRPCs and DAG-dependent activation of PKC upon receptor stimulation, as well as identification of the PKC isoform/s involved, would significantly improve understanding of the physiological relevance of such a mechanism in regulation of these channels. Of note, recent work from Minke's lab indicates that *Drosophila* TRP and TRPL channel activities are antagonized by PKC-dependent phosphorylation, but also phosphatase inhibition by calyculin A (calyA) abolishes channel activation, suggesting that a phosphorylation-dephosphorylation cycle regulates channel activity [129]. As of this writing, there appear to be no studies specifically addressing the role of protein phosphatases in the signaling mechanisms for the mammalian TRPC family.

In contrast to the findings reported for TRPC3, a positive modulatory role of PKC was proposed in TRPC1 regulation [130]. Pharmacological inhibition of PKC or expression of a kinase-defective mutant of PKC α markedly inhibited IP₃-induced TRPC1-mediated currents in human endothelial cells. Also, store depletion through either thrombin stimulation or thapsigargin induced a rapid and sustained phosphorylation of TRPC1. Channel phosphorylation required functional PKC α , as it was not observed in cells overexpressing the kinase-defective mutant of the enzyme. Although the data indicated that PKC α activity is required for signaling to TRPC1, the question remains whether the channel itself is the direct substrate for PKC α .

Protein kinase G (PKG) was also reported to have a role in negative regulation of human TRPC3 activity [131]. In vitro analysis of TRPC3 phosphorylation showed that the channel protein, either immunoprecipitated from HEK293 cells transiently expressing TRPC3 or synthesized in vitro, could be phosphorylated by active PKG α 1, and that a specific PKG inhibitor abolished this phosphorylation. Point mutations at two of the three potential PKG phosphorylation sites within the TRPC3 sequence (Thr-11 and Ser-263) reduced the phosphorylation signal in the in vitro PKG assays. In the same study, fluorescence monitoring of TRPC3-mediated calcium entry indicated that cGMP inhibited TRPC3-mediated calcium entry in a PKG-dependent fashion, an effect not observed in the TRPC3 mutants. The nature of the assay used in this study makes it likely that only constitutive activity of the channels was assessed. In ³²P-labeled platelets, TRPC6 was significantly phosphorylated by PKA and PKG, and associated with other unidentified phosphoproteins; in this study, channel phosphorylation did not appear to affect either thrombin- or OAG-induced TRPC6 activity [132].

The possibility that tyrosine kinases might play a role in regulation of TRPCs has recently received some attention. Hassock et al. [132] reported that in platelets, under conditions that provide widespread tyrosine phosphorylation of cellular proteins, endogenous TRPC6 and TRPC1 did not undergo tyrosine phosphorylation. A more recent report implicated the Src family kinase Fyn in epidermal growth factor (EGF)-dependent regulation of human TRPC6 [133]. In COS-7 cells overexpressing both Fyn and TRPC6, the channel was tyrosine phosphorylated following EGF treatment, and Fyn phosphorylated TRPC6 in an in vitro kinase assay. Fyn and TRPC6 were shown to interact through the SH2 domain of Fyn and the channel N-terminal domain of TRPC6 in a manner independent of channel phosphorylation. In inside-out patches excised from TRPC6-expressing HEK293 cells, Fyn-dependent phosphorylation of TRPC6 increased OAG-induced channel activity. These findings contrast with those from Hassock et al. [132] who observed no tyrosine phosphorylation of endogenous TRPC6 in human platelets.

Thus, despite some contradictory findings, the currently available evidence suggests that protein phosphorylation may

play an important regulatory role in signaling to TRPC channels. A number of kinases have been recently implicated in regulation of TRP channels outside the TRPC family. PKC and PKA modulate TRPV1 activity by direct channel phosphorylation [134], whereas Src is involved in regulation of TRPM7 [135], TRPV1 [136] and TRPV4 [137]. Thus, protein phosphorylation is emerging as a potential critical regulatory element within the signaling mechanisms to TRP superfamily channels in general.

5. Understanding the physiological functions of TRPC channels: future directions

In non-excitable cells, plasma membrane receptors coupled to PLC drive mobilization of Ca^{2+} from ER stores through IP_3 -mediated Ca^{2+} release, and Ca^{2+} entry across the plasma membrane through both store-operated (SOC) and/or non-store-operated (non-SOC) Ca^{2+} entry pathways. The multitude of cellular functions and responses that are dependent upon Ca^{2+} entry into the cell (reviewed in Refs. [56,138,139] requires a finely tuned regulation of the Ca^{2+} channels mediating the entry process. Currently, the molecular identity of SOCs and many other regulated channels, especially those regulated by PLC, remains uncertain. However, evidence continues to accrue indicating important roles for members of the TRPC family of cation channels in several different modes of Ca^{2+} entry. As discussed above, and despite the existence of controversial results, each member of the TRPC family has been implicated in both store-operated and store-independent Ca^{2+} entry in different cell lines. Importantly, TRPC channels share the common property of being activated by PLC-linked signals, an early and common event in G-protein-coupled receptor and receptor tyrosine kinase calcium signaling in a variety of physiological situations. Perhaps the most significant example of the potential physiological importance of these channels is as a component of the endogenous nonselective cation channels in smooth muscle cells. In smooth muscle cells, the biophysical properties of the endogenous channels as well as their sensitivity to DAG closely resemble ectopically expressed TRPC6 [50], pointing to TRPC6 as an important player in regulation of vascular tone.

The next era of research on TRPC channels should focus on several issues. First, it is important to resolve the many discrepancies in the literature that impede understanding of the true functions and modes of regulation of these channels. Understanding subtle effects of expression conditions and environment, for example differences in expression level, can help to resolve these inconsistencies. Second, it is important to resolve the current conundrum involving the role of TRPCs in store-operated channels. Although there are clearly examples of store-operated nonselective cation channels for which the TRPCs make excellent candidates, there is also evidence for a role of TRPCs in the more Ca^{2+} -selective SOCs, such as I_{crac} . How

can channels that consistently behave as nonselective cation channels when expressed ectopically form Ca^{2+} -selective channels in situ, or in fact do they? Finally we need to move from the logical first step of understanding the regulation of the channels in expression systems to investigations of their roles in real cells in real physiological systems. To this end, we will need to rely to a greater extent on knock-down strategies such as knock-out animals and RNAi gene silencing. Hopefully in the near future a clearer and more consistent picture of the true functions and regulatory mechanisms for TRPC channels will emerge.

References

- [1] X. Zhu, P.B. Chu, M. Peyton, L. Birnbaumer, Molecular cloning of a widely expressed human homologue for the *Drosophila* trp gene, *FEBS Lett.* 373 (1995) 193–198.
- [2] P.D. Wes, J. Chevesich, A. Jeromin, C. Rosenberg, G. Stetten, C. Montell, TRPC1, a human homolog of a *Drosophila* store-operated channel, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 9652–9656.
- [3] R.C. Hardie, B. Minke, Novel Ca^{2+} channels underlying transduction in *Drosophila* photoreceptors: implications for phosphoinositide-mediated Ca^{2+} mobilization, *Trends Neurosci.* 16 (1993) 371–376.
- [4] L. Birnbaumer, X. Zhu, M. Jiang, G. Boulay, M. Peyton, B. Vannier, D. Brown, D. Platano, H. Sadeghi, E. Stefani, M. Birnbaumer, On the molecular basis and regulation of cellular capacitative calcium entry: Roles for Trp proteins, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 15195–15202.
- [5] C. Montell, Visual transduction in *Drosophila*, *Annu. Rev. Cell Dev. Biol.* 15 (1999) 231–268.
- [6] D.E. Clapham, L.W. Runnels, C. Strübing, The TRP ion channel family, *Nat. Rev., Neurosci.* 2 (2002) 387–396.
- [7] R. Vennekens, T. Voets, J.M. Bindels, G. Droogmans, B. Nilius, Current understanding of mammalian TRP homologues, *Cell Calcium* 31 (2002) 253–264.
- [8] C. Zitt, C.R. Halaszovich, A. Lückhoff, The TRP family of cation channels: probing and advancing the concepts on receptor-activated calcium entry, *Prog. Neurobiol.* 66 (2002) 243–264.
- [9] C. Montell, L. Birnbaumer, V. Flockerzi, R.J. Bindels, E.A. Bruford, M.J. Caterina, D.E. Clapham, C. Harteneck, S. Heller, D. Julius, I. Kojima, Y. Mori, R. Penner, D. Prawitt, A.M. Scharenberg, G. Schultz, N. Shimizu, M.X. Zhu, A unified nomenclature for the superfamily of TRP cation channels, *Mol. Cell* 9 (2002) 229–231.
- [10] E.R. Liman, H. Innan, Relaxed selective pressure on an essential component of pheromone transduction in primate evolution, *Proc. Natl. Acad. Sci.* 100 (2003) 3328–3332.
- [11] B. Vannier, X. Zhu, D. Brown, L. Birnbaumer, The membrane topology of human transient receptor potential 3 as inferred from glycosylation-scanning mutagenesis and epitope immunocytochemistry, *J. Biol. Chem.* 273 (1998) 8675–8679.
- [12] Y. Dohke, Y.S. Oh, I.S. Ambudkar, R.J. Turner, Biogenesis and topology of the transient receptor potential Ca^{2+} channel TRPC1, *J. Biol. Chem.* 279 (2004) 12242–12248.
- [13] L.K. Mosavi, T.J. Cammett, D.C. Desrosiers, Z.y. Peng, The ankyrin repeat as molecular architecture for protein recognition, *Protein Sci.* 13 (2004) 1435–1448.
- [14] B.J. Wedel, G. Vazquez, R.R. McKay, G.St.J. Bird, J.W. Putney Jr., A calmodulin/inositol 1,4,5-trisphosphate (IP_3) receptor-binding region targets TRPC3 to the plasma membrane in a calmodulin/ IP_3 receptor-independent process, *J. Biol. Chem.* 278 (2003) 25758–25765.
- [15] C. Zitt, A. Zobel, A.G. Obukhov, C. Harteneck, F. Kalkbrenner, A. Lückhoff, G. Schultz, Cloning and functional expression of a human

- Ca²⁺-permeable cation channel activated by calcium store depletion, *Neuron* 16 (1996) 1189–1196.
- [16] M. Engelke, O. Friedrich, P. Budde, C. Schafer, U. Niemann, C. Zitt, E. Jüngling, O. Rocks, A. Lückhoff, J. Frey, Structural domains required for channel function of the mouse transient receptor potential protein homologue TRP1b, *FEBS Lett.* 523 (2002) 193–199.
- [17] A. Greka, B. Navarro, E. Oancea, A. Duggan, D.E. Clapham, TRPC5 is a regulator of hippocampal neurite length and growth cone morphology, *Nat. Neurosci.* 6 (2003) 837–845.
- [18] B.B. Singh, X. Liu, J. Tang, M.X. Zhu, I.S. Ambudkar, Calmodulin regulates Ca²⁺-dependent feedback inhibition of store-operated Ca²⁺ influx by interaction with a site in the C terminus of TrpC1, *Mol. Cell* 9 (2002) 739–750.
- [19] W.M. Krajewska, I. Maslowska, Caveolins: structure and function in signal transduction, *Cell. Mol. Biol. Lett.* 9 (2004) 195–220.
- [20] S.c. Brazer, B.B. Singh, X. Liu, W. Swaim, I.S. Ambudkar, Caveolin-1 Contributes to Assembly of Store-operated Ca²⁺ Influx Channels by Regulating Plasma Membrane Localization of TRPC1, *J. Biol. Chem.* 278 (2003) 27208–27215.
- [21] T.P. Lockwich, X. Liu, B.B. Singh, J. Jadlowiec, S. Wieland, I.S. Ambudkar, Assembly of Trp1 in a signaling complex associated with caveolin-scaffolding lipid raft domains, *J. Biol. Chem.* 275 (2000) 11934–11942.
- [22] T. Lockwich, B.B. Singh, X. Liu, I.S. Ambudkar, Stabilization of cortical actin induces internalization of transient receptor potential 3 (Trp3)-associated caveolar Ca²⁺ signaling complex and loss of Ca²⁺ influx without disruption of Trp3-inositol trisphosphate receptor association, *J. Biol. Chem.* 276 (2001) 42401–42408.
- [23] J. Couet, S.W. Li, T. Okamoto, T. Ikezu, M.P. Lisanti, Identification of peptide and protein ligands for the caveolin-scaffolding domain: Implications for the interaction of caveolin with caveolin-associated proteins, *J. Biol. Chem.* 272 (1997) 6525–6533.
- [24] C. Strübing, G. Krapivinsky, L. Krapivinsky, D.E. Clapham, Formation of novel TRPC channels by complex subunit interactions in embryonic brain, *J. Biol. Chem.* 278 (2003) 39014–39019.
- [25] T. Hofmann, M. Schaefer, G. Schultz, T. Gudermann, Subunit composition of mammalian transient receptor potential channels in living cells, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 7461–7466.
- [26] S. Jung, A. Mühle, M. Schaefer, R. Strotmann, G. Schultz, T.D. Plant, Lanthanides potentiate TRPC5 currents by an action at extracellular sites close to the pore mouth, *J. Biol. Chem.* 278 (2003) 3562–3571.
- [27] X. Liu, B.B. Singh, I.S. Ambudkar, TRPC1 is required for functional store-operated Ca²⁺ channels. Role of acidic amino acid residues in the S5–S6 region, *J. Biol. Chem.* 278 (2003) 11337–11343.
- [28] J.P. Yuan, K. Kiselyov, D.M. Shin, J. Chen, N. Sheyknikov, S.H. Kang, M.H. Dehoff, M.K. Schwarz, P.H. Seeburg, S. Muallem, P.F. Worley, Homer binds TRPC family channels and is required for gating of TRPC1 by IP(3) receptors, *Cell* 114 (2003) 777–789.
- [29] W.G. Sinkins, M. Goel, M. Estacion, W.P. Schilling, Association of immunophilins with mammalian TRPC channels, *J. Biol. Chem.* 279 (2004) 34521–34529.
- [30] B.B. Singh, X. Liu, I.S. Ambudkar, Expression of truncated transient receptor potential protein 1α (Trp1α). Evidence that the Trp1 C terminus modulates store-operated Ca²⁺ entry, *J. Biol. Chem.* 275 (2000) 36483–36486.
- [31] M. Schaefer, T.D. Plant, N. Stresow, N. Albrecht, G. Schultz, Functional differences between TRPC4 splice variants, *J. Biol. Chem.* 277 (2002) 3752–3759.
- [32] L. Mery, F. Magnino, K. Schmidt, K.-H. Krause, Alternative splice variants of hTrp4 differentially interact with the C-terminal portion of the inositol 1,4,5-trisphosphate receptors, *FEBS Lett.* 487 (2001) 377–383.
- [33] C. Trost, C. Bergs, N. Himmerkus, V. Flockerzi, The transient receptor potential, TRP4, cation channel is a novel member of the family of calmodulin binding proteins, *Biochem. J.* 355 (2001) 663–670.
- [34] Y. Tang, J. Tang, Z. Chen, C. Trost, V. Flockerzi, M. Li, V. Ramesh, M.X. Zhu, Association of mammalian Trp4 and phospholipase C isozymes with a PDZ domain-containing protein, NHERF, *J. Biol. Chem.* 275 (2000) 37559–37564.
- [35] L. Mery, B. Strauß, J.F. Dufour, K.H. Krause, M. Hoth, The PDZ-interacting domain of TRPC4 controls its localization and surface expression in HEK293 cells, *J. Cell. Sci.* 115 (2002) 3497–3508.
- [36] A. Dietrich, M. Schnitzler, J. Emmel, H. Kalwa, T. Hofmann, T. Gudermann, N-linked protein glycosylation is a major determinant for basal TRPC3 and TRPC6 channel activity, *J. Biol. Chem.* 278 (2003) 47842–47852.
- [37] T. Okada, R. Inoue, K. Yamazaki, A. Maeda, T. Kurosaki, T. Yamakuni, I. Tanaka, S. Shimizu, K. Ikenaka, K. Imoto, Y. Mori, Molecular and functional characterization of a novel mouse transient receptor potential protein homologue TRP7. Ca²⁺-permeable cation channel that is constitutively activated and enhanced by stimulation of G protein-coupled receptor, *J. Biol. Chem.* 274 (1999) 27359–27370.
- [38] C. Strübing, G. Krapivinsky, L. Krapivinsky, D.E. Clapham, TRPC1 and TRPC5 form a novel cation channel in mammalian brain, *Neuron* 29 (2001) 645–655.
- [39] B. Lintschinger, M. Balzer-Geldsetzer, T. Baskaran, W.F. Graier, C. Romanin, M.X. Zhu, K. Groschner, Coassembly of Trp1 and Trp3 proteins generates diacylglycerol- and Ca²⁺-sensitive cation channels, *J. Biol. Chem.* 275 (2000) 27799–27805.
- [40] M. Hoth, R. Penner, Depletion of intracellular calcium stores activates a calcium current in mast cells, *Nature* 355 (1992) 353–355.
- [41] P. Lucas, K. Ukhanov, T. Leinders-Zufall, F. Zufall, A diacylglycerol-gated cation channel in vomeronasal neuron dendrites is impaired in TRPC2 mutant mice: mechanism of pheromone transduction, *Neuron* 40 (2003) 551–561.
- [42] B. Vannier, M. Peyton, G. Boulay, D. Brown, N. Qin, M. Jiang, X. Zhu, L. Birnbaumer, Mouse trp2, the homologue of the human trpc2 pseudogene, encodes mTrp2, a store depletion-activated capacitative Ca²⁺ channel, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 2060–2064.
- [43] X. Zhu, M. Jiang, L. Birnbaumer, Receptor-activated Ca²⁺ influx via human Trp3 stably expressed in human embryonic kidney (HEK)293 cells. Evidence for a non-capacitative calcium entry, *J. Biol. Chem.* 273 (1998) 133–142.
- [44] T. Hofmann, A.G. Obukhov, M. Schaefer, C. Harteneck, T. Gudermann, G. Schultz, Direct activation of human TRPC6 and TRPC3 channels by diacylglycerol, *Nature* 397 (1999) 259–262.
- [45] K. Kiselyov, X. Xu, G. Mozhayeva, T. Kuo, I. Pessah, G. Mignery, X. Zhu, L. Birnbaumer, S. Muallem, Functional interaction between InsP₃ receptors and store-operated hTrp3 channels, *Nature* 396 (1998) 478–482.
- [46] K.-D. Preuß, J.K. Nöller, E. Krause, A. Göbel, I. Schulz, Expression and characterization of a trp1 homolog from rat, *Biochem. Biophys. Res. Commun.* 240 (1997) 167–172.
- [47] M. Trebak, G.St.J. Bird, R.R. McKay, L. Birnbaumer, J.W. Putney Jr., Signaling mechanism for receptor-activated TRPC3 channels, *J. Biol. Chem.* 278 (2003) 16244–16252.
- [48] M. Trebak, G.St.J. Bird, R.R. McKay, J.W. Putney Jr., Comparison of human TRPC3 channels in receptor-activated and store-operated modes. Differential sensitivity to channel blockers suggests fundamental differences in channel composition, *J. Biol. Chem.* 277 (2002) 21617–21623.
- [49] C.R. Halaszovich, C. Zitt, E. Jüngling, A. Lückhoff, Inhibition of TRP3 by lanthanides. Block from the cytosolic side of the plasma membrane, *J. Biol. Chem.* 275 (2000) 37423–37428.
- [50] R. Inoue, T. Okada, H. Onoue, Y. Hara, S. Shimizu, S. Naitoh, Y. Ito, Y. Mori, The transient receptor potential protein homologue TRP6 is the essential component of vascular α1-adrenoceptor-activated Ca²⁺-permeable cation channel, *Circ. Res.* 88 (2001) 325–332.
- [51] D.E. Clapham, L.W. Runnels, C. Strübing, The TRP ion channel family, *Nat. Rev., Neurosci.* 2 (2001) 387–396.

- [52] S. Philipp, A. Cavalié, M. Freichel, U. Wissenbach, S. Zimmer, C. Trost, A. Marguart, M. Murakami, V. Flockerzi, A mammalian capacitative calcium entry channel homologous to *Drosophila* TRP and TRPL, *EMBO J.* 15 (1996) 6166–6171.
- [53] M. Schaefer, T.D. Plant, A.G. Obukhov, T. Hofmann, T. Gudermann, G. Schultz, Receptor-mediated regulation of the nonselective cation channels TRPC4 and TRPC5, *J. Biol. Chem.* 275 (2000) 17517–17526.
- [54] S. Philipp, J. Hambrecht, L. Braslavski, G. Schroth, M. Freichel, M. Murakami, A. Cavalié, V. Flockerzi, A novel capacitative calcium entry channel expressed in excitable cells, *EMBO J.* 17 (1998) 4274–4282.
- [55] T. Okada, S. Shimizu, M. Wakamori, A. Maeda, T. Kurosaki, N. Takada, K. Imoto, Y. Mori, Molecular cloning and functional characterization of a novel receptor-activated TRP Ca^{2+} channel from mouse brain, *J. Biol. Chem.* 273 (1998) 10279–10287.
- [56] J.W. Putney Jr., *Capacitative Calcium Entry*, Landes Biomedical Publishing, Austin, TX, 1997.
- [57] E.S. Trepakova, P. Csutora, D.L. Hunton, R.B. Marchase, R.A. Cohen, V.M. Bolotina, Calcium influx factor (CIF) directly activates store-operated cation channels in vascular smooth muscle cells, *J. Biol. Chem.* 275 (2000) 26158–26163.
- [58] A.P. Albert, W.A. Large, A Ca^{2+} -permeable non-selective cation channel activated by depletion of internal Ca^{2+} stores in single rabbit portal vein myocytes, *J. Physiol. (Lond.)* 538 (2002) 717–728.
- [59] J.W. Putney Jr., L.M. Broad, F.-J. Braun, J.-P. Lièvreumont, G.St.J. Bird, Mechanisms of capacitative calcium entry, *J. Cell. Sci.* 114 (2001) 2223–2229.
- [60] K. Venkatachalam, D.B. van Rossum, R.L. Patterson, H.-T. Ma, D.L. Gill, The cellular and molecular basis of store-operated calcium entry, *Nat. Cell Biol.* 4 (2002) E263–E272.
- [61] D.J. Beech, S.Z. Xu, D. McHugh, R. Flemming, TRPC1 store-operated cationic channel subunit, *Cell Calcium* 33 (2003) 433–440.
- [62] X. Liu, W. Wang, B.B. Singh, T. Lockwich, J. Jadlowiec, B. O’Connell, R. Wellner, M.X. Zhu, I.S. Ambudkar, Trp1, a candidate protein for the store-operated Ca^{2+} influx mechanism in salivary gland cells, *J. Biol. Chem.* 275 (2000) 3403–3411.
- [63] X. Wu, G. Babnigg, M.L. Villereal, Functional significance of human trp1 and trp3 in store-operated Ca^{2+} entry in HEK-293 cells, *Am. J. Physiol.* 278 (2000) C526–C536.
- [64] Y. Tomita, S. Kaneko, M. Funayama, H. Kondo, M. Satoh, A. Akaike, Intracellular Ca^{2+} store-operated influx of Ca^{2+} through TRP-R, a rat homolog of TRP, expressed in *Xenopus* oocytes, *Neurosci. Lett.* 248 (1998) 195–198.
- [65] G.H. Brough, S. Wu, D. Cioffi, T.M. Moore, M. Li, N. Dean, T. Stevens, Contribution of endogenously expressed Trp1 to a Ca^{2+} -selective, store-operated Ca^{2+} entry pathway, *FASEB J.* 15 (2001) 1727–1738.
- [66] M. Sweeney, Y. Yu, O. Platoshyn, S. Zhang, S.S. McDaniel, J.X.J. Yuan, Inhibition of endogenous TRP1 decreases capacitative Ca^{2+} entry and attenuates pulmonary artery smooth muscle cell proliferation, *Am. J. Physiol.* 283 (2002) L144–L155.
- [67] Y. Mori, M. Wakamori, T. Miyakawa, M. Hermosura, Y. Hara, M. Nishida, K. Hirose, A. Mizushima, M. Kurosaki, E. Mori, K. Gotoh, T. Okada, A. Fleig, R. Penner, M. Iino, T. Kurosaki, Transient receptor potential 1 regulates capacitative Ca^{2+} entry and Ca^{2+} release from endoplasmic reticulum in B lymphocytes, *J. Exp. Med.* 195 (2002) 673–681.
- [68] S.Z. Xu, D.J. Beech, TrpC1 is a membrane-spanning subunit of store-operated Ca^{2+} channels in native vascular smooth muscle cells, *Circ. Res.* 88 (2001) 84–87.
- [69] J.A. Rosado, S.L. Brownlow, S.O. Sage, Endogenously expressed Trp1 is involved in store-mediated Ca^{2+} entry by conformational coupling in human platelets, *J. Biol. Chem.* 277 (2002) 42157–42163.
- [70] M.K. Jungnickel, H. Marreo, L. Birnbaumer, J.R. Lemos, H.M. Florman, Trp2 regulates entry of Ca^{2+} into mouse sperm triggered by egg ZP3, *Nat. Cell Biol.* 3 (2001) 499–502.
- [71] T. Hofmann, M. Schaefer, G. Schultz, T. Gudermann, Cloning, expression and subcellular localization of two splice variants of mouse transient receptor potential channel 2, *Biochem. J.* 351 (2000) 115–122.
- [72] P. Gailly, M. Colson-Van Schoor, Involvement of trp-2 protein in store-operated influx of calcium in fibroblasts, *Cell Calcium* 30 (2001) 157–165.
- [73] X. Zhu, M. Jiang, M. Peyton, G. Boulay, R. Hurst, E. Stefani, L. Birnbaumer, *trp*, a novel mammalian gene family essential for agonist-activated capacitative Ca^{2+} entry, *Cell* 85 (1996) 661–671.
- [74] M. Trebak, G. Vazquez, G.St.J. Bird, J.W. Putney Jr., The TRPC3/6/7 subfamily of cation channels, *Cell Calcium* 33 (2003) 451–461.
- [75] C. Zitt, A.G. Obukhov, C. Strübing, A. Zobel, F. Kalkbrenner, A. Lückhoff, G. Schultz, Expression of TRPC3 in Chinese hamster ovary cells results in calcium-activated cation currents not related to store depletion, *J. Cell Biol.* 138 (1997) 1333–1341.
- [76] H.-T. Ma, R.L. Patterson, D.B. van Rossum, L. Birnbaumer, K. Mikoshiba, D.L. Gill, Requirement of the inositol trisphosphate receptor for activation of store-operated Ca^{2+} channels, *Science* 287 (2000) 1647–1651.
- [77] R.R. McKay, C.L. Szmeczek-Seay, J.-P. Lièvreumont, G.St.J. Bird, C. Zitt, E. Jüngling, A. Lückhoff, J.W. Putney Jr., Cloning and expression of the human transient receptor potential 4 (TRP4) gene: localization and functional expression of human TRP4 and TRP3, *Biochem. J.* 351 (2000) 735–746.
- [78] G. Vazquez, J.-P. Lièvreumont, G.St.J. Bird, J.W. Putney Jr., Trp3 forms both inositol trisphosphate receptor-dependent and independent store-operated cation channels in DT40 avian B-lymphocytes, *Proc. Natl. Acad. Sci. U. S. A.* 98 (2001) 11777–11782.
- [79] G. Vazquez, B.J. Wedel, M. Trebak, G.St.J. Bird, J.W. Putney Jr., Expression level of TRPC3 channel determines its mechanism of activation, *J. Biol. Chem.* 278 (2003) 21649–21654.
- [80] K. Venkatachalam, H.-T. Ma, D.L. Ford, D.L. Gill, Expression of functional receptor-coupled TRPC3 channels in DT40 triple receptor *InsP3* knockout cells, *J. Biol. Chem.* 276 (2001) 33980–33985.
- [81] L. Yue, J.-B. Peng, M.A. Hediger, D.E. Clapham, CaT1 manifests the pore properties of the calcium release activated calcium channel, *Nature* 410 (2001) 705–709.
- [82] J.W. Putney Jr., The enigmatic TRPCs: multifunctional cation channels, *Trends Cell Biol.* 14 (2004) 282–286.
- [83] A. Riccio, C. Mattei, R.E. Kelsell, A.D. Medhurst, A.R. Calver, A.D. Randall, J.B. Davis, C.D. Benham, M.N. Pangalos, Cloning and functional expression of human short TRP7, a candidate protein for store-operated Ca^{2+} influx, *J. Biol. Chem.* 277 (2002) 12302–12309.
- [84] G. Boulay, X. Zhu, M. Peyton, M. Jiang, R. Hurst, E. Stefani, L. Birnbaumer, Cloning and expression of a novel mammalian homolog of *Drosophila* Transient Receptor Potential (Trp) involved in calcium entry secondary to activation of receptors coupled by the Gq class of G protein, *J. Biol. Chem.* 272 (1997) 29672–29680.
- [85] J. Wamat, S. Philipp, S. Zimmer, V. Flockerzi, A. Cavalié, Phenotype of a recombinant store-operated channel: highly selective permeation of Ca^{2+} , *J. Physiol. (Lond.)* 518 (1999) 631–638.
- [86] M. Kinoshita, A. Akaike, M. Satoh, S. Kaneko, Positive regulation of capacitative Ca^{2+} entry by intracellular Ca^{2+} in *Xenopus* oocytes expressing rat TRP4, *Cell Calcium* 28 (2000) 151–159.
- [87] S. Philipp, C. Trost, J. Wamat, J. Rautmann, N. Himmerkus, G. Schroth, O. Kretz, W. Nastainczyk, A. Cavalié, M. Hoth, V. Flockerzi, Trp4 (CCE1) protein is part of native calcium release-activated Ca^{2+} -like channels in adrenal cells, *J. Biol. Chem.* 275 (2000) 23965–23972.
- [88] M. Freichel, S.H. Suh, A. Pfeifer, U. Schweig, C. Trost, P. Weißgerber, M. Biel, S. Philipp, D. Freise, G. Droogmans, F. Hofmann, V. Flockerzi, B. Nilius, Lack of an endothelial store-operated Ca^{2+} current impairs agonist-dependent vasorelaxation in TRP4^{-/-} mice, *Nat. Cell Biol.* 3 (2001) 121–127.

- [89] C. Tiruppathi, M. Freichel, S.M. Vogel, B.C. Paria, D. Mehta, V. Flockerzi, A.B. Malik, Impairment of store-operated Ca^{2+} entry in TRPC4(–/–) mice interferes with increase in lung microvascular permeability, *Circ. Res.* 91 (2002) 70–76.
- [90] X. Wang, J.L. Pluznick, P. Wei, B.J. Padanilam, S.C. Sansom, TRPC4 forms store-operated Ca^{2+} channels in mouse mesangial cells, *Am. J. Physiol., Cell Physiol.* 287 (2004) C357–C364.
- [91] A.G. Obukhov, M.C. Nowicky, TRPC4 can be activated by G-protein-coupled receptors and provides sufficient Ca^{2+} to trigger exocytosis in neuroendocrine cells, *J. Biol. Chem.* 277 (2002) 16172–16178.
- [92] X. Wu, G. Babnigg, T. Zagranichnaya, M.L. Villereal, The role of endogenous human Trp4 in regulating carbachol-induced calcium oscillations in HEK-293 cells, *J. Biol. Chem.* 277 (2002) 13597–13608.
- [93] H. Kanki, M. Kinoshita, A. Akaike, M. Satoh, Y. Mori, S. Kaneko, Activation of inositol 1,4,5-trisphosphate receptor is essential for the opening of mouse TRP5 channel, *Mol. Pharmacol.* 60 (2001) 989–998.
- [94] R.F. Irvine, “Quantal” Ca^{2+} release and the control of Ca^{2+} entry by inositol phosphates—a possible mechanism, *FEBS Lett.* 263 (1990) 5–9.
- [95] M.J. Berridge, Capacitative calcium entry, *Biochem. J.* 312 (1995) 1–11.
- [96] L. Birnbaumer, G. Boulay, D. Brown, M. Jiang, A. Dietrich, K. Mikoshiba, X. Zhu, N. Qin, Mechanism of capacitative Ca^{2+} entry (CCE): interaction between IP3 receptor and TRP links the internal calcium storage compartment to plasma membrane CCE channels, *Recent Prog. Horm. Res.* 55 (2000) 127–161.
- [97] J. Tang, Y. Lin, Z. Zhang, S. Tikunova, L. Birnbaumer, M.X. Zhu, Identification of common binding sites for calmodulin and inositol 1,4,5-trisphosphate receptors on the carboxy termini of Trp channels, *J. Biol. Chem.* 276 (2001) 21303–21310.
- [98] J.A. Rosado, S.O. Sage, Coupling between inositol 1,4,5-trisphosphate receptors and human transient receptor potential channel 1 when intracellular Ca^{2+} stores are depleted, *Biochem. J.* 350 (2000) 631–635.
- [99] J.A. Rosado, S.O. Sage, Activation of store-mediated calcium entry by secretion-like coupling between the inositol 1,4,5-trisphosphate receptor type II and human transient receptor potential (hTrp1) channels in human platelets, *Biochem. J.* 356 (2001) 191–198.
- [100] R.L. Patterson, D.B. van Rossum, D.L. Gill, Store-operated Ca^{2+} entry: evidence for a secretion-like coupling model, *Cell* 98 (1999) 487–499.
- [101] S.L. Brownlow, S.O. Sage, Rapid agonist-evoked coupling of type II Ins(1,4,5)P3 receptor with human transient receptor potential (hTRPC1) channels in human platelets, *Biochem. J.* 375 (2003) 697–704.
- [102] D. Mehta, G.U. Ahmed, B.C. Paria, M. Holinstat, T. Voynoyasenetskaya, C. Tiruppathi, R.D. Minshall, A.B. Malik, RhoA interaction with inositol 1,4,5-trisphosphate receptor and transient receptor potential channel-1 regulates Ca^{2+} entry. Role in signaling increased endothelial permeability, *J. Biol. Chem.* 278 (2003) 33492–33500.
- [103] J.H. Brann, J.C. Dennis, E.E. Morrison, D.A. Fadool, Type-specific inositol 1,4,5-trisphosphate receptor localization in the vomeronasal organ and its interaction with a transient receptor potential channel, TRPC2, *J. Neurochem.* 83 (2002) 1452–1460.
- [104] A.P. Albert, W.A. Large, Synergism between inositol phosphates and diacylglycerol on native TRPC6-like channels in rabbit portal vein myocytes, *J. Physiol.* 552 (2003) 789–795.
- [105] K. Kiselyov, G.A. Mignery, M.X. Zhu, S. Muallem, The N-terminal domain of the IP3 receptor gates store-operated hTrp3 channels, *Mol. Cell* 4 (1999) 423–429.
- [106] G. Boulay, D.M. Brown, N. Qin, M. Jiang, A. Dietrich, M.X. Zhu, Z. Chen, M. Birnbaumer, K. Mikoshiba, L. Birnbaumer, Modulation of Ca^{2+} entry by TRP-binding polypeptides of the inositol 1,4,5-trisphosphate receptor (IP3R). Evidence for roles of TRP and the IP3R in store depletion-activated Ca^{2+} entry, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 14955–14960.
- [107] Z. Zhang, J. Tang, S. Tikunova, J.D. Johnson, Z. Chen, N. Qin, A. Dietrich, E. Stefani, L. Birnbaumer, M.X. Zhu, Activation of Trp3 by inositol 1,4,5-trisphosphate receptors through displacement of inhibitory calmodulin from a common binding domain, *Proc. Natl. Acad. Sci. U. S. A.* 98 (2001) 3168–3173.
- [108] T. Maruyama, T. Kanaji, S. Nakade, T. Kanno, K. Mikoshiba, 2APB, 2-aminoethoxydiphenyl borate, a membrane-penetrable modulator of Ins(1,4,5)P3-induced Ca^{2+} release, *J. Biochem.* 122 (1997) 498–505.
- [109] F.-J. Braun, L.M. Broad, D.L. Armstrong, J.W. Putney Jr., Stable activation of single CRAC-channels in divalent cation-free solutions, *J. Biol. Chem.* 276 (2001) 1063–1070.
- [110] D. Bakowski, M.D. Glitsch, A.B. Parekh, An examination of the secretion-like coupling model for the activation of the Ca^{2+} release-activated Ca^{2+} current Icrac in RBL-1 cells, *J. Physiol. (Lond.)* 532 (2001) 55–71.
- [111] R.B. Gregory, G. Rychkov, G.J. Barritt, Evidence that 2-aminoethyl diphenylborate is a novel inhibitor of store-operated Ca^{2+} channels in liver cells, and acts through a mechanism which does not involve inositol trisphosphate receptors, *Biochem. J.* 354 (2001) 285–290.
- [112] M. Prakriya, R.S. Lewis, Potentiation and inhibition of Ca^{2+} release-activated Ca^{2+} channels by 2-aminoethyl diphenyl borate (2-APB) occurs independently of IP3 receptors, *J. Physiol. (Lond.)* 536 (2001) 3–19.
- [113] Y. Dobrydneva, P. Blackmore, 2-Aminoethoxydiphenyl borate directly inhibits store-operated calcium entry channels in human platelets, *Mol. Pharmacol.* 60 (2001) 541–552.
- [114] H. Iwasaki, Y. Mori, Y. Hara, K. Uchida, H. Zhou, K. Mikoshiba, 2-Aminoethoxydiphenyl borate (2-APB) inhibits capacitative calcium entry independently of the function of inositol 1,4,5-trisphosphate receptors, *Recept. Channels* 7 (2001) 429–439.
- [115] J.M. Diver, S.O. Sage, J.A. Rosado, The inositol trisphosphate receptor antagonist 2-aminoethoxydiphenylborate (2-APB) blocks Ca^{2+} entry channels in human platelets: cautions for its use in studying Ca^{2+} influx, *Cell Calcium* 30 (2001) 323–329.
- [116] H.-S. Li, X.-Z.S. Xu, C. Montell, Activation of a TRPC3-dependent cation channel through the neurotrophin BDNF, *Neuron* 24 (1999) 261–273.
- [117] H. Sugawara, M. Kurosaki, M. Takata, T. Kurosaki, Genetic evidence for involvement of type 1, type 2 and type 3 inositol 1,4,5-trisphosphate receptors in signal transduction through the B-cell antigen receptor, *EMBO J.* 16 (1997) 3078–3088.
- [118] L.M. Broad, F.-J. Braun, J.-P. Lièvreumont, G.St.J. Bird, T. Kurosaki, J.W. Putney Jr., Role of the phospholipase C- inositol 1,4,5-trisphosphate pathway in calcium release-activated calcium current (Icrac) and capacitative calcium entry, *J. Biol. Chem.* 276 (2001) 15945–15952.
- [119] T.D. Plant, M. Schaefer, TRPC4 and TRPC5: receptor-operated Ca^{2+} -permeable nonselective cation channels, *Cell Calcium* 33 (2003) 441–450.
- [120] K. Venkatchalam, F. Zheng, D.L. Gill, Regulation of canonical transient receptor potential (TRPC) channel function by diacylglycerol and protein kinase C, *J. Biol. Chem.* 278 (2003) 29031–29040.
- [121] P. Delmas, N. Wanaverbecq, F.C. Abogadie, M. Mistry, D.A. Brown, Signaling microdomains define the specificity of receptor-mediated InsP3 pathways in Neurons, *Neuron* 34 (2002) 209–220.
- [122] Y. Saimi, C. Kung, Calmodulin as an ion channel subunit, *Annu. Rev. Physiol.* 64 (2002) 289–311.
- [123] L. Lan, M.J. Bawden, A.M. Auld, G.J. Barritt, Expression of *Drosophila* trpl cRNA in *Xenopus laevis* oocytes leads to the appearance of a Ca^{2+} channel activated by Ca^{2+} and calmodulin, and by guanosine 5′ [γ -thio]triphosphate, *Biochem. J.* 316 (Pt 3) (1996) 793–803.

- [124] C.G. Warr, L.E. Kelly, Identification and characterization of two distinct calmodulin-binding sites in the Trp1 ion-channel protein of *Drosophila melanogaster*, *Biochem. J.* 314 (Pt 2) (1996) 497–503.
- [125] G. Boulay, Ca²⁺-calmodulin regulates receptor-operated Ca²⁺ entry activity of TRPC6 in HEK-293 cells, *Cell Calcium* 320 (2002) 201–207.
- [126] M. Kamouchi, S. Philipp, V. Flockerzi, U. Wissenbach, A. Mamin, L. Raeymaekers, J. Eggermont, G. Droogmans, B. Nilius, Properties of heterologously expressed hTRP3 channels in bovine pulmonary artery endothelial cells, *J. Physiol. (Lond.)* 518 (1999) 345–358.
- [127] L. Zhang, D. Saffen, Muscarinic acetylcholine receptor regulation of TRP6 Ca²⁺ channel isoforms, *J. Biol. Chem.* 276 (2001) 13331–13339.
- [128] M. Estacion, S. Li, W.G. Sinkins, M. Gosling, P. Bahra, C. Poll, J. Westwick, W.P. Schilling, Activation of human TRPC6 channels by receptor stimulation, *J. Biol. Chem.* 279 (2004) 22047–22056.
- [129] K. Agam, S. Frechter, B. Minke, Activation of the *Drosophila* TRP and TRPL channels requires both Ca²⁺ and protein dephosphorylation, *Cell Calcium* 35 (2004) 87–105.
- [130] G.U. Ahmmed, D. Mehta, S. Vogel, M. Holinstat, B.C. Paria, C. Tiruppathi, A.B. Malik, Protein kinase C α phosphorylates the TRPC1 channel and regulates store-operated Ca²⁺ entry in endothelial cells, *J. Biol. Chem.* 279 (2004) 20941–20949.
- [131] H.Y. Kwan, Y. Huang, X. Yao, Regulation of canonical transient receptor potential isoform 3 (TRPC3) channel by protein kinase G, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 2625–2630.
- [132] S.R. Hassock, M.X. Zhu, C. Trost, V. Flockerzi, K.S. Authi, Expression and role of TRPC proteins in human platelets: evidence that TRPC6 forms the store-independent calcium entry channel, *Blood* 100 (2002) 2801–2811.
- [133] C. Hisatsune, Y. Kuroda, K. Nakamura, T. Inoue, T. Nakamura, T. Michikawa, A. Mizutani, K. Mikoshiba, Regulation of TRPC6 channel activity by tyrosine phosphorylation, *J. Biol. Chem.* 279 (2004) 18887–18894.
- [134] L.S. Premkumar, G.P. Ahern, Induction of vanilloid receptor channel by protein kinase C, *Nature* 408 (2000) 985–990.
- [135] X. Jiang, E.W. Newell, L.C. Schlichter, Regulation of a TRPM7-like current in rat brain microglia, *J. Biol. Chem.* 278 (2003) 42867–42876.
- [136] X. Jin, N. Morsy, J. Winston, P.J. Pasricha, K. Garrett, H.I. Akbarali, Modulation of TRPV1 by non-receptor tyrosine kinase, c-src kinase, *Am. J. Physiol., Cell Physiol.* 287 (2004) C558–C563.
- [137] H. Xu, H. Zhao, W. Tian, K. Yoshida, J.-B. Roullet, D.M. Cohen, Regulation of a transient receptor potential (TRP) channel by tyrosine phosphorylation, *J. Biol. Chem.* 278 (2003) 11520–11527.
- [138] D.E. Clapham, Calcium signaling, *Cell* 80 (1995) 259–268.
- [139] M.J. Berridge, P. Lipp, M.D. Bootman, The versatility and universality of calcium signalling, *Nat. Rev., Mol. Cell Biol.* 1 (2000) 11–21.
- [140] M. Kimura, *The Neutral Theory of Molecular Evolution*, Cambridge University Press, Cambridge, England, 1983.
- [141] N. Saitou, M. Nei, The neighbor-joining method: a new method for reconstructing phylogenetic trees, *Mol. Biol. Evol.* 4 (1987) 406–425.