

Available online at www.sciencedirect.com

Biochimica et Biophysica Acta 1768 (2007) 2343

www.elsevier.com/locate/bbamem

Erratum

Erratum to “Conformational behavior of oxygenated mycobacterial mycolic acids from *Mycobacterium bovis* BCG” [Biochim. Biophys. Acta 1768 (2007) 1717–1726]

Masumi Villeneuve^{a,*}, Mizuo Kawai^a, Motoko Watanabe^b, Yutaka Aoyagi^b,
Yukio Hitotsuyanagi^b, Koichi Takeya^b, Hiroaki Gouda^c, Shuichi Hirono^c,
David E. Minnikin^d, Hiroo Nakahara^a

^a Department of Chemistry, Faculty of Science, Saitama University, 255 Shimo-okubo, Sakura-ku, Saitama, 338-8570, Japan

^b School of Pharmacy, Tokyo University of Pharmacy and Life Science, 1432-1 Horinouchi, Hachioji, Tokyo, 192-0392, Japan

^c School of Pharmaceutical Sciences, Kitasato University, 5-9-1 Shirokane, Minato-ku, Tokyo, 108-8641, Japan

^d School of Biosciences, The University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

The publisher regrets that the last column of Table 2 was omitted. The corrected Table 2 appears here.

Table 2
Thickness of Langmuir monolayer estimated by ellipsometry

Origin	Keto-MA			MeO-MA			Deoxo-MA		
	T / °C	π /mN m ⁻¹	Thickness/nm	T / °C	π /mN m ⁻¹	Thickness/nm	T / °C	π /mN m ⁻¹	Thickness/nm
BCG	18	30	2.90±0.05	18	18	2.78±0.04	30	4	3.98±0.09
	32	20	2.80±0.08	32	30	5.62±0.18	30	20	4.81±0.11
							30	40	6.10±0.08
<i>M. tb</i>	18	30	2.90±0.07	18	18	2.91±0.05			
				32	30	4.96±0.88			

DOI of original article: [10.1016/j.bbamem.2007.04.003](https://doi.org/10.1016/j.bbamem.2007.04.003).

* Corresponding author.

E-mail address: villeneuve@chem.saitama-u.ac.jp (M. Villeneuve).