



Early dark energy from zero-point quantum fluctuations

Michele Maggiore*, Lukas Hollenstein, Maud Jaccard, Ermis Mitsou

Département de Physique Théorique and Center for Astroparticle Physics, Université de Genève, 24 quai Ansermet, CH-1211 Genève 4, Switzerland

ARTICLE INFO

Article history:

Received 14 June 2011

Received in revised form 24 July 2011

Accepted 4 September 2011

Available online 7 September 2011

Editor: L. Alvarez-Gaumé

Keywords:

Early dark energy

Cosmological constant

Vacuum fluctuations

ABSTRACT

We examine a cosmological model with a dark energy density of the form $\rho_{\text{DE}}(t) = \rho_X(t) + \rho_Z(t)$, where ρ_X is the component that accelerates the Hubble expansion at late times and $\rho_Z(t)$ is an extra contribution proportional to $H^2(t)$. This form of $\rho_Z(t)$ follows from the recent proposal that the contribution of zero-point fluctuations of quantum fields to the total energy density should be computed by subtracting the Minkowski-space result from that computed in the FRW space–time. We discuss theoretical arguments that support this subtraction. By definition, this eliminates the quartic divergence in the vacuum energy density responsible for the cosmological constant problem. We show that the remaining quadratic divergence can be reabsorbed into a redefinition of Newton's constant only under the assumption that $\nabla^\mu \langle 0|T_{\mu\nu}|0\rangle = 0$, i.e. that the energy–momentum tensor of vacuum fluctuations is conserved in isolation. However in the presence of an ultra-light scalar field X with $m_X < H_0$, as typical of some dark energy models, the gravity effective action depends both on the gravitational field and on the X field. In this case general covariance only requires $\nabla^\mu (T_{\mu\nu}^X + \langle 0|T_{\mu\nu}|0\rangle) = 0$. If there is an exchange of energy between these two terms, there are potentially observable consequences. We construct an explicit model with an interaction between ρ_X and ρ_Z and we show that the total dark energy density $\rho_{\text{DE}}(t) = \rho_X(t) + \rho_Z(t)$ always remains a finite fraction of the critical density at any time, providing a specific model of early dark energy. We discuss the implication of this result for the coincidence problem and we estimate the model parameters by means of a full likelihood analysis using current CMB, SNe Ia and BAO data.

© 2011 Elsevier B.V. Open access under [CC BY license](http://creativecommons.org/licenses/by/3.0/).

1. Introduction

Understanding the origin of dark energy is one of the most important challenges facing cosmology and theoretical physics (see e.g. [1–4]). One aspect of the problem is to understand what is the role of zero-point vacuum fluctuations in cosmology. In a Friedmann–Robertson–Walker (FRW) metric with Hubble parameter $H(t)$ the bare vacuum energy density takes the form

$$[\rho_{\text{bare}}(\Lambda_c)]_{\text{FRW}} = [\rho_{\text{bare}}(\Lambda_c)]_{\text{Mink}} + \mathcal{O}(H^2(t)\Lambda_c^2), \quad (1)$$

where $[\rho_{\text{bare}}(\Lambda_c)]_{\text{Mink}}$ is the bare vacuum energy density in Minkowski space, whose leading divergence is $\mathcal{O}(\Lambda_c^4)$, and we used for definiteness a momentum space cutoff Λ_c . In the usual treatment this Λ_c^4 divergence is reabsorbed into a renormalization of the cosmological constant, giving rise to the cosmological constant problem. The divergence $\propto H^2\Lambda_c^2$ is instead absorbed into a renormalization of Newton's constant G [5,6].

In this Letter, expanding on results presented in [7], we reexamine the role of vacuum energies in cosmology. First, we will propose theoretical arguments suggesting that the correct way of

computing the physical vacuum energy is to subtract the bare vacuum energy density of Minkowski space, $[\rho_{\text{bare}}(\Lambda_c)]_{\text{Mink}}$, from the FRW result given in Eq. (1), before renormalizing the result. By definition this subtraction eliminates the troublesome Λ_c^4 divergence and, therefore, the cosmological constant problem. Then we turn our attention to the left over term $H^2\Lambda_c^2$ which now becomes the leading term in the vacuum energy. It is usually believed that this quadratic divergence can be reabsorbed into a renormalization of G . We show that this is correct only under the assumption that vacuum expectation value (VEV) of the energy–momentum tensor is conserved in isolation (an assumption that was implicit in the literature). However, general covariance of General Relativity (GR) only implies the conservation of the *total* energy–momentum tensor $T_{\mu\nu} + \langle 0|T_{\mu\nu}|0\rangle$, including both the classical term $T_{\mu\nu}$ and the semiclassical term $\langle 0|T_{\mu\nu}|0\rangle$. The separate conservation of $\langle 0|T_{\mu\nu}|0\rangle$ only takes place if we can define an effective action which depends only on the gravitational field, by integrating out the matter degrees of freedom. This is possible only if the matter degrees of freedom are heavy with respect to the energy scale of the problem, and can then be integrated out. In a cosmological setting, this means that matter fields should satisfy $m > H_0$. If, in contrast, there is an ultra-light scalar fields with $m < H_0$, as is typical of some quintessence model, this field cannot be integrated out from the effective low energy action. We show that, as a

* Corresponding author.

E-mail address: michele.maggiore@unige.ch (M. Maggiore).

result, in general $\nabla^\mu T_{\mu\nu}^X = -\nabla^\mu \langle 0|T_{\mu\nu}|0\rangle \neq 0$. In this case the effect of the quadratically divergent term in the vacuum fluctuations cannot simply be absorbed into a renormalization of Newton's constant G , and gives rise to interesting and potentially detectable cosmological effects. We construct a specific coupled early dark energy model and test it against current observations.

We use natural units where $\hbar = c = 1$, $G = M_{\text{pl}}^{-2}$. If not specified otherwise, we work in a spatially flat FRW metric with signature $(-+++)$, cosmic time t , scale factor $a(t)$ and Hubble parameter $H(t) = (da/dt)/a$. Today, the Hubble parameter and the critical density take the values H_0 and $\rho_0 = 3H_0^2/(8\pi G)$, respectively.

2. Subtraction of the flat-space vacuum energy

In Minkowski space the divergence in the vacuum energy density is usually dealt with by normal ordering the Hamiltonian, which gives by definition a vanishing result for the physical vacuum energy density. However, it is useful to realize that the problem can be treated more generally in the context of renormalization theory, which rather allows us to fix the renormalized vacuum energy density to any observed value. In the standard language of renormalization, divergences in a generic N -point Green's function are cured by adding the corresponding counterterms to the Lagrangian density. The same procedure can be applied to vacuum energy, i.e. to the $N = 0$ Green's function: one simply adds a constant counterterm $-\rho_{\text{count}}(\Lambda_c)$ to the Lagrangian density. This corresponds to adding a term $+\rho_{\text{count}}(\Lambda_c)$ to the Hamiltonian density. Hence the renormalized, physical vacuum energy density is given by $\rho_{\text{ren}} = \rho_{\text{bare}}(\Lambda_c) + \rho_{\text{count}}(\Lambda_c)$. As always in renormalization theory, the counterterm ρ_{count} is chosen so to cancel the divergences in ρ_{bare} and leave us with the desired finite part that is fixed by comparison with the experiment.

Using the language of renormalization theory is useful in this context because it makes clear that the cosmological constant problem is not that quantum field theory (QFT) gives a *wrong* prediction for the cosmological constant (as it is sometimes incorrectly said). Strictly speaking QFT makes no prediction for the cosmological constant, just as it does not predict the electron mass nor the fine structure constant. Rather, it is a problem of naturalness, in the sense that the counterterm $\rho_{\text{count}}(\Lambda_c)$ must be fine-tuned to exceeding accuracy, in order to cancel the Λ_c^4 divergence in ρ_{bare} , leaving a physical vacuum energy density that, if one identifies Λ_c with the Planck mass, is about $\mathcal{O}(10^{120})$ times smaller than Λ_c^4 .

Posing the problem in terms of a cancellation between $\rho_{\text{bare}}(\Lambda_c)$ and $\rho_{\text{count}}(\Lambda_c)$ can also give a first hint for a possible solution. First of all, one should appreciate that neither the bare vacuum energy $\rho_{\text{bare}}(\Lambda_c)$ nor the counterterm $\rho_{\text{count}}(\Lambda_c)$ have a physical meaning and only their sum is an observable. Thus, this kind of cancellation is different from a fine-tuning between observable quantities. Indeed, the Casimir effect is a well-known example where a rather similar cancellation takes place. In that case the physical vacuum energy density of a quantum field in a finite volume is found by taking the difference between the bare vacuum energy density computed in this finite volume and the bare vacuum energy density in an infinite volume. Regularizing with a cutoff Λ_c both terms diverge as Λ_c^4 , but their difference is finite and depends only on the physical size of the system. This might suggest that, similarly, to obtain the physical effect of the vacuum energy density in cosmology, one should compute the vacuum energy density in a FRW space-time and subtract from it the value computed in a reference geometry, which could be naturally taken as Minkowski space, leading to a sort of “cosmological Casimir effect”.

Before taking this analogy with the Casimir effect seriously, one must however face the obvious objection that in special relativity the zero of the energy can be chosen arbitrarily, and only energy differences with respect to the ground state are relevant.¹ In contrast, in GR we cannot choose the zero of the energy arbitrarily. One typically expects that “every form of energy gravitates”, so the contribution of Minkowski space cannot just be dropped.

While it is certainly true that in GR the choice of the zero for the energy is not arbitrary, the point that we wish to make here is that what is the correct choice can be a non-trivial issue. As a first example, consider the definition of energy for asymptotically flat space-times. This is obtained from the Hamiltonian formulation of GR, which goes back to the classic paper by Arnowitt, Deser and Misner (ADM) [8,9]. To properly define the Hamiltonian of a given field configuration in GR one must work at first in a finite three-dimensional volume V . Then the Hamiltonian takes the form $H_{\text{GR}} = H_{\text{bulk}} + H_{\text{boundary}}$, where H_{bulk} is given by an integral over the spatial volume V at fixed time, while H_{boundary} is given by an integral over the two-dimensional boundary ∂V . When one evaluates H_{bulk} on any classical solution of the equations of motion one finds a vanishing result (since H_{bulk} is proportional to the constraint equations of GR), so the whole contribution comes from the boundary term. On the other hand, H_{boundary} diverges for any asymptotically flat metric $g_{\mu\nu}$ (including flat space-time), when the boundary is finally taken to infinity. The solution proposed by ADM is to subtract from this boundary term the same term computed in Minkowski space $\eta_{\mu\nu}$. Accordingly, the energy E associated with a classical asymptotically flat metric $g_{\mu\nu}$ is obtained by defining

$$E = H_{\text{GR}}[g_{\mu\nu}] - H_{\text{GR}}[\eta_{\mu\nu}]. \quad (2)$$

This provides the standard definition of mass in GR, and reproduces the expected properties of asymptotically flat space-times. For instance, when applied to the Schwarzschild space-time, it correctly gives the mass that appears in the Schwarzschild metric. This underlines that our intuition that any form of energy gravitates according to GR is not entirely correct: Eq. (2) tells us that the energy associated to Minkowski space does not gravitate.

Similar subtractions also hold for non-asymptotically flat space-times, and can be performed either by subtracting the contribution of some reference space-time whose boundary has the same induced metric as the background under consideration [10–13], or even without introducing a reference background but just by adding some local counterterms to the boundary action, given by a coordinate-invariant functional of the intrinsic boundary geometry [14,15]. The latter prescription is particularly appealing for asymptotically AdS space-times. In fact, in the context of the AdS/CFT correspondence, this way of removing divergences in the gravitational action on the AdS side corresponds to the renormalization of the UV divergences in the conformal QFT that lives on the boundary [14–18].

These examples show that, already in classical GR, the energy that actually acts as a source of gravity can be obtained from a Hamiltonian only after performing an appropriate subtraction. It is quite natural to assume that the same should hold at the quantum level, so in particular for zero-point fluctuations of quantum fields in curved space. To understand what is the appropriate subtraction for the FRW metric, we consider the Friedmann equation that results from the Einstein equations sourced by $T_\nu^\mu + \langle 0|T_\nu^\mu|0\rangle$,

¹ Equivalently, one may observe that in the Casimir effect one actually measures the force between the plates, i.e. not the energy density itself but only its derivative w.r.t. the size of the system L . Since the divergence Λ_c^4 is independent of L , it can simply be dropped.

where $T_\nu^\mu = \text{diag}(-\rho, p, p, p)$ is the ordinary contribution of matter, radiation, etc., and $\langle 0|T_\nu^\mu|0\rangle = \text{diag}(-\rho_{\text{vac}}, p_{\text{vac}}, p_{\text{vac}}, p_{\text{vac}})$ is the corresponding contribution of zero-point fluctuations:

$$H^2(t) = \frac{8\pi G}{3}(\rho + \rho_{\text{vac}}). \quad (3)$$

We then require that Minkowski space, $H(t) = 0$, should be a solution in the limit $\rho \rightarrow 0$. This implies that all terms in $[\rho_{\text{vac}}]_{\text{FRW}}$ that do not vanish for $H \rightarrow 0$ must be subtracted. In other words, we must subtract the vacuum energy computed in Minkowski space.² In contrast, all terms proportional to H^2 or H^4 are consistent with this requirement and thus generically allowed. This procedure eliminates the Λ_c^4 term (as well as flat-space terms that appear for massive fields, such as $m^2\Lambda_c^2$ and $m^4 \ln \Lambda_c$, and are also much larger than the observed vacuum energy, for all known massive particles). Thus, this treatment of vacuum fluctuations solves the cosmological constant problem by definition, at least in its “old” form, i.e. it explains why the observed vacuum energy is many orders of magnitude smaller than M_{pl}^4 .

We next turn to the quadratic divergence left over after the subtraction. We will see that it could teach us something about the more recent form of the cosmological constant problem, namely explaining why the observed dark energy (DE) density is just of the order of the critical density of the universe today, the coincidence problem.

3. Non-interacting zero-point fluctuations and renormalization of G

Let us now discuss the fate of the quadratic divergence $\propto \Lambda_c^2 H^2(t)$ in Eq. (1). To understand what is the structure of the renormalized VEV of the energy–momentum tensor, consider the semiclassical Einstein equations for the renormalized quantities,

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi G(T_{\mu\nu} + \langle 0|T_{\mu\nu}|0\rangle), \quad (4)$$

in which the vacuum expectation value of $T_{\mu\nu}$ is added as an additional source term. Together with the Bianchi identities the above equation implies

$$\nabla^\mu(T_{\mu\nu} + \langle 0|T_{\mu\nu}|0\rangle) = 0. \quad (5)$$

This equation is therefore a consequence of the general covariance of the renormalized theory.³ If $T_{\mu\nu}$ and $\langle 0|T_{\mu\nu}|0\rangle$ are separately conserved we further have $\nabla^\mu T_{\mu\nu} = \nabla^\mu \langle 0|T_{\mu\nu}|0\rangle = 0$.

The stronger condition $\nabla^\mu \langle 0|T_{\mu\nu}|0\rangle = 0$ can indeed be derived by using the effective action for gravity, which is obtained by treating the metric $g_{\mu\nu}$ as a classical background and integrating out the matter degrees of freedom (see e.g. Refs. [21–23]). The VEV of the renormalized energy–momentum tensor is then obtained by taking the functional derivative $(2/\sqrt{-g})\delta/\delta g_{\mu\nu}$ of the effective action. One can perform the calculation of the effective action using regularizations, such as dimensional regularization or point-splitting, that preserve general covariance explicitly. The effective

² This conclusion also fits nicely with that of Ref. [19] where the authors considered the QFT of a large number \mathcal{N} of fields in Minkowski space, and found that the vacuum fluctuations collapse to black holes on scales smaller than $\mathcal{O}(\mathcal{N}^{1/4}l_{\text{pl}})$. They conclude that Minkowski space would therefore be unstable to black hole formation unless either the length-scale where quantum gravity sets in, for a theory with \mathcal{N} fields, is of order $\mathcal{N}^{1/4}l_{\text{pl}}$, or the vacuum fluctuations in Minkowski space do not gravitate.

³ Observe that general covariance holds, for renormalized quantities, independently of the regularization scheme employed, as it should. If one uses a scheme that breaks general covariance at the level of regularization, such as a momentum space cutoff, then to recover it at the level of renormalized quantity one must also add non-covariant counterterms. See [20] for an extended discussion of this point, and comparison with the literature.

action is therefore explicitly generally covariant, and the VEV of the energy–momentum tensor derived from it is automatically covariantly conserved. In this case, all divergences in $\langle 0|T_{\mu\nu}|0\rangle$ can be reabsorbed into generally covariant counterterms in the effective action. In particular, the divergence $\propto H^2(t)\Lambda_c^2$ is cured by a counterterm proportional to the Einstein–Hilbert action. In fact, taking the variation of the Einstein–Hilbert term gives the Einstein tensor $G_{\mu\nu}$, and G_{00} , specialized to the FRW metric, is proportional to $H^2(t)$. This shows that a VEV $\langle 0|T_{00}|0\rangle \propto H^2(t)$ is obtained, in the effective action language, by an additional term proportional to $\int d^4x \sqrt{-g}R$, and therefore is reabsorbed into a renormalization of Newton’s constant, as it is well known [5,6].

Such an effective action approach, however, assumes that we can integrate out all matter fields, i.e. that they are sufficiently massive with respect to the scale of interest. In our cosmological context this means that we are implicitly assuming that all fields have a mass m bigger than the Hubble parameter $H(t)$ at the time of interest, which is the quantity that fixes the relevant scale. Equivalently, we are assuming that the wavelength $1/m$ is smaller than the horizon size $H^{-1}(t)$. It is however interesting to consider the case in which in the spectrum there is a scalar particle X with a mass $m_X < H_0$. Such an ultra-light scalar field, with the mass protected against radiative correction by demanding that it is a pseudo Nambu–Goldstone boson, provides in fact a typical realization of quintessence [24]. In this case we cannot integrate out this field, and the low-energy effective action necessarily depends both on the metric and on this scalar field. Then the above derivation giving $\nabla^\mu \langle 0|T_{\mu\nu}|0\rangle = 0$ no longer goes through. By taking the variation of this action with respect to the metric we get the total energy–momentum tensor, including both $\langle 0|T_{\mu\nu}|0\rangle$ and the energy–momentum tensor $T_{\mu\nu}^X$ of this scalar field, and general covariance now only implies $\nabla^\mu (\langle 0|T_{\mu\nu}|0\rangle + T_{\mu\nu}^X) = 0$.

Whether these two terms are separately conserved is now a dynamical question, and depends on whether there is an interaction among them. One can imagine mechanisms by which vacuum fluctuations can exchange energy with other forms of matter. Typical examples are the amplification of vacuum fluctuations [25, 26], or the change in a large-scale scalar field due to the continuous horizon-crossing of small-scale quantum fluctuations of the same scalar field, which is also at the basis of stochastic inflation [27]. If $\nabla^\mu \langle 0|T_{\mu\nu}|0\rangle = -\nabla^\mu T_{\mu\nu}^X \neq 0$, it is no longer possible to reabsorb the effect of $\langle 0|T_{\mu\nu}|0\rangle$ into a renormalization of the Einstein–Hilbert term, nor of any other generally covariant local operator in the effective action. In fact, taking the functional derivative $(2/\sqrt{-g})\delta/\delta g_{\mu\nu}$ of a generally covariant term, we necessarily obtain a covariantly conserved tensor, so we can never obtain a quantity $\langle 0|T_{\mu\nu}|0\rangle$ that satisfies $\nabla^\mu \langle 0|T_{\mu\nu}|0\rangle \neq 0$.

4. Cosmology with zero-point fluctuations

We now explore the cosmological consequences of the hypothesis that $\langle 0|T_{\mu\nu}|0\rangle$ and $T_{\mu\nu}$ are conserved in conjunction, as in Eq. (5), but not separately. In this case, as shown above, the term $\Lambda_c^2 H^2$ cannot be reabsorbed into a renormalization of G , and we rather expect that it will give a genuine contribution to the total energy density $\rho_Z(t) = \mathcal{O}(H^2(t)M^2)$, where M is the UV scale where new physics sets in (so that M could be typically given by the Planck mass M_{pl} , or by the string mass). Recalling that the critical density is $\rho_c(t) = 3H^2(t)M_{\text{pl}}^2/(8\pi)$ we see that, for M of order M_{pl} , $\rho_Z(t)$ is of order of the critical density $\rho_c(t)$ at any time t . Thus we write

$$\rho_Z(t) = \Omega_Z \rho_c(t) = \Omega_Z \rho_0 H^2(t)/H_0^2, \quad (6)$$

where ρ_0 is the present value of the critical density. The value of Ω_Z is fixed by the renormalization procedure to the observed

value, so we keep it as a free parameter. The same is true for the equation of state (EOS) parameter w_Z defined by $p_Z = w_Z \rho_Z$, that can in principle be a function of time (see [20] for an extended discussion of this point).

The late time acceleration of the Universe cannot be explained only by a DE density that scales like $H^2(t)$ [7,28,29], basically because observations tell us that the total DE density is at least approximately constant in the recent cosmological epoch. Therefore we assume $\rho_Z(t)$ only to provide a part of the total DE density that we write as $\rho_{DE}(t) = \rho_X(t) + \rho_Z(t)$. Here $\rho_X(t)$ is a second dynamical DE component that is the one dominating the energy budget at the current epoch. In this approach the physical origin of ρ_X and ρ_Z can a priori be completely different. For instance, ρ_X could be due to a scalar field, as in quintessence models. In particular, we will take ρ_X to be due to an ultra-light scalar field with $m < H_0$ as discussed in Section 3. We define the EOS parameter of the X-component by $w_X = p_X/\rho_X$. Since ρ_X and ρ_Z have different origins, w_X and w_Z can in principle be different and we will see that only for $w_X \neq w_Z$ a tracking mechanism for DE emerges.

Mechanisms such as the amplification of vacuum fluctuations can produce an interaction between ρ_Z and scalar fields, but do not lead to interactions with photons nor massless fermions, that are not amplified in the FRW space-time because of conformal invariance. As we saw in Section 3, massive particles with $m > H_0$ can be integrated out in the effective action and therefore cannot contribute to the violation of the condition $\nabla^\mu \langle 0|T_{\mu\nu}|0\rangle = 0$. Thus, in our context it is natural to consider an interaction between ρ_Z and ρ_X , while non-relativistic matter and radiation are conserved in isolation, so they scale in the standard way, $\rho_M \sim a^{-3}$ and $\rho_R \sim a^{-4}$. In contrast, ρ_Z and ρ_X satisfy the coupled conservation equation

$$\dot{\rho}_Z + \dot{\rho}_X + 3(1 + w_Z)H\rho_Z + 3(1 + w_X)H\rho_X = 0. \quad (7)$$

In terms of the total DE density, $\rho_{DE}(t) = \rho_X(t) + \rho_Z(t)$, it reads

$$\dot{\rho}_{DE} + 3(1 + w_X)H\rho_{DE} = 3(w_X - w_Z)H\rho_Z. \quad (8)$$

We insert $\rho_Z(t) = \Omega_Z \rho_0 H^2/H_0^2$ on the r.h.s. and use the Friedmann equation, that now reads $H^2(t)/H_0^2 = \Omega_R a^{-4} + \Omega_M a^{-3} + \rho_{DE}(t)/\rho_0$. With the definitions

$$w_0 \equiv w_X + \epsilon, \quad \epsilon \equiv \Omega_Z(w_Z - w_X), \quad (9)$$

and using $d/dt = aH(d/da)$, we obtain

$$\left[a \frac{d}{da} + 3(1 + w_0) \right] \frac{\rho_{DE}}{\rho_0} = -3\epsilon \left[\frac{\Omega_R}{a^4} + \frac{\Omega_M}{a^3} \right]. \quad (10)$$

Note that the background evolution of the total DE density is fully determined by the two parameters w_0 and ϵ which can in principle be functions of time. We assume these functions to be constant for the scope of this work, i.e. we assume w_Z and w_X to be constant. A more general analysis will be presented in [20]. Then the evolution can be solved analytically,

$$\frac{\rho_{DE}}{\rho_0} = \frac{C}{a^{3(1+w_0)}} + \epsilon \left[\frac{1}{\frac{1}{3} - w_0} \frac{\Omega_R}{a^4} - \frac{1}{w_0} \frac{\Omega_M}{a^3} \right], \quad (11)$$

where C is the integration constant that is fixed by the condition $\rho_{DE}(a=1)/\rho_0 = \Omega_{DE} = 1 - (\Omega_R + \Omega_M)$. Observe that the model reduces to w CDM when $w_X = w_Z$ and the deviation from w CDM only depends on ϵ , so we are dealing with a one-parameter extension of w CDM, that we will call w ZCDM.

A very interesting feature of (11) is that ρ_{DE} always scales as the dominant energy component, for $\epsilon \neq 0$. In Fig. 1 we show the ratio $\rho_{DE}(a)/\rho_{tot}(a)$, where $\rho_{tot}(a) = \rho_{DE}(a) + \rho_0 \Omega_M a^{-3} + \rho_0 \Omega_R a^{-4}$. Deep into the RD phase, as well as in the MD phase,




Fig. 1. The energy fraction $\rho_{DE}(a)/\rho_{tot}(a)$, for $w_0 = -1$ and $\epsilon = 0.1$ (solid line) and $\epsilon = -0.1$ (dashed).




Fig. 2. The equation of state parameter $w_{DE}(a)$ for the same models as in Fig. 1.

$\rho_{DE}(a)/\rho_{tot}(a) = \mathcal{O}(\epsilon)$, while today it becomes $\mathcal{O}(1)$. Compared to Λ CDM, where the ratio ρ_Λ/ρ_{tot} is of order one today, but goes to zero as a^3 during MD and as a^4 during RD, the coincidence problem is sensibly alleviated.

We cannot claim that w ZCDM entirely solves the coincidence problem; in this model, in fact, the transition between the regime where $\rho_{DE}/\rho_{tot} = \mathcal{O}(\epsilon)$ to the regime where $\rho_{DE}/\rho_{tot} = \mathcal{O}(1)$ takes place at the present epoch simply because we have fixed the integration constant C in (11) by the requirement that $\rho_{DE}(a=1)/\rho_0 = \Omega_{DE}$. Nevertheless the coincidence problem is certainly alleviated, compared to Λ CDM, where ρ_{DE} is parametrically different from ρ_{tot} , and the ratio ρ_{DE}/ρ_{tot} evolves from $\mathcal{O}(10^{-120})$ at a Planck time to ~ 0.7 today.

It is also interesting to note that our model provides a different theoretical justification for parameterizations of early DE models that have been proposed in the literature. To make the relation explicit, it is useful to define $w_{DE} = p_{DE}/\rho_{DE}$ and derive it from the total DE continuity equation, $\dot{\rho}_{DE} + 3H(\rho_{DE} + p_{DE}) = 0$. Inserting the explicit solution for $\rho_{DE}(t)$ found in (11) we get

$$w_{DE}(t) = w_0 + \epsilon \left[\frac{\rho_M(t) + \rho_R(t)}{\rho_{DE}(t)} \right]. \quad (12)$$

This function is shown in Fig. 2, setting $w_0 = -1$ for definiteness. For $\epsilon > 0$ it evolves smoothly from a value $w_{DE} \simeq 1/3$ during RD, to $w_{DE} \simeq 0$ during MD and finally goes asymptotically to $w_{DE} \simeq w_0$. This EOS is quite similar to that obtained in a commonly used parameterization of early dark energy [30,31]. Note that, for $\epsilon < 0$, w_{DE} goes through infinity, as a consequence of the fact that ρ_{DE} goes through zero with $\dot{\rho}_{DE} \neq 0$, see Fig. 1, but the pressure $w_{DE}\rho_{DE}$ stays finite, and the background evolution is regular.

Our model has some similarities, as well as important differences, with other DE models studied in the literature. In particular,




Fig. 3. Left: The marginalized posterior probabilities of w_0 (upper panel) and ϵ (lower panel). Right: 1σ and 2σ mean-likelihood contours in the (ϵ, w_0) plane, after marginalization over all other parameters. Red is SNe only, blue is CMB only, and black is CMB + SNe + BAO. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this Letter.)

in [32–35] a model was proposed where the DE density has the form $\rho_\Lambda(t) = n_0 + n_1 H^2(t)$, inspired by the idea that the cosmological constant could run under renormalization group. In this case, however, the two components n_0 and $n_1 H^2(t)$ necessarily have the same EOS parameter. More closely related is the Λ XCDM model proposed in [36] in which DE has two components, an energy density $\rho_\Lambda = n_0 + n_1 H^2(t)$ associated to a running of the cosmological constant interacting with an unspecified dynamical “cosmon” field, although in our case the interaction is rather between ρ_χ (that, for $w_\chi \simeq -1$, plays basically the role of n_0) and $\rho_Z \sim H^2$, which has a different EOS.

We have performed a detailed comparison of our model with current observations of CMB, SNe Ia and BAO using modified versions of CAMB and CosmoMC [37,38], treating perturbations in the DE by modeling it as a perfect fluid (without anisotropic stress) with the EOS parameter w_{DE} given in (12) and a unit rest-frame sound speed. Full details will be reported in [20]. In Fig. 3 we give a sample of our results. The plots on the left are the one-dimensional posterior probabilities marginalized over all parameters except w_0 or ϵ , respectively, while the plot on the right shows the two-dimensional posterior probability marginalized over all parameters except the pair (ϵ, w_0) . In particular, we find the marginalized limits $-1.25 < w_0 < -0.908$ and $-0.0201 < \epsilon < 0.0460$ at 95% C.L., consistent with Λ CDM. The means of the marginalized posteriors are at $\langle w_0 \rangle = -1.07$ and $\langle \epsilon \rangle = 0.0104$ and the standard deviations are $\sigma_{w_0} = 0.0873$ and $\sigma_\epsilon = 0.0167$, respectively. Thus, our w ZCDM model is consistent with current data, and its deviations from w CDM, expressed by the parameter ϵ , are constrained at the level $\mathcal{O}(10^{-2})$. Future data will be able to set more stringent limits or to detect a non-vanishing value of ϵ .

Acknowledgements

We thank R. Brustein, S. Foffa, M. Frommert, R.K. Jain, M. Kunz, M. Shaposhnikov, R. Rattazzi and A. Riotto for very useful discussions. This work is supported by the Fonds National Suisse.

References

- [1] S. Weinberg, *Rev. Mod. Phys.* 61 (1989) 1.
- [2] P.J.E. Peebles, B. Ratra, *Rev. Mod. Phys.* 75 (2003) 559.
- [3] T. Padmanabhan, *Phys. Rep.* 380 (2003) 235.
- [4] E.J. Copeland, M. Sami, S. Tsujikawa, *Int. J. Mod. Phys. D* 15 (2006) 1753.
- [5] S.A. Fulling, L. Parker, *Ann. Phys.* 87 (1974) 176.
- [6] N. Birrell, P. Davies, *Quantum Fields in Curved Space*, Cambridge University Press, 1982, 340 pp.
- [7] M. Maggiore, *Phys. Rev. D* 83 (2011) 063514.
- [8] R.L. Arnowitt, S. Deser, C.W. Misner, in: L. Witten (Ed.), *Gravitation: An Introduction to Current Research*, John Wiley & Sons, New York, London, 1962, pp. 227–265.
- [9] E. Poisson, *A Relativist's Toolkit. The Mathematics of Black-Hole Mechanics*, Cambridge University Press, 2004.
- [10] G.W. Gibbons, S.W. Hawking, *Phys. Rev. D* 15 (1977) 2752.
- [11] J.D. Brown, J.W. York Jr., *Phys. Rev. D* 47 (1993) 1407.
- [12] J.D. Brown, J. Creighton, R.B. Mann, *Phys. Rev. D* 50 (1994) 6394.
- [13] S.W. Hawking, G.T. Horowitz, *Class. Quantum Grav.* 13 (1996) 1487.
- [14] V. Balasubramanian, P. Kraus, *Commun. Math. Phys.* 208 (1999) 413.
- [15] P. Kraus, E. Larsen, R. Siebelink, *Nucl. Phys. B* 563 (1999) 259.
- [16] S.R. Lau, *Phys. Rev. D* 60 (1999) 104034.
- [17] R.B. Mann, *Phys. Rev. D* 60 (1999) 104047.
- [18] R. Emparan, C.V. Johnson, R.C. Myers, *Phys. Rev. D* 60 (1999) 104001.
- [19] R. Brustein, D. Eichler, S. Foffa, D.H. Oaknin, *Phys. Rev. D* 65 (2002) 105013.
- [20] L. Hollenstein, M. Jaccard, M. Maggiore, E. Mitsou, in preparation (2011).
- [21] A.O. Barvinsky, G.A. Vilkovisky, *Phys. Rep.* 119 (1985) 1.
- [22] I.L. Buchbinder, S.D. Odintsov, I.L. Shapiro, *Effective Action in Quantum Gravity*, Institute of Physics, Bristol, UK, 1992.
- [23] I.L. Shapiro, *Class. Quantum Grav.* 25 (2008) 103001.
- [24] J.A. Frieman, C.T. Hill, A. Stebbins, I. Waga, *Phys. Rev. Lett.* 75 (1995) 2077.
- [25] L.P. Grishchuk, *Sov. Phys. JETP* 40 (1975) 409.
- [26] A.A. Starobinsky, *JETP Lett.* 30 (1979) 682.
- [27] A.A. Starobinsky, in: H.J. De Vega, N. Sanchez (Eds.), *Field Theory, Quantum Gravity and Strings*, Springer-Verlag, 1986, pp. 107–126.
- [28] S. Basilakos, M. Plionis, J. Sola, *Phys. Rev. D* 80 (2009) 083511.
- [29] J. Grande, J. Sola, S. Basilakos, M. Plionis, arXiv:1103.4632 [astro-ph.CO], 2011.
- [30] M. Doran, G. Robbers, *JCAP* 0606 (2006) 026.
- [31] L. Hollenstein, D. Sapone, R. Crittenden, B.M. Schaefer, *JCAP* 0904 (2009) 012.

- [32] I.L. Shapiro, J. Sola, JHEP 0202 (2002) 006.
- [33] I.L. Shapiro, J. Sola, C. Espana-Bonet, P. Ruiz-Lapuente, Phys. Lett. B 574 (2003) 149.
- [34] C. Espana-Bonet, P. Ruiz-Lapuente, I.L. Shapiro, J. Sola, JCAP 0402 (2004) 006.
- [35] I.L. Shapiro, J. Sola, Phys. Lett. B 682 (2009) 105.
- [36] J. Grande, J. Sola, H. Stefancic, JCAP 0608 (2006) 011.
- [37] A. Lewis, A. Challinor, A. Lasenby, Astrophys. J. 538 (2000) 473.
- [38] A. Lewis, S. Bridle, Phys. Rev. D 66 (2002) 103511.