

## Report

# The Spinster Homolog, Two of Hearts, Is Required for Sphingosine 1-Phosphate Signaling in Zebrafish

Nick Osborne,<sup>1,7</sup> Koroboshka Brand-Arzamendi,<sup>1</sup>  
Elke A. Ober,<sup>1,3</sup> Suk-Won Jin,<sup>1,4</sup> Heather Verkade,<sup>1,5</sup>  
Nathalia Glickman Holtzman,<sup>2,6</sup> Deborah Yelon,<sup>2</sup>  
and Didier Y.R. Stainier<sup>1,\*</sup>

<sup>1</sup>Department of Biochemistry and Biophysics  
Programs in Developmental Biology, Genetics and Human  
Genetics

Cardiovascular Research Institute  
University of California, San Francisco  
1550 Fourth Street  
San Francisco, CA 94143  
USA

<sup>2</sup>Developmental Genetics Program and Department of Cell  
Biology

Kimmel Center for Biology and Medicine  
Skirball Institute of Biomolecular Medicine  
New York University School of Medicine  
New York, NY 10016  
USA

<sup>3</sup>National Institute for Medical Research  
Division of Developmental Biology  
The Ridgeway, Mill Hill  
London NW71AA  
UK

<sup>4</sup>Department of Cell and Molecular Physiology and the  
Carolina Cardiovascular Biology Center  
University of North Carolina  
Chapel Hill, NC 27599  
USA

<sup>5</sup>School of Biological Sciences  
Monash University  
Clayton, VIC 3800  
Australia

<sup>6</sup>Department of Biology  
Queens College, CUNY  
65-30 Kissena Boulevard  
Flushing, NY 11367  
USA

## Summary

The bioactive lipid sphingosine 1-phosphate (S1P) and its G protein-coupled receptors play critical roles in cardiovascular, immunological, and neural development and function [1–6]. Despite its importance, many questions remain about S1P signaling, including how S1P, which is synthesized intracellularly, is released from cells. Mutations in the zebrafish gene encoding the S1P receptor Miles Apart (Mil)/S1P<sub>2</sub> disrupt the formation of the primitive heart tube [5]. We find that mutations of another zebrafish locus, *two of hearts* (*toh*), cause phenotypes that are morphologically indistinguishable from those seen in *mil/s1p2* mutants. Positional

cloning of *toh* reveals that it encodes a member of the Spinster-like family of putative transmembrane transporters. The biological functions of these proteins are poorly understood, although phenotypes of the *Drosophila spinster* and zebrafish *not really started* mutants suggest that these proteins may play a role in lipid trafficking [7, 8]. Through gain- and loss-of-function analyses, we show that *toh* is required for signaling by S1P<sub>2</sub>. Further evidence indicates that Toh is involved in the trafficking or cellular release of S1P.

## Results and Discussion

The lysophospholipid sphingosine 1-phosphate (S1P) has emerged as a key cellular signaling molecule. Many of the relevant signaling properties of S1P are mediated via its interaction with a family of G protein-coupled receptors (GPCRs). The interaction of these receptors with S1P is known to affect numerous processes, including the development and function of the vertebrate cardiovascular system [1, 3–5]. In addition, S1P has been identified as a mediator of immunological function [6]. The S1P analog FTY720 has recently been shown to function as a potent immunological modulator, promising to improve the treatment of organ-transplant recipients [9], as well as those suffering from pathological immunological responses following infection [10]. Despite the importance of S1P signaling, little is known about the processes that make this lipid available to bind its receptors.

Previously, we have shown that the *miles apart* (*mil*) gene, which is required for the formation of the primitive heart tube in zebrafish, encodes the ortholog of the mammalian receptor S1P<sub>2</sub>, initially called Edg5 and also known as S1PR2 [5, 11]. Screening mutagenized lines of zebrafish, we have identified another recessive mutation, *two of hearts* (*toh*), which causes phenotypes indistinguishable from those caused by *mil/s1p2* mutations.

At 36 hr postfertilization (hpf), wild-type (WT) embryos have a functioning heart (Figure 1A), whereas *toh* (Figure 1D) and *mil/s1p2* (Figure 1G) mutants exhibit pericardial edema (arrows), indicating circulatory defects. Circulatory failure and consequential pericardial edema in *toh* and *mil/s1p2* mutants is, in fact, the result of a defect in early heart-tube formation. In zebrafish, as in all vertebrates, the primitive heart tube is formed from bilateral groups of anterior mesodermal cells. These two cell populations migrate to the embryonic midline and fuse to form a single heart tube [12]. In WT zebrafish embryos at 19 hpf, the ring-shaped primitive heart tube has formed (Figure 1B). At 19 hpf, the myocardial cells of *toh* (Figure 1E) and *mil/s1p2* (Figure 1H) mutants have not migrated to the midline, resulting in the formation of bilateral heart-like structures, a phenotype called cardia bifida. Despite the cardia bifida, differentiation of the myocardial cells in *toh* mutants appears unaffected, because the bifid heart structures have WT-like chamber-specific gene expression and are infiltrated by endocardial cells (data not shown). However, the bifid heart structures in *toh* and *mil/s1p2* mutants are not appropriately connected to the vasculature and thus cannot support circulation.

\*Correspondence: didier\_stainier@biochem.ucsf.edu

<sup>7</sup>Present address: Department of Genetics, University of North Carolina, Chapel Hill, NC 27599, USA




Figure 1. *two of hearts* and *miles apart* Mutant Phenotypes

Comparison of WT (A–C), *toh* mutant (D–F), and *mill/s1p2* mutant (G–I) embryos.

(A, D, and G) Lateral brightfield images, anterior to the left, at 36 hpf show pericardial edema (arrow) and epidermal blisters (insets, arrowheads) in the tails of *toh*<sup>sk12</sup> (D) and *mill/s1p2*<sup>mn93</sup> (G) mutants.

(B, E and H) Examination of *cmlc2* expression at 19 hpf shows heart-ring formation in WT embryos (B) and a failure in precardiic-mesoderm migration in *toh*<sup>s420</sup> (E) and *mill/s1p2*<sup>mn93</sup> (H) mutant embryos. Dorsal views with anterior upwards.

(C, F and I) Visualization of the anterior endoderm by *Tg(-0.7her5:EGFP)*<sup>ne2067</sup> expression at 18 hpf. In embryos injected with *toh* (F) and *mill/s1p2* (I) MOs, numerous gaps (arrowheads) appear in the endodermal sheet, which is also irregularly shaped. The most anterior region of *mill/s1p2* morphants lacks GFP<sup>+</sup> endodermal cells at the midline (asterisk). Dorsal views with anterior upwards.

In addition to cardiac defects, *toh* and *mill/s1p2* mutants also display blistering in the tip of the tail, first evident around 26 hpf (Figures 1D and 1G, respectively; arrowheads). This tail-blister phenotype is not shared with other cardia bifida mutants in zebrafish [12], suggesting that *toh* and *mill/s1p2* may function in the same pathway. Because *mill/s1p2* mutant embryos have also been shown to have defects in the morphogenesis of the anterior endoderm [5], we analyzed the endoderm in *toh* mutants. The anterior endoderm of WT embryos, visualized at 18 hpf by the expression of the *-0.7her5:EGFP* transgene [13], forms a contiguous sheet across the embryonic midline (Figure 1C). Embryos lacking *toh* function display holes in their anterior endodermal sheet (Figure 1F), similar to *mill/s1p2* mutants (Figure 1I). We also observed these endodermal morphogenesis defects in *toh* and *mill/s1p2* mutants by examining the expression of the endodermal markers *foxa1* and *foxa2* (data not shown). Given that the endoderm is required for precardiic mesoderm migration [14–17], the cardia bifida phenotype seen in *toh* and *mill/s1p2* mutants is probably a result of these endodermal defects.

Because of the similarities between the *toh* and *mill/s1p2* mutant phenotypes, we hypothesized that Toh was involved in signaling by Mii/S1P<sub>2</sub>. In order to test this hypothesis, we isolated the *toh* gene (Figure 2A). Positional cloning of *toh* (Figure 2A) is described in detail in the Experimental Procedures section. *toh* encodes a 12-pass transmembrane-domain protein of the major facilitator superfamily (MFS) of non-ATP-dependent transporters (Figure 2B). The protein is a predicted 504-amino-acid member of the Spinster-like family of proteins (Figure S1, available online). This family is named after the *Drosophila spinster* gene, also called *benchwarmer* [18, 19]. The lesions found in the *spinster-like* gene in the *toh* mutant alleles are described in Figure S2.

The identification of the *spinster-like* gene as the *toh* locus was further verified by loss- and gain-of-function analyses. First, injection of a morpholino antisense oligonucleotide (MO) blocking *toh* mRNA splicing at the boundary between exon 4 and intron 4 resulted in the phenotypes seen in *toh* mutants (Figure 2C). Second, mRNA encoding the putative Toh protein rescued migration of the precardiic mesoderm when injected into maternal-zygotic *toh*<sup>ss</sup> mutants (*MZtoh*<sup>ss</sup>;

generation of these embryos is described in the Experimental Procedures section) and zygotic *toh*<sup>s420</sup> mutants (data not shown), leading to functional hearts. Overexpression of the *toh* mRNA had no effect on WT development and did not rescue *mill/s1p2* mutants. These loss- and gain-of-function experiments, together with tight genetic linkage between the cardiac phenotype and the *spinl2* locus and the presence of molecular lesions in the *spinl2* gene, show that we have isolated the *toh* gene. Interestingly, precardiic-mesoderm migration in *toh* mutants cannot be rescued by overexpression of *Drosophila spinster* (*spin*) or either of the two additional *spinster-like* genes found in the zebrafish genome (Figure S1), *not really started* (*nrs*) [20] and a gene that we here name *spinster-like 3* (*spinl3*). These data indicate that the Spinster-like proteins have acquired divergent functions.

The *toh* gene is expressed dynamically during early development (Figure 3 and Figure S3). *toh* mRNA is maternally provided (Figure S3A) and appears to be distributed ubiquitously during cleavage stages (Figure S3B). At the onset of gastrulation, the expression of *toh* changes. Cells that have undergone involution appear to express heightened levels of *toh* (Figure S3C, arrowhead). In addition, transcripts can be seen in the yolk syncytial layer, an extraembryonic tissue (Figures 3A and 3B, arrowhead). At the conclusion of gastrulation, *toh* is expressed strongly in tissues adjacent to the yolk cell and remains evident in the yolk syncytial layer (YSL) (Figures 3C and 3D, arrowhead). This expression continues through early somitogenesis. However, as somitogenesis proceeds, expression domains of *toh* become evident in the somitic mesoderm and in the endoderm adjacent to the yolk extension. By 24 hpf, *toh* is strongly expressed in a distinct compartment of the somites (Figures S3D and S3E, arrowhead), in the endoderm (Figure S3D), and in the heart (Figures S3F and S3G).

Because *mill/s1p2* and *toh* mutants have clear defects in the morphogenesis of the anterior endoderm and because both genes are expressed in the developing endoderm, we hypothesized that they might function cell-autonomously in the anterior endoderm to regulate precardiic-mesoderm migration. To test this hypothesis, we performed endoderm transplantation [17] in *mill/s1p2* and *toh* MO-injected embryos (morphants). This technique allows one to populate the endoderm of




Figure 2. Isolation of the *two of hearts* Gene

(A) Positional cloning of *toh*. Direction of the chromosomal walk is indicated with black arrows above the marker names. Markers used for mapping are indicated above the line representing the genomic region. The numbers of recombination events out of 2394 meioses found at each marker are indicated below the genomic region. A magnification of the critical region depicts portions of six open-reading frames (open arrows) in the identified genetic interval, including the previously cloned locus *liebeskummer/reptin*. The critical region contains two *spinster-like* genes, one of which was identified as *toh* (gray filled arrow) and the other of which we here name *spinl3*.

(B) Schematic diagram of Toh proteins produced from each allele. Arrows point to the site affected in the mutant alleles. Transmembrane domains are indicated in blue, and the predicted translated intronic sequence in the *s8* allele is indicated in red. Numbers also identify the individual transmembrane domains.

(C) Injection of a splice blocking MO against *toh* into WT embryos phenocopies the *toh* mutations, leading to pericardial edema (arrow) and blistering in the tail (arrowhead).

injection of equivalent amounts of *nrs* mRNA into the YSL never rescued precardiac-mesoderm migration. Altogether, these data indicate that Toh function in the YSL is necessary and sufficient for precardiac-mesoderm migration.

Little is known about the mechanism by which Spinster-like proteins carry out their function. However, the common phenotypes of *toh* and *mil/s1p2* mutants suggest that *toh* and *mil/s1p2* function in the same pathway. To test this hypothesis, we injected suboptimal amounts of *mil/s1p2* and *toh* MOs alone and in combination. When injected singly, 0.4 ng of *mil/s1p2* MO and 0.2 ng of *toh* MO caused cardia bifida in 2.5% (n = 40) and 2.4% (n = 85) of the embryos, respectively. However, when embryos were coinjected with 0.4 ng of *mil/s1p2* and 0.2 ng of *toh* MO, 85.7% (n = 70) of the embryos exhibited cardia bifida. Therefore, partial loss of function of both Mil/S1P<sub>2</sub> and Toh

morphants with WT cells while leaving other tissues untouched. In four cases, we were able to rescue the *mil/s1p2* mutant heart phenotype (data not shown). In these cases, the majority of the anterior endoderm had been replaced with WT cells. In cases of partial replacement of the anterior endoderm or replacement of only the posterior endoderm, no rescue was observed. However, in the case of *toh* morphants, rescue was never observed, regardless of the level of endoderm replacement. Altogether, these data indicate that whereas *mil/s1p2* functions cell-autonomously in the endoderm to regulate precardiac-mesoderm migration, *toh* functions in another cell type.

*toh* is clearly expressed in the YSL (Figures 3A–3D), a tissue previously implicated in precardiac-mesoderm migration in zebrafish [21]. In order to test whether the YSL expression of *toh* regulates precardiac-mesoderm migration, we injected *toh* MO into the YSL. Injection of *toh* MO (Figure 3F), but not of mock carrier solution (Figure 3E), into the YSL resulted in cardia bifida at a high frequency (84%, n = 56). We also injected 200 pg of *toh* mRNA into the YSL of *toh* morphants. YSL injection of *toh* mRNA was able to rescue precardiac-mesoderm migration in some *toh* morphants (38%, n = 32). However,

caused cardia bifida much more frequently than did the partial loss of either protein alone. We also observed that *mil/s1p2*; *toh* double mutants have no additional phenotypes compared to *mil* or *toh* single mutants. Together, these data suggest that Mil/S1P<sub>2</sub> and Toh may function in a common genetic pathway.

In order to test directly whether *toh* is required for Mil/S1P<sub>2</sub> signaling, we overexpressed mRNA encoding Mil/S1P<sub>2</sub> in WT and *toh* mutant embryos. Injection of 100 pg of *mil/s1p2* mRNA at the one-cell stage caused severe morphological defects in WT embryos (89%, n = 208; Figure 4A), including cyclopia (asterisk) and disorganized body-axis formation. Defects caused by *mil/s1p2* overexpression can be traced to problems with cell movements during gastrulation; embryos overexpressing *mil/s1p2* exhibited defects in convergence, extension, and epiboly movements (Figure 4D). It is likely that the gastrulation defects seen in *mil/s1p2*-overexpressing embryos are due to the antagonism between *mil/s1p2* and *silberblick/wnt11* signaling [22], given that mutations in *slb/wnt11* cause defects in convergence and extension movements similar in nature to those seen in *mil/s1p2*-overexpressing embryos [23, 24]. If Toh function is required for signaling by Mil/S1P<sub>2</sub>, one would expect that loss of Toh function would




Figure 3. *Toh* Function in the YSL Is Required for Precardiac-Mesoderm Migration

(A–D) *toh* expression at 6 (A and B) and 8 (C and D) hpf. (A) Animal-pole view with dorsal to the right, 6 hpf, showing *toh* expression around the margin and diffusely throughout the YSL. (B) Magnified view of the box in (A) showing *toh* expression around the YSL nuclei (arrowhead). (C) Lateral view with dorsal to the right, 8 hpf, showing continued *toh* expression in cells that have involuted, as well as in the YSL. (D) Magnified view of the box in (C) showing pronounced *toh* expression in the YSL (arrowhead).

(E and F) Dorsal images of *Tg(cmlc2:GFP)*<sup>fl</sup> embryos injected into the YSL with mock solution (E) or with 4 ng *toh* MO (F) and visualized at 30 hpf. Mock-injected embryos have a single heart tube (E, arrow), whereas embryos with loss of *Toh* function in the YSL very frequently (84%, n = 56) display cardia bifida (F; arrows).


suppress the phenotypes caused by *mil/s1p2* overexpression. Indeed, *toh* mutants or morphants overexpressing *mil/s1p2* only rarely showed morphological defects similar to those seen in control *mil/s1p2*-overexpressing embryos, as assessed at both 30 hpf (7%, n = 195; Figure 4B) and 8 hpf (0%, n = 40; Figure 4E). Therefore, *Toh* function is required for Mil/S1P<sub>2</sub> signaling in this overexpression assay. Note that these *mil/s1p2*-overexpressing embryos still, however, display the phenotypes seen in *toh* mutants or morphants, including cardia bifida and blister formation in the tail, indicating that *mil/s1p2* mRNA is incapable of rescuing loss of *toh* function.

One manner by which *Toh* might affect the function of the Mil/S1P<sub>2</sub> receptor is by regulating the availability of the receptor's ligand, S1P. To determine whether the phenotypes of Mil/S1P<sub>2</sub> overexpression are dependent on receptor-ligand interaction, we generated a mutant form of Mil/S1P<sub>2</sub>. The E129A mutation in Mil is analogous to mutations that have been shown, for other S1P receptors, to block the receptor's ability

to interact with S1P without affecting the receptor's stability or ability to interact with downstream signaling components [25]. Embryos injected with 100 or 200 pg of *milE129A* mRNA gastrulated normally and displayed none of the phenotypes observed in embryos injected with 100 pg of WT *mil/s1p2* mRNA. Thus, it appears that the effects seen in Mil/S1P<sub>2</sub>-overexpressing embryos require an interaction between the exogenous receptor and S1P.

Because the yolk cell contains nutrients as well as some developmental signals necessary for embryonic development [26], we hypothesized that *Toh* in the YSL was required for S1P availability from the yolk cell to the embryo. One prediction of this hypothesis is that the Mil/S1P<sub>2</sub> receptor should be capable of responding to exogenously applied S1P even in the absence of *Toh* function. To test this prediction, we injected embryos with *mil/s1p2* mRNA alone or in combination with *toh* MO. At the onset of gastrulation (~5 hpf), the embryos were also injected at the animal pole with a mock carrier solution or a carrier solution containing 10 μM S1P. Embryos injected with S1P alone showed no effect (Figure 4F). However, embryos injected with *mil/s1p2* mRNA showed immediate morphogenetic effects from the application of exogenous S1P, regardless of whether or not they had intact *Toh* expression (Figures 4H and 4J). These effects were S1P-dependent, because *mil/s1p2*-overexpressing embryos injected with the carrier solution alone did not show this response (Figures 4G and 4I). The presence or absence of *Toh* function had no effect on the severity of a *mil/s1p2*-overexpressing embryo's response to exogenously supplied S1P. In both groups of overexpressing embryos, gastrulation movements paused and the cells of the embryo retreated back to the animal pole, causing a thickening of the embryo (Figures 4H and 4J, asterisks). These data suggest that the reason that *toh* mutants or morphants do not exhibit any defects upon *mil/s1p2* overexpression is due to a lack of interaction between overexpressed Mil/S1P<sub>2</sub> and endogenous S1P.

The above results and the fact that *Toh* is a member of a superfamily of transporters raise the possibility that *Toh* functions as a transporter of S1P. However, *Drosophila* embryos with mutations in the *toh* homolog *spin* display a dramatic expansion of the acidified compartment of the cell, as visualized by LysoTracker staining [7], accompanied by inappropriate accumulations of lipids and sugars in those cellular compartments [19, 27]. Therefore, *toh* mutations might be affecting S1P release by affecting the storage and trafficking of many lipids, not just S1P or its precursors. We examined the levels of LysoTracker staining in *toh* morphants and *MZtoh*<sup>se</sup> mutants, and they appeared equivalent to those of WT embryos (data not shown), indicating normal lysosomal structure in *toh* mutants. We also examined the early and recycling endocytic compartments of WT and *toh* MO-injected embryos to determine whether the earlier stages of the endocytic pathway might be disrupted in *toh* mutants. The architecture of the early and recycling endosomal compartments appeared indistinguishable between WT and *toh* MO-injected embryos when visualized with a yellow fluorescent protein (YFP)-tagged Rab5c protein [28] (data not shown). Therefore, in contrast to *Drosophila spin* mutants and zebrafish *nrs* heterozygotes [7, 8], *toh* mutants do not appear to exhibit gross defects in lipid or carbohydrate trafficking. These findings, together with the failure of *Drosophila spin* and zebrafish *nrs* genes to rescue *toh* mutants, indicate that vertebrate *spinster-like* genes have diverged in function and that *Toh* may have a specific role in the trafficking or release of S1P.



**Figure 4. Exogenous S1P Substitutes for Toh Function in Mil/S1P<sub>2</sub>-Overexpressing Embryos** (A and B) Lateral views with anterior to the left, of embryos overexpressing *mil/s1p2* at 30 hpf. WT embryos injected with 100 pg of *mil/s1p2* RNA (A) display a shortened body axis and cyclopia (asterisk). *MZtoh*<sup>sh</sup> mutants injected with 100 pg of *mil/s1p2* RNA have no body-axis defects (B) but develop pericardial edema (arrow) and tail blisters (arrowheads).

(C–E) Visualization of axial mesoderm and endoderm by expression of *foxa2* at late gastrula stages, dorsal views with anterior upwards. WT embryos overexpressing *mil/s1p2* (D) have broadened axial mesoderm (solid line) compared to uninjected embryos (C). Progression of epiboly (dashed line) is also impaired in embryos overexpressing *mil/s1p2* as compared to uninjected siblings. *toh* MO-injected embryos overexpressing *mil/s1p2* (E) are indistinguishable from uninjected siblings (C).

(F–J) Lateral views of 5.5 hpf embryos injected with mock carrier solution or 10 µM S1P. Control embryos show no response to exogenous S1P (F). Embryos overexpressing Mil/S1P<sub>2</sub> do not show more severe phenotypes when injected with mock carrier solution at 5 hpf (G). Mil/S1P<sub>2</sub>-overexpressing embryos injected with a 10 µM S1P solution at 5 hpf exhibit severe phenotypes (H), including an exaggerated thickened layer of cells at the animal pole (asterisk). Mil/S1P<sub>2</sub>-overexpressing embryos coinjected with *toh* MO show no phenotypes when injected with mock solution (I) and resemble controls (F). When injected with a 10 µM S1P solution, Mil/S1P<sub>2</sub>-overexpressing embryos lacking Toh function show severe defects (J), including a thickened layer of cells at the animal pole (asterisk).

absence of Mtx1 would lead to an absence of Toh function in the YSL and, therefore, a lack of S1P release from the yolk. Lack of S1P release would, in turn, prevent Mil/S1P<sub>2</sub> signaling and

In this study, we identify Toh as a novel component of signaling via the zebrafish S1P<sub>2</sub> ortholog, Mil. The combination of shared phenotypes between *toh* and *mil/s1p2* mutants, the fact that *toh* loss of function suppresses the deleterious effects of *mil/s1p2* overexpression, and the synergistic effects of suboptimal *mil/s1p2* and *toh* MO injections suggest that *toh* is a critical component of the Mil/S1P<sub>2</sub> signaling pathway. Furthermore, we have shown that the Mil/S1P<sub>2</sub> receptor is capable of signaling in the absence of Toh function when exogenous S1P is applied. Therefore, Toh appears to be a novel contributor to the biosynthesis, trafficking, or release of S1P.

Previous analysis has shown that the YSL is critical for precardiac-mesoderm migration [21]. This study suggested that the transcription-factor gene *mtx1*, which is expressed exclusively in the YSL, regulates the deposition at the embryonic midline of Fibronectin (FN), an extracellular-matrix component critical for precardiac-mesoderm migration in mice [29] and zebrafish [30]. Interestingly, the deposition of FN in *mil/s1p2* morphants is also deficient (Figure S4). Furthermore, injection of FN into the midline of *mil/s1p2*-deficient embryos appears to rescue precardiac-mesoderm migration [31]. Thus, *mtx1* might regulate *toh* expression or function in the YSL. The

the downstream deposition of FN required for endoderm and precardiac-mesoderm morphogenesis.

The exact biochemical function of Toh and other Spinster-like proteins remains unclear. The homology of these proteins to small solute transporters suggests the interesting possibility that Toh is involved in the trafficking or actual cellular release of S1P and that *toh* mutations affect Mil/S1P<sub>2</sub> signaling by limiting available S1P. Our data showing that overexpressed Mil/S1P<sub>2</sub> is capable of responding to exogenously supplied S1P strongly supports the idea that the defect observed in animals lacking a functional Toh/Spinl2 transporter is due to a reduction in endogenous ligand production or release. Thus, Toh is a strong candidate for a transmembrane transporter that, as previously claimed for ABCC1 [32], may be capable of moving S1P across cellular membranes to make it available for receptor-ligand interactions.

## Conclusions

As the relevance of S1P signaling to both basic and clinical sciences becomes more evident, it is critical that the signaling partners of S1P receptors be identified. S1P<sub>2</sub> signaling in

mammalian systems has been shown to play a vital role in activation of mast cells [33], a cell type thought to contribute to the pathogenesis of asthma [34]. In addition, S1P<sub>2</sub>-receptor function is known to affect vascular tone [35] and may contribute to the protective effects that S1P demonstrates during ischemic challenge to the heart during myocardial infarction [36]. The identification of Toh as a component of S1P<sub>2</sub> signaling in zebrafish suggests that the mammalian orthologs of *toh* may have similar functions in S1P-mediated signaling. Therefore, Toh and its orthologs may represent new targets for the manipulation of specific S1P signaling pathways in both normal and pathological states.

## Experimental Procedures

### Zebrafish Strains and Care

Adult and embryonic zebrafish were raised and cared for with the use of standard laboratory procedures [37]. We used the following zebrafish mutant and transgenic strains: *toh<sup>ss</sup>*, *toh<sup>s220</sup>*, *toh<sup>sk12</sup>*, *toh<sup>s420</sup>*, *mil<sup>m93</sup>*, *Tg(-0.7her5:egfp)<sup>ne2067</sup>* [13], and *Tg(cmlc2:EGFP)<sup>1</sup>* [38]. Maternal-zygotic *s8* (*MZtoh<sup>ss</sup>*) mutant embryos were generated via the mating of heterozygous *toh<sup>ss</sup>* fish. Escaping homozygous *toh<sup>ss</sup>* embryos were then raised to adulthood.

### Immunohistochemistry, Fluorescence Microscopy, and Confocal Analysis

Embryos were fixed at room temperature for 1 hr in 4% Paraformaldehyde or overnight at 4°C in 2% Paraformaldehyde in PBS. LysoTracker DND-99 (Molecular Probes) was diluted 1:200 in 1/10 Hanks Basic Salt Solution, and staining was carried out on live embryos for 1 hr at 28°C, after which embryos were washed three times, for 5 min each, with 1/10 Hanks Basic Salt Solution, then fixed and processed as described above. Images were acquired with a Zeiss LSM5 Pascal confocal microscope. Whole-mount fluorescence microscopy was performed with a Zeiss SteREO Lumar.V12 microscope.

### Endoderm Transplantation

Endodermal cell transplantation was carried out as described by Stafford et al. [17], with 100 pg *cas/sox32* mRNA used to force donor cells into the endodermal lineage. Host embryos were injected with 2 ng *mil/s1p2* MO or 1 ng *toh* MO. These host embryos were also coinjected with 1 ng *cas/sox32* MO for depletion of their endoderm.

### In Situ Hybridization

Whole-mount in situ hybridization was carried out as described by Alexander et al. [39], with the following probes used: *cmlc2* [40], *foxa2* [41], and *toh*. The *toh* probe was generated with the use of primers 5'-TTG GAGCCATCAC ATGTGTGA-3' and 5'-TACTTGTGGCGCTTTGT-3' for PCR of a 516 bp fragment of *toh*. The primers were engineered with T3 and T7 promoters, respectively, allowing the generation of sense and antisense probes directly from the PCR product.

### RNA Overexpression and Morpholino-Oligonucleotide Generation

All capped mRNA used for injection was generated with mMessage mMachine kits (Ambion). *mil* mRNA was generated as described by Kupperman et al. [5]. pCS2+ *mil E129A* was generated with the QuikChange II Mutagenesis kit (Stratagene). mRNA encoding *toh* was generated by linearization of pCS2+ *toh* with NotI and transcription with SP6 polymerase. Zebrafish *spin13* mRNA was generated by subcloning of the coding region of *spin13* into pCS2+, linearization of this construct with NotI, and transcription with SP6 polymerase. *Drosophila spinster* mRNA was generated by subcloning of *spinster-RFP* from pUAS *spinster-RFP* (gift of Sean Sweeney and Graeme Davis) into pCS2+, linearization with NotI, and transcription with SP6 polymerase. Zebrafish *nrs* was generated from pCS2+ *nrs* [20]. *rab5c-YFP* mRNA was generated from pCS2+ *Rab5c-YFP* (gift of C.-P. Heisenberg) as described by Ulrich et al. [28], and embryos were injected with 100 pg of mRNA. The *toh* MO (5'-GCAGCTTACCCTCAGTGCCAGT-3') was designed by Gene Tools. YSL injections of 4 ng of *toh* MO were carried out as described by Amack et al. [42].

### Exogenous S1P Application

S1P (Sigma Aldrich catalog no. S9666) was dissolved in 100% methanol at 1 mg/mL. This stock solution was further diluted to 10 μM concentration into a carrier solution of 250 mM potassium chloride containing 0.5% w/v Fatty Acid Free Bovine Serum Albumin (Calbiochem/EMD catalog no. 126575). Embryos were injected with either 100 pg *mil/s1p2* mRNA or *mil/s1p2* mRNA plus 1 ng *toh* MO at the one-cell stage. Controls were not injected at this stage. At 5 hpf, the embryos were then injected into the animal pole, among the cells of the embryos, with 2.3 nL of 10 μM S1P solution or the carrier solution. They were aged for 30 min at 28°C and then imaged.

### Cloning of the *toh* Locus

Via bulk-segregant analysis, the *toh* locus was localized to zebrafish chromosome 5. We used 1197 diploid *s220 toh* mutant embryos to narrow the affected locus to a 295 kb region between the CA repeat marker z9419 and a single-nucleotide polymorphism (SNP) in a homolog of the mammalian *ankhzn* gene. This SNP was designated *ankdCAP4* and was amplified with primers designed by the dCAPS 2.0 web-based program and detected with Hinfl cutting [43]. Zebrafish CA-repeat microsatellite primers were obtained from the Massachusetts General Hospital MGH/CVRC Zebrafish Server website (<http://zebrafish.mgh.harvard.edu>). The 295 kb region is covered entirely by three bacterial artificial chromosomes (BACs): CHORI-211 134D21, CHORI-211 138A6, and DanioKey 7B17. These BACs have been sequenced and assembled by the Sanger Centre *Danio rerio* Sequencing Project. Full-length sequences are available at <ftp://ftp.sanger.ac.uk/pub/sequences/zebrafish>. Six putative open-reading frames (ORFs) were found between z9419 and *ankdCAP4* via a combination of the GENSCAN exon-prediction software [44] and an analysis of conservation of synteny between zebrafish, mouse, and human genomic sequences. These ORFs were sequenced from cDNA and genomic DNA in WT and mutant samples.

### Supplemental Data

Supplemental Data include four figures and can be found with this paper online at [http://www.current-biology.com/supplemental/S0960-9822\(08\)01431-0](http://www.current-biology.com/supplemental/S0960-9822(08)01431-0).

### Acknowledgments

We thank Holly Field and Jonathan Alexander for assistance with initial studies of *toh*; Courtney Babbitt for assistance with phylogenetic analysis; Atsuo Kawahara for discussions and sharing of unpublished data; and Jason Cyster, Ian Scott, and Courtney Griffin for helpful comments on the manuscript. This work was supported in part by the National Science Foundation (N.O.), the American Heart Association (E.A.O., S-W.J., N.G.H.), the Human Frontier Science Program (H.V.), the National Institutes of Health (D.Y., D.Y.R.S.), and the Packard foundation (D.Y.R.S.).

Received: April 18, 2006

Revised: October 19, 2008

Accepted: October 21, 2008

Published online: December 4, 2008

### References

1. Liu, Y., Wada, R., Yamashita, T., Mi, Y., Deng, C.X., Hobson, J.P., Rosenfeldt, H.M., Nava, V.E., Chae, S.S., Lee, M.J., et al. (2000). Edg-1, the G protein-coupled receptor for sphingosine-1-phosphate, is essential for vascular maturation. *J. Clin. Invest.* 106, 951–961.
2. Chun, J. (2005). Lysophospholipids in the nervous system. *Prostaglandins Other Lipid Mediat.* 77, 46–51.
3. Sanna, M.G., Liao, J., Jo, E., Alfonso, C., Ahn, M.Y., Peterson, M.S., Webb, B., Lefebvre, S., Chun, J., Gray, N., and Rosen, H. (2004). Sphingosine 1-phosphate (S1P) receptor subtypes S1P1 and S1P3, respectively, regulate lymphocyte recirculation and heart rate. *J. Biol. Chem.* 279, 13839–13848.
4. Wendler, C.C., and Rivkees, S.A. (2006). Sphingosine-1-phosphate inhibits cell migration and endothelial to mesenchymal cell transformation during cardiac development. *Dev. Biol.* 291, 264–277.
5. Kupperman, E., An, S., Osborne, N., Waldron, S., and Stainier, D.Y. (2000). A sphingosine-1-phosphate receptor regulates cell migration during vertebrate heart development. *Nature* 406, 192–195.
6. Chiba, K., Yanagawa, Y., Masubuchi, Y., Kataoka, H., Kawaguchi, T., Ohtsuki, M., and Hoshino, Y. (1998). FTY720, a novel immunosuppressant,

- induces sequestration of circulating mature lymphocytes by acceleration of lymphocyte homing in rats. I. FTY720 selectively decreases the number of circulating mature lymphocytes by acceleration of lymphocyte homing. *J. Immunol.* **160**, 5037–5044.
7. Sweeney, S.T., and Davis, G.W. (2002). Unrestricted synaptic growth in spinster—a late endosomal protein implicated in TGF- $\beta$ -mediated synaptic growth regulation. *Neuron* **36**, 403–416.
  8. Kishi, S., Bayliss, P.E., Uchiyama, J., Koshimizu, E., Qi, J., Nanjappa, P., Imamura, S., Islam, A., Neuberger, D., Amsterdam, A., and Roberts, T.M. (2008). The identification of zebrafish mutants showing alterations in senescence-associated biomarkers. *PLoS Genet* **4**, e1000152.
  9. Chun, J., and Rosen, H. (2006). Lysophospholipid receptors as potential drug targets in tissue transplantation and autoimmune diseases. *Curr. Pharm. Des.* **12**, 161–171.
  10. Premenko-Lanier, M., Moseley, N.B., Pruett, S.T., Romagnoli, P.A., and Altman, J.D. (2008). Transient FTY720 treatment promotes immune-mediated clearance of a chronic viral infection. *Nature* **454**, 894–898.
  11. Osborne, N., and Stainier, D.Y. (2003). Lipid receptors in cardiovascular development. *Annu. Rev. Physiol.* **65**, 23–43.
  12. Stainier, D.Y. (2001). Zebrafish genetics and vertebrate heart formation. *Nat. Rev. Genet.* **2**, 39–48.
  13. Tallafuss, A., and Bally-Cuif, L. (2003). Tracing of her5 progeny in zebrafish transgenics reveals the dynamics of midbrain-hindbrain neurogenesis and maintenance. *Development* **130**, 4307–4323.
  14. Kikuchi, Y., Trinh, L.A., Reiter, J.F., Alexander, J., Yelon, D., and Stainier, D.Y. (2000). The zebrafish bonnie and clyde gene encodes a Mix family homeodomain protein that regulates the generation of endodermal precursors. *Genes Dev.* **14**, 1279–1289.
  15. Reiter, J.F., Alexander, J., Rodaway, A., Yelon, D., Patient, R., Holder, N., and Stainier, D.Y. (1999). Gata5 is required for the development of the heart and endoderm in zebrafish. *Genes Dev.* **13**, 2983–2995.
  16. Alexander, J., Rothenberg, M., Henry, G.L., and Stainier, D.Y. (1999). casanova plays an early and essential role in endoderm formation in zebrafish. *Dev. Biol.* **215**, 343–357.
  17. Stafford, D., White, R.J., Kinkel, M.D., Linville, A., Schilling, T.F., and Prince, V.E. (2006). Retinoids signal directly to zebrafish endoderm to specify insulin-expressing beta-cells. *Development* **143**, 949–956.
  18. Yamamoto, D., and Nakano, Y. (1999). Sexual behavior mutants revisited: molecular and cellular basis of *Drosophila* mating. *Cell. Mol. Life Sci.* **56**, 634–646.
  19. Dermaut, B., Norga, K.K., Kania, A., Verstreken, P., Pan, H., Zhou, Y., Callaerts, P., and Bellen, H.J. (2005). Aberrant lysosomal carbohydrate storage accompanies endocytic defects and neurodegeneration in *Drosophila* benchwarmer. *J. Cell Biol.* **170**, 127–139.
  20. Young, R.M., Marty, S., Nakano, Y., Wang, H., Yamamoto, D., Lin, S., and Allende, M.L. (2002). Zebrafish yolk-specific not really started (nrs) gene is a vertebrate homolog of the *Drosophila* spinster gene and is essential for embryogenesis. *Dev. Dyn.* **223**, 298–305.
  21. Sakaguchi, T., Kikuchi, Y., Kuroiwa, A., Takeda, H., and Stainier, D.Y. (2006). The yolk syncytial layer regulates myocardial migration by influencing extracellular matrix assembly in zebrafish. *Development* **133**, 4063–4072.
  22. Kai, M., Heisenberg, C.P., and Tada, M. (2008). Sphingosine-1-phosphate receptors regulate individual cell behaviours underlying the directed migration of prechordal plate progenitor cells during zebrafish gastrulation. *Development* **135**, 3043–3051.
  23. Heisenberg, C.P., and Nusslein-Volhard, C. (1997). The function of silberblick in the positioning of the eye anlage in the zebrafish embryo. *Dev. Biol.* **184**, 85–94.
  24. Heisenberg, C.P., Tada, M., Rauch, G.J., Saude, L., Concha, M.L., Geisler, R., Stemple, D.L., Smith, J.C., and Wilson, S.W. (2000). Silberblick/Wnt11 mediates convergent extension movements during zebrafish gastrulation. *Nature* **405**, 76–81.
  25. Parrill, A.L., Wang, D., Bautista, D.L., Van Brocklyn, J.R., Lorincz, Z., Fischer, D.J., Baker, D.L., Liliom, K., Spiegel, S., and Tigyi, G. (2000). Identification of Edg1 receptor residues that recognize sphingosine 1-phosphate. *J. Biol. Chem.* **275**, 39379–39384.
  26. Chen, S., and Kimelman, D. (2000). The role of the yolk syncytial layer in germ layer patterning in zebrafish. *Development* **127**, 4681–4689.
  27. Nakano, Y., Fujitani, K., Kurihara, J., Ragan, J., Usui-Aoki, K., Shimoda, L., Lukacsovich, T., Suzuki, K., Sezaki, M., Sano, Y., et al. (2001). Mutations in the novel membrane protein spinster interfere with programmed cell death and cause neural degeneration in *Drosophila melanogaster*. *Mol. Cell. Biol.* **21**, 3775–3788.
  28. Ulrich, F., Krieg, M., Schotz, E.M., Link, V., Castanon, I., Schnabel, V., Taubenberger, A., Mueller, D., Puech, P.H., and Heisenberg, C.P. (2005). Wnt11 functions in gastrulation by controlling cell cohesion through Rab5c and E-cadherin. *Dev. Cell* **9**, 555–564.
  29. George, E.L., Georges-Labouesse, E.N., Patel-King, R.S., Rayburn, H., and Hynes, R.O. (1993). Defects in mesoderm, neural tube and vascular development in mouse embryos lacking fibronectin. *Development* **119**, 1079–1091.
  30. Trinh, L.A., and Stainier, D.Y. (2004). Fibronectin regulates epithelial organization during myocardial migration in zebrafish. *Dev. Cell* **6**, 371–382.
  31. Matsui, T., Raya, A., Callol-Massot, C., Kawakami, Y., Oishi, I., Rodriguez-Esteban, C., and Belmonte, J.C. (2007). miles-apart-Mediated regulation of cell-fibronectin interaction and myocardial migration in zebrafish. *Nat. Clin. Pract. Cardiovasc. Med.* **4** (Suppl 1), S77–S82.
  32. Mitra, P., Oskeritzian, C.A., Payne, S.G., Beaven, M.A., Milstien, S., and Spiegel, S. (2006). Role of ABCG1 in export of sphingosine-1-phosphate from mast cells. *Proc. Natl. Acad. Sci. USA* **103**, 16394–16399.
  33. Jolly, P.S., Bektas, M., Olivera, A., Gonzalez-Espinosa, C., Proia, R.L., Rivera, J., Milstien, S., and Spiegel, S. (2004). Transactivation of sphingosine-1-phosphate receptors by Fc $\epsilon$ RI triggering is required for normal mast cell degranulation and chemotaxis. *J. Exp. Med.* **199**, 959–970.
  34. Chiappara, G., Gagliardo, R., Siena, A., Bonsignore, M.R., Bousquet, J., Bonsignore, G., and Vignola, A.M. (2001). Airway remodelling in the pathogenesis of asthma. *Curr. Opin. Allergy Clin. Immunol.* **1**, 85–93.
  35. Ohmori, T., Yatomi, Y., Osada, M., Kazama, F., Takafuta, T., Ikeda, H., and Ozaki, Y. (2003). Sphingosine 1-phosphate induces contraction of coronary artery smooth muscle cells via S1P2. *Cardiovasc. Res.* **58**, 170–177.
  36. Karliner, J.S. (2004). Mechanisms of cardioprotection by lysophospholipids. *J. Cell. Biochem.* **92**, 1095–1103.
  37. Westerfield, M. (2000). *The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish (Danio rerio)* (Eugene, OR: University of Oregon Press).
  38. Huang, C.J., Tu, C.T., Hsiao, C.D., Hsieh, F.J., and Tsai, H.J. (2003). Germ-line transmission of a myocardium-specific GFP transgene reveals critical regulatory elements in the cardiac myosin light chain 2 promoter of zebrafish. *Dev. Dyn.* **228**, 30–40.
  39. Alexander, J., Stainier, D.Y., and Yelon, D. (1998). Screening mosaic F1 females for mutations affecting zebrafish heart induction and patterning. *Dev. Genet.* **22**, 288–299.
  40. Yelon, D., Horne, S.A., and Stainier, D.Y. (1999). Restricted expression of cardiac myosin genes reveals regulated aspects of heart tube assembly in zebrafish. *Dev. Biol.* **214**, 23–37.
  41. Odenthal, J., and Nusslein-Volhard, C. (1998). fork head domain genes in zebrafish. *Dev. Genes Evol.* **208**, 245–258.
  42. Amack, J.D., Wang, X., and Yost, H.J. (2007). Two T-box genes play independent and cooperative roles to regulate morphogenesis of ciliated Kupffer's vesicle in zebrafish. *Dev. Biol.* **310**, 196–210.
  43. Neff, M.M., Turk, E., and Kalishman, M. (2002). Web-based primer design for single nucleotide polymorphism analysis. *Trends Genet.* **18**, 613–615.
  44. Burge, C., and Karlin, S. (1997). Prediction of complete gene structures in human genomic DNA. *J. Mol. Biol.* **268**, 78–94.