

REVIEW

10.1111/j.1469-0691.2006.01512.x

Post-operative aspergillosis

A. C. Pasqualotto and D. W. Denning

School of Medicine, The University of Manchester and Wythenshawe Hospital, Manchester, UK

ABSTRACT

While invasive aspergillosis occurs typically in severely immunocompromised patients, cases of surgical site infection have been reported in immunocompetent individuals. The Medline, LILACS and EMBASE databases were searched for descriptions of cases of post-operative aspergillosis, and references from relevant articles and conference abstracts were reviewed. More than 500 cases of post-operative aspergillosis were found. Cardiac surgery ($n = 188$), ophthalmological surgery ($n > 90$) and dental surgery ($n > 100$) were associated with the majority of cases. Other cases involved wound infections ($n = 22$), bronchial infections ($n = 30$), mediastinitis ($n = 11$), pleural aspergillosis ($n = 1$), infections following orthopaedic surgery ($n = 42$), vascular prosthetic surgery ($n = 22$), breast surgery ($n = 5$), abdominal surgery ($n = 10$) and neurosurgery ($n = 25$). In most patients, the source was presumed to be airborne infection during the surgical procedure. Prevention of these infections requires special care of the ventilation system in the operating room. Successful treatment requires rapid diagnosis, surgical debridement and antifungal therapy, often with voriconazole. In order to improve the outcome, better diagnostic methods are needed, particularly for cases of endocarditis and aortitis.

Keywords *Aspergillus*, neurosurgical infection, nosocomial infection, post-operative infection, review, surgical site infection

Accepted: 22 December 2005

Clin Microbiol Infect 2006; 12: 1060–1076

INTRODUCTION

Fungi belonging to the genus *Aspergillus* are ubiquitous and occur in soil, water and decaying vegetation [1,2]. Like other filamentous fungi, *Aspergillus* is acquired primarily from an inanimate reservoir, usually by the inhalation of small airborne spores [3]. While nosocomial aspergillosis typically affects immunocompromised patients [4], post-operative aspergillosis occurs mainly in immunocompetent patients, including some patients who received corticosteroids temporarily. This review is based on a search of the world literature to identify cases of post-operative aspergillosis occurring after surgical procedures that were not secondary to disseminated infection or previous known colonisation by *Aspergillus* spp.

PATIENTS AND METHODS

Search strategy

The Medline, LILACS and EMBASE databases were searched and references from relevant articles were reviewed. Search terms were 'aspergillus', 'surgery', 'surgical site infections', 'cardiovascular surgical procedures', 'endocarditis', 'aortitis', 'mediastinitis', 'breast', 'laparotomy', 'arthroplasty', 'orthopaedic procedures', 'neurosurgery', 'transplantation', 'ophthalmologic surgical procedures', 'osteomyelitis', 'prosthesis', 'treatment' and 'prevention'. Conference abstracts were reviewed using the *Aspergillus* web-site (<http://www.aspergillus.man.ac.uk>). Apart from papers published in Japanese, Russian, Czech or Polish, no language restrictions were applied. Non-surgical cases of primary cutaneous aspergillosis and infections associated with intravascular devices were not considered for this review.

Definitions

Only cases of proven or probable infection were reviewed, which required that all cases included were laboratory-confirmed by culture and/or histology and/or microscopy. These criteria were based on standard definitions for invasive fungal infections in immunocompromised patients for clinical research [5]. As the incubation period of aspergillosis is not well-established, this review was not limited by the CDC definitions of surgical site infection (i.e., infections occurring within 30 days of a surgical procedure if no implant is left in

Corresponding author and reprint requests: D. W. Denning, Education and Research Centre, Wythenshawe Hospital, Southmoor Road, Manchester M23 9LT, UK
E-mail: ddenning@manchester.ac.uk

place, or within 1 year if an implant is present) [6]. Wound aspergillosis was defined as infections involving only the skin or subcutaneous tissue of the incision. Infections that involved both superficial and deep incision sites were classified as deep surgical site infections, according to the site affected. For classification as a wound infection, in addition to a positive culture of *Aspergillus* spp. taken directly from the wound, the wound itself had to be non-healing following treatment with standard antibiotics, with other pathogens absent or minimally present.

LITERATURE REVIEW

From the initial search, *c.* 450 papers were excluded for several reasons, including cases of aspergillosis unrelated to surgery and disseminated

infections from pulmonary or other sources. Other review articles with no new contributions were also excluded. More than 500 patients with post-operative aspergillosis were included in this review, including cases following heart surgery (188), neurosurgery (25), dental surgery (>100), ophthalmological surgery (>90), orthopaedic surgery (42), vascular prosthetic surgery (22), abdominal surgery (10) and breast surgery (5). Reports of wound infections (22), bronchial infections (30), mediastinitis (11) and pleural aspergillosis (1) were also included. Some cases were reported more than once, *i.e.*, [10] and [11]; [16] and [129]. Table 1 summarises the key aspects of

Table 1. Summary of main clinical presentations, treatment and prophylaxis of cases of post-operative aspergillosis

	Comments
Clinical presentation	
Endocarditis/aortitis (<i>n</i> = 174)	Marked male predominance (<i>c.</i> 70%) Absence of immunosuppression for almost all patients Aortic valve mostly involved (60.5%) Median time from surgery to diagnosis = 2.7 months <i>Aspergillus fumigatus</i> : aetiology in only 58.7% of cases Low ante-mortem diagnosis (43.5%), usually following surgical intervention Blood cultures rarely positive (6.4%) Propensity for late embolisation Mortality >80% despite intervention (overall mortality 92.7%)
Neurosurgery infections (<i>n</i> = 25)	Frequently associated with the use of steroids (52.0%) All proven cases caused by <i>A. fumigatus</i> Median time from surgery to diagnosis = 3 months Higher ante-mortem diagnosis rate (68.0%) than for endocarditis Mortality much lower than observed for immunocompromised patients with aspergillosis disseminated to the central nervous system
Dental surgery infections (<i>n</i> > 100)	Many cases of sinus aspergillosis associated with endodontic treatment Obturing pastes within the maxillary antrum probably a risk-factor
Wound infections (<i>n</i> = 22)	Most patients immunosuppressed Median time from surgery to diagnosis = 17 days <i>A. flavus</i> accounts for 41.2% of cases confirmed by culture Risk of dissemination
Ophthalmological surgery infections (<i>n</i> > 90)	Usually presents as keratitis Diagnosis requires sampling at the site of infection
Vascular prosthesis infections (<i>n</i> = 22)	Almost all patients are male and immunocompetent Median time from surgery to diagnosis = 8 months Clinical picture indistinguishable from <i>Staphylococcus epidermidis</i> infections Usually occurs close to suture lines of previous aortotomy High incidence of embolisation to vertebral disks
Other cases (<i>n</i> = 113)	Frequent association with prosthesis insertion
Treatment	The optimal treatment has not been specifically studied Survival seemingly dependent on excision of the infected tissue; in cases involving a prosthesis, the new prosthesis should be placed in a non-infected field Concomitant use of a systemic antifungal agent is also crucial. Although there are no data favouring any one antifungal drug, the response rates to voriconazole in immunocompromised patients were superior to those in response to amphotericin B. Caspofungin and other echinocandins may also be useful. Longer-term oral extension with itraconazole or voriconazole is appropriate for some patients The optimal duration of therapy is unknown and is likely to differ between patients, but once as much infected tissue as possible has been removed and any immunocompromising factors, such as corticosteroid therapy, are minimised, treatment for 3 months beyond the last evidence of active disease may be appropriate
Prevention	Requires special care with the ventilation system in the operating theatre Proper disinfection and storage of surgical materials

post-operative aspergillosis following different surgical procedures.

The first case reported in the literature

In 1933, a female aged 14 years underwent surgery for an abdominal tumour [7]. After 16 days, the dressing was removed and an ulcer was observed. There were no systemic manifestations of the infection, but *Aspergillus niger* grew in the surgical dressings, which were covered with a 'dark powder'. It was speculated that contamination may have occurred in the operating room, and the authors commented on two previous similar cases of wound aspergillosis in the same medical centre, each exhibiting similar pustulation confined to the area of the gauze dressing, and with rapid healing after local treatment with iodine and alcohol.

Endocarditis, aortitis and other presentations of aspergillosis following cardiac surgery

Table S1 (see Supplementary material) summarises some clinical details for 124 patients with *Aspergillus* endocarditis or aortitis following heart surgery [4,8–91]. The median age of these patients was 43.5 (range 0.8–71) years. Most of the surgical procedures involved the aortic (60.5%) or mitral (30.6%) valve, and no particular prosthesis was associated with the infection. Male patients predominated (69.9%), which may be related to the higher prevalence of coronary and aortic valve surgery in males than in females. The incubation period was variable, with a mean time from surgery to the diagnosis (often following autopsy) of 2.7 (range <1 to >12) months. Two cases [43,88] had possible incubation periods of several years; silent *Aspergillus* fungaemia with late endocarditis is possible in such cases, although there is no direct evidence for such a diagnosis. Forty additional cases were cited in other studies, but without detailed information [92–111], and at least ten additional cases were reported in languages other than English [112–120].

Notably, almost none of the patients were immunosuppressed, and there was no evidence of bronchopulmonary aspergillosis. The post-operative course of these patients was usually consistent with culture-negative endocarditis. Unusual presentations of aspergillosis following heart surgery included osteochondritis and/or

osteomyelitis after sternotomy [78,121–125], and saphenous vein thrombosis [126] or rupture [127] after coronary by-pass surgery.

Aspergillus fumigatus was the aetiological agent in only 58.7% of cases confirmed by culture. Other important species were *Aspergillus terreus* (12.5%), which is resistant to amphotericin B, *A. niger* (11.2%) and *Aspergillus flavus* (11.2%). Most reports associated the infection with contaminated grafts, contaminated sutures or intra-operative dispersion of spores. The prevention of such infections will be considered below.

The difficulty in establishing the diagnosis of *Aspergillus* endocarditis is well-known, since the fungus is isolated infrequently from blood cultures. Only eight cases of post-operative aspergillosis associated with a positive blood culture (6.4% of cases) were found in the literature [11,12,14,36,38,46,61,87]. Pre-autopsy diagnosis was made in only 43.5% of cases. In the vast majority of cases, surgical intervention was crucial in obtaining a definitive diagnosis, often following examination of either the valve (23.4%) or emboli to large peripheral arteries (16.9%). Unfortunately, some surgical specimens were not sent for culture. The diagnosis of *Aspergillus* aortitis is also made more difficult because of the localisation of the infection and the absence of endocarditis vegetation; this condition is therefore often undetected by both trans-thoracic and trans-oesophageal echocardiography. Computed tomography scan, magnetic resonance imaging and angiography might be more helpful for the diagnosis of such cases.

The overall mortality rate was 92.7%. In addition, 83.0% of patients who had an ante-mortem diagnosis died, and a high mortality rate (80.9%) existed even among patients treated with a surgical intervention in the affected valve. This might be related to a delayed diagnosis, which gives the fungus the opportunity to establish large vegetations, to cause extensive local necrosis, and eventually to embolise to large arteries. Many of the patients who underwent valve replacement died during surgery or a few days later. These results indicate an unequivocal need for better diagnostic methods for this condition. At the present time, the role of antibody tests and molecular tests of blood samples has not been properly investigated. In one report, a false-negative galactomannan result occurred for a

patient with non-surgical *Aspergillus* endocarditis, despite a positive blood culture for *A. fumigatus* [128].

Wound infections

For patients with post-operative wound infection caused by *Aspergillus* spp., the interval between surgery and infection is measured in days, rather than the months required for endocarditis (Table S2, see Supplementary material) [5,87,129–143]. In addition, many patients with wound aspergillosis were immunosuppressed, with half of the cases occurring in solid-organ transplant recipients. As this condition can progress to profound or disseminated infection, all patients need to be treated aggressively with combined medical and surgical therapy. *A. flavus* is a particularly important species in wound aspergillosis, accounting for 41.2% of cases confirmed by culture.

Chronic lung disease was the only independent risk-factor for *A. fumigatus* sternal wound infection after open-heart surgery in a case-control study [144]. *A. fumigatus* was grown simultaneously from the bronchial washings of one patient, suggesting that colonised patients may be at increased risk for these infections. In another study, a cluster of four cases of surgical and burn wound aspergillosis was associated with the outside packages of dressing supplies, which had become contaminated during hospital construction work [145]. This resulted in patients with large exposed vulnerable surface areas being inoculated directly with *Aspergillus* spores. When outbreaks of cutaneous aspergillosis occur, wound dressing and tape should be cultured [146,147]. Cases of *A. flavus* sternal wound infection coinciding with hospital renovation activities have also been reported [148].

Neurosurgery

Involvement of the central nervous system in aspergillosis may occur via haematogenous dissemination (usually from the lungs), or by direct extension from nasal or paranasal foci [149]. Table S3 (see Supplementary material) summarises the published literature concerning aspergillosis following neurosurgical procedures [150–168]. In contrast to *Aspergillus* endocarditis, male patients were not over-represented in neurosurgical infections (44.0%). Previous treatment with cortico-

steroids was mentioned in 52.0% of the reports, including patients receiving regimens of pulse therapy, high-dose regimens for periods of weeks to months, and even physiological steroid replacement therapy. The median time from surgery to diagnosis was 3 (range <1 to >12) months. All proven cases were caused by *A. fumigatus*. As shown in Table S3, different presentations of the infection can occur, including meningitis, central nervous system abscesses, mycotic aneurysms and cerebral infarction. Many cases were associated with *Aspergillus* contamination from paranasal sinuses, particularly following trans-sphenoidal surgery.

An ante-mortem diagnosis was performed in 68% of cases, which was a proportion considerably higher than that for cases of *Aspergillus* endocarditis. Most of these cases were diagnosed after abscess drainage (36%), and 20% of patients had a cerebrospinal fluid culture positive for *Aspergillus*. In some cases, a culture positive for *Aspergillus* was considered to be contamination, which was associated with delayed treatment, sometimes with a fatal outcome. All patients who survived received combined aggressive clinical and surgical therapy. The overall mortality rate was 68.0%, which is surprisingly low in comparison with mortality rates of >90% for immunosuppressed patients with disseminated aspergillosis involving the central nervous system. This may reflect an earlier diagnosis, facilitated by imaging techniques, in association with aggressive neurosurgical intervention. Furthermore, dexamethasone therapy could be withdrawn for many of these patients, unlike the situation for patients with more permanent immunosuppression.

Dental surgery

A number of studies have reported a connection between endodontic treatment and non-invasive sinus aspergillosis, mainly in the presence of obturating pastes within the maxillary antrum [169–183]. Experiments with fungal cultures have revealed accelerated growth of *Aspergillus* spp. in the presence of zinc oxide in the culture medium [182]. With respect to treatment, it is necessary to remove surgically the fungal masses in the sinuses, and especially all zinc-containing radiopaque material, and to ensure establishment of adequate sinus drainage and aeration [178,181].

Systemic antifungal agents are generally not recommended to complement the surgical procedure unless there are signs of local invasion [181].

Ophthalmological surgery

Ocular aspergillosis usually presents as keratitis, and rarely as endophthalmitis [92,184–187]. Post-operative infections have been described following penetrating keratoplasty [188–190], radial keratotomy [191–194], excimer laser photorefractive keratectomy [195], laser-assisted in-situ keratomileusis [196–199], pterygium excision [200,201], hydroxyapatite orbital implant surgery [202], cataract surgery [179,184,203–212], scleral buckling procedures [213,214], sutureless surgery [215,216] and trabeculectomy [204,217]. Sampling at the site of infection provides the best chance for obtaining a positive culture [196,218–220]. While some cases of *Aspergillus* endophthalmitis and keratitis have been associated with hospital construction [209,221], others have been linked to contaminated irrigating fluids used during surgery [208]. Many cases of *Aspergillus* keratitis and scleritis following non-surgical corneal trauma have also been reported [209,222–248], frequently in farmers. The efficacy of the antifungal therapy depends to a great extent on the ability of the compound used to penetrate the eye [219,249].

Orthopaedic surgery

A male aged 64 years underwent hip replacement surgery and was admitted after a 9-year interval because of progressive hip pain [250]. *A. niger* grew from aspirated hip fluid, and the patient was treated with amphotericin B and prosthesis replacement. Two cases of aspergillosis following laminectomy were also reported in the same publication. In another report, *A. fumigatus* spondylodiskitis occurred 2.5 months after lumbar disk surgery, and the patient was treated with itraconazole in combination with surgical debridement [251]. Other cases following lumbar surgery have also been reported [252–254].

In a prospective study of 658 consecutive total hip arthroplasty procedures in which cultures were taken at surgery, *Aspergillus* was recovered in 4.1% of cases, and *Aspergillus* was also cultivated from the air in the operating theatre [255]. In another report, a male aged 22 years developed *Aspergillus* osteomyelitis after a closed fracture of

the femur and subsequent surgery [256]. Another male, aged 16 years, developed infection with *A. flavus* following surgery for a fracture of the femur. The infection was associated with pressure and necrosis caused by casts [257]. The first case of aspergillosis complicating total knee arthroplasty was described in a male aged 80 years who was receiving chronic corticosteroid therapy [258]. This patient presented with a popliteal cyst 27 months after surgery, and was cured with a 12-week course of amphotericin B in combination with surgical debridement and prosthesis removal. Other similar cases have been described in the literature [259,260].

Bronchopulmonary surgery

Bronchial stump aspergillosis, i.e., *Aspergillus* infection of bronchial granulation tissue surrounding endobronchial sutures after pulmonary resection, is a rare clinical presentation of *Aspergillus* infection [261–267]. This condition is more common when silk thread is used as the suture material rather than stainless steel [268,269], and should be considered in the differential diagnosis of haemoptysis, coughing and dyspnoea occurring up to several years after lung surgery, although asymptomatic presentation is also possible [262–264,269]. *Aspergillus* suture granuloma may simulate recurrent neoplastic disease in patients undergoing resection for pulmonary cancer [270]. Although no comparative studies exist, removal of the silk suture is probably the key therapeutic intervention for this condition.

An association between the presence of saprophytic fungi in the bronchial anastomosis and the occurrence of serious airway complications following lung transplantation has been proposed [271]. The ischaemic and necrotic airway debris at the bronchial anastomosis provides a fertile environment for *Aspergillus* proliferation. Several reports have also described an increased risk of complications as a consequence of post-transplantation saprophytic *Aspergillus* disease [272–275]. The source of these infections has not been well-studied, but may be related to either colonisation of the diseased or transplanted lungs, or to direct perioperative acquisition.

Pleural aspergillosis has been reported as an iatrogenic complication of lung adenocarcinoma resection [276]. A 14-day course of intravenous caspofungin failed to sterilise the cavity, and

the patient was treated with a combination of intra-pleural and systemic amphotericin B lipid complex.

Infection of vascular prosthesis

Glotzbach [277] seems to have been the first to describe *Aspergillus* infection of a vascular prosthesis in 1982, and this was followed by the report of an immunocompetent patient, aged 73 years, with aortic pseudoaneurysm and contiguous vertebral osteomyelitis, who was treated with graft resection and amphotericin B [278]; as in the previous case [277], this patient presented with back pain *c.* 2 years after surgery. In a study of 14 cases of fungal graft infections [279], only four were caused by *Aspergillus* spp.

Aspergillus graft infections usually occur on the suture line of a previous aortotomy [280]. Definitive diagnostic procedures for these patients generally involve culture of the excised aortic graft or the peripheral embolus, and biopsy of the contiguously affected vertebral disk space [277,281–284]. As with endocarditis, almost all cases have involved immunocompetent males [277,278,281–290]. The median time from the placement of the graft to diagnosis is 8 months, which is a longer period than that for *Candida* graft infections (usually <6 weeks) [279]. Fever is absent in about one-half of the cases. One report associated in-situ replacements or treatment without excision of the infected graft with recurrence and death in all cases, whereas all patients who underwent an extra-anatomical by-pass through a clean field survived [279]. No difference seems to exist between the clinical presentations of fungal and bacterial vascular prosthesis infections, particularly when *Staphylococcus epidermidis* is involved [291].

Abdominal surgery

Multiple abdominal visceral infection by *A. fumigatus* occurred in a female aged 37 years after laparostomy for Crohn's disease [292]. The patient was treated successfully with a 4-week course of amphotericin B. Sampling of air from the patient's environment yielded one isolate of *A. fumigatus* that matched the patient's isolates, suggesting that this patient acquired the infection from air in the intensive care unit. Peritoneal aspergillosis has also been reported following laparotomy in non-trans-

plant patients [4,293,294], and *Aspergillus* cholangitis complicating portoenterostomy for a biliary atresia has also been reported [295]. A case of *A. flavus* renal pelvis infection, following a pyelolithotomy, has been described, with *Aspergillus* around the suture material [296]. Post-operative cases of wound infection [87,138,139,141–143,297], peritonitis [297,298] and renal allograft infection [299] have also been described following solid-organ transplantation. Although not always apparent, it is possible that some of these patients had occult foci from which the *Aspergillus* may have disseminated [299–325].

Mediastinitis

Aspergillus mediastinitis in surgical patients has been described in association with deep sternal wound infection [4,123,125,326], heart transplantation [326–329], aortitis [4,46] and patch infection after repaired tetralogy of Fallot [113]. Two immunocompetent patients were also affected during an outbreak of aspergillosis related to a contaminated air-handling system [4]. The importance of considering aspergillosis in the differential diagnosis of slowly progressive, but destructive, wound infections and culture-negative mediastinitis after cardiac surgery has been emphasised.

Breast surgery

Fungal infections of the soft tissue have been reported in association with prosthetic breast implants [330–333]. A healthy female, aged 39 years, with bilateral inflatable silicone breast implants was colonised by *A. niger* as a possible consequence of massive contamination during surgery [331]. *A. flavus* was also cultured from a saline-filled implant removed from a healthy young patient who developed unilateral breast inflammation 4 years after surgery [332]. It has been shown that fungi can grow and reproduce in a saline-only environment, and that their survival periods differ among species [334–336]. Colonisation of a tissue expander by *A. niger*, resulting in mechanical obstruction of the device, has also been described [337].

Linking the infection with the operating theatre

Many studies have associated the occurrence of post-operative aspergillosis with the dissemin-

ation of *Aspergillus* spores in the operating theatre. In one investigation, pigeon excreta and moss in the immediate vicinity of the ventilator intake port were found to harbour large numbers of *Aspergillus* spores [14]. Similarly, *Aspergillus* was isolated from all areas of the theatre suite in another study [38], except in a room fitted with laminar airflow. Air-conditioner cooling coils and pigeon droppings on the ledges outside the suite were found to harbour large numbers of *Aspergillus* spores.

Environmental studies conducted during an outbreak of *A. flavus* sternal infection [123] revealed massive fungal contamination in some areas of the surgical ward [338]. When isolates from patients and from the environment were typed, the same genotype was detected, implicating a single clonal source for the environmental contamination and intra-operative acquisition of *A. flavus*. Diaz-Guerra *et al.* [339] reported the simultaneous isolation of an *A. flavus* isolate from the aortic prosthesis of a heart surgery patient and two isolates from a dual-reservoir cooler-heater used in the operating theatre where this patient received surgery. Genetic typing of these isolates by random amplified polymorphic DNA analysis revealed identical genotypes, indicating the nosocomial origin of the strain. Similar results were reported in another investigation [51], and during an epidemiological investigation of cases of *Aspergillus* aortitis following cardiac surgery [46], high concentrations of *A. fumigatus* were found in the ventilation system and on the floor of the surgical suite.

Prevention

Most modern operating theatres have conventional ventilation and filters that remove airborne particles $\geq 5 \mu\text{m}$ [340]. However, these filters are not sufficient to remove *Aspergillus* spores (2.5–3.0 μm in size). Laminar airflow systems and HEPA filtration will remove airborne particles $\geq 0.3 \mu\text{m}$ with 99.97% efficiency. These systems are generally used in orthopaedic and other implant surgery [341], as well as in protective environments for high-risk patients [342–347]. Theoretically, these systems are more effective in preventing post-operative aspergillosis, but data demonstrating a survival benefit for patients are lacking [348,349]. Furthermore, these systems are limited by the expense of construction and maintenance, and by a lack

of consensus concerning the level of airborne conidia at which the risk can be numerically defined for aspergillosis [340].

Since outbreaks of post-operative aspergillosis have occurred occasionally in units with HEPA filters [78], periodic maintenance of ventilation systems seems to be warranted [38,346,350]. Some of these incidents were caused by contamination of insulation in variable airflow volume units that had deteriorated after becoming wet. As fungi can inhabit water distribution systems and may cause nosocomial infection, tap water should not be used on surgical wounds. However, larger controlled studies are needed to determine the role of water in the transmission of aspergillosis [2,3]. The patients may also be their own source of infection in some circumstances [159], as has been demonstrated for staphylococcal infections. Prevention of post-operative aspergillosis requires proper disinfection techniques, as well as appropriate storage of surgical materials. A single proven post-operative case of aspergillosis should be sufficient to initiate epidemiological investigations.

CONCLUSIONS

Post-operative aspergillosis seems to be an underappreciated problem, particularly when compared with other extra-pulmonary manifestations of invasive aspergillosis. Mortality is high for non-cutaneous infections. Several organs and surgical procedures may be involved, depending on the surgery performed. These infections are usually indolent, in some cases occurring several months after surgery. In most patients, the source of the spores seems to have originated from the air during surgery, but contamination from paranasal sinuses, bronchopulmonary lesions, haematogenous dissemination and contaminated grafts is also possible. In order to improve the outcome, better diagnostic methods are needed. Infected patients should receive aggressive combined medical and surgical therapy. Prevention of these infections should include special care of the ventilation system in the operating theatre and adequate sterilisation techniques.

ACKNOWLEDGEMENTS

This work was presented, in part, at the 2nd Trends in Medical Mycology meeting (Berlin, 2005).

SUPPLEMENTARY MATERIAL

The following supplementary material is available for this article online at <http://www.blackwell-synergy.com>:

Table S1. Case reports of post-operative endocarditis/aortitis caused by *Aspergillus* spp.

Table S2. Case reports of wound infections with *Aspergillus* spp. following surgical procedures.

Table S3. Case reports of central nervous system infections with *Aspergillus* spp. following neurosurgical procedures.

REFERENCES

- Walsh TJ, Dixon DM. Nosocomial aspergillosis: environmental microbiology, hospital epidemiology, diagnosis and treatment. *Eur J Epidemiol* 1989; **5**: 131–142.
- Anaissie EJ, Stratton SL, Dignani MC *et al.* Pathogenic *Aspergillus* species recovered from a hospital water system: a 3-year prospective study. *Clin Infect Dis* 2002; **34**: 780–789.
- Soubani AO, Chandrasekar PH. The clinical spectrum of pulmonary aspergillosis. *Chest* 2002; **121**: 1988–1999.
- Lutz BD, Jin J, Rinaldi MG, Wickes BL, Huycke MM. Outbreak of invasive *Aspergillus* infection in surgical patients, associated with a contaminated air-handling system. *Clin Infect Dis* 2003; **37**: 786–793.
- Ascioglu S, Rex JH, de Pauw B *et al.* Defining opportunistic invasive fungal infections in immunocompromised patients with cancer and hematopoietic stem cell transplants: an international consensus. *Clin Infect Dis* 2002; **34**: 7–14.
- Garner JS, Jarvis WR, Emori TG, Horan TC, Hughes JM. CDC definitions for nosocomial infections. In: Olmsted RN, ed., *APIC infection control and applied epidemiology: principles and practice*. St Louis: Mosby, 1996; A1–A20.
- Frank L, Alton OM. Aspergillosis: a case of post-operative skin infection. *JAMA* 1933; **100**: 2007–2008.
- Amoury RA, Bowman FO, Malm JR. Endocarditis associated with intracardiac prostheses. Diagnosis, management, and prophylaxis. *J Thorac Cardiovasc Surg* 1966; **51**: 36–48.
- Leffert RL, Hackett RL. *Aspergillus* aortitis following replacement of aortic valve. *J Thorac Cardiovasc Surg* 1967; **53**: 866–874.
- Jones T, Meshel L, Rubin IL. *Aspergillus* endocarditis superimposed on aortic valve prosthesis. *NY State J Med* 1969; **69**: 1923–1928.
- Hairston P, Lee WH. Mycotic (fungal) endocarditis after cardiovascular surgery. *Am Surg* 1969; **35**: 135–143.
- Hairston P, Lee WH. Management of infected prosthetic heart valves. *Ann Thorac Surg* 1970; **9**: 229–237.
- Aslam PA, Gourley R, Eastridge CE, Pate JW. *Aspergillus* endocarditis after aortic valve replacement. *Int Surg* 1970; **53**: 91–96.
- Gage AA, Dean DC, Schimert G, Minsley N. *Aspergillus* infection after cardiac surgery. *Arch Surg* 1970; **101**: 384–387.
- Chaudhuri MR. Fungal endocarditis after valve replacement. *J Thorac Cardiovasc Surg* 1970; **60**: 207–214.
- Ostermiller WE, Dye WS, Weinberg M. Fungal endocarditis following cardiovascular surgery. *J Thorac Cardiovasc Surg* 1971; **61**: 670–675.
- Lawrence T, Shockman AT, MacVaugh H. *Aspergillus* infection of prosthetic aortic valves. *Chest* 1971; **60**: 406–414.
- Carrizosa J, Levison ME, Lawrence T, Kaye D. Cure of *Aspergillus ustus* endocarditis on a prosthetic valve. *Arch Intern Med* 1974; **133**: 486–490.
- Doshi R. *Aspergillus fumigatus* endocarditis of an aortic homograft with aneurysm of the ascending aorta. *J Pathol* 1971; **103**: 263–265.
- Harthorne JW, Castleman B. Case records of the Massachusetts General Hospital. Weekly clinicopathological exercises. Case 24-1973. *N Engl J Med* 1973; **288**: 1290–1296.
- Dayal Y, Weindling HK, Price DL. Cerebral infarction due to fungal embolus. A complication of *Aspergillus* infection on an aortic valve prosthesis. *Neurology* 1974; **24**: 76–79.
- Foxworthy D, Gunnar R. Complications of aortic valve replacement. *Postgrad Med* 1974; **56**: 146–151.
- Rubinstein E, Noriega ER, Simberkoff MS, Holzman R, Rahal JJ. Fungal endocarditis: analysis of 24 cases and review of the literature. *Medicine (Baltimore)* 1975; **54**: 331–334.
- Uitley JR, Mills J, Roe BB. The role of valve replacement in the treatment of fungal endocarditis. *J Thorac Cardiovasc Surg* 1975; **69**: 255–258.
- Wheeler EO, Robboy SJ. Case records of the Massachusetts General Hospital. Weekly clinicopathological exercises. Case 39-1976. *N Engl J Med* 1976; **295**: 718–724.
- Petheram IS, Seal RM. *Aspergillus* prosthetic valve endocarditis. *Thorax* 1976; **31**: 380–390.
- Irvin RG, Sade RM. Endocarditis and musculoskeletal manifestations. *Ann Intern Med* 1978; **88**: 578.
- Drexler L, Rytel M, Keelan M, Bonchek LI, Olinger GN. *Aspergillus terreus* infective endocarditis on a porcine heterograft valve. *J Thorac Cardiovasc Surg* 1980; **79**: 269–274.
- Middleton J, Chmel H, Tecson F, Sarkaria JS, Neville WE. Aortotomy site infections: case presentation and review of the literature. *Am J Med Sci* 1980; **279**: 105–109.
- Vo NM, Russell JC, Becker DR. Mycotic emboli of the peripheral vessels: analysis of forty-four cases. *Surgery* 1981; **90**: 541–545.
- Rozo L, Cesbron JP, de Lajarte AY, Marjolet M, Dupon H, Nicolas G. *Aspergillus* endocarditis following valve replacement. Apropos of 5 cases. *Arch Mal Coeur Vaiss* 1983; **76**: 822–830.
- Moore RS, Hasleton PS, Lawson RA, Stanbridge TN. *Aspergillus niger* endocarditis complicating aortic tissue valve replacement. *Thorax* 1984; **39**: 76–77.
- Farina C, Filice C, Carnevale G, Maserati R, Alberici F. *Aspergillus* endocarditis: a clinical report. *G Malatt Infett Parassitar* 1985; **37**: 205–208.
- John V, Shuhaiber HJ, Chugh TD, Ghosh DK. *Aspergillus fumigatus* endocarditis on a Carpentier-Edwards xenograft. *J Kuwait Med Assoc* 1987; **21**: 210–213.
- Corrigan C, Horner SM. *Aspergillus* endocarditis in association with a false aortic aneurysm. *Clin Cardiol* 1988; **11**: 430–432.

36. Verghese S, Mullasari A, Padmaja P, Sudha P, Sapna MC, Cherian KM. Fungal endocarditis following cardiac surgery. *Ind Heart J* 1998; **50**: 418–422.
37. Vivas C. Endocarditis caused by *Aspergillus niger*: case report. *Clin Infect Dis* 1998; **27**: 1322–1323.
38. Mehta G. *Aspergillus* endocarditis after open heart surgery: an epidemiological investigation. *J Hosp Infect* 1990; **15**: 245–253.
39. Jenckes GA. *Aspergillus* aortitis. *J Thorac Cardiovasc Surg* 1990; **99**: 375–376.
40. Ledesma D, William HP. Septic (*Aspergillus*) embolus. *N Engl J Med* 2000; **342**: 1015.
41. Kocazeybek B, Sonmez B, Sarman K *et al.* Diagnosis at first glance: hairy black colonies in resected prosthetic aortic valve. *Clin Microbiol Infect* 2000; **6**: 617–618.
42. Galrinho A, Tavares J, Caria R, Veiga C. Fungal endocarditis with atypical localization in patients with aortic prosthesis. *Rev Port Cardiol* 2000; **19**: 1045–1047.
43. Plaza Martínez A, Díaz López M, Riera Vázquez R *et al.* Bilateral popliteal embolism secondary to the fungal infection of the scar tissue resulting from aortotomy. *Angiologia* 2002; **54**: 455–459.
44. Ramos AO, Medeiros AR, Paulista PP, Abboud CS, Meneghelo ZM. *Aspergillus* infection in the ascending aorta of a patient with aortic and mitral valve prostheses. *Arq Bras Cardiol* 2003; **81**: 419–420.
45. López-Gómez M, López-Ruz MA, Jiménez-Alonso JF. Mycotic aneurysm of the superior mesenteric artery and intestinal infarction due to *Aspergillus*. *Enferm Infecc Microbiol Clin* 2003; **21**: 530.
46. Sanchez-Recalde A, Mate I, Merino JL, Simon RS, Sobrino JA. *Aspergillus* aortitis after cardiac surgery. *J Am Coll Cardiol* 2003; **41**: 152–156.
47. El-Hamamsy I, Dürreleman N, Stevens LM *et al.* A cluster of cases of *Aspergillus* endocarditis after cardiac surgery. *Ann Thorac Surg* 2004; **77**: 2184–2186.
48. Reis LJ, Barton TD, Pochettino A *et al.* Successful treatment of *Aspergillus* prosthetic valve endocarditis with oral voriconazole. *Clin Infect Dis* 2005; **41**: 752–753.
49. Lónyai T, Fehér C, Timár K. *Aspergillus* endocarditis following aortic valve replacement. *Acta Chir Acad Sci Hung* 1973; **14**: 43–51.
50. Villate JI, Aldamiz-Echevarria G, Gaztelurrutia L, Barrenechea JI, Gonzalez de Zarate P. Cardiac aspergillosis. *J Thorac Cardiovasc Surg* 2000; **119**: 403–404.
51. Veinot JP, Lorimer JW, Walley VM, Turek M, Saginur R, Rubens F. Aortitis with multiple aneurysms mimicking infective endocarditis. *Can J Cardiol* 1999; **15**: 105–109.
52. Malivi TA, Webb HM, Dixon CD, Boone JA. Systemic aspergillosis caused by *Aspergillus niger* after open-heart surgery. *JAMA* 1968; **203**: 520–522.
53. Anonymous. A case of fungal endocarditis. *BMJ* 1969; **3**: 765–779.
54. Malcolm AD, Bakerspigel A, Enriquez AA. *Aspergillus flavus* endocarditis following aortic valvotomy. *Thorax* 1971; **26**: 435–442.
55. Horten BC, Abbott GF, Porro RS. Fungal aneurysms of intracranial vessels. *Arch Neurol* 1976; **33**: 577–579.
56. Kleiman MB. Echocardiography in *Aspergillus* endocarditis. *Pediatrics* 1982; **69**: 252–253.
57. Koshi G, Cherian KM. *Aspergillus terreus*, an uncommon fungus causing aortic root abscess and pseudoaneurysm. *Ind Heart J* 1995; **47**: 265–267.
58. Hadorn W. Aortic rupture caused by *Aspergillus* infection following operation in aortic stenosis. Endocarditis polyposa mycotica. *Schweiz Med Wochenschr* 1960; **90**: 929–934.
59. Bortolotti U, Thiene G, Milano A, Panizzon G, Valente M, Gallucci V. Pathological study of infective endocarditis on Hancock porcine bioprostheses. *J Thorac Cardiovasc Surg* 1981; **81**: 934–942.
60. Laham MN, Carpenter JL. *Aspergillus terreus*, a pathogen capable of causing infective endocarditis, pulmonary mycetoma, and allergic bronchopulmonary aspergillosis. *Am Rev Respir Dis* 1982; **125**: 769–772.
61. Newman WH, Cordell RA. *Aspergillus* endocarditis after open-heart surgery. Report of a case and review of literature. *J Thorac Cardiovasc Surg* 1964; **48**: 652–660.
62. Darrell RW. Endogenous *Aspergillus* uveitis following heart surgery. *Arch Ophthalmol* 1967; **78**: 354–357.
63. Morea M, Girelli L, Benintendi V, Scaravilli F. *Aspergillus* endocarditis following cardiocirculatory operation. *Chir Ital* 1967; **19** (suppl 2): 173–185.
64. Khan TH, Kane EG, Dean DC. *Aspergillus* endocarditis of mitral prosthesis. *Am J Cardiol* 1968; **22**: 277–280.
65. Chelloul P, Adotti F, Caulin C, Manicacci M, Cornu P, Dentan M. Cardio-aortic aspergillosis. *Sem Hop* 1971; **47**: 825–831.
66. Schelbert HR, Muller OF. Detection of fungal vegetations involving a Starr–Edwards mitral prosthesis by means of ultrasound. *Vasc Surg* 1972; **6**: 20–25.
67. Harford CG. Postoperative fungal endocarditis. Fungemia, embolism, and therapy. *Arch Intern Med* 1974; **134**: 116–120.
68. Kammer RB, Utz JP. *Aspergillus* species endocarditis. The new face of a not so rare disease. *Am J Med* 1974; **56**: 506–521.
69. Gallego Page JC, Marquez Montes J, Ortigosa Aso FJ *et al.* Endocarditis by *Aspergillus*: a case report. *Clin Cardiovasc* 1998; **16**: 12–14.
70. Chow WH, Ng TH, Chow TC, Tai YT, Cheung KL. Free-floating *Aspergillus* vegetation in left atrium: identification by two-dimensional echocardiography. *J Clin Ultrasound* 1991; **19**: 315–317.
71. Verghese S, Maria CF, Mullasari AS, Asha M, Padmaja P, Padhye AA. *Aspergillus* endocarditis presenting as femoral artery embolism. *Mycosis* 2004; **47**: 252–256.
72. Tomaru A, Horiguchi M. Complications of valvular surgery—5 case reports with special reference to the conduction system. *Jpn Circ J* 1975; **39**: 1225–1240.
73. Mershon JC, Samuelson DR, Layman TE. Left ventricular ‘fibrous body’ aneurysm caused by aspergillus endocarditis. *Am J Cardiol* 1968; **22**: 281–285.
74. Wagner DK, Werner PH, Bonchek LI, Shimshak T, Rytel MW. Successful treatment of post-mitral valve annuloplasty *Aspergillus flavus* endocarditis. *Am J Med* 1985; **79**: 777–780.
75. Robbins JA, Mohanty PK, Mohanty LB. Aortic aspergiloma resulting in supraventricular aortic stenosis after aortocoronary bypass surgery. *Am Heart J* 1983; **106**: 163–166.
76. Gray R, Kaplan L, Matloff J *et al.* Aortic pseudo-aneurysm with *Aspergillus* aortitis: an unusual complication of coronary bypass surgery. *Chest* 1986; **89**: 306–308.
77. Opal SM, Reller LB, Harrington G, Cannady P. *Aspergillus clavatus* endocarditis involving a normal aortic valve

- following coronary artery surgery. *Rev Infect Dis* 1986; **8**: 781–785.
78. Weber SF, Washburn RG. Invasive *Aspergillus* infections complicating coronary artery bypass grafting. *South Med J* 1990; **83**: 584–588.
 79. Kaneda T, Iemura J, Oka H *et al.* Treatment of deep infection following thoracic aorta graft replacement without graft removal. *Ann Vasc Surg* 2001; **15**: 430–434.
 80. Choyke PL, Edmonds PR, Markowitz RI, Kleinman CS, Laks H. Mycotic pulmonary artery aneurysm: complication of *Aspergillus* endocarditis. *Am J Roentgenol* 1982; **138**: 1172–1175.
 81. Shrivastava S, Radhakrishnan S. Infective endocarditis following patch closure of ventricular septal defect: a cross-sectional Doppler echocardiographic study. *Int J Cardiol* 1989; **25**: 27–31.
 82. Navabi MA, Ajami H, Amirghofran A, Peyravian F. *Aspergillus* endocarditis: rare but serious *Aspergillus* ball obstructing the pulmonary artery. *Eur J Cardiothorac Surg* 1998; **14**: 530–532.
 83. Barst RJ, Prince AS, Neu HC. *Aspergillus* endocarditis in children: case report and review of the literature. *Pediatrics* 1981; **68**: 73–78.
 84. Shoar MG, Zomorodian K, Saadat F, Hashemi MJ, Tarazoei B. Fatal endocarditis due to *Aspergillus flavus* in Iran. *J Pak Med Assoc* 2004; **54**: 485–486.
 85. Doughten RM, Pearson HA. Disseminated intravascular coagulation associated with *Aspergillus* endocarditis. Fatal outcome following heparin therapy. *J Pediatr* 1968; **73**: 576–582.
 86. Hosking MC, MacDonald NE, Cornel G. Liposomal amphotericin B for postoperative *Aspergillus fumigatus* endocarditis. *Ann Thorac Surg* 1995; **59**: 1015–1017.
 87. Khan ZU, Chugh TD. Invasive fungal infections in Kuwait: a retrospective study. *Ind J Chest Dis Allied Sci* 2000; **42**: 279–287.
 88. Fujioka S, Karashima K, Inoue A, Takami H, Nishikawa N, Saito Y. Case of infectious endocarditis predicted by orbital color Doppler imaging. *Jpn J Ophthalmol* 2005; **49**: 46–48.
 89. Thamlikitkul V, Prachuabmoh K, Sukroongreung S, Danchaiwijitr S. *Aspergillus terreus* endocarditis—a case report. *J Med Assoc Thai* 1983; **66**: 723–726.
 90. Barbone A, Pini D, Grossi P *et al.* *Aspergillus* left ventricular assist device endocarditis. *Ital Heart J* 2004; **5**: 876–880.
 91. Escribano Subias P, Lopez Rios F, Delgado Jimenez JF, Sotelo Rodriguez T, Aguado JM, Rodriguez Hernandez E. Mycotic aneurysm caused by *Aspergillus* of the aortic suture line after heart transplantation. *Rev Esp Cardiol* 2000; **53**: 1403–1405.
 92. Rao NA, Hidayat A. A comparative clinicopathologic study of endogenous mycotic endophthalmitis: variations in clinical and histopathologic changes in candidiasis compared to aspergillosis. *Trans Am Ophthalmol Soc* 2000; **98**: 183–193.
 93. Atkinson JB, Connor DH, Robinowitz M, McAllister HA, Virmani R. Cardiac fungal infections: review of autopsy findings in 60 patients. *Hum Pathol* 1984; **15**: 935–942.
 94. Walsh TJ, Hutchins GM, Bulkley BH, Mendelsohn G. Fungal infections of the heart: analysis of 51 autopsy cases. *Am J Cardiol* 1980; **45**: 357–366.
 95. Wain W, Ahmed M, Thompson R, Yacoub M. The role of chemotherapy in the management of fungal endocarditis following homograft valve replacement. *Postgrad Med J* 1979; **55**: 629–631.
 96. Raghavan R, Krishnaswami H, Date A. Experience with fungal infections affecting the heart in necropsies at an Indian hospital. *Trans R Soc Trop Med Hyg* 1989; **83**: 563–564.
 97. Mencl K, Otcenasek M, Spacek J, Rehulova E. *Aspergillus restrictus* and *Candida parapsilosis*—agents of endocarditis following heart valve replacement. *Mykosen* 1985; **28**: 127–133.
 98. Stein PD, Harken DE, Dexter L. The nature and prevention of prosthetic valve endocarditis. *Am Heart J* 1966; **71**: 393–407.
 99. Cherubin CE, Neu HC. Infective endocarditis at the Presbyterian Hospital in New York City from 1938–1967. *Am J Med* 1938; **51**: 83–96.
 100. Choudhary SK, Talwar S, Kumar AS. Bentall operation with valved homograft conduit. *Tex Heart Inst J* 2000; **27**: 366–368.
 101. Puig LB, Verginelli G, Kawabe L *et al.* Four years experience with dura mater cardiac valves. *J Cardiovasc Surg (Torino)* 1977; **18**: 247–255.
 102. Vogeser M, Wanders A, Haas A, Ruckdeschel G. A four-year review of fatal aspergillosis. *Eur J Clin Microbiol Infect Dis* 1999; **18**: 42–45.
 103. Boyd AD, Spencer FC, Isom OW *et al.* Infective endocarditis. An analysis of 54 surgically treated patients. *J Thorac Cardiovasc Surg* 1977; **73**: 23–30.
 104. Arnett EN, Roberts WC. Prosthetic valve endocarditis: clinicopathologic analysis of 22 necropsy patients with comparison observations in 74 necropsy patients with active infective endocarditis involving natural left-sided cardiac valves. *Am J Cardiol* 1976; **38**: 281–292.
 105. Anonymous. News and notes, epidemiology: mycosis. *BMJ* 1970; **ii**: 185.
 106. Clarkson PM, Barratt-Boyes BG. Bacterial endocarditis following homograft replacement of the aortic valve. *Circulation* 1970; **42**: 987–991.
 107. Kalil RAK, Macedo RG, Prates PR *et al.* Re-operations following Bentall–De Bono operation for anulo-aortic ectasia. *Rev Brasil Cirurg Cardiovasc* 1988; **3**: 93–100.
 108. Chen SC, Sorrell TC, Dwyer DE, Collignon PJ, Wright EJ. Endocarditis associated with prosthetic cardiac valves. *Med J Aust* 1990; **152**: 458–463.
 109. Semb BK, Hall KV, Nitter-Hauge S, Abdelnoor M. A 5-year follow-up of the Medtronic–Hall valve: survival and thromboembolism. *Thorac Cardiovasc Surg* 1983; **31**: 61–65.
 110. Coutlee F, Carceller AM, Deschamps L, Kratz C, Lapointe JR, Davignon A. The evolving pattern of pediatric endocarditis from 1960 to 1985. *Can J Cardiol* 1990; **6**: 164–170.
 111. Stanulla H. Mycotic prosthetic valve endocarditis. *Zentralbl Allgem Pathol Pathol Anat* 1983; **127**: 97–103.
 112. Pospisil K, Strata K, Otcenasek M *et al.* Mycotic ulcerative aortitis after replacement of the aortic valve caused by the fungus *Aspergillus restrictus*. *Vnitr Lek* 1984; **30**: 292–297.
 113. Kawasaki S, Naitoh Y, Takagaki Y, Komai Y, Noguchi Y. Two cases of *Aspergillus* endocarditis after cardiac surgery in childhood. *Kyobu Geka* 1996; **49** (suppl 8): 673–676.
 114. Katayama Y, Okabe M, Mizutani T, Yada I, Yuasa H, Kusagawa M. Treatment of prosthetic valve endo-

- carditis—analysis of eleven cases. *Nippon Kyobu Geka Gakkai Zasshi* 1989; **37**: 2111–2116.
115. Otto T, Halweg H, Pietraszek A, Kaminski Z. *Aspergillus* endocarditis following aortic valve replacement. *Kardiol Pol* 1979; **22**: 221–226.
 116. Nakanishi M, Teramoto S, Doi T, Mihara Y, Nasu M. Case of *Aspergillus* endocarditis following aortic valve replacement. *Kyobu Geka* 1973; **26**: 432–436.
 117. Grabowski H, Holak L, Musiatowicz B. Case of *Aspergillus* endocarditis in a patient following surgery for Fallot's tetralogy. *Pol Tyg Lek* 1971; **26**: 520–521.
 118. Tomsova M. Mycotic diseases of the heart. *Cesk Patol* 2001; **37**: 172–176.
 119. Dudkiewicz B, Szewczyk E. Etiology of bacterial endocarditis in materials from Cardiology and Cardiac Surgery Clinics of the Lodz Academy. *Med Dosw Mikrobiol* 1993; **45**: 357–359.
 120. Sakashita I, Otani S, Terashima M, Equchi A, Nakamura C. A study on the relationship between prosthetic valve replacement and infective endocarditis before and after surgery. Part 2. With special reference to prosthetic valve endocarditis. *Nippon Kyobu Geka Gakkai Zasshi* 1979; **27**: 1537–1543.
 121. Wellens F, Potvlieghe C, Deuvaert FE, Primo G. *Aspergillus* osteochondritis after median sternotomy. Combined operative treatment and drug therapy with amphotericin B. *Thorac Cardiovasc Surg* 1982; **30**: 322–324.
 122. Sachs MK, Paluzzi RG, Moore JH, Fraimow HS, Ost D. Amphotericin-resistant *Aspergillus* osteomyelitis controlled by itraconazole. *Lancet* 1990; **335**: 1475.
 123. Vandecasteele SJ, Boelaert JR, Verrelst P, Graulus E, Gordts BZ. Diagnosis and treatment of *Aspergillus flavus* sternal wound infections after cardiac surgery. *Clin Infect Dis* 2002; **35**: 887–890.
 124. Elahi MM, Mitra A, Spears J, McClurken JB. Recalcitrant chest wall *Aspergillus fumigatus* osteomyelitis after coronary artery bypass grafting: successful radical surgical and medical management. *Ann Thorac Surg* 2005; **79**: 1057–1059.
 125. Barzaghi N, Emmi V, Mencherini S, Minzioni G, Marone P, Minoli L. Sternal osteomyelitis due to *Aspergillus fumigatus* after cardiac surgery. *Chest* 1994; **105**: 1275–1277.
 126. Whiting RB, Barner HB, Leone P, Westura EE. Aspergilloma: an unusual cause of late failure of aortocoronary bypass graft. *Chest* 1973; **63**: 1030–1033.
 127. Draganov J, Klein HM, Ghodsizad A, Gehrke M, Gams E. Rupture of a saphenous vein coronary artery bypass graft due to *Aspergillus* necrotizing vasculitis. *Ann Thorac Surg* 2005; **80**: 724–726.
 128. Pemán J, Ortíz R, Osseyran F *et al.* Non-surgical endocarditis by *Aspergillus* with mould isolation in blood culture and negative galactomannan. *Mycoses* 2005; **48** (suppl 2): 45.
 129. Frank KA, Merz WG, Hutchins GM. Sclerotium formation in an *Aspergillus flavus* wound infection. *Mycopathologia* 1988; **102**: 185–188.
 130. Sawyer RG, Schenk WG, Adams RB, Pruett TL. *Aspergillus flavus* wound infection following repair of a ruptured duodenum in a non-immunocompromised host. *Scand J Infect Dis* 1992; **24**: 805–809.
 131. Yuen JC, Puri SK, Feng Z. Scalp necrotizing fasciitis with osteomyelitis of the skull from *Aspergillus*. *J Craniofac Surg* 2002; **13**: 762–764.
 132. Falsey AR, Goldsticker D, Ahem MJ. Fatal subcutaneous aspergillosis following necrotizing fasciitis: a case report. *Yale J Biol Med* 1990; **63**: 9–13.
 133. Sergi C, Weitz J, Hofmann WJ *et al.* *Aspergillus* endocarditis, myocarditis and pericarditis complicating necrotizing fasciitis. Case report and subject review. *Virchows Arch* 1996; **429**: 177–180.
 134. Johnson MA, Lyle G, Hanly M, Yeh KA. *Aspergillus*: a rare primary organism in soft-tissue infections. *Am Surg* 1998; **64**: 122–126.
 135. Lai CS, Lin SD, Chou CK, Lin HJ. Aspergillosis complicating the grafted skin and free muscle flap in a diabetic. *Plast Reconstr Surg* 1993; **92**: 532–536.
 136. Anderson LL, Giandoni MB, Keller RA, Grabski WJ. Surgical wound healing complicated by *Aspergillus* infection in a nonimmunocompromised host. *Dermatol Surg* 1995; **21**: 799–801.
 137. Attah CA, Cerruti MM. *Aspergillus* osteomyelitis of sternum after cardiac surgery. *NY State J Med* 1979; **79**: 1420–1421.
 138. Langlois RP, Flegel KM, Meakins JL, Morehouse DD, Robson HG, Guttman RD. Cutaneous aspergillosis with fatal dissemination in a renal transplant recipient. *Can Med Assoc J* 1980; **120**: 673–676.
 139. Nampoory MR, Khan ZU, Johny KV *et al.* Invasive fungal infections in renal transplant recipients. *J Infect* 1996; **33**: 95–101.
 140. Grossi P, Farina C, Fiocchi R, Dalla Gasperina D. Prevalence and outcome of invasive fungal infections in 1,963 thoracic organ transplant recipients: a multi-center retrospective study. *Transplantation* 2000; **70**: 112–116.
 141. Le Conte P, Blanloeil Y, Michel P, Francois T, Paineau J. Cutaneous aspergillosis in a patient with orthotopic hepatic transplantation. *Transplantation* 1992; **53**: 1153–1154.
 142. Plá MP, Berenguer J, Arzuaga JA, Bafiares R, Polo JR, Bouza E. Surgical wound infection by *Aspergillus fumigatus* in liver transplant recipients. *Diagn Microbiol Infect Dis* 1992; **15**: 703–706.
 143. Pegues DA, Lasker BA, McNeil MM, Hamm PM, Lundal JL, Kubak BM. Cluster of cases of invasive aspergillosis in a transplant intensive care unit: evidence of person-to-person airborne transmission. *Clin Infect Dis* 2002; **34**: 412–416.
 144. Richet HM, McNeil MM, Davis BJ *et al.* *Aspergillus fumigatus* sternal wound infections in patients undergoing open heart surgery. *Am J Epidemiol* 1992; **135**: 48–58.
 145. Bryce EA, Walker M, Scharf S *et al.* An outbreak of cutaneous aspergillosis in a tertiary-care hospital. *Infect Control Hosp Epidemiol* 1996; **17**: 170–172.
 146. Marples RR. Contaminated first-aid dressings: report of a Working Party of the PHLS. *J Hyg* 1983; **90**: 241–252.
 147. Patterson JE, Barden GE, Bia FJ. Hospital-acquired gangrenous mucormycosis. *Yale J Biol Med* 1986; **59**: 453–459.
 148. Florio M, Marroni M, Morosi S, Stagni G. Nosocomial *Aspergillus flavus* wound infections following cardiac surgery. *Infez Med* 2004; **12**: 270–273.
 149. Siddiqui AA, Shah AA, Bashir SH. Craniocerebral aspergillosis of sinonasal origin in immunocompetent patients: clinical spectrum and outcome in 25 cases. *Neurosurgery* 2004; **55**: 602–613.

150. Visudhiphan P, Bunyaratavej S, Khantanaphar S. Cerebral aspergillosis. Report of three cases. *J Neurosurg* 1973; **38**: 472–476.
151. Shapiro K, Tabaddor K. Cerebral aspergillosis. *Surg Neurol* 1975; **4**: 465–471.
152. Feely M, Steinberg M. *Aspergillus* infection complicating transphenoidal yttrium-90 pituitary implant. Report of two cases. *J Neurosurg* 1977; **46**: 530–532.
153. Morioka T, Tashima T, Nagata S, Fukui M, Hasuo K. Cerebral aspergillosis after burr-hole surgery for chronic subdural hematoma: case report. *Neurosurgery* 1990; **26**: 332–335.
154. Galassi E, Pozzati E, Poppi M, Vinci A. Cerebral aspergillosis following intracranial surgery. Case report. *J Neurosurg* 1978; **49**: 308–311.
155. Hajjar J, Brunon J, Jaubert J *et al.* Cerebral aspergillosis. Apropos of 4 cases. *Neurochirurgie* 1987; **33**: 142–147.
156. Mielke B, Weir B, Oldring D, von Westarp C. Fungal aneurysm: case report and review of the literature. *Neurosurgery* 1981; **9**: 578–582.
157. Ouammou A, el Ouarzazi A, Belghmaidi M, el Faidouzi M. Cerebral aspergillosis and encephalomeningocele. *Childs Nerv Syst* 1986; **2**: 216–218.
158. Piotrowski WP, Pilz P, Chuang IH. Subarachnoid hemorrhage caused by a fungal aneurysm of the vertebral artery as a complication of intracranial aneurysm clipping. Case report. *J Neurosurg* 1990; **73**: 962–964.
159. Kim DG, Hong SC, Kim HJ *et al.* Cerebral aspergillosis in immunologically competent patients. *Surg Neurol* 1993; **40**: 326–331.
160. Komatsu Y, Narushima K, Kobayashi E, Tomono Y, Nose T. *Aspergillus* mycotic aneurysm—case report. *Neurol Med Chir (Tokyo)* 1991; **31**: 346–350.
161. Darras-Joly C, Veber B, Bedos JP, Gachot B, Regnier B, Wolff M. Nosocomial cerebral aspergillosis: a report of 3 cases. *Scand J Infect Dis* 1996; **28**: 317–319.
162. Sharma RR, Gurusinge NT, Lynch PG. Cerebral infarction due to *Aspergillus* arteritis following glioma surgery. *Br J Neurosurg* 1992; **6**: 485–490.
163. Beal MF, O'Carroll CP, Kleinman GM, Grossman RI. Aspergillosis of the nervous system. *Neurology* 1982; **32**: 473–479.
164. Viriyavejakul P, Phudhichareonrat S, Boonpasat Y *et al.* Intracerebral hemorrhage due to nosocomial aspergillosis following neurosurgery. *SE Asian J Trop Med Pub Hlth* 1999; **30**: 154–156.
165. Letscher V, Herbrecht R, Gaudias J *et al.* Post-traumatic intracranial epidural *Aspergillus fumigatus* abscess. *J Med Vet Mycol* 1997; **35**: 279–282.
166. Takeshita M, Izawa M, Kubo O *et al.* Aspergillotic aneurysm formation of cerebral artery following neurosurgical operation. *Surg Neurol* 1992; **38**: 146–151.
167. Cuccia V, Galarza M, Monges J. Cerebral aspergillosis in children. Report of three cases. *Pediatr Neurosurg* 2000; **33**: 43–48.
168. Endo T, Numagami Y, Jokura H, Ikeda H, Shirane R, Yoshimoto T. *Aspergillus* parasellar abscess mimicking radiation-induced neuropathy. Case report. *Surg Neurol* 2001; **56**: 195–200.
169. Theaker ED, Rushton VE, Corcoran JP, Hatton P. Chronic sinusitis and zinc-containing endodontic obturating pastes. *Br Dent J* 1995; **179**: 64–68.
170. Legent F, Beauvillain C, Mercier J, Hofmann B, Wesoluch M. Relations between sinus aspergillosis, primary rhinoliths and dental pathology. *Ann Otolaryngol Chir Cervicofac* 1982; **99**: 541–545.
171. Beck-Mannagetta J, Necek D, Grasserbauer M. Dental aspects of solitary maxillary sinus aspergillosis. A clinical, microanalytical and experimental study. *Z Stomatol* 1986; **83**: 283–315.
172. Legent F, Billet J, Beauvillain C, Bonnet J, Miegerville M. The role of dental canal fillings in the development of *Aspergillus* sinusitis. A report of 85 cases. *Arch Otorhinolaryngol* 1989; **246**: 318–320.
173. De Foer C, Fossion E, Vaillant JM. Sinus aspergillosis. *J Craniomaxillofac Surg* 1990; **18**: 33–40.
174. Bader G. *Aspergillus* sinusitis of dental origin. *Rev Odontostomatol (Paris)* 1989; **18**: 345–353.
175. Buffe P, Nottet JB, Poncet JL. Maxillary *Aspergillus* sinusitis. Comments apropos of 20 cases. *Rev Laryngol Otol Rhinol (Bord)* 1992; **113**: 197–200.
176. Khongkhunthian P, Reichart PA. Aspergillosis of the maxillary sinus as a complication of overfilling root canal material into the sinus: report of two cases. *J Endod* 2001; **27**: 476–478.
177. Rodriguez V, Bardwil JM, Bodey GP. Primary aspergillosis cured with amphotericin B. *South Med J* 1971; **64**: 396–398.
178. Beck-Mannagetta J, Necek D. Radiologic findings in aspergillosis of the maxillary sinus. *Oral Surg Oral Med Oral Pathol* 1986; **62**: 345–349.
179. Krennmair G, Lenglinger F, Laine FJ. Maxillary sinus aspergillosis: diagnosis and differentiation of the pathogenesis based on computed tomography densitometry of sinus concretions. *J Oral Maxillofac Surg* 1995; **53**: 657–664.
180. Legend F, Desnos J, Beauvillain C. *Aspergillus* sinusitis. *J Fr Otorhinolaryngol Audiophonol Chirur Maxillofac* 1980; **29**: 39–49.
181. Martins WD, Ribeiro Rosa EA. Aspergillosis of the maxillary sinus: review and case report. *Scand J Infect Dis* 2004; **36**: 758–761.
182. Beck-Mannagetta J, Necek D, Grasserbauer M. Solitary aspergillosis of maxillary sinus, a complication of dental treatment. *Lancet* 1983; **ii**: 1260.
183. Gonty AA, Page LR. Aspergillosis of the maxillary sinus. Review of the literature and report of a case. *Oral Surg Oral Med Oral Pathol* 1977; **43**: 350–356.
184. Brar GS, Ram J, Kaushik S, Chakraborti A, Dogra MR, Gupta A. *Aspergillus niger* endophthalmitis after cataract surgery. *J Cataract Refract Surg* 2002; **28**: 1882–1883.
185. Mendicute J, Orbegozo J, Ruiz M, Saiz A, Eder F, Aramberri J. Keratomycosis after cataract surgery. *J Cataract Refract Surg* 2000; **26**: 1660–1666.
186. Kermani NK, Aggarwal SP. Isolated post-operative *Aspergillus niger* endophthalmitis. *Eye* 2000; **14**: 114–116.
187. Das T, Vyas P, Sharma S. *Aspergillus terreus* postoperative endophthalmitis. *Br J Ophthalmol* 1993; **77**: 386–387.
188. Kloess PM, Stulting RD, Waring GO, Wilson LA. Bacterial and fungal endophthalmitis after penetrating keratoplasty. *Am J Ophthalmol* 1993; **115**: 309–316.
189. Weichel ED, Bower KS, Ward TP, Hidayat A. Epithelial aspergilloma after penetrating keratoplasty. *Cornea* 2002; **21**: 825–827.
190. Cameron JA, Antonios SR, Cotter JB, Habash NR. Endophthalmitis from contaminated donor corneas following

- penetrating keratoplasty. *Arch Ophthalmol* 1991; **109**: 54–59.
191. Heidemann DG, Dunn SP, Watts JC. *Aspergillus* keratitis after radial keratotomy. *Am J Ophthalmol* 1995; **120**: 254–256.
 192. Panda A, Das GK, Vanathi M, Kumar A. Corneal infection after radial keratotomy. *J Cataract Refractive Surg* 1998; **24**: 331–334.
 193. Maskin SL, Alfonso E. Fungal keratitis after radial keratotomy. *Am J Ophthalmol* 1992; **114**: 369–370.
 194. de Sousa LB, Cronemberger MF, Campos M, Nose W. Penetrating keratoplasty in the treatment of fungal keratitis after radial keratotomy: report of 3 cases. *Arquivos Brasil Oftalmol* 1995; **58**: 47–49.
 195. Dunphy D, Andrews D, Seamone C, Ramsey M. Fungal keratitis following excimer laser photorefractive keratectomy. *Can J Ophthalmol* 1999; **34**: 286–289.
 196. Kuo IC, Margolis TP, Cevallos V, Hwang DG. *Aspergillus fumigatus* keratitis after laser in situ keratomileusis. *Cornea* 2001; **20**: 342–344.
 197. Sridhar MS, Garg P, Bansal AK, Gopinathan U. *Aspergillus flavus* keratitis following laser in situ keratomileusis. *Am J Ophthalmol* 2000; **129**: 802–804.
 198. Ritterband D, Kelly J, McNamara T, Kresloff M, Koplin R, Seedor J. Delayed-onset multifocal polymicrobial keratitis after laser in situ keratomileusis. *J Cataract Refractive Surg* 2002; **28**: 898–899.
 199. Tehrani M, Dick HB, Daeschlein G, Kramer A. Microbial keratitis following laser in situ keratomileusis. Prevention, differential diagnosis, and therapy. *Ophthalmologie* 2005; **102**: 514–519.
 200. Margo CE, Polack FM, Hood CI. *Aspergillus* panophthalmitis complicating treatment of pterygium. *Cornea* 1988; **7**: 285–289.
 201. Hsiao CH, Chen JJY, Huang SCM, Ma HK, Chen PYF, Tsai RJF. Intrasceral dissemination of infectious scleritis following pterygium excision. *Br J Ophthalmol* 1998; **82**: 29–34.
 202. Oestreicher JH, Bashour M, Jong R, Chiu B. *Aspergillus* mycetoma in a secondary hydroxyapatite orbital implant: a case report and literature review. *Ophthalmology* 1999; **106**: 987–991.
 203. Mendicute J, Ondarra A, Eder F, Ostolaza JI, Salaberria M, Lamsfus JM. The use of collagen shields impregnated with amphotericin B to treat *Aspergillus* keratomycosis. *CLAO J* 1995; **21**: 252–255.
 204. Bernauer W, Allan BDS, Dart JK. Successful management of *Aspergillus* scleritis by medical and surgical treatment. *Eye* 1998; **12**: 311–316.
 205. Carlson AN, Foulks GN, Perfect JR, Kim JH. Fungal scleritis after cataract surgery. Successful outcome using itraconazole. *Cornea* 1992; **11**: 151–154.
 206. Oshitari K, Hirakata A, Okada AA *et al*. Vitrectomy for endophthalmitis after cataract surgery. *Nippon Ganka Gakkai Zasshi* 2003; **107**: 590–596.
 207. Valenton M. Wound infection after cataract surgery. *Jpn J Ophthalmol* 1996; **40**: 447–455.
 208. Narang S, Gupta A, Gupta V *et al*. Fungal endophthalmitis following cataract surgery: clinical presentation, microbiological spectrum, and outcome. *Am J Ophthalmol* 2001; **132**: 609–617.
 209. Tabbara KF, al Jabarti AL. Hospital construction-associated outbreak of ocular aspergillosis after cataract surgery. *Ophthalmology* 1998; **105**: 522–526.
 210. Gupta A, Gupta V, Gupta A *et al*. Spectrum and clinical profile of post cataract surgery endophthalmitis in North India. *Ind J Ophthalmol* 2003; **51**: 139–145.
 211. Tasaka Y, Fujii S. Fungal endophthalmitis following intraocular lens implantation. A case report. *Folia Ophthalmol Jpn* 1990; **41**: 1338–1344.
 212. Mahajan VM. Postoperative ocular infections. An analysis of laboratory data on 750 cases. *Ann Ophthalmol* 1984; **16**: 847–848.
 213. Kondo T, Yoshida H. Fungal infection following scleral buckling procedures. *Folia Ophthalmol Jpn* 1974; **25**: 781–787.
 214. Newton JC, Fowler BJ. *Aspergillus* infection of a scleral implant. *Ann Ophthalmol* 1980; **12**: 952–954.
 215. Garg P, Mahesh S, Bansal AK, Gopinathan U, Rao GN. Fungal infection of sutureless self-sealing incision for cataract surgery. *Ophthalmology* 2003; **110**: 2173–2177.
 216. Oxford K, Abbott RL, Fung WE, Ellis DS. *Aspergillus* endophthalmitis after sutureless cataract surgery. *Am J Ophthalmol* 1995; **120**: 534–535.
 217. Krzystolik MG, Ciulla TA, Topping TM, Baker AS. Exogenous *Aspergillus niger* endophthalmitis in a patient with a filtering bleb. *Retina* 1997; **17**: 461–462.
 218. Alexandris G, Haimovici R, Miller D *et al*. Corneal biopsy in the management of progressive microbial keratitis. *Am J Ophthalmol* 2000; **129**: 571–577.
 219. Thomas PA. Current perspectives on ophthalmic mycoses. *Clin Microbiol Rev* 2003; **16**: 730–797.
 220. Ishibashi Y, Kaufman HE. Corneal biopsy in the diagnosis of keratomycosis. *Am J Ophthalmol* 1986; **101**: 288–293.
 221. Burt B, Pappas G, Simcock P. Hospital acquired *Aspergillus* keratitis. *Br J Ophthalmol* 2003; **87**: 923.
 222. Fahad B, McKellar M, Armstrong M, Denning D, Tullo A. *Aspergillus* keratitis following corneal foreign body. *Br J Ophthalmol* 2004; **88**: 847–848.
 223. Klotz SA, Penn CC, Negvesky GJ, Butrus SI. Fungal and parasitic infections of the eye. *Clin Microbiol Rev* 2000; **13**: 662–685.
 224. Rosa RH, Miller D, Alfonso EC. The changing spectrum of fungal keratitis in South Florida. *Ophthalmology* 1994; **101**: 1005–1013.
 225. Wong TY, Ng TP, Fong KS, Tan DT. Risk factors and clinical outcome between fungal and bacterial keratitis. A comparative study. *CLAO J* 1997; **23**: 275–281.
 226. Armstrong RA. The microbiology of the eye. *Ophthalmic Physiol Opt* 2000; **20**: 429–441.
 227. Gopinathan U, Garg P, Fernandes M, Sharma S, Athmanathan S, Rao GN. The epidemiological features and laboratory results of fungal keratitis: a 10-year review at a referral eye care center in South India. *Cornea* 2002; **21**: 555–559.
 228. Karimov MK. Mycotic infections as possible complications in penetrating eye injuries. *Oftalmol Zh* 1981; **36**: 214–216.
 229. Anonymous. Direct fungal infection of the eye and its prevention. *Br J Ophthalmol* 1976; **60**: 605–606.
 230. Hanish SJ, Perlmutter JC, Boucher C, Bohigian G, Grand MG. Exogenous *Aspergillus* endophthalmitis. *Ann Ophthalmol* 1984; **16**: 417–419.
 231. Ho PC, Tolentino FI, Baker AS. Successful treatment of exogenous *Aspergillus* endophthalmitis: a case report. *Br J Ophthalmol* 1984; **68**: 412–415.

232. Kaushik S, Ram J, Brar GS, Jain AK, Chakraborti A, Gupta A. Intracameral amphotericin B: initial experience in severe keratomycosis. *Cornea* 2001; **20**: 715–719.
233. Abu el-Asrar AM, al-Amro SA, al-Mosallam AA, al-Obeidan S. Post-traumatic endophthalmitis: causative organisms and visual outcome. *Eur J Ophthalmol* 1999; **9**: 21–31.
234. Carranza R, Lopez N, Gonzalez F, Dominguez J, Amigo A, Martinez F. Ocular perforation in the agrarian environment. *Enferm Infecc Microbiol Clin* 1998; **16**: 377–378.
235. Hirose H, Terasaki H, Awaya S, Yasuma T. Treatment of fungal corneal ulcers with amphotericin B ointment. *Am J Ophthalmol* 1997; **124**: 836–838.
236. Kulshrestha M, Palimar P. Corneal ulcers in paddy workers in south India. *Occup Med (Lond)* 1996; **46**: 320.
237. Rodriguez-Ares MT, De Rojas Silva MV, Pereiro M, Fente Sampayo B, Gallegos Chamas GS, Salorio M. *Aspergillus fumigatus* scleritis. *Acta Ophthalmol Scand* 1995; **73**: 467–469.
238. Ariyasu RG, Kumar S, LaBree LD, Wagner DG, Smith RE. Microorganisms cultured from the anterior chamber of ruptured globes at the time of repair. *Am J Ophthalmol* 1995; **119**: 181–188.
239. Upadhyay MP, Karmacharya PC, Koirala S *et al.* Epidemiologic characteristics, predisposing factors, and etiologic diagnosis of corneal ulceration in Nepal. *Am J Ophthalmol* 1991; **111**: 92–99.
240. Lubniewski A, Olk RJ, Grand MG. Ocular dangers in the garden. A new menace—nylon line lawn trimmers. *Ophthalmology* 1988; **95**: 906–910.
241. Srinivasan M, Gonzales CA, George C *et al.* Epidemiology and aetiological diagnosis of corneal ulceration in Madurai, south India. *Br J Ophthalmol* 1997; **81**: 965–971.
242. Gugnani HC, Talwar RS, Njoku-Obi AN, Kodilinye HC. Mycotic keratitis in Nigeria. A study of 21 cases. *Br J Ophthalmol* 1976; **60**: 607–613.
243. Koul RL, Pratap VB. Keratomycosis in Lucknow. *Br J Ophthalmol* 1975; **59**: 47–51.
244. O'Day DM, Moore TE, Aronson SB. Deep fungal corneal abscess. Combined corticosteroid therapy. *Arch Ophthalmol* 1971; **86**: 414–419.
245. Vemuganti GK, Garg P, Gopinathan U *et al.* Evaluation of agent and host factors in progression of mycotic keratitis: a histologic and microbiologic study of 167 corneal buttons. *Ophthalmology* 2002; **109**: 1538–1546.
246. Wood TO, Tuberville AW, Monnett R. Keratomycosis and amphotericin B. *Trans Am Ophthalmol Soc* 1985; **83**: 397–409.
247. Gonawardena SA, Ranasinghe KP, Arseculeratne SN, Seimon CR, Ajello L. Survey of mycotic and bacterial keratitis in Sri Lanka. *Mycopathologia* 1994; **127**: 77–81.
248. Giobbia M, Rossi MC, Conti E *et al.* Post-traumatic *Aspergillus fumigatus* keratitis. *J Mycol Med* 2002; **12**: 32–33.
249. Hariprasad SM, Mieler WF, Holz ER *et al.* Determination of vitreous, aqueous, and plasma concentration of orally administered voriconazole in humans. *Arch Ophthalmol* 2004; **122**: 42–47.
250. Tack KJ, Rhame FS, Brown B, Thompson RC. *Aspergillus* osteomyelitis. Report of four cases and review of the literature. *Am J Med* 1982; **73**: 295–300.
251. Peters-Christodoulou MN, de Beer FC, Bots GT, Ottenhoff TM, Thompson J, van't Wout JW. Treatment of post-operative *Aspergillus fumigatus* spondylodiscitis with itraconazole. *Scand J Infect Dis* 1991; **23**: 373–376.
252. Mawk JR, Erickson DL, Chou SN, Seljeskog EL. *Aspergillus* infections of the lumbar disc spaces. Report of three cases. *J Neurosurg* 1983; **58**: 270–274.
253. Dabrowski LC, Marchou B, Dabrowski A. Spondylodiscites aspergillaires: à propos de 3 cas et revue de la littérature. *Med Mal Infect* 2003; **33**: 258–265.
254. Lenzi J, Agrillo A, Santoro A, Marotta N, Cantore GP. Postoperative spondylodiscitis from *Aspergillus fumigatus* in immunocompetent subjects. *J Neurosurg Sci* 2004; **48**: 81–85.
255. Fitzgerald RH, Peterson LF, Washington JA, Van Scoy RE, Coventry MB. Bacterial colonization of wounds and sepsis in total hip arthroplasty. *J Bone Joint Surg Am* 1973; **55**: 1242–1250.
256. Cimerman M, Gunde-Cimerman N, Zalar P, Perkovic T. Femur osteomyelitis due to a mixed fungal infection in a previously healthy man. *J Clin Microbiol* 1999; **37**: 1532–1535.
257. Brook I, Aaron RK. Hospital-acquired *Aspergillus flavus* infection in a surgical wound: a case report. *Orthopedics* 1980; **3**: 1215–1216.
258. Austin KS, Testa NN, Luntz RK, Greene JB, Smiles S. *Aspergillus* infection of total knee arthroplasty presenting as a popliteal cyst. Case report and review of the literature. *J Arthroplasty* 1992; **7**: 311–314.
259. Baumann PA, Cunningham B, Patel NS, Finn HA. *Aspergillus fumigatus* infection in a mega prosthetic total knee arthroplasty: salvage by staged reimplantation with 5-year follow-up. *J Arthroplasty* 2001; **16**: 498–503.
260. Langer P, Kassim RA, Macari GS, Saleh KJ. *Aspergillus* infection after total knee arthroplasty. *Am J Orthop* 2003; **32**: 402–404.
261. Kelemen S, Novak E. Mycotic stitch abscess after lobectomy (*Aspergillus* infection) cured by antibiotic therapy. *Beitr Klin Tuberk Spezif Tuberkuloseforsch* 1960; **121**: 654–658.
262. Roig J, Ruiz J, Puig X, Carreras A, Morera J. Bronchial stump aspergillosis four years after lobectomy. *Chest* 1993; **104**: 295–296.
263. Noppen M, Claes I, Maillet B, Meysman M, Monsieur I, Vincken W. Three cases of bronchial stump aspergillosis: unusual clinical presentations and beneficial effect of oral itraconazole. *Eur Respir J* 1995; **8**: 477–480.
264. Le Rochais JP, Icard P, Simon T, Poirier P, Evrard C. Bronchial stump aspergillosis. *Ann Thorac Surg* 2000; **70**: 302–304.
265. Zajackowska J, Halweg H, Pirozynski M, Olechnowicz H, Podsiadlo B, Krakowka P. Further studies on the occurrence of mycoses in the bronchial stump and bronchial anastomosis or trachea. *Pneumonol Pol* 1989; **57**: 104–113.
266. Filippov VP, Lebedev KM, Sofronii SV. Mycotic lesions of the stump of the resected bronchus. *Grudn Khir* 1980; **5**: 72–74.
267. Krakowka P, Halweg H, Nowicki J, Zajackowska J. Aspergillosis of the bronchial stump following resection of pulmonary tissue. *Pol Przegl Chir* 1969; **41**: 242–247.
268. Franke H, Ammen K, Haering R. On the problem of the suture for bronchial closure: observation on mycoses of the bronchial stump. *Thoraxchirurgie* 1961; **8**: 621–626.

269. Sawasaki H, Horie K, Yamada M *et al.* Bronchial stump aspergilliosis. Experimental and clinical study. *J Thorac Cardiovasc Surg* 1969; **58**: 198–208.
270. Katsura H, Iuchi K, Matsumura A. Pulmonary suture granuloma with *Aspergillus* after partial resection for lung cancer; report of a case. *Kyobu Geka* 2005; **58**: 169–171.
271. Nunley DR, Gal AA, Veja JD, Perlino C, Smith P, Lawrence EC. Saprophytic fungal infections and complications involving the bronchial anastomosis following human lung transplantation. *Chest* 2002; **122**: 1185–1191.
272. Kramer MR, Denning DW, Marshall SE *et al.* Ulcerative tracheobronchitis after lung transplantation: a new form of invasive aspergilliosis. *Am Rev Respir Dis* 1991; **144**: 552–556.
273. Bertocchi M, Thevenet F, Bastien O *et al.* Fungal infections in lung transplant recipients. *Transplant Proc* 1995; **27**: 1695.
274. Guillemain R, Lavarde V, Amrein C, Chevalier P, Guinvarc'h A, Glotz D. Invasive aspergilliosis after transplantation. *Transplant Proc* 1995; **27**: 1307–1309.
275. Patterson DL, Singh N. Invasive aspergilliosis in transplant recipients. *Medicine (Baltimore)* 1999; **78**: 123–138.
276. Tezer Y, Memikoglu O, Kart H. Iatrogenic pleural aspergilliosis treated with combination of systemic and intrapleural amphotericin B. *Mycoses* 2005; **48** (suppl 2): 43.
277. Glotzbach RE. *Aspergillus terreus* infection of pseudoaneurysm of aortofemoral vascular graft with contiguous vertebral osteomyelitis. *Am J Clin Pathol* 1982; **77**: 224–227.
278. Brandt SJ, Thompson RL, Wenzel RP. Mycotic pseudoaneurysm of an aortic bypass graft and contiguous vertebral osteomyelitis due to *Aspergillus fumigatus*. *Am J Med* 1985; **79**: 259–262.
279. Doscher W, Krishnasastri KV, Deckoff SL. Fungal graft infections: case report and review of the literature. *J Vasc Surg* 1987; **6**: 398–402.
280. Aguado JM, Valle R, Arjona R, Ferreres JC, Gutierrez JA. Aortic bypass graft infection due to *Aspergillus*: report of a case and review. *Clin Infect Dis* 1992; **14**: 916–921.
281. Echene B, Brunel D, Astruc J, Perez C. Aspergillose disséminée chez un enfant porteur d'une prosthèse aortique. *Med Mal Infect* 1980; **10**: 263–266.
282. Anderson J, Kron IL. Treatment of *Aspergillus* infection of the proximal aortic prosthetic graft with associated vertebral osteomyelitis. *J Vasc Surg* 1984; **1**: 579–581.
283. Hargrove WC, Edmunds LH. Management of infected thoracic aortic prosthetic grafts. *Ann Thorac Surg* 1984; **37**: 72–77.
284. Nussaume O, Elzaabi M, Icard P, Branger C, Bouttier S, Andreassian B. False aneurysm infected by *Aspergillus fumigatus*: an unusual complication of aortofemoral bypass graft. *Ann Vasc Surg* 1990; **4**: 388–392.
285. Collazos J, Mayo J, Martinez E, Ibarra S. Prosthetic vascular graft infection due to *Aspergillus* species: case report and literature review. *Eur J Clin Microbiol Infect Dis* 2001; **20**: 414–417.
286. Lifschultz BD, Leestma JE, Stryker S. Multiple mycotic aneurysms and transverse myelopathy complicating repair of aortic coarctation. *Ann Thorac Surg* 1982; **33**: 192–196.
287. Motte S, Bellens B, Rickaert F, Serruys E, Thys JP, Dereume JP. Vascular graft infection caused by *Aspergillus* species: case report and review of the literature. *J Vasc Surg* 1993; **17**: 607–612.
288. Sánchez Lastres J, Rodríguez Núñez A, Alvez González F *et al.* Infección por *Aspergillus* de prótesis vascular en niño con coartación aórtica. Implicaciones evolutivas y terapéuticas. *An Esp Pediatr* 1993; **38**: 363–364.
289. Wandschneider W, Deutsch M. Fatal fungal infection of an ascending aortic graft. *Thorac Cardiovasc Surg* 1995; **43**: 217–219.
290. García Fuster R, Clará A, di Stefano S, Legarra J, Sarraide JA. An unusual vascular graft infection by *Aspergillus*. A case report and literature review. *Angiology* 1999; **50**: 169–173.
291. Marroni M, Cao A, Repetto A, Prattichizzo L, Parlani G, Fiorio M. *Aspergillus flavus* infection of an aortic bypass. *Eur J Clin Microbiol Infect Dis* 2001; **20**: 439–441.
292. Carlson GL, Mughal MM, Birch M, Denning DW. *Aspergillus* wound infection following laparostomy. *J Infect* 1996; **33**: 119–121.
293. Nenoff P, Horn LC, Schwenke H, Mierzwa M, Rieske K, Haustein UF. Invasive mould infections in the University Clinics of Leipzig during the period from 1992 to 1994. *Mycoses* 1996; **39** (suppl 1): 107–112.
294. Vairani G, Rebeschini R, Barbazza R. Hepatic and subcutaneous abscesses due to aspergilliosis. Initial diagnosis of a case by intraoperative fine needle aspiration cytology. *Acta Cytol* 1990; **34**: 891–894.
295. Erdman SH, Barber BJ, Barton LL. *Aspergillus* cholangitis: a late complication after Kasai portoenterostomy. *J Pediatr Surg* 2002; **37**: 923–925.
296. Davies SP, Webb WJ, Patou G, Murray WK, Denning DW. Renal aspergilloma—a case illustrating the problems of medical therapy. *Nephrol Dial Transplant* 1987; **2**: 568–572.
297. Kusne S, Torre-Cisneros J, Manez R *et al.* Factors associated with invasive lung aspergilliosis and the significance of positive *Aspergillus* culture after liver transplantation. *J Infect Dis* 1992; **166**: 1379–1383.
298. Sartin JS, Wilhelm MP, Keating MR, Batts K, Krom RA. A case of *Aspergillus fumigatus* peritonitis complicating liver transplantation. *Eur J Clin Microbiol Infect Dis* 1994; **13**: 25–28.
299. Marañes A, Portoles J, Blanco J *et al.* *Aspergillus* infection of a renal allograft without evidence of a site of origin. *Nephrol Dial Transplant* 1996; **11**: 1639–1642.
300. Woods GL, Wood RP, Shaw BW. *Aspergillus* endocarditis in patients without prior cardiovascular surgery: report of a case in a liver transplant recipient and review. *Rev Infect Dis* 1989; **11**: 263–272.
301. Paterson DL, Dominguez EA, Chang FY, Snyderman DR, Singh N. Infective endocarditis in solid organ transplant recipients. *Clin Infect Dis* 1998; **26**: 689–694.
302. Kaplan R, Duncalf D, Cizmar S. *Aspergillus* pancarditis and cardiac arrest during anesthesia. *Anesth Analg* 1981; **60**: 440–444.
303. Viertel A, Ditting T, Pistorius K, Geiger H, Scheuermann EH, Just-Nubling G. An unusual case of *Aspergillus* endocarditis in a kidney transplant recipient. *Transplantation* 1999; **68**: 1812–1813.
304. Marín P, Garcia-Martos P, Garcia-Doncel A *et al.* Endocarditis by *Aspergillus fumigatus* in a renal transplant. *Mycopathologia* 1999; **145**: 127–129.
305. Myerowitz RL, Friedman R, Grossman WL. Mycotic 'mycotic aneurysm' of the aorta due to *Aspergillus fumigatus*. *Am J Clin Pathol* 1971; **55**: 241–246.

306. Haddad SH, Arabi YM, Memish ZA, Al-Shimemeri AA. Nosocomial infective endocarditis in critically ill patients: a report of three cases and review of the literature. *Int J Infect Dis* 2004; **8**: 210–216.
307. Elzi L, Laifer G, Bremerich J, Vosbeck J, Mayr M. Invasive aspergillosis with myocardial involvement after kidney transplantation. *Nephrol Dial Transplant* 2005; **20**: 631–634.
308. Alvarez L, Calvo E, Abril C. Articular aspergillosis: case report. *Clin Infect Dis* 1995; **20**: 457–460.
309. Franco M, Van Elslande L, Robino C *et al.* *Aspergillus* arthritis of the shoulder in a renal transplant recipient. Failure of itraconazole therapy. *Rev Rhum Engl Ed* 1995; **62**: 215–218.
310. Cassuto-Viguiere E, Mondain JR, Van Elslande L *et al.* Fatal outcome of *Aspergillus fumigatus* arthritis in a renal transplant recipient. *Transplant Proc* 1995; **27**: 2461.
311. Ingwer I, McLeish KR, Tight RR, White AC. *Aspergillus fumigatus* epidural abscess in a renal transplant recipient. *Arch Intern Med* 1978; **138**: 153–154.
312. Salvalaggio PR, Bassetti M, Lorber MI *et al.* *Aspergillus* vertebral osteomyelitis after simultaneous kidney–pancreas transplantation. *Transpl Infect Dis* 2003; **5**: 187–190.
313. Taillandier J, Alemanni M, Cerrina J, Le Roy Ladurie F, Dartevielle P. *Aspergillus* osteomyelitis after heart–lung transplantation. *J Heart Lung Transplant* 1997; **16**: 436–438.
314. Kaneko J, Sugawara Y, Makuuchi M. *Aspergillus* osteomyelitis after liver transplantation. *Liver Transpl* 2002; **8**: 1073–1075.
315. Tang TJ, Janssen HL, van der Vlies CH *et al.* *Aspergillus* osteomyelitis after liver transplantation: conservative or surgical treatment? *Eur J Gastroenterol Hepatol* 2000; **12**: 123–126.
316. Williams RL, Fukui MB, Meltzer CC, Swarnkar A, Johnson DW, Welch W. Fungal spinal osteomyelitis in the immunocompromised patient: MR findings in three cases. *Am J Neuroradiol* 1999; **20**: 381–385.
317. Cortet B, Richard R, Deprez X *et al.* *Aspergillus* spondylodiscitis: successful conservative treatment in 9 cases. *J Rheumatol* 1994; **21**: 1287–1291.
318. Hummel M, Schuler S, Weber U *et al.* Aspergillosis with *Aspergillus* osteomyelitis and diskitis after heart transplantation: surgical and medical management. *J Heart Lung Transplant* 1993; **12**: 599–603.
319. Castelli C, Benazzo F, Minoli L, Marone P, Seghezzi R, Carlizzi CN. *Aspergillus* infection of the L3–L4 disc space in an immunosuppressed heart transplant patient. *Spine* 1990; **15**: 1369–1373.
320. De Vuyst D, Surmont I, Verhaegen J, Vanhaecke J. Tibial osteomyelitis due to *Aspergillus flavus* in a heart transplant patient. *Infection* 1992; **20**: 48–49.
321. Bridwell KH, Campbell JW, Barenkamp SJ. Surgical treatment of hematogenous vertebral aspergillus osteomyelitis. *Spine* 1990; **15**: 281–285.
322. Byrd BF, Weiner MH, McGee ZA. *Aspergillus* spinal epidural abscess. *JAMA* 1982; **248**: 3138–3139.
323. Grossman M. Aspergillosis of bone. *Br J Radiol* 1975; **48**: 57–59.
324. Sonin AH, Stern SH, Levi E. Primary *Aspergillus* osteomyelitis in the tibia of an immunosuppressed man. *Am J Roentgenol* 1996; **166**: 1277–1279.
325. Delmas Y, Merville P, Dousset V, Dervau-Durieux L, Morel D, Potaux L. A renal transplant recipient with acute paraparesis due to an *Aspergillus* epidural abscess. *Nephrol Dial Transplant* 1997; **12**: 2185–2187.
326. Jeevanandam V, Smith CR, Rose EA, Malm JR, Hugo NE. Single-stage management of sternal wound infections. *J Thorac Cardiovasc Surg* 1990; **99**: 256–262.
327. Byl B, Jacobs F, Antoine M *et al.* Mediastinitis caused by *Aspergillus fumigatus* with ruptured aortic pseudoaneurysm in a heart transplant recipient: case study. *Heart Lung* 1993; **22**: 145–147.
328. Levin T, Suh B, Beltramo D, Samuel R. *Aspergillus* mediastinitis following orthotopic heart transplantation: case report and review of the literature. *Transpl Infect Dis* 2004; **6**: 129–131.
329. Forestier E, Remy V, Lesens O *et al.* A case of *Aspergillus* mediastinitis after heart transplantation successfully treated with liposomal amphotericin B, caspofungin and voriconazole. *Eur J Clin Microbiol Infect Dis* 2005; **24**: 347–349.
330. Truppman ES, Ellenby JD, Schwartz BM. Fungi in and around implants after augmentation mammoplasty. *Plast Reconstr Surg* 1979; **64**: 804–806.
331. Williams K, Walton RL, Bunkis J. *Aspergillus* colonization associated with bilateral silicone mammary implants. *Plast Reconstr Surg* 1983; **71**: 260–261.
332. Rosenblatt WB, Pollock A. *Aspergillus flavus* cultured from a saline-filled implant. *Plast Reconstr Surg* 1997; **99**: 1470–1472.
333. Héry G, Hu W, Jestin N, Lagarde N, Le Flohic AM. *Aspergillus fumigatus* infection associated with mammary prosthesis. *J Mycol Med* 2002; **12**: 193–195.
334. Saray A, Kilic D, Kaygusuz S, Boyunaga H, Ozluk O. Fungal growth inside saline-filled implants and the role of injection ports in fungal translocation: in vitro study. *Plast Reconstr Surg* 2004; **114**: 1170–1178.
335. Brown MH, Markus YM, Belchetz B, Vearncombe M, Semple JL. Microbial growth in saline breast implants and saline tissue expanders. *Plast Reconstr Surg* 2002; **109**: 2242–2244.
336. Young VL, Hertl MC, Murray PR, Jensen J, Witt H, Schorr MW. Microbial growth inside saline-filled breast implants. *Plast Reconstr Surg* 1997; **100**: 182–196.
337. Coady MSE, Gaylor J, Knight SL. Fungal growth within a silicone tissue expander: Case report. *Br J Plast Surg* 1995; **48**: 428–430.
338. Heinemann S, Symoens F, Gordts B, Jannes B, Nolard N. Environmental investigations and molecular typing of *Aspergillus flavus* during an outbreak of post-operative infections. *J Hosp Infect* 2004; **57**: 149–155.
339. Diaz-Guerra TM, Mellado E, Cuenca-Estrella M, Gaztelurrutia L, Navarro JI, Tudela JL. Genetic similarity among one *Aspergillus flavus* strain isolated from a patient who underwent heart surgery and two environmental strains obtained from the operating room. *J Clin Microbiol* 2000; **38**: 2419–2422.
340. Dharan S, Pittet D. Environmental controls in operating theatres. *J Hosp Infect* 2002; **51**: 79–84.
341. Babb JR, Lynam P, Ayliffe GAJ. Risk of airborne transmission in an operating theater containing four ultraclean air units. *J Hosp Infect* 1995; **31**: 159–168.
342. Barnes RA, Rogers TR. Control of an outbreak of nosocomial aspergillosis by laminar air-flow isolation. *J Hosp Infect* 1989; **14**: 89–94.
343. Rhame FS. Prevention of nosocomial aspergillosis. *J Hosp Infect* 1991; **18** (suppl A): 466–472.

344. Walmsley S, Devi S, King S, Schneider R, Richardson S, Ford-Jones L. Invasive *Aspergillus* infections in a pediatric hospital: a ten-year review. *Pediatr Infect Dis* 1993; **12**: 673–682.
345. Pittet D, Duce G. Infectious risk factors related to operating rooms. *Infect Control Hosp Epidemiol* 1994; **15**: 456–462.
346. Sehulster LM, Chinn RYW, Arduino MJ *et al*. *Guidelines for environmental infection control in health-care facilities. Recommendations from CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC)*. Chicago IL: American Society for Healthcare Engineering/American Hospital Association, 2004.
347. Faure O, Fricker-Hidalgo H, Lebeau B, Mallaret MR, Ambroise-Thomas P, Grillot R. Eight-year surveillance of environmental fungal contamination in hospital operating rooms and haematological units. *J Hosp Infect* 2002; **50**: 155–160.
348. Roy M-C. The operating theater: a special environmental area. In: Wenzel RP, ed., *Prevention and control of nosocomial infections*, 3rd edn. Baltimore, MD: William & Wilkins, 1997; 515–538.
349. Humphreys H. Positive-pressure isolation and the prevention of invasive aspergillosis. What is the evidence? *J Hosp Infect* 2004; **56**: 93–100.
350. Kelkar U, Bal AM, Kulkarni S. Fungal contamination of air conditioning units in operating theatres in India. *J Hosp Infect* 2005; **60**: 81–84.