



Egyptian Mathematical Society
Journal of the Egyptian Mathematical Society

www.etms-eg.org
www.elsevier.com/locate/joems



ORIGINAL ARTICLE

Characterization through distributional properties of dual generalized order statistics

A.H. Khan ^{a,*}, Imtiyaz A. Shah ^a, M. Ahsanullah ^b

^a Department of Statistics and Operations Research, Aligarh Muslim University, Aligarh 202 002, India

^b Department of Management Sciences, Rider University, Lawrenceville, NJ 08648-3099, USA

Received 25 June 2011; revised 30 June 2012

Available online 6 December 2012

KEYWORDS

Order statistics;
 Lower record statistics;
 Upper record statistics;
 Generalized order statistics;
 Dual generalized order statistics;
 Contraction;
 Dilation;
 Characterization of distributions;
 Generalized exponential;
 Generalized Pareto;
 Generalized power function;
 Gumbel;
 Weibull;
 Inverse Weibull;
 Exponential;
 Power function;
 Pareto;
 Distributions

Abstract Distributional properties of two non-adjacent dual generalized order statistics have been used to characterize distributions. Further, one sided contraction and dilation for the dual generalized order statistics are discussed and then the results are deduced for generalized order statistics, order statistics, lower record statistics, upper record statistics and adjacent dual generalized order statistics.

© 2012 Egyptian Mathematical Society. Production and hosting by Elsevier B.V.
 Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

Kamps [6] introduced the concept of generalized order statistics (*gos*) as follows:

Let X_1, X_2, \dots, X_n be a sequence of independent and identically distributed (*iid*) random variables (*rv*) with the absolutely continuous distribution function (*df*) $F(x)$ and the probability density function (*pdf*) $f(x)$, $x \in (\alpha, \beta)$. Let $n \in \mathbb{N}$, $n \geq 2$, $k > 0$, $\vec{m} = (m_1, m_2, \dots, m_{n-1}) \in \mathfrak{R}^{n-1}$, $M_r = \sum_{j=r}^{n-1} m_j$,

* Corresponding author.

E-mail address: ahamidkhan@rediffmail.com (A.H. Khan).

Peer review under responsibility of Egyptian Mathematical Society.



Production and hosting by Elsevier

such that $\gamma_r = k + (n - r) + M_r > 0$ for all $r \in \{1, 2, \dots, n - 1\}$. If $m_1 = m_2 = \dots = m_{n-1} = m$, then $X(r, n, m, k)$ is called the r th m -gos and its pdf is given as:

$$f_{X(r,n,m,k)}(x) = \frac{c_{r-1}}{(r-1)!} [\bar{F}(x)]^{\gamma_r-1} \left[\frac{1 - [\bar{F}(x)]^{m+1}}{m+1} \right]^{r-1} f(x),$$

$$\alpha < x < \beta, \quad (1.1)$$

where $\gamma_r = k + (n - r)(m + 1)$ and $c_{r-1} = \prod_{i=1}^r \gamma_i$.

Based on the generalized order statistics (gos), Burkschat et al. [4] introduced the concept of the dual generalized order statistics (dgos) where the pdf of the r th m -dgos $X^*(r, n, m, k)$ is given as

$$f_{X^*(r,n,m,k)}(x) = \frac{c_{r-1}}{(r-1)!} [F(x)]^{\gamma_r-1} \left[\frac{1 - [F(x)]^{m+1}}{m+1} \right]^{r-1} f(x),$$

$$\alpha < x < \beta,$$

which is obtained just by replacing $\bar{F}(x) = 1 - F(x)$ by $F(x)$.

Ahsanullah [1] has characterized uniform distribution under random contraction for adjacent dgos. Khan and Shah [7] have characterized distributions using distributional properties of non-adjacent lower records, upper records and order statistics. In this paper, distributional properties of the dgos have been used to characterize a general form of distributions for non-adjacent dgos under random translation, dilation and contraction, thus generalizing the results of Ahsanullah [1]. Further, results in terms of lower records, upper records and order statistics are deduced. One may also refer to Alzaid and Ahsanullah [2], Beutner and Kamps [3], Wesolowski and Ahsanullah [8] and Castaño-Martínez et al. [5] for the related results.

Remark 1.1. It may be seen that if Y is a measurable function of X with the relation $Y = h(X)$, then $Y^*(r, n, m, k) = h(X^*(r, n, m, k))$ and $Y(r, n, m, k) = h(X(r, n, m, k))$, if h is increasing function (e.g., $Y_{r:n} = h(X_{r:n})$ and $Y_{L(r)} = h(X_{L(r)})$, where $X_{r:n}$ and $X_{L(r)}$ are the r th order statistic and lower record, respectively). Moreover, $Y(r, n, m, k) = h(X^*(r, n, m, k))$ and $Y^*(r, n, m, k) = h(X(r, n, m, k))$, if h is decreasing function (e.g., $Y_{n-r+1:n} = h(X_{r:n})$ and $Y_{U(r)} = h(X_{L(r)})$, where $Y_{U(r)}$ is the r th upper record).

Remark 1.2. The following elementary facts will be needed in the next section:

- (i) if $Y = \log X \sim Gum(\alpha)$ (i.e., $F_Y(y) = e^{-e^{-\alpha y}}$, $-\infty < y < \infty, \alpha > 0$), then $X \sim in W(\alpha)$ (i.e., $F_X(x) = e^{-x^{-\alpha}}$, $0 < x < \infty, \alpha > 0$).
- (ii) if $-\log X \sim Gum(\alpha)$, then $X \sim Wei(\alpha)$, (i.e., $F_X(x) = 1 - e^{-x^\alpha}, 0 < x < \infty, \alpha > 0$).
- (iii) if $Y = \log X \sim exp(\alpha)$ (i.e., $F_Y(y) = 1 - e^{-\alpha y}, 0 < y < \infty, \alpha > 0$), then $X \sim Par(\alpha)$, (i.e., $F_X(x) = 1 - x^{-\alpha}, 1 < x < \infty, \alpha > 0$).
- (iv) if $-\log X \sim exp(\alpha)$, then $X \sim pow(\alpha)$, (i.e., $F_X(x) = x^\alpha, 0 < x < 1, \alpha > 0$).
- (v) if $Y = \log X \sim genexp(\alpha)$ (i.e., $F_Y(y) = [1 - (m + 1)e^{-\alpha y}]^{\frac{1}{m+1}}, \frac{1}{\alpha} \log(m + 1) < y < \infty, \alpha > 0$), then $X \sim gen-Par(\alpha)$ (i.e., $F_X(x) = [1 - (m + 1)x^{-\alpha}]^{\frac{1}{m+1}}, (m + 1)^{\frac{1}{2}} < x < \infty, \alpha > 0$).
- (vi) if $-\log X \sim genexp(\alpha)$, then $X \sim genpow(\alpha)$ (i.e., $F_X(x) = 1 - [1 - (m + 1)x^\alpha]^{\frac{1}{m+1}}, 0 < x < (m + 1)^{\frac{1}{2}}, \alpha > 0$).

2. Characterizing results

Theorem 2.1. Let $X^*(s, n, m, k)$ be the s th m -dgos from a sample of size n drawn from a continuous population with the $pdf f(x)$ and the $df F(x)$, then for $1 \leq r < s \leq n$,

$$X^*(r + j, n, m, k) \stackrel{d}{=} X^*(s, n, m, k) + Y_{s-r-j:s-1}, \quad j = 0, 1, \quad (2.1)$$

where $Y_{s-r-j:s-1}$ is the $(s - r - j)$ th order statistic from a sample of size $(s - 1)$ drawn from $exp(\alpha)$ distribution and is independent of $X^*(s, n, m, k)$ if and only if $X_1 \sim genexp(\alpha)$ and $X \stackrel{d}{=} Y$ denotes that X and Y have the same df .

Proof. To prove the necessary part, let the moment generating function (mgf) of $X^*(r, n, m, k)$ be $M_{X^*(r)}(t)$, then $X^*(r, n, m, k) \stackrel{d}{=} X^*(s, n, m, k) + Y$, implies that $M_{X^*(r)}(t) = M_{X^*(s)}(t) \cdot M_Y(t)$.

Since for the $genexp(\alpha)$ distribution, we have

$$M_{X^*(r)}(t) = \frac{C_{r-1}}{(r-1)!} \frac{1}{(m+1)^{r-\frac{1}{2}}} \frac{\Gamma(r - \frac{1}{2}) \Gamma(\frac{\gamma_r}{m+1})}{\Gamma(r - \frac{1}{2} + \frac{\gamma_r}{m+1})}.$$

Therefore,

$$M_Y(t) = \frac{M_{X^*(r)}(t)}{M_{X^*(s)}(t)} = \frac{\Gamma(s)}{\Gamma(r)} \frac{\Gamma(r - \frac{1}{2})}{\Gamma(s - \frac{1}{2})}.$$

But this is the mgf of $Y_{s-r:s-1}$, which is the $(s - r)$ th order statistic from a sample of size $(s - 1)$ drawn from $exp(\alpha)$.

To prove the sufficiency part, we have for $s \geq r + 1$,

$$f_{X^*(r,n,m,k)}(x) = \int_0^x f_{X^*(s,n,m,k)}(y) \cdot f_{Y_{s-r:s-1}}(x - y) dy$$

$$= \frac{\alpha(s-1)!}{(r-1)!(s-r-1)!} \int_0^x [e^{-\alpha(x-y)}]^r [1 - e^{-\alpha(x-y)}]^{s-r-1} \times f_{X^*(s,n,m,k)}(y) dy. \quad (2.2)$$

Differentiate both the sides of (2.2) w.r.t. x , to get

$$\frac{d}{dx} f_{X^*(r,n,m,k)}(x) = \alpha r [f_{X^*(r+1,n,m,k)}(x) - f_{X^*(r,n,m,k)}(x)]$$

$$\text{or, } f_{X^*(r,n,m,k)}(x) = \alpha r [F_{X^*(r+1,n,m,k)}(x) - F_{X^*(r,n,m,k)}(x)]$$

Now, since (Ahsanullah [1])

$$[F_{X^*(r+1,n,m,k)}(x) - F_{X^*(r,n,m,k)}(x)] = \frac{F(x)}{\gamma_{r+1} f(x)} f_{X^*(r+1,n,m,k)}(x).$$

Therefore, we have $\frac{(m+1)[F(x)]^{m+1} f(x)}{[1 - (F(x))^{m+1}]} = \alpha$, which implies that

$$F(x) = [1 - (m + 1)e^{-\alpha x}]^{\frac{1}{m+1}}. \text{ Hence the proof. } \square$$

Remark 2.1. Let $X_{r:n}$ be the r th order statistic from a sample of size n drawn from a continuous population with the $pdf f(x)$ and the $df F(x)$, then for $1 \leq r < s \leq n$,

$$X_{s-j:n} \stackrel{d}{=} X_{r:n} + X_{s-r-j:n-r}, \quad j = 0, 1, \quad (2.3)$$

where $X_{s-r-j:n-r}$ is independent of $X_{r:n}$ if and only if $X_1 \sim exp(\alpha)$.

This can be established by noting that order statistic appear in the generalized order statistics (*gos*) model as well as in dual generalized order statistics (*dgos*) model. Therefore at $m = 0$, (2.1) may be written as

$$X_{n-r-j+1:n} \stackrel{d}{=} X_{n-s+1:n} + X_{s-r-j:s-1}, \quad j = 0, 1; \quad 1 \leq r < s \leq n,$$

which implies

$$X_{s-j:n} \stackrel{d}{=} X_{r:n} + X_{s-r-j:n-r}, \quad j = 0, 1; \quad 1 \leq r < s \leq n,$$

obtained by replacing $(n - s + 1)$ by r and $(n - r + 1)$ by s as given by Khan and Shah [7].

Remark 2.2. Alzaid and Ahsanullah [2] have proved that

$$X_{r:n} \stackrel{d}{=} X_{r-1:n} + V$$

where $V \sim \exp(n - r + 1)$ if and only if $X_1 \sim \exp(1)$.

Remark 2.3. Castaño-Martínez *et al.* [5] have shown that

$$X_{s:n} \stackrel{d}{=} X_{r:n} + V$$

where $V \stackrel{d}{=} -\log W$ with $W \sim Be(n - s + 1, s - r)$ if and only if $X_1 \sim \exp(1)$.

Remark 2.4. As $m \rightarrow -1$, $genexp(x)$ tends to the $Gum(x)$ and $X^*(r, n, m, k)$ to $X_{L(r)}$, the r th lower records. Therefore, we have

$$X_{L(r+j)} \stackrel{d}{=} X_{L(s)} + Y_{s-r-j:s-1}, \quad j = 0, 1; \quad 1 \leq r < s,$$

where $Y_{s-r-j:s-1}$ is the $(s - r - j)$ th order statistic from a sample of size $(s - 1)$ drawn from $\exp(x)$ distribution and is independent of $X_{L(s)}$ if and only if $X_1 \sim Gum(x)$, as obtained by Khan and Shah [7].

Remark 2.5. Alzaid and Ahsanullah [2] have shown that

$$X_{L(r)} \stackrel{d}{=} X_{L(r+1)} + V$$

where $V \sim \exp(r)$ if and only if $X_1 \sim Gum(1)$.

Corollary 2.1. Let $X^*(s, n, m, k)$ be the s th *m-dgos* from a sample of size n drawn from a continuous population with the pdf $f(x)$ and the df $F(x)$, then for $1 \leq r < s \leq n$,

$$X^*(r + j, n, m, k) \stackrel{d}{=} X^*(s, n, m, k) \cdot Y_{s-r-j:s-1}, \quad j = 0, 1, \quad (2.4)$$

where $Y_{s-r-j:s-1}$ is the $(s - r - j)$ th order statistic from a sample of size $(s - 1)$ drawn from $Par(\alpha)$ distribution and is independent of $X^*(s, n, m, k)$ if and only if $X_1 \sim genPar(\alpha)$.

Proof. Here the product $X^*(s, n, m, k) \cdot Y_{s-r-j:s-1}$ in (2.4) is called random dilation of $X^*(s, n, m, k)$ (Beutner and Kamps [3]). Note that if

$$\log X^*(r, n, m, k) \stackrel{d}{=} \log X^*(s, n, m, k) + \log Y_{s-r:s-1}$$

then

$$X^*(r, n, m, k) \stackrel{d}{=} X^*(s, n, m, k) \cdot Y_{s-r:s-1}$$

in view of Remarks 1.1 and 1.2 and the result follows. \square

Remark 2.6. In case of ordinary order statistics, i.e., at $m = 0$, we have

$$X_{s-j:n} \stackrel{d}{=} X_{r:n} \cdot X_{s-r-j:n-r}, \quad j = 0, 1; \quad 1 \leq r < s \leq n,$$

where $X_{s-r-j:n-r}$ is independent of $X_{r:n}$ if and only if $X_1 \sim Par(\alpha)$, as obtained by Castaño-Martínez *et al.* [5] and Khan and Shah Imtiyaz [7].

Remark 2.7. As $m \rightarrow -1$, we get

$$X_{L(r+j)} \stackrel{d}{=} X_{L(s)} \cdot Y_{s-r-j:s-1}, \quad j = 0, 1; \quad 1 \leq r < s,$$

where $Y_{s-r-j:s-1}$ is the $(s - r - j)$ th order statistic from a sample of size $(s - 1)$ drawn from the $Par(\alpha)$ distribution and is independent of $X_{L(s)}$, the s th lower records if and only if $X_1 \sim inW(\alpha)$.

Corollary 2.2. Let $X(s, n, m, k)$ be the s th *m-gos* from a sample of size n drawn from a continuous population with the pdf $f(x)$ and the df $F(x)$, then for $1 \leq r < s \leq n$,

$$X(r + j, n, m, k) \stackrel{d}{=} X(s, n, m, k) \cdot Y_{r+j:s-1}, \quad j = 0, 1, \quad (2.5)$$

where $Y_{r+j:s-1}$ is the $(r + j)$ th order statistic from a sample of size $(s - 1)$ drawn from $pow(\alpha)$ distribution and is independent of $X(s, n, m, k)$ if and only if $X_1 \sim genpow(\alpha)$.

Proof. Here the product $X(s, n, m, k) \cdot Y_{r+j:s-1}$ in (2.5) is called random contraction of $X(s - j, n, m, k)$ (Beutner and Kamps [3]). Since

$$-\log X^*(r, n, m, k) \stackrel{d}{=} -\log X^*(s, n, m, k) - \log Y_{s-r:s-1}$$

implies

$$X(r, n, m, k) \stackrel{d}{=} X(s, n, m, k) \cdot Y_{r:s-1}$$

in view of Remarks 1.1 and 1.2 and the result follows. \square

Remark 2.8. Beutner and Kamps [3] have shown that for adjacent generalized order statistics

$$X(r, n, m, k) \stackrel{d}{=} X(r + 1, n, m, k) \cdot V$$

where $V \sim pow(r\alpha)$ if and only if $X_1 \sim genpow(\alpha)$.

Remark 2.9. We can get the corresponding characterizing results for the order statistics at $m = 0$ as:

Let $X_{r:n}$ be the r th order statistic from a sample of size n drawn from continuous population with the pdf $f(x)$ and the df $F(x)$, then for $1 \leq r < s \leq n$,

$$X_{r+j:n} \stackrel{d}{=} X_{s:n} \cdot X_{r+j:s-1}, \quad j = 0, 1,$$

where $X_{r+j:s-1}$ is independent of $X_{s:n}$ if and only if $X_1 \sim pow(\alpha)$, as given by Khan and Shah [7].

For adjacent order statistics one may also refer to Ahsanullah [1] and Wesolowski and Ahsanullah [8].

Remark 2.10. The corresponding result for the lower records as $m \rightarrow -1$ is:

Let $X_{U(s)}$ be the s th upper record from a continuous population with the *pdf* $f(x)$ and the *df* $F(x)$, then

$$X_{U(r+j)} \stackrel{d}{=} X_{U(s)} \cdot Y_{r+j:s-1}, \quad j = 0, 1; 1 \leq r < s,$$

where $Y_{r+j:s-1}$ is the $(r + j)$ th order statistic from a sample of size $(s - 1)$ drawn from *pow*(α) distribution and is independent of $X_{U(s)}$ if and only if $X_1 \sim \text{Wei}(\alpha)$, as obtained by Khan and Shah [7].

Acknowledgment

The authors are thankful to the Referees and the Editor for their fruitful suggestions and comments.

References

- [1] M. Ahsanullah, A characterization of the uniform distribution by dual generalized order statistics, *Commun. Statist. Theor. Meth.* 33 (2004) 2921–2928.

- [2] A.A. Alzaid, M. Ahsanullah, A characterization of the Gumbel distribution based on record values, *Commun. Statist. Theor. Meth.* 32 (2003) 2101–2108.
- [3] E. Beutner, U. Kamps, Random contraction and random dilation of generalized order statistics, *Commun. Statist. Theor. Meth.* 37 (2008) 2185–2201.
- [4] M. Burkschat, E. Cramer, U. Kamps, Dual generalized order statistics, *Metron LXI* (2003) 13–26.
- [5] A. Castaño-Martínez, F. López-Blázquez, B. Salamanca-Miño, Random translations, contractions and dilations of order statistics and records, *Statistics* (2010) 1–11.
- [6] U. Kamps, *A Concept of Generalized Order Statistics*, Teubner, Stuttgart, 1995.
- [7] A.H. Khan, Imtiaz A. Shah, Distributional properties of order statistics and record statistics, *Pak. J. Statist. Oper. Res.* 8 (3) (2012) 293–301.
- [8] J. Wesolowski, M. Ahsanullah, Switching order statistics through random power contractions, *Aust. N.Z.J. Statist.* 46 (2) (2004) 297–303.