



An elementary introduction to the homotopy perturbation method

Ji-Huan He^{a,b,*}

^a College of Mathematics & Physics, Nanjing University of Information Science and Technology, Nanjing 210044, PR China

^b Modern Textile Institute, Donghua University, 1882 Yan'an Xilu Road, Shanghai 200051, PR China

ARTICLE INFO

Article history:

Received 30 December 2007

Accepted 19 June 2008

Keywords:

Homotopy perturbation method

Parameter-expansion method

Axisymmetric flow

Asymptotic solution

Lindstedt–Poincaré method

ABSTRACT

This paper is an elementary introduction to the concepts of the homotopy perturbation method. Particular attention is paid to giving an intuitive grasp for the solution procedure throughout the paper.

© 2008 Elsevier Ltd. All rights reserved.

This short note is actually a comment on Ariel's publication [1] and a reply to El-Mistikawy's comment [2].

Ariel [1] considered the following coupled equations

$$F''' + (F + G)F'' - F'^2 = 0 \quad (1)$$

$$G''' + (F + G)G'' - G'^2 = 0 \quad (2)$$

$$F'(0) = 1, \quad G'(0) = \beta, \quad F'(\infty) = 0, \quad G'(\infty) = 0, \quad \text{and} \quad F(0) + G(0) = 0 \quad (3)$$

where prime denotes the derivative with respect to the similarity coordinate η .

Ariel constructed the following homotopy equations

$$F''' - b^2F' + p \left\{ (F + G)F'' - F'^2 - b^2F' \right\} = 0 \quad (4)$$

$$G''' - b^2G' + p \left\{ (F + G)G'' - G'^2 - b^2G' \right\} = 0. \quad (5)$$

The obtained results are of high accuracy even for the first-order approximate solution, showing that the solution procedure is acceptable.

El-Mistikawy [2] considers that the homotopy perturbation cannot obtain higher-order approximation except first-order approximation. Actually El-Mistikawy might misapprehend the method, which always tends to solve first-order approximations but the solution procedure can continue without any difficulty to obtain higher-order approximations, see Refs. [3–7].

* Corresponding address: Modern Textile Institute, Donghua University, 1882 Yan'an Xilu Road, Shanghai 200051, PR China.

E-mail address: jhhe@dhu.edu.cn.

Let us outline the basic solution procedure of the homotopy perturbation method:

1. Qualitative sketch/trial function solution

From physical understandings and boundary conditions, we choose such initial guesses:

$$F'(\eta) = e^{-a\eta} \tag{6}$$

and

$$G'(\eta) = \beta e^{-b\eta} \tag{7}$$

where a and b are unknown constants to be further identified. It is obvious that Eqs. (6) and (7) satisfy the boundary condition $F'(0) = 1, F'(\infty) = 0$ and $G'(0) = \beta, G'(\infty) = 0$ respectively.

2. Construction of homotopy equations

According to the initial guesses, Eqs. (6) and (7), homotopy equations can be constructed as follows

$$F''' - a^2F' + p \{ (F + G)F'' - F'^2 - a^2F' \} = 0 \tag{8}$$

$$G''' - b^2G' + p \{ (F + G)G'' - G'^2 - b^2G' \} = 0. \tag{9}$$

When $p = 0$, we can obtain the initial guesses; When $p = 1$, Eqs. (8) and (9) turn out to be the original ones.

Our homotopy system turns out to be Ariel's system when we choose $a = b$ for simplicity.

3. Solution procedure similar to that of classical perturbation methods

Now we can use the homotopy parameter, p , as an expanding parameter used in classic perturbation methods. The simplest way is the method of straightforward expansion:

$$F = F_0 + pF_1 + p^2F_2 + \dots \tag{10}$$

$$G = G_0 + pG_1 + p^2G_2 + \dots \tag{11}$$

Generally we stop before second iteration. Setting $p = 1$, we obtain the first-order approximation which reads

$$F(\eta) = F_0(\eta) + F_1(\eta) \tag{12}$$

$$G(\eta) = G_0(\eta) + G_1(\eta). \tag{13}$$

4. Optimal identification of the unknown parameter in the trial functions

There are many approaches to identify the unknown parameters in the obtained approximation, for example, the method of weighted residuals, especially the least squares method.

For the present problem, we set

$$\int_0^{+\infty} e^{-a\eta} \{ F''' + (F + G)F'' - F'^2 \} d\eta = 0 \tag{14}$$

and

$$F(0) + G(0) = 0 \tag{15}$$

to identify the unknown constants a and b . Ariel's identification of the unknown constant is equivalent to Eq. (14).

5. Higher-order approximations

If first-order approximation cannot meet the requirement of engineering applications, higher-order approximations should be solved.

As it is well known the straightforward expansion leads to secular terms [8] and various methods, such as Lindstedt–Poincare method [9–11], are suggested to eliminate the secular terms. One of the advantages of the homotopy perturbation method is that it can take full advantage of various known perturbation methods.

El-Mistikawy [2] actually uses the Lindstedt–Poincare method, of course, El-Mistikawy's solution procedure is valid for higher-order approximations.

In Ref. [9], the present author suggested a modified Lindstedt–Poincare method, which is also very useful and effective to solve higher-order approximations. We re-write Eqs. (1) and (2) in the forms

$$F''' + 0 \cdot F' + p \left\{ (F + G)F'' - F'^2 \right\} = 0 \quad (16)$$

$$G''' + 0 \cdot G' + p \left\{ (F + G)G'' - G'^2 \right\} = 0. \quad (17)$$

The zero coefficient in Eq. (16) can be expressed in [6,9,12]

$$0 = -a^2 + a_1p + a_2p^2 + \dots \quad (18)$$

The zero coefficient in Eq. (17) can be expanded in a similar way

$$0 = -b^2 + b_1p + b_2p^2 + \dots \quad (19)$$

The constants a_i and b_i can be identified in view of no η terms, i.e., no $\eta e^{-a\eta}$ term in F and no $\eta e^{-b\eta}$ term in G .

Why such expansion is valid is explained in Ref. [12].

The above solution procedure can be achieved by the parameter-expansion method [6]. We re-write Eqs. (1) and (2) in the forms:

$$F''' + 0 \cdot F' + 1 \cdot \left\{ (F + G)F'' - F'^2 \right\} = 0 \quad (20)$$

$$G''' + 0 \cdot G' + 1 \cdot \left\{ (F + G)G'' - G'^2 \right\} = 0. \quad (21)$$

We expand F , G and zero as before, and expand unit in Eqs. (20) and (21), respectively, as

$$1 = c + c_1p + c_2p^2 + \dots \quad (22)$$

and

$$1 = d + d_1p + d_2p^2 + \dots \quad (23)$$

For detailed solution procedure for parameter-expansion method, please refer to Refs. [6,13–18].

For detailed description of the homotopy perturbation method, please refer to Refs. [6,19].

References

- [1] P.D. Ariel, The three-dimensional flow past a stretching sheet and the homotopy perturbation method, *Comput. Math. Appl.* 54 (2007) 920–925.
- [2] T.M.A. El-Mistikawy, Comment on the three-dimensional flow past a stretching sheet and the homotopy perturbation method, *Comput. Math. Appl.* 57 (3) (2009) 404–406.
- [3] P.D. Ariel, T. Hayat, S. Asghar, Homotopy perturbation method and axisymmetric flow over a stretching sheet, *Int. J. Nonlinear Sci. Num.* 7 (2006) 399–406.
- [4] A. Belendez, A. Hernandez, T. Belendez, et al., Application of He's homotopy perturbation method to the Duffing-harmonic oscillator, *Int. J. Nonlinear Sci. Num.* 8 (2007) 79–88.
- [5] D.D. Ganji, A. Sadighi, Application of He's homotopy-perturbation method to nonlinear coupled systems of reaction-diffusion equations, *Int. J. Nonlinear Sci. Num.* 7 (2006) 411–418.
- [6] J.H. He, Some asymptotic methods for strongly nonlinear equations, *Internat. J. Modern Phys. B.* 20 (2006) 1141–1199.
- [7] A.M. Siddiqui, R. Mahmood, Q.K. Ghori, Thin film flow of a third grade fluid on a moving belt by He's homotopy perturbation method, *Int. J. Nonlinear Sci. Num.* 7 (2006) 7–14.
- [8] J.H. He, Modified Lindstedt–Poincare methods for some strongly nonlinear oscillations Part III: Double series expansion, *Int. J. Nonlinear Sci. Num.* 2 (2001) 317–320.
- [9] J.H. He, Modified Lindstedt–Poincare methods for some strongly non-linear oscillations Part I: Expansion of a constant, *Internat. J. Nonlinear Mech.* 37 (2002) 309–314.
- [10] J.H. He, Modified Lindstedt–Poincare methods for some strongly non-linear oscillations Part II: A new transformation, *Internat. J. Nonlinear Mech.* 37 (2002) 315–320.
- [11] J.H. He, Modified straightforward expansion, *Meccanica* 34 (1999) 287–289.
- [12] J.H. He, New interpretation of homotopy perturbation method, *Internat. J. Modern Phys. B* 20 (2006) 2561–2568.
- [13] D.H. Shou, J.H. He, Application of parameter-expanding method to strongly nonlinear oscillators, *Int. J. Nonlinear Sci. Num.* 8 (2007) 121–124.
- [14] S.Q. Wang, J.H. He, Nonlinear oscillator with discontinuity by parameter-expansion method, *Chaos Solitons Fractals* 35 (2008) 688–691.
- [15] L. Xu, Application of He's parameter-expansion method to an oscillation of a mass attached to a stretched elastic wire, *Phys. Lett. A* 368 (2007) 259–262.
- [16] L. Xu, He's parameter-expanding methods for strongly nonlinear oscillators, *J. Comput. Appl. Math.* 207 (2007) 148–154.
- [17] L. Xu, Determination of limit cycle by He's parameter-expanding method for strongly nonlinear oscillators, *J. Sound Vib.* 302 (2007) 178–184.
- [18] L.N. Zhang, L. Xu, Determination of the limit cycle by He's parameter-expansion for oscillators in a $u(3)/(1 + u(2))$ potential, *Z. Naturforsch. A* 62 (2007) 396–398.
- [19] J.H. He, *Nonperturbative methods for strongly nonlinear problems*, Dissertation.de-Verlag im Internet GmbH 2006.